


Notation: Sets


# Notation: Sets

\_\_\_\_\_→  $x$


# Notation: Sets




# Notation: Sets




# Notation: Sets




# Notation: Sets



# Notation: Sets




# Notation: Sets






# Notation: Sets

Point:  $(1, 1)$




# Notation: Sets

Point:  $(1, 1) \in \mathbb{R}^2$




# Notation: Sets

Point:  $(1, 2) \in \mathbb{R}^2$




# Notation: Sets

Point:  $(1, 2.5) \in \mathbb{R}^2$




# Notation: Sets


Point:  $(3, 2) \in \mathbb{R}^2$




# Notation: Sets



# Notation: Sets




# Notation: Sets






# Notation: Sets

$$\{(1, 1), (1, 2), (1, 3)\}$$




# Notation: Sets

$$S := \{(1, 1), (1, 2), (1, 3)\}$$





# Notation: Sets

$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$





# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line,

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane


# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

# Notation: Sets




$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$\{???\}$$

# Notation: Sets




$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by **points**  $(x, y)$  in the plane so that  $y = x - 1.7$

$$\{(x, y)\}$$



# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$\{(x, y) \in \mathbb{R}^2\}$$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$\{(x, y) \in \mathbb{R}^2 \mid \}$$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$\{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $\{x \in \mathbb{R}\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $\{x \in \mathbb{R} \mid \}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $\{x \in \mathbb{R} \mid \alpha < x\}$



# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $\{x \in \mathbb{R} \mid \alpha < x < \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $:= \{x \in \mathbb{R} \mid \alpha < x < \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

- 1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
- 2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

- 1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
- 2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $\{x \in \mathbb{R}\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

- 1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
- 2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
- 3.  $\{x \in \mathbb{R} \mid \}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $\{x \in \mathbb{R} \mid \alpha < x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $\{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$



# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $\{x \in \mathbb{R}\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $\{x \in \mathbb{R} \mid \}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $\{x \in \mathbb{R} \mid \alpha \leq x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$


$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$



# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $\{x \in \mathbb{R}\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $\{x \in \mathbb{R} \mid \}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $\{x \in \mathbb{R} \mid \alpha \leq x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $\{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $\{x \in \mathbb{R}\}$



# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $\{x \in \mathbb{R} \mid \}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $\{x \in \mathbb{R} \mid \alpha < x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $:= \{x \in \mathbb{R} \mid \alpha < x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, := \{x \in \mathbb{R} \mid \alpha < x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$
7.  $\{x \in \mathbb{R}\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$
7.  $\{x \in \mathbb{R} \mid \}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$


$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$
7.  $\{x \in \mathbb{R} \mid x \leq \beta\}$



# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$


A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$
7.  $:= \{x \in \mathbb{R} \mid x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$

$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$
7.  $(-\infty, \beta] := \{x \in \mathbb{R} \mid x \leq \beta\}$

# Notation: Sets



$$S := \{(1, 1), (1, 2), (1, 3)\} \subset \mathbb{R}^2$$

A line, defined by points  $(x, y)$  in the plane so that  $y = x - 1.7$



$$C := \{(x, y) \in \mathbb{R}^2 \mid y = x - 1.75\}$$

## Examples.

1.  $\mathbb{R}$  : set of all real numbers.  $2, \pi$  etc  $\in \mathbb{R}$
2.  $(\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha < x < \beta\}$
3.  $(\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha < x \leq \beta\}$
4.  $[\alpha, \beta) := \{x \in \mathbb{R} \mid \alpha \leq x < \beta\}$
5.  $[\alpha, \beta] := \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$
6.  $(\alpha, \infty) := \{x \in \mathbb{R} \mid \alpha < x\}$
7.  $(-\infty, \beta] := \{x \in \mathbb{R} \mid x \leq \beta\}$


# Notation: Sets

{???




# Notation: Sets

$$\{(x,y)\}$$




# Notation: Sets

$$\{(x,y) \in \mathbb{R}^2\}$$




# Notation: Sets

$$\{(x,y) \in \mathbb{R}^2 \mid \}$$



# Notation: Sets


$$\{(x, y) \in \mathbb{R}^2 \mid y = x\}$$






# Notation: Sets

$$\{(x, y) \in \mathbb{R}^2 \mid y = x, 0 < x\}$$




# Notation: Sets

$$\{(x, y) \in \mathbb{R}^2 \mid y = x, 0 < x < 3\}$$




# Notation: Sets

$$L := \{(x, y) \in \mathbb{R}^2 \mid y = x, 0 < x < 3\}$$



# Notation: Sets

$$L := \{(x, y) \in \mathbb{R}^2 \mid y = x, x \in (0, 3)\}$$



# Notation: Functions

# Notation: Functions

$\mathbb{R}$

# Notation: Functions

$$\mathbb{R} \rightarrow \mathbb{R}$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$



# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$\mathbb{R}^2$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$\mathbb{R}^2 \rightarrow \mathbb{R}$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f((x, y)) = x - y$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$



# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : [0, 2\pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Notation: Functions**

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve”

**Notation: Functions**

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta)$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes



# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2\}$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), \}$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

- $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty)$



# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

- $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

- $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$   
 $\gamma(t) = (t, \frac{7t+3}{4})$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,

$$\gamma(t) = (f_1(t), f_2(t)), \text{ for planes}$$

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in P$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$   
 $\gamma(t) = (t, \frac{7t+3}{4}) \in P$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$   
 $\gamma(t) = (t, \frac{7t+3}{4}) \in P$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$   
 $\gamma$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,

$$\gamma(t) = (f_1(t), f_2(t)), \text{ for planes}$$

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in P$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

$$\gamma : (-\infty, \infty)$$

## Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,

$$\gamma(t) = (f_1(t), f_2(t)), \text{ for planes}$$

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

## Examples.

$$1. \ L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in P$$

$$2. \ P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

## Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,

$$\gamma(t) = (f_1(t), f_2(t)), \text{ for planes}$$

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

## Examples.

$$1. \ L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in P$$

$$2. \ P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t^2, t)$$



## Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,

$$\gamma(t) = (f_1(t), f_2(t)), \text{ for planes}$$

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

## Examples.

$$1. \ L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in P$$

$$2. \ P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t^2, t) \in P$$

# Notation: Functions

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x - 1.75$$

$$g : \{1, 2\} \rightarrow \{1, 2, 3\}$$

$$g(1) = 3$$

$$g(2) = 1$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$f(x, y) = x - y$$

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (\cos(t), \sin(t))$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,

$$\gamma(t) = (f_1(t), f_2(t)), \text{ for planes}$$

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

## Examples.

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in P$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t^2, t) \in P$$