

Theorem. *Given,*

- 1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$,*

Theorem. *Given,*

- 1. functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t ,*

Theorem. *Given,*

- 1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*

Theorem. *Given,*

- 1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
- 2. *$p \in \mathbb{R}^3$*

Theorem. *Given,*

- 1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
- 2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
- 3. *an orthonormal basis $\{\mathbf{E}_1, \}$*

Theorem. *Given,*

- 1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
- 2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
- 3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \}$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p ,*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization,

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$,*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$,*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*

2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*

3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*

2. *$p = \gamma(t_0)$*

3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{T}}(t) &= 0\mathbf{T}(t) + \kappa(t)\mathbf{N}(t) + 0\mathbf{B}(t) \\ \dot{\mathbf{N}}(t) &= -\kappa(t)\mathbf{T}(t) + 0\mathbf{N}(t) + \tau(t)\mathbf{B}(t) \\ \dot{\mathbf{B}}(t) &= 0\mathbf{T}(t) - \tau(t)\mathbf{N}(t) + 0\mathbf{B}(t)\end{aligned}$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*

2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*

3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*

2. *$p = \gamma(t_0)$*

3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations,

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

By the theory of differential equations, always has a solution, □

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$\mathbf{e}_1(t_0)$, $\mathbf{e}_2(t_0)$, and $\mathbf{e}_3(t_0)$.

□

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$\mathbf{e}_1(t_0) = \mathbf{E}_1$, $\mathbf{e}_2(t_0)$, and $\mathbf{e}_3(t_0)$.

□

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$\mathbf{e}_1(t_0) = \mathbf{E}_1$, $\mathbf{e}_2(t_0) = \mathbf{E}_2$, and $\mathbf{e}_3(t_0)$.

□

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

□

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$$

□

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0 \text{ (exercise!)}$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ □

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! □)

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration □)

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!) □

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*

2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*

3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*

2. *$p = \gamma(t_0)$*

3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!) □

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\begin{aligned}\kappa(t)\mathbf{N}(t) &= \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t) \\ \text{So, } \mathbf{N}(t) &= \mathbf{e}_2(t)\end{aligned}$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$$\mathbf{B}(t)$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$$\mathbf{B}(t) \text{ unit}$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t)$ and $\mathbf{N}(t)$. □

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\dot{\mathbf{e}}_1(t) = 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_2(t) = -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t)$$

$$\dot{\mathbf{e}}_3(t) = 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$. □

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$.

$$\mathbf{e}_3(t)$$

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$.

$\mathbf{e}_3(t)$ unit

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$.

$\mathbf{e}_3(t)$ unit and orthogonal

□

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$.

$\mathbf{e}_3(t)$ unit and orthogonal to $\mathbf{e}_1(t)$ and $\mathbf{e}_2(t)$. □

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0)$, $\mathbf{E}_2 = \mathbf{N}(t_0)$, $\mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$.

$\mathbf{e}_3(t)$ unit and orthogonal to $\mathbf{e}_1(t)$ and $\mathbf{e}_2(t)$. only two choices, and negatives of each other. So one is $\mathbf{e}_1(t) \times \mathbf{e}_2(t)$, other is $-\mathbf{e}_1(t) \times \mathbf{e}_2(t)$. \square

Theorem. *Given,*

1. *functions $\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}$, $\tilde{\kappa}(t) > 0$ for all t , and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}$*
2. *$p \in \mathbb{R}^3$, $t_0 \in (\alpha, \beta)$,*
3. *an orthonormal basis $\{\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_3\}$ based at p , such that $\mathbf{E}_3 = \mathbf{E}_1 \times \mathbf{E}_2$.*

there is a unit speed parametrization, $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$, so that,

1. *$\tilde{\kappa} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its curvature function and $\tilde{\tau} : (\alpha, \beta) \rightarrow \mathbb{R}^3$ is its torsion function*
2. *$p = \gamma(t_0)$*
3. *$\mathbf{E}_1 = \mathbf{T}(t_0), \mathbf{E}_2 = \mathbf{N}(t_0), \mathbf{E}_3 = \mathbf{B}(t_0)$*

Proof.

$$\begin{aligned}\dot{\mathbf{e}}_1(t) &= 0\mathbf{e}_1(t) + \kappa(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_2(t) &= -\kappa(t)\mathbf{e}_1(t) + 0\mathbf{e}_2(t) + \tau(t)\mathbf{e}_3(t) \\ \dot{\mathbf{e}}_3(t) &= 0\mathbf{e}_1(t) - \tau(t)\mathbf{e}_2(t) + 0\mathbf{e}_3(t)\end{aligned}$$

By the theory of differential equations, always has a solution, unique if we fix,

$$\mathbf{e}_1(t_0) = \mathbf{E}_1, \mathbf{e}_2(t_0) = \mathbf{E}_2, \text{ and } \mathbf{e}_3(t_0) = \mathbf{E}_3.$$

$(\mathbf{e}_i \cdot \mathbf{e}_j)' = 0$ (exercise!), so constant, so $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is orthonormal for *all* t

Can find $\gamma(t)$, so that, $\mathbf{T}(t) = \dot{\gamma}(t) = \mathbf{e}_1(t)$ and $\gamma(t_0) = p$ (exercise!! Integration (anti-derivative)!!)

$$\kappa(t)\mathbf{N}(t) = \dot{\mathbf{T}}(t) = \dot{\mathbf{e}}_1(t) = \kappa(t)\mathbf{e}_2(t)$$

$$\text{So, } \mathbf{N}(t) = \mathbf{e}_2(t)$$

$\mathbf{B}(t)$ unit and orthogonal to $\mathbf{T}(t) = \mathbf{e}_1(t)$ and $\mathbf{N}(t) = \mathbf{e}_2(t)$.

$\mathbf{e}_3(t)$ unit and orthogonal to $\mathbf{e}_1(t)$ and $\mathbf{e}_2(t)$. only two choices, and negatives of each other. So one is $\mathbf{e}_1(t) \times \mathbf{e}_2(t)$, other is $-\mathbf{e}_1(t) \times \mathbf{e}_2(t)$. Can distinguish using dot

products...

