

Chain rule for mult-variable functions

f

Chain rule for mult-variable functions

$$f : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = f_x(x(t_0), y(t_0))x'(t_0) + f_y(x(t_0), y(t_0))y'(t_0)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = f_x(x(t_0), y(t_0))x'(t_0) + f_y(x(t_0), y(t_0))y'(t_0) = (f_x(x, y), f_y(x, y)) \cdot \dot{\gamma}(t_0)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = f_x(x(t_0), y(t_0))x'(t_0) +$$

$$f_y(x(t_0), y(t_0))y'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0),$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0),$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{(f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0)},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$[f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0)],$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

f

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a γ

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta)$

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S$

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.
 $\sigma : U \rightarrow S$ a surface patch.

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t))$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(x(t), y(t)) + y'(t)\sigma_y(x(t), y(t))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(x(t), y(t)) + y'(t)\sigma_y(x(t), y(t))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(p) + y'(t)\sigma_y(p)$$

The tangent vectors at $p \in S \subset \mathbb{R}^3$ always belong to the span of $\sigma_x(p)$ and $\sigma_y(p)$.

Exercise. Show that any vector that belongs to the span of $\sigma_x(p)$ and $\sigma_y(p)$, is a tangent vector.

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(p) + y'(t)\sigma_y(p)$$

The tangent vectors at $p \in S \subset \mathbb{R}^3$ always belong to the span of $\sigma_x(p)$ and $\sigma_y(p)$.

Exercise. Show that any vector that belongs to the span of $\sigma_x(p)$ and $\sigma_y(p)$, is a tangent vector.

Exercise. Show that σ is regular at p if and only if the tangent vectors at p form a two dimensional subspace of \mathbb{R}^3 .

Inverse function theorem

f

Inverse function theorem

$$f : \mathbf{R}^3$$

Inverse function theorem

$$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$$

Inverse function theorem

$$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$$

i.e.

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x,y,z) = (f_1(x,y,z), f_2(x,y,z), f_3(x,y,z))$

define,

$$J(f)(p)$$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x,y,z) = (f_1(x,y,z), f_2(x,y,z), f_3(x,y,z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \end{pmatrix}$$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x,y,z) = (f_1(x,y,z), f_2(x,y,z), f_3(x,y,z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \end{pmatrix}$$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x,y,z) = (f_1(x,y,z), f_2(x,y,z), f_3(x,y,z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix}$$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x,y,z) = (f_1(x,y,z), f_2(x,y,z), f_3(x,y,z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem.

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f :$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$
and

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$
and $J(f)$ is invertible

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3.$

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$
and $J(f)$ is invertible (equivalently, $\det(J(f)(p)) \neq 0$

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$
*and $J(f)$ is invertible (equivalently, $\det(J(f)(p)) \neq 0$,
or the rows are linearly independent)*

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$
and $J(f)$ is invertible (equivalently, $\det(J(f)(p)) \neq 0$, or the rows are linearly independent), then f is invertible

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$
and $J(f)$ is invertible (equivalently, $\det(J(f)(p)) \neq 0$, or the rows are linearly independent), then f is invertible and has a smooth inverse

Inverse function theorem

$f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$.

i.e. $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$

define,

$$J(f)(p) = \begin{pmatrix} \frac{\partial f_1}{\partial x}(p) & \frac{\partial f_1}{\partial y}(p) & \frac{\partial f_1}{\partial z}(p) \\ \frac{\partial f_2}{\partial x}(p) & \frac{\partial f_2}{\partial y}(p) & \frac{\partial f_2}{\partial z}(p) \\ \frac{\partial f_3}{\partial x}(p) & \frac{\partial f_3}{\partial y}(p) & \frac{\partial f_3}{\partial z}(p) \end{pmatrix} = \begin{pmatrix} \nabla(f_1)(p) \\ \nabla(f_2)(p) \\ \nabla(f_3)(p) \end{pmatrix}$$

Theorem (Inverse function theorem). $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ and $J(f)$ is invertible (equivalently, $\det(J(f)(p)) \neq 0$, or the rows are linearly independent), then f is invertible and has a smooth inverse in a neighbourhood of p .

Applications of the inverse function theorem

Given $f(x, y, z) = 0$

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$?

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$?

Equivalently, can we find,

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$?

Equivalently, can we find,

$\theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$?

Equivalently, can we find,

$F(x, y, t) := (x, y, \theta(x, y))$ so that $f(x, y, \theta(x, y)) = t$

Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?


Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$?

Equivalently, can we find,

$F(x, y, t) := (x, y, \theta(x, y))$ so that $f(x, y, \theta(x, y)) = t$



Applications of the inverse function theorem

Given $f(x, y, z) = 0$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = 0$?

Making it stronger works!

Given $f(x, y, z) = t$, can we “solve for” z in terms of x and y ?

Can we write $z = \theta(x, y)$ so that $f(x, y, \theta(x, y)) = t$?



Equivalently, can we find,

$F(x, y, t) := (x, y, \theta(x, y))$ so that $f(x, y, \theta(x, y)) = t$

Equivalently, can we define an inverse of

$$G(x, y, z) := (x, y, f(x, y, z))$$

so that $G(F(x, y, z)) = (x, y, z)$



$$F_x = (1, 0, f_x(x, y, z))$$

$$F_y = (0, 1, f_y(x, y, z))$$

$$F_z = (0, 0, f_z(x, y, z))$$

linearly independent if and only if $f_z(x, y, z) \neq 0$

Can “solve for z” if and only if $f_z(x, y, z) \neq 0$

$$F_x = (1, 0, f_x(x, y, z))$$

$$F_y = (0, 1, f_y(x, y, z))$$

$$F_z = (0, 0, f_z(x, y, z))$$

linearly independent if and only if $f_z(x, y, z) \neq 0$

Can “solve for z” if and only if $f_z(x, y, z) \neq 0$

Can “solve for y” if and only if $f_y(x, y, z) \neq 0$

$$F_x = (1, 0, f_x(x, y, z))$$

$$F_y = (0, 1, f_y(x, y, z))$$

$$F_z = (0, 0, f_z(x, y, z))$$

linearly independent if and only if $f_z(x, y, z) \neq 0$

Can “solve for z” if and only if $f_z(x, y, z) \neq 0$

Can “solve for y” if and only if $f_y(x, y, z) \neq 0$

Can “solve for x” if and only if $f_x(x, y, z) \neq 0$

$$F_x = (1, 0, f_x(x, y, z))$$

$$F_y = (0, 1, f_y(x, y, z))$$

$$F_z = (0, 0, f_z(x, y, z))$$

linearly independent if and only if $f_z(x, y, z) \neq 0$

Can “solve for z” if and only if $f_z(x, y, z) \neq 0$

Can “solve for y” if and only if $f_y(x, y, z) \neq 0$

Can “solve for x” if and only if $f_x(x, y, z) \neq 0$

Can solve for either x , y , or z if and only if
 $\nabla(f)(x, y, z) \neq 0$

Understanding the cross products

Understanding the cross products

$$\mathbf{T}(t) \times \mathbf{N}(t) = \mathbf{B}(t)$$

Understanding the cross products

$$\mathbf{T}(t) \times \mathbf{N}(t) = \mathbf{B}(t)$$

$$\mathbf{B}(t) \times \mathbf{T}(t) = \mathbf{N}(t)$$

Understanding the cross products

$$\mathbf{T}(t) \times \mathbf{N}(t) = \mathbf{B}(t)$$

$$\mathbf{B}(t) \times \mathbf{T}(t) = \mathbf{N}(t)$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t)$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t)))$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

If $\kappa_s(t) = \kappa,$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

If $\kappa_s(t) = \kappa$,

$$\theta'(t) = \kappa$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

If $\kappa_s(t) = \kappa$,

$$\theta'(t) = \kappa$$

$$\theta(t) = \kappa t$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

If $\kappa_s(t) = \kappa$,

$$\theta'(t) = \kappa$$

$$\theta(t) = \kappa t + \theta_0$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

$$\text{If } \kappa_s(t) = \kappa,$$

$$\theta'(t) = \kappa$$

$$\theta(t) = \kappa t + \theta_0$$

$$\gamma'(t) = \mathbf{T}(t) = (\cos(at + \theta_0), \sin(at + \theta_0))$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

$$\text{If } \kappa_s(t) = \kappa,$$

$$\theta'(t) = \kappa$$

$$\theta(t) = \kappa t + \theta_0$$

$$\gamma'(t) = \mathbf{T}(t) = (\cos(at + \theta_0), \sin(at + \theta_0))$$

$$\gamma(t) = (\frac{1}{\kappa} \sin(\kappa t + \theta_0) + p_1, -\frac{1}{\kappa} \cos(\kappa t + \theta_0) + p_2)$$

$$\mathbf{T}(t) = (\cos(\theta(t)), \sin(\theta(t)))$$

$$\dot{\mathbf{T}}(t) = \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t)$$

$$\kappa_s(t) = \theta'(t)$$

$$\text{If } \kappa_s(t) = \kappa,$$

$$\theta'(t) = \kappa$$

$$\theta(t) = \kappa t + \theta_0$$

$$\gamma'(t) = \mathbf{T}(t) = (\cos(at + \theta_0), \sin(at + \theta_0))$$

$$\gamma(t) = (\frac{1}{\kappa} \sin(\kappa t + \theta_0) + p_1, -\frac{1}{\kappa} \cos(\kappa t + \theta_0) + p_2)$$

$$\gamma(t) = (\frac{1}{\kappa} \sin(\kappa t + \theta_0), -\frac{1}{\kappa} \cos(\kappa t + \theta_0)) + p$$

$$\begin{aligned}\mathbf{T}(t) &= (\cos(\theta(t)), \sin(\theta(t))) \\ \dot{\mathbf{T}}(t) &= \theta'(t)(-\sin(\theta(t)), \cos(\theta(t))) = \theta'(t)\mathbf{N}(t) \\ \kappa_s(t) &= \theta'(t)\end{aligned}$$

$$\begin{aligned}\text{If } \kappa_s(t) &= \kappa, \\ \theta'(t) &= \kappa \\ \theta(t) &= \kappa t + \theta_0\end{aligned}$$

$$\begin{aligned}\gamma'(t) &= \mathbf{T}(t) = (\cos(at + \theta_0), \sin(at + \theta_0)) \\ \gamma(t) &= (\tfrac{1}{\kappa} \sin(\kappa t + \theta_0) + p_1, -\tfrac{1}{\kappa} \cos(\kappa t + \theta_0) + p_2) \\ \gamma(t) &= (\tfrac{1}{\kappa} \sin(\kappa t + \theta_0), -\tfrac{1}{\kappa} \cos(\kappa t + \theta_0)) + p \\ \text{where } p &= (p_1, p_2)\end{aligned}$$

Exercise. If $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is a unit speed parametrization with constant curvature 0, then $\gamma(t)$ lies on a line for all t .

Solution. $\ddot{\gamma}(t) = 0$ □

Exercise. If $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is a unit speed parametrization with constant curvature 0, then $\gamma(t)$ lies on a line for all t .

Solution. $\ddot{\gamma}(t) = 0$

$\dot{\gamma}(t) = \mathbf{v}$ □

Exercise. If $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is a unit speed parametrization with constant curvature 0, then $\gamma(t)$ lies on a line for all t .

Solution. $\ddot{\gamma}(t) = 0$

$\dot{\gamma}(t) = \mathbf{v}$ for some $\mathbf{v} \in \mathbb{R}^2$ □

Exercise. If $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is a unit speed parametrization with constant curvature 0, then $\gamma(t)$ lies on a line for all t .

Solution. $\ddot{\gamma}(t) = 0$

$\dot{\gamma}(t) = \mathbf{v}$ for some $\mathbf{v} \in \mathbb{R}^2$

$\gamma(t) = t\mathbf{v} + p$ □

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) = 0$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t)\mathbf{N}(t).\dot{\mathbf{N}}(t) = 0$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t)\mathbf{N}(t).\dot{\mathbf{N}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t) = 0$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t)\mathbf{N}(t).\dot{\mathbf{N}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) = -\kappa_s(t)$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t)\mathbf{N}(t).\dot{\mathbf{N}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) = -\kappa_s(t)$$

$$\mathbf{N}(t).\mathbf{N}(t) = 0$$

$$\mathbf{N}(t).\mathbf{T}(t) = 0$$

$$(\mathbf{N}(t).\mathbf{T}(t))' = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t)\mathbf{N}(t).\dot{\mathbf{N}}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{T}(t) = -\kappa_s(t)$$

$$\mathbf{N}(t).\mathbf{N}(t) = 0$$

$$\dot{\mathbf{N}}(t).\mathbf{N}(t) = 0$$


So,

$$\begin{aligned}\mathbf{N}(t).\mathbf{T}(t) &= 0 \\ (\mathbf{N}(t).\mathbf{T}(t))' &= 0 \\ \dot{\mathbf{N}}(t).\mathbf{T}(t) + \mathbf{N}(t).\dot{\mathbf{T}}(t) &= 0 \\ \dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t)\mathbf{N}(t).\dot{\mathbf{N}}(t) &= 0 \\ \dot{\mathbf{N}}(t).\mathbf{T}(t) + \kappa_s(t) &= 0 \\ \dot{\mathbf{N}}(t).\mathbf{T}(t) &= -\kappa_s(t)\end{aligned}$$

$$\begin{aligned}\mathbf{N}(t).\mathbf{N}(t) &= 0 \\ \dot{\mathbf{N}}(t).\mathbf{N}(t) &= 0\end{aligned}$$

So,

$$\dot{\mathbf{N}}(t) = -\kappa(t)\mathbf{T}(t)$$



$$P = \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\}$$

$$\begin{aligned} P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\ &= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \end{aligned}$$

$$\begin{aligned}
P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
&= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
&= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
\end{aligned}$$

$$\begin{aligned}
 P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
 &= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
 &= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
 \end{aligned}$$

$\gamma :$

$$\begin{aligned}
 P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
 &= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c).(x, y, z) = d\} \\
 &= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n}.\mathbf{v} = d\}
 \end{aligned}$$

$$\gamma : (\alpha, \beta)$$

$$\begin{aligned}
P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
&= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
&= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
\end{aligned}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$$

$$\begin{aligned}
 P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
 &= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
 &= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
 \end{aligned}$$

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ parametrizes a curve

$$\begin{aligned}
P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
&= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
&= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
\end{aligned}$$

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ parametrizes a curve that lies on the plane, P ,

$$\begin{aligned}
P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
&= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
&= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
\end{aligned}$$

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ parametrizes a curve that lies on the plane, P , if and only if

$$\mathbf{n} \cdot \gamma(t) = d$$

$$\begin{aligned}
P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
&= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
&= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
\end{aligned}$$

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ parametrizes a curve that lies on the plane, P , if and only if

$$\mathbf{n} \cdot \gamma(t) = d$$

for all t

$$\begin{aligned}
P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\
&= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\
&= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\}
\end{aligned}$$

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ parametrizes a curve that lies on the plane, P , if and only if

$$\mathbf{n} \cdot \gamma(t) = d$$

for all $t \in (\alpha, \beta)$.

Alternatively,

$$\begin{aligned} P &= \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\} \\ &= \{(x, y, z) \in \mathbb{R}^3 \mid (a, b, c) \cdot (x, y, z) = d\} \\ &= \{\mathbf{v} \in \mathbb{R}^3 \mid \mathbf{n} \cdot \mathbf{v} = d\} \end{aligned}$$

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ parametrizes a curve that lies on the plane, P , if and only if

$$\mathbf{n} \cdot \gamma(t) = d$$

for all $t \in (\alpha, \beta)$.