

**Definition.**

$\gamma$

**Definition.**

$$\gamma : (\alpha, \beta)$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector*

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t)$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = s'(t)$



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = s'(t)\mathbf{T}(t)$ .

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \mathbf{T}(t)$ .

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}} .$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}} .$

*Solution.*  $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \square$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}}$ .

*Solution.*  $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \|\dot{\gamma}(t)\|$  □

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}} .$

*Solution.*  $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \|\dot{\gamma}(t)\| = \mathbf{T}(t) \times \square$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

Show that  $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}}$ .

*Solution.*  $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \|\dot{\gamma}(t)\| = \mathbf{T}(t) \times s'(t) \quad \square$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta)$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma}$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta})$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t})$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\dot{\tilde{\gamma}}(\tilde{t})$$



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t}))\end{aligned}$$



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If  $g(f(t)) = t$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then

$$g'(f(t))f'(t) = 1,$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore,



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

□

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

□

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.  
 $\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.  
 $\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.  
Show that,  
 $\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,  
If  $g(f(t)) = t$ , then  
 $g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned} \dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ?? \end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)}$$

□



**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\gamma'(t)\|}$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\gamma'(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))}$$

□

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

□

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

□

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned} \dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|} \end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$  is the *unit tangent vector* at  $t$ .

**Exercise.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$  a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

*Solution.*  $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned} \dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|} \\ &= \mathbf{T}(s^{-1}(\tilde{t})) \end{aligned}$$

□

Recall,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

**Exercise.**

If  $\|\dot{\gamma}(t)\|$

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ ,



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant.

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$s(t) =$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$s(t) = \int$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$s(t) = \int_{t_0}$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$s(t) = \int_{t_0}^t$$





**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$s(t) = \int_{t_0}^t \|\dot{\gamma}(t)\|$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$s(t) = \int_{t_0}^t \|\dot{\gamma}(t)\| dt$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned} s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \end{aligned}$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So,  $s(t)/c$  □

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So,  $s(t)/c = t - t_0$  □

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So,  $s(t)/c = t - t_0$

$\implies$   $\square$

**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So,  $s(t)/c = t - t_0$

$\implies s(t)/c - t$





**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So,  $s(t)/c = t - t_0$

$$\implies s(t)/c - t = -t_0$$



**Exercise.**

If  $\|\dot{\gamma}(t)\|$  is a constant  $c$ , show that  $t - s(t)/c$  is a constant. Here  $s(t)$  is the arc-length function “measured from (fixed)  $t_0$ ”.

*Solution.*

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

$$\begin{aligned}\text{So, } s(t)/c &= t - t_0 \\ \implies s(t)/c - t &= -t_0\end{aligned}$$

□

# Curvature

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization  
 $\|\ddot{\gamma}(t)\|$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$



# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$\gamma(t)$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t), r \sin(t))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t), r \sin(t))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t)$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (-r/r \sin(t/r), r/r \cos(t/r))$$



# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (-r/r \sin(t/r), r/r \cos(t/r))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), r/r \cos(t/r))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t)$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), r/r \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \cos(t/r), -1/r \sin(t/r))$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\|\ddot{\gamma}(t)\| = 1/r$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\|\ddot{\gamma}(t)\| = 1/r$$

# Curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$



# Curvature

**Exercise.** Show that the curvature

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

# Curvature

**Exercise.** Show that the curvature at any point of any line

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

# Curvature

**Exercise.** Show that the curvature at any point of any line is 0.

**Definition.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ , *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$  is the curvature, denoted  $\kappa(t)$ , at  $\gamma(t)$

**Example.**

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and

## Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),



## Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t)$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) =$$

## Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t)$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) =$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 =$$



# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) =$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) =$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) =$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 +$$



# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

# Deriving a general formula for curvature

$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$\kappa(t)$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$   
and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$\begin{aligned}s'(t) &= \|\dot{\gamma}(t)\| \\ (s'(t))^2 &= \dot{\gamma}(t) \cdot \dot{\gamma}(t) \\ 2(s'(t))s''(t) &= 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t) \\ 2\|\dot{\gamma}(t)\|s''(t) &= 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t) \\ s''(t) &= \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}\end{aligned}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\kappa(t) \; = \; \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\kappa(t) \; = \; \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|} = \ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\kappa(t) \; = \; \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 = \ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

# Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,  $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),  $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 = \ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\begin{aligned} \kappa(t) &= \|\ddot{\tilde{\gamma}}(s(t))\| \\ &= \left\| \frac{\ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}}{\|\dot{\gamma}(t)\|^2} \right\| \end{aligned}$$

## Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

$$\text{Equivalently, } \gamma(t) = \tilde{\gamma}(s(t))$$

$$\text{and (by definition), } \kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t) \cdot \dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 = \ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\begin{aligned} \kappa(t) &= \|\ddot{\tilde{\gamma}}(s(t))\| \\ &= \left\| \frac{\ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}}{\|\dot{\gamma}(t)\|^2} \right\| \\ &= \left\| \frac{\ddot{\gamma}(t) - \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}}{\|\dot{\gamma}(t)\|^2} \right\| \end{aligned}$$