

**Definition.** A “parametrized plane curve”

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta)$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2\}$



**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), \}$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:  
Image  $\gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$

**Examples.**

- 1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty)$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$   
 $\gamma(t) = (t, \frac{7t+3}{4})$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$   
 $\gamma(t) = (t, \frac{7t+3}{4}) \in L$



**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in L$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in L$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

$$\gamma$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

$$\begin{aligned} 1. \quad L &:= \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\} \\ \gamma &: (-\infty, \infty) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (t, \frac{7t+3}{4}) \in L \end{aligned}$$

$$\begin{aligned} 2. \quad P &:= \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\} \\ \gamma &: (-\infty, \infty) \end{aligned}$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$   
 $\gamma(t) = (t, \frac{7t+3}{4}) \in L$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$   
 $\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in L$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t^2, t)$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in L$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t^2, t) \in P$$

**Definition.** A “parametrized plane curve” is a function,  
 $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ .

Explicitly,  
 $\gamma(t) = (f_1(t), f_2(t))$ , for planes

Set of points on the curve:

$$\text{Image } \gamma = \{(x, y) \in \mathbb{R}^2 \mid (x, y) = \gamma(t), t \in \mathbb{R}\}$$

**Examples.**

1.  $L := \{(x, y) \in \mathbb{R}^2 \mid 4y = 7x + 3\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (t, \frac{7t+3}{4}) \in L$$

2.  $P := \{(x, y) \in \mathbb{R}^2 \mid y^2 = x\}$

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2$$


$$\gamma(t) = (t^2, t) \in P$$

# Parametrizing a circle







# Parametrizing a circle



# Parametrizing a circle




# Parametrizing a circle




# Parametrizing a circle

$$\gamma : (0, \pi/18) \rightarrow \mathbb{R}^2$$



# Parametrizing a circle


$$\gamma : (0, \pi/18) \rightarrow \mathbb{R}^2$$
$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$



# Parametrizing a circle

$$\gamma : (0, 2\pi/18) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$



# Parametrizing a circle


$$\gamma : (0, 3\pi/18) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$



# Parametrizing a circle

$$\gamma : (0, 6\pi/18) \rightarrow \mathbb{R}^2$$
$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$






# Parametrizing a circle

$$\gamma : (0, 9\pi/18) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$



# Parametrizing a circle

$$\gamma : (0, 12\pi/18) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$



# Parametrizing a circle


$$\gamma : (0, 18\pi/18) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$



# Parametrizing a circle


$$\gamma : (0, 24\pi/18) \rightarrow \mathbb{R}^2$$
$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$




# Parametrizing a circle

$$\gamma : (0, 27\pi/18) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$




# Parametrizing a line



$$\gamma : (0, 27\pi/18) \rightarrow \mathbb{R}^2$$
$$\gamma(t) := (2 \cos(t), 2 \sin(t))$$

# Parametrizing a line


$$\gamma : (-5, -4) \rightarrow \mathbb{R}^2$$



# Parametrizing a line

$$\gamma : (-5, -4) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (t, t)$$






# Parametrizing a line

$$\gamma : (-5, -3) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, -2) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, -1) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, 0) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, 1) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, 2) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, 3) \rightarrow \mathbb{R}^2$$


$$\gamma(t) := (t, t)$$



# Parametrizing a line

$$\gamma : (-5, 4) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (t, t)$$






# Parametrizing a line

$$\gamma : (-5, 5) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (t, t)$$



# Quick review: Derivative

# Quick review: Derivative

Example.  $f : \mathbb{R} \rightarrow \mathbb{R}$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \{$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \end{cases}$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \end{cases}$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$



# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x)$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x)$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^{\textcolor{red}{2}} & x > 5 \end{cases}$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$



# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

# Quick review: Derivative

Example.  $f(x) = x^2$

Example.  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

Example.

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$$

$f$  is “continuous”.

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$$

$f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$$

$f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$\begin{aligned} f(5) &= 0 \\ \lim_{x \rightarrow 5^-} f(x) &= 5^2 \\ \lim_{x \rightarrow 5^+} f(x) &= 5^3 \end{aligned}$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$   
 $\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$   
 $f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists,



# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$$

$f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, then  $f$  is “differentiable”

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$f(5) = 0$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2$$

$$\lim_{x \rightarrow 5^+} f(x) = 5^3$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$

$$\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$$

$f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, then  $f$  is “differentiable” and the limit is the derivative

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$\begin{aligned} f(5) &= 0 \\ \lim_{x \rightarrow 5^-} f(x) &= 5^2 \\ \lim_{x \rightarrow 5^+} f(x) &= 5^3 \end{aligned}$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$   
 $\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$   
 $f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, then  $f$  is “differentiable” and the limit is the derivative of  $f$

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$\begin{aligned} f(5) &= 0 \\ \lim_{x \rightarrow 5^-} f(x) &= 5^2 \\ \lim_{x \rightarrow 5^+} f(x) &= 5^3 \end{aligned}$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$   
 $\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$   
 $f$  is “continous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, then  $f$  is “differentiable” and the limit is the derivative of  $f$  at  $x$ ,

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$\begin{aligned} f(5) &= 0 \\ \lim_{x \rightarrow 5^-} f(x) &= 5^2 \\ \lim_{x \rightarrow 5^+} f(x) &= 5^3 \end{aligned}$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$   
 $\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$   
 $f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, then  $f$  is “differentiable” and the limit is the derivative of  $f$  at  $x$ , denoted  $f'(x)$  or  $\frac{df}{dx}$ .

# Quick review: Derivative

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} x^2 & x < 5 \\ 0 & x = 5 \\ x^3 & x > 5 \end{cases}$$

$$\begin{aligned} f(5) &= 0 \\ \lim_{x \rightarrow 5^-} f(x) &= 5^2 \\ \lim_{x \rightarrow 5^+} f(x) &= 5^3 \end{aligned}$$

**Example.**

$$f(x) = \begin{cases} x^2 & x \neq 5 \\ 0 & x = 5 \end{cases}$$

$$\lim_{x \rightarrow 5^-} f(x) = 5^2 = \lim_{x \rightarrow 5^+} f(x)$$

Can say,  $\lim_{x \rightarrow 5} f(x) = 5^2$

**Example.**  $f(x) = x^2$   
 $\lim_{x \rightarrow 5} f(x) = 5^2 = f(5)$   
 $f$  is “continuous”.

**Definition** (Continuous function).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if  $\lim_{x \rightarrow a} f(x) = f(a)$

**Definition** (Derivative). If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is such that

$$f'(x) := \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists, then  $f$  is “differentiable” and the limit is the derivative of  $f$  at  $x$ , denoted  $f'(x)$  or  $\frac{df}{dx}$ .