

Example. If the distance covered in time t is $f(t)$ for some function f , then $f'(t)$ is the velocity at time t .

Example. If the distance covered in time t is $f(t)$ for some function f , then $f'(t)$ is the velocity at time t .

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\gamma'(t) := (f_1'(t), f_2'(t), f_3'(t))$ is called the velocity of γ .

Example. If the distance covered in time t is $f(t)$ for some function f , then $f'(t)$ is the velocity at time t . $f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\gamma'(t) := (f_1'(t), f_2'(t), f_3'(t))$ is called the velocity of γ .

Example. If the distance covered in time t is $f(t)$ for some function f , then $f'(t)$ is the velocity at time t . $f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\gamma'(t) := (f_1'(t), f_2'(t), f_3'(t))$ is called the velocity of γ . $\gamma''(t)$ is called the acceleration.

Example. If the distance covered in time t is $f(t)$ for some function f , then $f'(t)$ is the velocity at time t . $f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f_1'(t), f_2'(t), f_3'(t))$ is called the velocity of γ . $\gamma''(t)$ is called the acceleration.

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f'_1(t), f'_2(t), f'_3(t))$ is called the velocity of γ . $\gamma''(t)$ is called the acceleration.

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f_1'(t), f_2'(t), f_3'(t))$ is called the velocity of γ . $\ddot{\gamma}(\mathbf{t}) := \gamma''(t)$ is called the acceleration.

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $\ddot{\mathbf{f}}(\mathbf{t}) := f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f'_1(t), f'_2(t), f'_3(t))$ is called the velocity of γ . $\ddot{\gamma}(\mathbf{t}) := \gamma''(t)$ is called the acceleration.

Different notations

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $\ddot{\mathbf{f}}(\mathbf{t}) := f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f'_1(t), f'_2(t), f'_3(t))$ is called the velocity of γ . $\ddot{\gamma}(\mathbf{t}) := \gamma''(t)$ is called the acceleration.

Different notations

$$\frac{d\gamma}{dt}(t) := \dot{\gamma}(t) := \gamma'(t)$$

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $\ddot{\mathbf{f}}(\mathbf{t}) := f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f'_1(t), f'_2(t), f'_3(t))$ is called the velocity of γ . $\ddot{\gamma}(\mathbf{t}) := \gamma''(t)$ is called the acceleration.

Different notations

$$\begin{aligned} \frac{d\gamma}{dt}(t) &:= \dot{\gamma}(t) := \gamma'(t) \\ \frac{d^2\gamma}{dt^2}(t) &:= \ddot{\gamma}(t) := \gamma''(t) \end{aligned}$$

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $\ddot{\mathbf{f}}(\mathbf{t}) := f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f'_1(t), f'_2(t), f'_3(t))$ is called the velocity of γ . $\ddot{\gamma}(\mathbf{t}) := \gamma''(t)$ is called the acceleration.

Different notations

$$\begin{aligned} \frac{d\gamma}{dt}(t) &:= \dot{\gamma}(t) := \gamma'(t) \\ \frac{d^2\gamma}{dt^2}(t) &:= \ddot{\gamma}(t) := \gamma''(t) \\ \frac{d^3\gamma}{dt^3}(t) &:= \dddot{\gamma}(t) := \gamma'''(t) \end{aligned}$$

Example. If the distance covered in time t is $f(t)$ for some function f , then $\dot{\mathbf{f}}(\mathbf{t}) := f'(t)$ is the velocity at time t . $\ddot{\mathbf{f}}(\mathbf{t}) := f''(t)$ is the acceleration.

Definition. $\gamma(t) = (f_1(t), f_2(t), f_3(t))$, then $\dot{\gamma}(\mathbf{t}) := \gamma'(t) := (f'_1(t), f'_2(t), f'_3(t))$ is called the velocity of γ . $\ddot{\gamma}(\mathbf{t}) := \gamma''(t)$ is called the acceleration.

Different notations

$$\begin{aligned} \frac{\mathrm{d}\gamma}{\mathrm{d}t}(t) &:= \dot{\gamma}(t) := \gamma'(t) \\ \frac{\mathrm{d}^2\gamma}{\mathrm{d}t^2}(t) &:= \ddot{\gamma}(t) := \gamma''(t) \\ \frac{\mathrm{d}^3\gamma}{\mathrm{d}t^3}(t) &:= \dddot{\gamma}(t) := \gamma'''(t) \\ \dots \end{aligned}$$

Derivative facts

Derivative facts

1. $c' = 0$

Derivative facts

1. $c' = 0$

2. $(x^n)' = nx^{n-1}$

Derivative facts

1. $c' = 0$

2. $(x^n)' = nx^{n-1}$

3. $(\sin(x))' = \cos(x)$

Derivative facts

1. $c' = 0$

2. $(x^n)' = nx^{n-1}$

3. $(\sin(x))' = \cos(x)$

4. $(\cos(x))' = -\sin(x)$

Derivative facts

1. $c' = 0$

2. $(x^n)' = nx^{n-1}$

3. $(\sin(x))' = \cos(x)$

4. $(\cos(x))' = -\sin(x)$

5. $(\tan(x))' = \sec^2(x)$

Derivative facts

1. $c' = 0$

2. $(x^n)' = nx^{n-1}$

3. $(\sin(x))' = \cos(x)$

4. $(\cos(x))' = -\sin(x)$

5. $(\tan(x))' = \sec^2(x)$

6. $(\log(x))' = \frac{1}{x}$

Derivative facts

1. $c' = 0$

2. $(x^n)' = nx^{n-1}$

3. $(\sin(x))' = \cos(x)$

4. $(\cos(x))' = -\sin(x)$

5. $(\tan(x))' = \sec^2(x)$

6. $(\log(x))' = \frac{1}{x}$

7. $(e^x)' = e^x$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' =$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' =$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' =$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' =$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' =$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' =$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' =$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' =$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) +$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' =$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' =$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' =$	

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' = f'(g(x))g'(x)$	$(\sin(x^2))' =$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' = f'(g(x))g'(x)$	$(\sin(x^2))' = \cos(x^2)2x$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Example.

$$(x^2 \sin(x^3) + \cos(x))' =$$

Rule

$$(cf)' = cf',$$

where $c \in \mathbb{R}$

$$(f + g)' = f' + g'$$

$$(f - g)' = f' - g'$$

$$(fg)' = f'g + fg'$$

$$(f/g)' = \frac{f'g - fg'}{g^2}$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

Example

$$(2 \sin(x))' = 2 \cos(x)$$

$$(\sin(x) + x^3)' = \cos(x) + 3x^2$$

$$(\sin(x) - x^3)' = \cos(x) - 3x^2$$

$$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$$

$$\left(\frac{\sin(x)}{x^2}\right)' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$$

$$(\sin(x^2))' = \cos(x^2)2x$$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Example.

$$(x^2 \sin(x^3) + \cos(x))' = 2x \sin(x^3) + x^2(\sin(x^3))'$$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' = f'(g(x))g'(x)$	$(\sin(x^2))' = \cos(x^2)2x$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Example.

$$\begin{aligned}(x^2 \sin(x^3) + \cos(x))' &= 2x \sin(x^3) + x^2(\sin(x^3))' \\ &\quad - \sin(x) \\ &= \end{aligned}$$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' = f'(g(x))g'(x)$	$(\sin(x^2))' = \cos(x^2)2x$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Example.

$$\begin{aligned}(x^2 \sin(x^3) + \cos(x))' &= 2x \sin(x^3) + x^2(\sin(x^3))' \\ &\quad - \sin(x) \\ &= 2x \sin(x^3) + x^2(3x^2 \cos(x^3))\end{aligned}$$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' = f'(g(x))g'(x)$	$(\sin(x^2))' = \cos(x^2)2x$

Derivative facts

- 1. $c' = 0$
- 2. $(x^n)' = nx^{n-1}$
- 3. $(\sin(x))' = \cos(x)$
- 4. $(\cos(x))' = -\sin(x)$
- 5. $(\tan(x))' = \sec^2(x)$
- 6. $(\log(x))' = \frac{1}{x}$
- 7. $(e^x)' = e^x$

Example.

$$\begin{aligned}(x^2 \sin(x^3) + \cos(x))' &= 2x \sin(x^3) + x^2(\sin(x^3))' \\ &\quad - \sin(x) \\ &= 2x \sin(x^3) + x^2(3x^2 \cos(x^3)) \\ &\quad - \sin(x)\end{aligned}$$

Rule	Example
$(cf)' = cf'$, where $c \in \mathbb{R}$	$(2 \sin(x))' = 2 \cos(x)$
$(f + g)' = f' + g'$	$(\sin(x) + x^3)' = \cos(x) + 3x^2$
$(f - g)' = f' - g'$	$(\sin(x) - x^3)' = \cos(x) - 3x^2$
$(fg)' = f'g + fg'$	$(x^2 \sin(x))' = 2x \sin(x) + x^2 \cos(x)$
$(f/g)' = \frac{f'g - fg'}{g^2}$	$(\frac{\sin(x)}{x^2})' = \frac{\cos(x)x^2 - \sin(x)2x}{x^4}$
$(f(g(x)))' = f'(g(x))g'(x)$	$(\sin(x^2))' = \cos(x^2)2x$

Example

Example

Example

γ :

Example

$$\gamma : (-\pi,$$

Example

$$\gamma : (-\pi, \pi)$$

Example

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

Example

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t)$$

Example

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (r \cos(t), r \sin(t))$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t)\end{aligned}$$

Example

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (r \cos(t), r \sin(t))$$

$$\dot{\gamma}(t) := (-r \sin(t), r \cos(t))$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, r \cos(t))\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\sqrt{v_1(t)^2 + v_2(t)^2}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\sqrt{v_1(t)^2 + v_2(t)^2} = \sqrt{(-r \sin(t))^2 + (r \cos(t))^2}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2((\sin(t))^2 + (\cos(t))^2)}\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1}\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2}\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Example

$$\gamma : (-\pi, \pi) \rightarrow \mathbb{R}^2$$

$$\gamma(t) := (r \cos(t), r \sin(t))$$

$$\dot{\gamma}(t) := (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})$$

The “speed” at time t is defined as

$$\begin{aligned} \sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r \end{aligned}$$

Can we find a parametrization of the circle

Example

Making the “speed” 1

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\gamma :$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\gamma : (-r\pi,$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\gamma : (-r\pi, r\pi)$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\gamma : (-r\pi, r\pi) \rightarrow \mathbb{R}^2$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r))\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (-r(1/r) \sin(t/r),\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (-r(1/r) \sin(t/r), r(1/r) \cos(t/r))\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (-\sin(t/r), \cos(t/r))\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \cos(t/r))\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\sqrt{v_1(t)^2 + v_2(t)^2}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\sqrt{v_1(t)^2 + v_2(t)^2} = \sqrt{(-\sin(t/r))^2 + (\cos(t/r))^2}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t/r))^2 + (\cos(t/r))^2} \\ &= \sqrt{\sin^2(t/r) + \cos^2(t/r)}\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t))^2 + (\cos(t))^2} \\ &= \sqrt{\sin^2(t) + \cos^2(t)} \\ &= 1\end{aligned}$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t))^2 + (\cos(t))^2} \\ &= \sqrt{\sin^2(t) + \cos^2(t)} \\ &= 1\end{aligned}$$

Acceleration of such a “unit speed parametrization” is,

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t))^2 + (\cos(t))^2} \\ &= \sqrt{\sin^2(t) + \cos^2(t)} \\ &= 1\end{aligned}$$

Acceleration of such a “unit speed parametrization” is,

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t))^2 + (\cos(t))^2} \\ &= \sqrt{\sin^2(t) + \cos^2(t)} \\ &= 1\end{aligned}$$

Acceleration of such a “unit speed parametrization” is,

$$\ddot{\gamma}(t)$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t))^2 + (\cos(t))^2} \\ &= \sqrt{\sin^2(t) + \cos^2(t)} \\ &= 1\end{aligned}$$

Acceleration of such a “unit speed parametrization” is,

$$\ddot{\gamma}(t) := (-1/r \cos(t/r),$$

Example

$$\begin{aligned}\gamma &: (-\pi, \pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t), r \sin(t)) \\ \dot{\gamma}(t) &:= (\underbrace{-r \sin(t)}_{v_1(t)}, \underbrace{r \cos(t)}_{v_2(t)})\end{aligned}$$

The “speed” at time t is defined as

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-r \sin(t))^2 + (r \cos(t))^2} \\ &= \sqrt{r^2 \underbrace{((\sin(t))^2 + (\cos(t))^2)}_1} \\ &= \sqrt{r^2} \\ &= r\end{aligned}$$

Can we find a parametrization of the circle to ensure the speed is 1?

Making the “speed” 1

$$\begin{aligned}\gamma &: (-r\pi, r\pi) \rightarrow \mathbb{R}^2 \\ \gamma(t) &:= (r \cos(t/r), r \sin(t/r)) \\ \dot{\gamma}(t) &:= (\underbrace{-\sin(t/r)}_{v_1(t)}, \underbrace{\cos(t/r)}_{v_2(t)})\end{aligned}$$

The “speed” is

$$\begin{aligned}\sqrt{v_1(t)^2 + v_2(t)^2} &= \sqrt{(-\sin(t))^2 + (\cos(t))^2} \\ &= \sqrt{\sin^2(t) + \cos^2(t)} \\ &= 1\end{aligned}$$

Acceleration of such a “unit speed parametrization” is,

$$\ddot{\gamma}(t) := (-1/r \cos(t/r), -1/r \sin(t/r))$$

A function is smooth

A function is smooth if it can be differentiated any number of times.

Example.

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$
 $f(x)' = \cos(x)$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$$\dot{\gamma}(t)$$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$$\dot{\gamma}(t) := \gamma'(t)$$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t)).$$

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

A function is smooth if it can be differentiated any number of times.

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

Note. We will study only smooth parametrizations.

From now on, all parametrizations will be assumed to be smooth.

A function is smooth if it can be differentiated any number of times. **Examples.**

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

1. $\gamma = (t, t)$

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

Note. We will study only smooth parametrizations.

From now on, all parametrizations will be assumed to be smooth.

A function is smooth if it can be differentiated any number of times. **Examples.**

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

1. $\gamma = (t, t)$ is a smooth parametrization.

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

Note. We will study only smooth parametrizations.

From now on, all parametrizations will be assumed to be smooth.

A function is smooth if it can be differentiated any number of times. **Examples.**

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

Note. We will study only smooth parametrizations.

From now on, all parametrizations will be assumed to be smooth.

1. $\gamma = (t, t)$ is a smooth parametrization. $\dot{\gamma}(t) = (1, 1)$.

2. $\gamma = (\cos(t), \sin(t))$

A function is smooth if it can be differentiated any number of times. **Examples.**

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

Note. We will study only smooth parametrizations.

From now on, all parametrizations will be assumed to be smooth.

1. $\gamma = (t, t)$ is a smooth parametrization. $\dot{\gamma}(t) = (1, 1)$.

2. $\gamma = (\cos(t), \sin(t))$ is a smooth parametrization.

A function is smooth if it can be differentiated any number of times. **Examples.**

Example. $f(x) = \sin(x)$

$$f(x)' = \cos(x)$$

$$f(x)'' = -\sin(x)$$

$$f(x)''' = -\cos(x)$$

$$f(x)'''' = \sin(x)$$

...

Definition. A parametrized plane curve $\gamma(t) := (f_1(t), f_2(t))$ is differentiable if $f_1(t)$ and $f_2(t)$ are differentiable.

$\dot{\gamma}(t) := \gamma'(t) := (f_1'(t), f_2'(t))$. It is smooth if it can be differentiated any number of times.

Note. We will study only smooth parametrizations.

From now on, all parametrizations will be assumed to be smooth.

1. $\gamma = (t, t)$ is a smooth parametrization. $\dot{\gamma}(t) = (1, 1)$.

2. $\gamma = (\cos(t), \sin(t))$ is a smooth parametrization. $\dot{\gamma}(t) = (-\sin(t), \cos(t))$.