

Chain rule for mult-variable functions

f

Chain rule for mult-variable functions

$$f : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = f_x(x(t_0), y(t_0))x'(t_0) + f_y(x(t_0), y(t_0))y'(t_0)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = f_x(x(t_0), y(t_0))x'(t_0) + f_y(x(t_0), y(t_0))y'(t_0) = (f_x(x, y), f_y(x, y)) \cdot \dot{\gamma}(t_0)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = f_x(x(t_0), y(t_0))x'(t_0) +$$

$$f_y(x(t_0), y(t_0))y'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0),$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$(f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0),$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{(f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0)},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \dot{\gamma}(t_0)},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(x(t_0), y(t_0)) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

f

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v)$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,

$\mathbf{v} = \dot{\gamma}(t_0)$,

and $p = (x(t_0), y(t_0))$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a γ

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta)$

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S$

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v},$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

Chain rule for mult-variable functions

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R} \end{aligned}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

where $\nabla(f)(x, y) = (f_x(x, y), f_y(x, y))$,
 $\mathbf{v} = \dot{\gamma}(t_0)$,
and $p = (x(t_0), y(t_0))$

$$\begin{aligned} f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R} \end{aligned}$$

$$\begin{aligned} &(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0) \end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.
 $\sigma : U \rightarrow S$ a surface patch.

Chain rule for mult-variable functions

$$\begin{aligned}f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}\end{aligned}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\begin{aligned}\text{where } \nabla(f)(x, y) &= (f_x(x, y), f_y(x, y)), \\ \mathbf{v} &= \dot{\gamma}(t_0), \\ \text{and } p &= (x(t_0), y(t_0))\end{aligned}$$

$$\begin{aligned}f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}\end{aligned}$$

$$\begin{aligned}(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)\end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t))$

Chain rule for mult-variable functions

$$\begin{aligned}f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \gamma(t) &= (x(t), y(t)) \\ f \circ \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}\end{aligned}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\begin{aligned}\text{where } \nabla(f)(x, y) &= (f_x(x, y), f_y(x, y)), \\ \mathbf{v} &= \dot{\gamma}(t_0), \\ \text{and } p &= (x(t_0), y(t_0))\end{aligned}$$

$$\begin{aligned}f &: \mathbb{R}^2 \rightarrow \mathbb{R} \\ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ \gamma(u, v) &= (x(u, v), y(u, v)) \\ f \circ \gamma &: \mathbb{R}^2 \rightarrow \mathbb{R}\end{aligned}$$

$$\begin{aligned}(f \circ \gamma)_u(u_0, v_0) \\ &= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0) \\ &+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)\end{aligned}$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(x(t), y(t)) + y'(t)\sigma_y(x(t), y(t))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(x(t), y(t)) + y'(t)\sigma_y(x(t), y(t))$$

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(p) + y'(t)\sigma_y(p)$$

The tangent vectors at $p \in S \subset \mathbb{R}^3$ always belong to the span of $\sigma_x(p)$ and $\sigma_y(p)$.

Exercise. Show that any vector that belongs to the span of $\sigma_x(p)$ and $\sigma_y(p)$, is a tangent vector.

Chain rule for mult-variable functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma(t) = (x(t), y(t))$$

$$f \circ \gamma : (\alpha, \beta) \rightarrow \mathbb{R}$$

$$\boxed{f_{\mathbf{v}}(x(t_0), y(t_0)) := (f \circ \gamma)'(t_0) = \nabla(f)(p) \cdot \mathbf{v}},$$

$$\text{where } \nabla(f)(x, y) = (f_x(x, y), f_y(x, y)),$$

$$\mathbf{v} = \dot{\gamma}(t_0),$$

$$\text{and } p = (x(t_0), y(t_0))$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$\gamma(u, v) = (x(u, v), y(u, v))$$

$$f \circ \gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$(f \circ \gamma)_u(u_0, v_0)$$

$$= f_x(x(u_0, v_0), y(u_0, v_0))x_u(u_0, v_0)$$

$$+ f_y(x(u_0, v_0), y(u_0, v_0))y_u(u_0, v_0)$$

Curves on surfaces

Definition. $\mathbf{v} \in \mathbb{R}^3$ is a tangent vector of the surface S at a point p , if there is a $\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ so that $p = \gamma(t)$ and $\mathbf{v} = \dot{\gamma}(t)$

$\gamma : (\alpha, \beta) \rightarrow S \subset \mathbb{R}^3$ is a curve.

$\sigma : U \rightarrow S$ a surface patch.

So, $\gamma(t) = \sigma(x(t), y(t)) = p \in S$

$$\dot{\gamma}(t) = x'(t)\sigma_x(p) + y'(t)\sigma_y(p)$$

The tangent vectors at $p \in S \subset \mathbb{R}^3$ always belong to the span of $\sigma_x(p)$ and $\sigma_y(p)$.

Exercise. Show that any vector that belongs to the span of $\sigma_x(p)$ and $\sigma_y(p)$, is a tangent vector.

Exercise. Show that σ is regular at p if and only if the tangent vectors at p form a two dimensional subspace of \mathbb{R}^3 .