

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta]$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,



$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$f(t_1)(t_2 - t_1) +$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$f(t_1)(t_2 - t_1) + f(t_2)(t_3 - t_2) +$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$f(t_1)(t_2 - t_1) + f(t_2)(t_3 - t_2) + \dots$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - t_1) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots + ||\dot{\gamma}(t_{n-1})|| (t_n - t_{n-1})$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2-t_1}_a)+||\dot{\gamma}(t_2)|| (t_3-t_2)+\dots+||\dot{\gamma}(t_{n-1})||(t_n-t_{n-1})$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

Better and better approximations “converge”.

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

Better and better approximations “converge”. Denoted,

$$||\dot{\gamma}(t)||$$



$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

Better and better approximations “converge”. Denoted,

$$||\dot{\gamma}(t)||dt$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

Better and better approximations “converge”. Denoted,

$$\int ||\dot{\gamma}(t)|| dt$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a} ||\dot{\gamma}(t)|| \mathrm{d}t$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)|| (t_2 - \underbrace{t_1}_a) + ||\dot{\gamma}(t_2)|| (t_3 - t_2) + \dots +$$

$$||\dot{\gamma}(t_{n-1})|| (\underbrace{t_n}_a - t_{n-1})$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)|| dt$$

# Arc length

## Definition.

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)|| dt$$

## Arc length

### Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!)

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)|| dt$$

## Arc length

### Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.



$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

## Arc length

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ ,

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

## Arc length

### Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

## Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t)$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

## Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

## Exercise.

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

## Arc length

### Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

### Exercise.

$$s_{\alpha}(t)$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

## Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

## Exercise.

$$s_{\alpha}(t) := \int_{t_{\alpha}}^t ||\dot{\gamma}(u)||du$$

$$s_{\beta}(t)$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

**Definition.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

**Exercise.**

$$s_{\alpha}(t) := \int_{t_{\alpha}}^t ||\dot{\gamma}(u)||du$$

$$s_{\beta}(t) := \int_{t_{\beta}}^t ||\dot{\gamma}(u)||du$$



$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$f(t) = ||\dot{\gamma}(t)||$$

$$f : [\alpha, \beta] \rightarrow \mathbb{R} \text{ continuous}$$

The arc-length,  $s$  from  $t = a$  to  $t = b$  is approximated by,

$$||\dot{\gamma}(t_1)||(\underbrace{t_2 - t_1}_a) + ||\dot{\gamma}(t_2)||(\underbrace{t_3 - t_2}_a) + \dots + ||\dot{\gamma}(t_{n-1})||(\underbrace{t_n - t_{n-1}}_a)$$

Better and better approximations “converge”. Denoted,

$$\int_{t=a}^{t=b} ||\dot{\gamma}(t)||dt$$

# Arc length

## Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular (of course!) parametrization.

Arc-length beginning at  $t_0$ , denoted  $s(t)$ ,

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

## Exercise.

$$s_{\alpha}(t) := \int_{t_{\alpha}}^t ||\dot{\gamma}(u)||du$$

$$s_{\beta}(t) := \int_{t_{\beta}}^t ||\dot{\gamma}(u)||du$$

Prove that  $s_{\beta}(t) - s_{\alpha}(t)$  is a constant.

**Theorem** (First Fundamental theorem of calculus).

$f$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$F(t)$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u)du$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u)du$$

*then,*

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) \mathrm{d}u$$

*then,  $F'(t) = f(t)$*

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) \mathrm{d}u$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$



**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) \mathrm{d}u$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  *is a smooth and regular*

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) \mathrm{d}u$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  *is a smooth and regular parametrization.*

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) \mathrm{d}u$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  *is a smooth and regular parametrization.*

*and  $s(t)$  its arc-length function beginning at  $t_0$  then,*

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = ||\dot{\gamma}(t)||$$

*Proof.*

$$s(t)$$



**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) du$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  *is a smooth and regular parametrization.*

*and  $s(t)$  its arc-length function beginning at  $t_0$  then,*

$$s'(t) = ||\dot{\gamma}(t)||$$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)|| du$$

□

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) du$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  *is a smooth and regular parametrization.*

*and  $s(t)$  its arc-length function beginning at  $t_0$  then,*

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus, □

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  *continuous*

$$F(t) := \int_{t_0}^t f(u) du$$

*then,  $F'(t) = f(t)$*

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  *is a smooth and regular parametrization.*

*and  $s(t)$  its arc-length function beginning at  $t_0$  then,*

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$  □

**Theorem** (First Fundamental theorem of calculus).  
 $f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = \|\dot{\gamma}(t)\|$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** *The arc length function*



**Theorem** (First Fundamental theorem of calculus).  
 $f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = \|\dot{\gamma}(t)\|$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) =$

$$\|\dot{\gamma}(t)\| \quad \square$$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*

$\square$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth  $\square$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why?  $\square$ )

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!).  $\square$

Observe,

If  $g(f(t)) = t$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!).  $\square$

Observe,

If  $g(f(t)) = t$ , then

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!)  $\square$

Observe,

If  $g(f(t)) = t$ , then

$$g'(f(t))f'(t) = 1,$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!).  $\square$

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore,



**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!).  $\square$

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!).  $\square$

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u) du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,

$$s'(t) = \|\dot{\gamma}(t)\|$$

*Proof.*

$$s(t) := \int_{t_0}^t \|\dot{\gamma}(u)\| du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = \|\dot{\gamma}(t)\|$   $\square$

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = \|\dot{\gamma}(t)\|$  which is itself smooth (Why? exercise!).  $\square$

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)}$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{||\gamma'(t)||}$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{||\dot{\gamma}(t)||}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))}$$



**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{||\gamma'(t)||}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

**Theorem** (First Fundamental theorem of calculus).

$f : [\alpha, \beta] \rightarrow \mathbb{R}$  continuous

$$F(t) := \int_{t_0}^t f(u)du$$

then,  $F'(t) = f(t)$

**Corollary.**  $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  is a smooth and regular parametrization.

and  $s(t)$  its arc-length function beginning at  $t_0$  then,  
 $s'(t) = ||\dot{\gamma}(t)||$

*Proof.*

$$s(t) := \int_{t_0}^t ||\dot{\gamma}(u)||du$$

by the First Fundamental Theorem of Calculus,  $s'(t) = ||\dot{\gamma}(t)||$  □

**Corollary.** The arc length function  $s(t)$  is smooth.

*Proof.*  $s'(t) = ||\dot{\gamma}(t)||$  which is itself smooth (Why? exercise!) □

Observe,

If  $g(f(t)) = t$ , then

$g'(f(t))f'(t) = 1$ , therefore, if  $f'(t) \neq 0$  for any  $t$ ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking  $f(t) = s(t)$  and  $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{||\gamma'(t)||}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$\gamma :$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} :$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$



$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t})$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t})$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$\text{If } \phi(t) = s^{-1}(t),$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$\text{If } \phi(t) = s^{-1}(t),$$

$$\tilde{\gamma}'(\tilde{t})$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$\text{If } \phi(t) = s^{-1}(t),$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$\text{If } \phi(t) = s^{-1}(t),$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})||$$



$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$\text{If } \phi(t) = s^{-1}(t),$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$\gamma :$

$$\begin{aligned}\gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma} &: (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma}(\tilde{t}) &= \gamma(\phi(\tilde{t})) \\ \tilde{\gamma}'(\tilde{t}) &= \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})\end{aligned}$$

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$$\gamma : (\alpha, \beta) \rightarrow$$

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$$

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})$$

$$\text{If } \phi(t) = s^{-1}(t),$$

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular smooth parametrization

$$\begin{aligned} \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma} &: (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma}(\tilde{t}) &= \gamma(\phi(\tilde{t})) \\ \tilde{\gamma}'(\tilde{t}) &= \gamma'(\phi(\tilde{t}))\phi'(\tilde{t}) \end{aligned}$$

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular smooth parametrization  $s(t)$ ,

$$\begin{aligned}\gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma} &: (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma}(\tilde{t}) &= \gamma(\phi(\tilde{t})) \\ \tilde{\gamma}'(\tilde{t}) &= \gamma'(\phi(\tilde{t}))\phi'(\tilde{t})\end{aligned}$$

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular smooth parametrization  $s(t)$ , the arc-length from  $t_0$  to  $t$

$$\begin{aligned} \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma} &: (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma}(\tilde{t}) &= \gamma(\phi(\tilde{t})) \\ \tilde{\gamma}'(\tilde{t}) &= \gamma'(\phi(\tilde{t}))\phi'(\tilde{t}) \end{aligned}$$

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular smooth parametrization  $s(t)$ , the arc-length from  $t_0$  to  $t$  (where  $t, t_0 \in (\alpha, \beta)$ )

$$\begin{aligned} \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma} &: (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma}(\tilde{t}) &= \gamma(\phi(\tilde{t})) \\ \tilde{\gamma}'(\tilde{t}) &= \gamma'(\phi(\tilde{t}))\phi'(\tilde{t}) \end{aligned}$$

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

Proved that,

**Theorem.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular smooth parametrization  $s(t)$ , the arc-length from  $t_0$  to  $t$  (where  $t, t_0 \in (\alpha, \beta)$ )  
 $\tilde{\gamma}(t) = \gamma(s^{-1}(t))$ , then

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$



$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{  \gamma'(s^{-1}(\tilde{t}))  }$
---

Proved that,

**Theorem.**

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$  a regular smooth parametrization  
 $s(t)$ , the arc-length from  $t_0$  to  $t$  (where  $t, t_0 \in (\alpha, \beta)$ )  
 $\tilde{\gamma}(t) = \gamma(s^{-1}(t))$ , then  
 $\tilde{\gamma}$  is a unit speed re-parametrization.

$$\begin{aligned} \gamma &: (\alpha, \beta) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma} &: (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2 \\ \tilde{\gamma}(\tilde{t}) &= \gamma(\phi(\tilde{t})) \\ \tilde{\gamma}'(\tilde{t}) &= \gamma'(\phi(\tilde{t}))\phi'(\tilde{t}) \end{aligned}$$

If  $\phi(t) = s^{-1}(t)$ ,

$$\tilde{\gamma}'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))(s^{-1})'(\tilde{t}) = \gamma'(s^{-1}(\tilde{t}))\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||}$$

$$||\tilde{\gamma}'(\tilde{t})|| = ||\gamma'(s^{-1}(\tilde{t}))||\frac{1}{||\gamma'(s^{-1}(\tilde{t}))||} = 1$$