

Definition. An n -clique

Definition. An n -*clique* is an n -graph in which each pair

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

Definition. An *n-clique* is an *n*-graph in which each pair from its *n*-vertices are joined by exactly one edge.

Generalized Pigeonhole principle
A

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f :$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A$

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow$

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B$

Definition. An n -*clique* is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}.$$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n -$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$,

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Example.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Example. In a group of 6 people,

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Example. In a group of 6 people, there is either a subset of 3

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Example. In a group of 6 people, there is either a subset of 3 who know each other,

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Definition. An *n-clique* is an *n*-graph in which each pair from its *n*-vertices are joined by exactly one edge.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3

Generalized Pigeonhole principle

A and *B* finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}.$

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some *i*.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set,

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A :=$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\}$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any b_i, b_j

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise,

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple

□

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,

then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other

and form the triple

□

Example.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,

then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other

and form the triple □

Example. Consider a graph with 6 vertices,

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Definition. An n -clique is an n -graph in which each pair from its n -vertices are joined by exactly one edge.

Generalized Pigeonhole principle

A and B finite sets

$f : A \rightarrow B := \{b_1, b_2, \dots, b_n\}$.

k_1, k_2, \dots, k_n are positive integers

If $|A| \geq k_1 + k_2 + \dots + k_n - (n - 1)$, then

$|f^{-1}(b_i)| \geq k_i$ for some i .

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices,

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be,

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V :=$

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.
 $16 = 3 \times 5 + 1,$

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other,
then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other
and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 ,

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise,

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours,

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple □

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above □

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

Solution.

Denote the set, $A := \{a_1, a_2, \dots, a_6\}$

a_1 either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$ (pigeonhole principle)

Assume a_1 knows $\{b_1, b_2, b_3\}$

If any $b_i, b_j \in \{b_1, b_2, b_3\}$ know each other, then $\{a_1, b_i, b_j\}$ form the triple.

Otherwise, none among $\{b_1, b_2, b_3\}$ know each other and form the triple \square

Example. Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution. Same as above \square

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number).

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Definition (Ramsay Number). $R(p, q)$ the smallest number n

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue,

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red, or a q -clique which is entirely blue.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red, or a q -clique which is entirely blue.

Example.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red, or a q -clique which is entirely blue.

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red, or a q -clique which is entirely blue.

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red, or a q -clique which is entirely blue.

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Definition (Ramsay Number). $R(p, q)$ the smallest number n such that a graph with n vertices, and each pair of vertices is joined by exactly one edge, and each coloured by either red or blue, will have either a p -clique which is entirely red, or a q -clique which is entirely blue.

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V :=$

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$
 v_1 is joined with $n - 1$ vertices.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1$

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) -$

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$
 v_1 is joined with $n - 1$ vertices.

$$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1),$$

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,

Either $R(p - 1, q)$ edges are red,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,

Either $R(p - 1, q)$ edges are red, or $R(p, q - 1)$ edges are blue.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,

Either $R(p - 1, q)$ edges are red, or $R(p, q - 1)$ edges are blue.

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_m}\}$

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.
 $16 = 3 \times 5 + 1$, so by pigeonhole principle,
 At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1
 Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$
 If even one of them are joined by the colour C_1 , we are done.
 Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$
 v_1 is joined with $n - 1$ vertices.
 $n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,
 Either $R(p - 1, q)$ edges are red, or $R(p, q - 1)$ edges are blue.
 Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_m}\}$
 They either have a q blue clique,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,

Either $R(p - 1, q)$ edges are red, or $R(p, q - 1)$ edges are blue.

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_m}\}$

They either have a q blue clique, then we are done.

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$

v_1 is joined with 16 vertices.

$16 = 3 \times 5 + 1$, so by pigeonhole principle,

At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$

If even one of them are joined by the colour C_1 , we are done.

Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. \square

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$

v_1 is joined with $n - 1$ vertices.

$n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,

Either $R(p - 1, q)$ edges are red, or $R(p, q - 1)$ edges are blue.

Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_m}\}$

They either have a q blue clique, then we are done. Or a $p - 1$ red clique,

Example. Consider a graph with 17 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, blue, or green. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_{17}\}$
 v_1 is joined with 16 vertices.
 $16 = 3 \times 5 + 1$, so by pigeonhole principle,
 At least $5 + 1 = 6$ edges containing v_1 are of the same colour C_1
 Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_6}\}$
 If even one of them are joined by the colour C_1 , we are done.
 Otherwise, the 6 vertices are joined by only two colours, reducing to the previous example. □

Example. Consider a graph with $n = R(p - 1, q) + R(p, q - 1)$ vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there is a red p -clique or a blue q -clique.

Solution.

Let the set of vertices be, $V := \{v_1, v_2, \dots, v_n\}$
 v_1 is joined with $n - 1$ vertices.
 $n - 1 = R(p - 1, q) + R(p, q - 1) - (2 - 1)$, so by pigeonhole principle,
 Either $R(p - 1, q)$ edges are red, or $R(p, q - 1)$ edges are blue.
 Let those edges be $\{v_{i_1}, v_{i_2}, \dots, v_{i_m}\}$
 They either have a q blue clique, then we are done. Or a $p - 1$ red clique, combined with v_1 , forms a p red clique. □