

Definition.

Definition (A path). A graph P

Definition (A path). A graph $P = (V(P),$

Definition (A path). A graph $P = (V(P), E(P))$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) =$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1}$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. A graph, G ,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length*

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation:



Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , \square

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , □

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

□

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n □

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. \square

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n 's neighbours. P 's length $\geq d(x_n)$ \square

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form, $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Definition.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P ’s length $\geq d(x_n) \geq \delta(G)$ □

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph C

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) =$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, \textcolor{red}{x_nx_0}\}$.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G ,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth).

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G ,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$,

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 .

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition.

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G ,*

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle,*

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .

$Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

Definition (A path). A graph $P = (V(P), E(P))$ on a set of vertices $V(P) = \{x_0, x_1, \dots, x_n\}$ is called a path of length n linking the “ends”, x_0 and x_n , if its set of edges is of the form,
 $E(P) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n\}$.

Notation:

$P := x_0x_1x_2 \dots x_n$ denotes a path linking x_0 with x_n .
 $Q := Px_{n+1} = x_0x_1x_2 \dots x_nx_{n+1}$

Proposition. *A graph, G , contains a path with length $\delta(G)$.*

Proof. Main observation: If a path P ends at x_n , but does not contain a “neighbour”, x_{n+1} of x_n , it can be made longer by including x_{n+1} to give Px_{n+1}

A path P with the largest length ending at x_n must contain all of x_n s neighbours. P 's length $\geq d(x_n) \geq \delta(G)$ □

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

$$g(G) \leq 2\text{diam } G + 1$$

Proof.

□ **Definition** (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
$$E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}.$$


Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

$$g(G) \leq 2\text{diam } G + 1$$



Proof.

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.


Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G . □

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

$$g(G) \leq 2\text{diam } G + 1$$



Proof.

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.


Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G . □

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

$$g(G) \leq 2\text{diam } G + 1$$



Proof.

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.

Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.


Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

□

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

$$g(G) \leq 2\text{diam } G + 1$$



Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.


Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*

$$g(G) \leq 2\text{diam } G + 1$$



Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square

Definition (A cycle). A graph $C = (V(C), E(C))$ on a set of vertices $V(C) = \{x_0, x_1, \dots, x_n\}$ is called a cycle of length $n + 1$ if its set of edges is of the form,
 $E(C) = \{x_0x_1, x_1x_2, \dots, x_{n-1}x_n, x_nx_0\}$.


Exercise. A graph, G , with at least one cycle contains a cycle with length *at least* $\delta(G) + 1$.

Definition (girth). The girth, $g(G)$, of a graph G is the minimum length of a cycle contained in G .

Definition. Given a graph G , the distance between two vertices $x_0, x_1 \in V(G)$, denoted $d_G(x_0, x_1)$, is the length of the shortest path joining x_0 and x_1 . The diameter of G , $\text{diam } G := \max\{d_G(x, y) \mid x, y \in V(G)\}$

Proposition. *Every graph, G , that contains at least one cycle, satisfies,*


$$g(G) \leq 2\text{diam } G + 1$$



Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square


Definition.



Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square


Definition. A (non-empty) graph, G , is connected



Proof.


Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square

Definition. A (non-empty) graph, G , is connected if any two of its vertices are linked by a path.



Proof.


Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square



Definition. A (non-empty) graph, G , is connected if any two of its vertices are linked by a path.
 $U \subset V(G)$ is connected

Proof.


Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square



Definition. A (non-empty) graph, G , is connected if any two of its vertices are linked by a path.
 $U \subset V(G)$ is connected if $G[U]$ is connected.

Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square




Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square

Definition. A (non-empty) graph, G , is connected if any two of its vertices are linked by a path.
 $U \subset V(G)$ is connected if $G[U]$ is connected.

Definition.




Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square

Definition. A (non-empty) graph, G , is connected if any two of its vertices are linked by a path.
 $U \subset V(G)$ is connected if $G[U]$ is connected.

Definition (component).




Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square

Definition. A (non-empty) graph, G , is connected if any two of its vertices are linked by a path.
 $U \subset V(G)$ is connected if $G[U]$ is connected.

Definition (component). A maximal connected subgraph is called a component




Definition. Graphs, G_1 and G_2 are called isomorphic if

Proof.


Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. □

Definition. Graphs, G_1 and G_2 are called isomorphic if there is a bijection



Proof.


Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square



Definition. Graphs, G_1 and G_2 are called isomorphic if there is a bijection $\phi : V(G_1) \rightarrow V(G_2)$ satisfying,

Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square



Proof.

Any cycle C has girth k , then it has at least two vertices with distance $k/2$. If $k/2 \geq \text{diam } G + 1$, then there is a path P joining the two points that is not in C . One can form cycle with smaller girth.. \square

Definition. Graphs, G_1 and G_2 are called isomorphic if there is a bijection $\phi : V(G_1) \rightarrow V(G_2)$ satisfying,
 $\{x, y\} \in E(G_1) \iff \{\phi(x), \phi(y)\} \in E(G_2)$