

Generating functions

Generating functions

Definition.

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) ,

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x)$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) :=$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 \dots$$

Example.

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 \dots$$

Example. Sequence:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$1 + x + x^2 + x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example.

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$1 + 2x + 3x^2 + 4x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example.

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $((\binom{n}{0}), (\binom{n}{1}), (\binom{n}{2}), \dots, (\binom{n}{n}), 0, 0, \dots)$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $((\binom{n}{0}), (\binom{n}{1}), (\binom{n}{2}), \dots, (\binom{n}{n}), 0, 0, \dots)$

Generating function:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$\binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Generating functions

Example.

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Generating functions

Example. Sequence:

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Generating functions

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Generating functions

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively,

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1 - x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1 - x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1 + x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1 - x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1 - x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1 + x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1 - x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1 - x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $((\binom{n}{0}), (\binom{n}{1}), (\binom{n}{2}), \dots, (\binom{n}{n}), 0, 0, \dots)$

Generating function:

$$(1 + x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1 - x)^n} = 1 + nx + \frac{n(n + 1)}{2!}x^2 + \frac{n(n + 1)(n + 2)}{3!}x^3 + \dots$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$(x^2 + x^4 + x^6 + \dots)^5$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \end{aligned}$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \end{aligned}$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$

Generating function:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$

Generating function:

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$

Generating function:

$$(1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right) \end{aligned}$$

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$
Generating function:

$$\frac{1}{1 - x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$
Generating function:

$$\frac{1}{(1 - x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, \dots)$
Generating function:

$$(1 + x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1 - x)^n} = 1 + nx + \frac{n(n + 1)}{2!}x^2 + \frac{n(n + 1)(n + 2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} &(x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1 - x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5 + k - 1}{k}x^{2k} + \dots \right) \end{aligned}$$

Answer:

Generating functions

Definition. Given a sequence (a_0, a_1, a_2, \dots) , its generating function is the power series,

$$A(x) := a_0 + a_1x + a_2x^2 + \dots$$

Example. Sequence: $(1, 1, 1, \dots)$
Generating function:

$$\frac{1}{1 - x} = 1 + x + x^2 + x^3 + \dots$$

Example. Sequence: $(1, 2, 3, \dots)$
Generating function:

$$\frac{1}{(1 - x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Example. Sequence: $((n \choose 0), (n \choose 1), (n \choose 2), \dots, (n \choose n), 0, 0, \dots)$
Generating function:

$$(1 + x)^n = {n \choose 0} + {n \choose 1}x + {n \choose 2}x^2 + {n \choose 3}x^3 + \dots$$

Example. Sequence: $(1, {n \choose 1}, {n+1 \choose 2}, {n+2 \choose 3}, \dots, {n+k-1 \choose k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1 - x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} &(x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1 - x^2)^5} \\ &= x^{10} \left(1 + {5 \choose 1}x^2 + \dots + {5+k-1 \choose k}x^{2k} + \dots \right) \end{aligned}$$

Answer: ${14 \choose 10}$

$$(a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots)$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
 Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
 Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right) \end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \cdots \end{aligned}$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
 Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
 Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right) \end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned}
& (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\
&= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \\
&c(a_0 + a_1x + a_2x^2 \dots)
\end{aligned}$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned}
& (x^2 + x^4 + x^6 + \dots)^5 \\
&= x^{10}(1 + x^2 + x^4 + \dots)^5 \\
&= x^{10} \frac{1}{(1-x^2)^5} \\
&= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)
\end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \dots) \\ &= ca_0 + ca_1x + ca_2x^2 \dots \end{aligned}$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
 Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
 Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right) \end{aligned}$$

Answer: $\binom{14}{10}$

$$(a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\ = (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots$$

$$c(a_0 + a_1x + a_2x^2 \dots) \\ = ca_0 + ca_1x + ca_2x^2 \dots$$

$$(a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots)$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
 Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$

Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$(x^2 + x^4 + x^6 + \dots)^5 \\ = x^{10}(1 + x^2 + x^4 + \dots)^5 \\ = x^{10} \frac{1}{(1-x^2)^5} \\ = x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)$$

Answer: $\binom{14}{10}$

$$\begin{aligned}
& (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\
&= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \\
& c(a_0 + a_1x + a_2x^2 \dots) \\
&= ca_0 + ca_1x + ca_2x^2 \dots \\
& (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\
&= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots
\end{aligned}$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned}
& (x^2 + x^4 + x^6 + \dots)^5 \\
&= x^{10}(1 + x^2 + x^4 + \dots)^5 \\
&= x^{10} \frac{1}{(1-x^2)^5} \\
&= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)
\end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned}
& (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\
&= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \\
& c(a_0 + a_1x + a_2x^2 \dots) \\
&= ca_0 + ca_1x + ca_2x^2 \dots \\
& (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\
&= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots
\end{aligned}$$

Example.

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned}
& (x^2 + x^4 + x^6 + \dots)^5 \\
&= x^{10}(1 + x^2 + x^4 + \dots)^5 \\
&= x^{10} \frac{1}{(1-x^2)^5} \\
&= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)
\end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \dots) \\ &= ca_0 + ca_1x + ca_2x^2 \dots \end{aligned}$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\ &= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots \end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
 Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
 Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right) \end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \dots) \\ &= ca_0 + ca_1x + ca_2x^2 \dots \end{aligned}$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\ &= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots \end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
 Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
 Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned} & (x^2 + x^4 + x^6 + \dots)^5 \\ &= x^{10}(1 + x^2 + x^4 + \dots)^5 \\ &= x^{10} \frac{1}{(1-x^2)^5} \\ &= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right) \end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned}
& (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\
&= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \\
& c(a_0 + a_1x + a_2x^2 \dots) \\
&= ca_0 + ca_1x + ca_2x^2 \dots \\
& (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\
&= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots
\end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2}$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned}
& (x^2 + x^4 + x^6 + \dots)^5 \\
&= x^{10}(1 + x^2 + x^4 + \dots)^5 \\
&= x^{10} \frac{1}{(1-x^2)^5} \\
&= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)
\end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned}
& (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\
&= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \\
& c(a_0 + a_1x + a_2x^2 \dots) \\
&= ca_0 + ca_1x + ca_2x^2 \dots \\
& (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\
&= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots
\end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned}
& (x^2 + x^4 + x^6 + \dots)^5 \\
&= x^{10}(1 + x^2 + x^4 + \dots)^5 \\
&= x^{10} \frac{1}{(1-x^2)^5} \\
&= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)
\end{aligned}$$

Answer: $\binom{14}{10}$

$$\begin{aligned}
& (a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\
&= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots \\
& c(a_0 + a_1x + a_2x^2 \dots) \\
&= ca_0 + ca_1x + ca_2x^2 \dots \\
& (a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\
&= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots
\end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\begin{aligned}
& \frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \dots \\
& \frac{2}{(1-x)^3}
\end{aligned}$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$
Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$\begin{aligned}
& (x^2 + x^4 + x^6 + \dots)^5 \\
&= x^{10}(1 + x^2 + x^4 + \dots)^5 \\
&= x^{10} \frac{1}{(1-x^2)^5} \\
&= x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)
\end{aligned}$$

Answer: $\binom{14}{10}$

$$(a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\ = (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots$$

$$c(a_0 + a_1x + a_2x^2 \dots) \\ = ca_0 + ca_1x + ca_2x^2 \dots$$

$$(a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\ = a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \dots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$

Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$(x^2 + x^4 + x^6 + \dots)^5 \\ = x^{10}(1 + x^2 + x^4 + \dots)^5 \\ = x^{10} \frac{1}{(1-x^2)^5} \\ = x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)$$

Answer: $\binom{14}{10}$

$$(a_0 + a_1x + a_2x^2 \dots) + (b_0 + b_1x + b_2x^2 \dots) \\ = (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \dots$$

$$c(a_0 + a_1x + a_2x^2 \dots) \\ = ca_0 + ca_1x + ca_2x^2 \dots$$

$$(a_0 + a_1x + a_2x^2 \dots)(b_0 + b_1x + b_2x^2 \dots) \\ = a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \dots$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \dots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \dots$$

Example. Sequence: $(1, \binom{n}{1}, \binom{n+1}{2}, \binom{n+2}{3}, \dots, \binom{n+k-1}{k}, \dots)$
Alternatively, $(1, n, \frac{n(n+1)}{2!}, \frac{n(n+1)(n+2)}{3!}, \dots)$

Generating function:

$$\frac{1}{(1-x)^n} = 1 + nx + \frac{n(n+1)}{2!}x^2 + \frac{n(n+1)(n+2)}{3!}x^3 + \dots$$

Example. Find the coefficient of x^{30} in $(x^2 + x^4 + x^6 + \dots)^5$.

Solution.

$$(x^2 + x^4 + x^6 + \dots)^5 \\ = x^{10}(1 + x^2 + x^4 + \dots)^5 \\ = x^{10} \frac{1}{(1-x^2)^5} \\ = x^{10} \left(1 + \binom{5}{1}x^2 + \dots + \binom{5+k-1}{k}x^{2k} + \dots \right)$$

Answer: $\binom{14}{10}$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \cdots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \cdots) \\ &= ca_0 + ca_1x + ca_2x^2 \cdots \end{aligned}$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots)(b_0 + b_1x + b_2x^2 \cdots) \\ &= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \cdots \end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \cdots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \cdots$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \cdots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \cdots) \\ &= ca_0 + ca_1x + ca_2x^2 \cdots \end{aligned}$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots)(b_0 + b_1x + b_2x^2 \cdots) \\ &= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \cdots \end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \cdots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \cdots$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution.

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \cdots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \cdots) \\ &= ca_0 + ca_1x + ca_2x^2 \cdots \end{aligned}$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots)(b_0 + b_1x + b_2x^2 \cdots) \\ &= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \cdots \end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \cdots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \cdots$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \cdots (1 + x + x^2 + \cdots)$

$$(a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots) \\ = (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \cdots$$

$$c(a_0 + a_1x + a_2x^2 \cdots) \\ = ca_0 + ca_1x + ca_2x^2 \cdots$$

$$(a_0 + a_1x + a_2x^2 \cdots)(b_0 + b_1x + b_2x^2 \cdots) \\ = a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \cdots$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \cdots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \cdots$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

$$\textit{Solution.} \quad (1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \cdots (1 + x + x^2 + \cdots) \\ = (1 + x + x^2 + \cdots)^n$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots) + (b_0 + b_1x + b_2x^2 \cdots) \\ &= (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 \cdots \end{aligned}$$

$$\begin{aligned} & c(a_0 + a_1x + a_2x^2 \cdots) \\ &= ca_0 + ca_1x + ca_2x^2 \cdots \end{aligned}$$

$$\begin{aligned} & (a_0 + a_1x + a_2x^2 \cdots)(b_0 + b_1x + b_2x^2 \cdots) \\ &= a_0b_0 + (a_1b_0 + a_0b_1)x + (a_2b_0 + a_1b_1 + a_0b_2)x^2 \cdots \end{aligned}$$

Example. Find the generating function for the sequence $(1^2, 2^2, 3^2, \dots, n^2, \dots)$

Solution.

$$\frac{1}{(1-x)^2} = 1 + x + 2x^2 + 3x^3 + \cdots$$

$$\frac{2}{(1-x)^3} = 1 + 2.2x + 3.3x^2 + \cdots$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

$$\begin{aligned} & \textit{Solution.} \quad (1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \cdots (1 + x + x^2 + \cdots) \\ &= (1 + x + x^2 + \cdots)^n \\ &= \frac{1}{(1-x)^{n+1}} \end{aligned}$$

Midsem question

What is the minimum number of numbers

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

$$\begin{aligned} \textit{Solution.} \quad & (1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \cdots (1 + \\ & x + x^2 + \cdots) \\ &= (1 + x + x^2 + \cdots)^n \\ &= \frac{1}{(1-x)^n} \end{aligned}$$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

$$\begin{aligned} \textit{Solution.} \quad & (1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots) \\ &= (1 + x + x^2 + \dots)^n \\ &= \frac{1}{(1-x)^n} \end{aligned}$$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution.

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

$$\begin{aligned} \textit{Solution.} \quad & (1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots) \\ &= (1 + x + x^2 + \dots)^n \\ &= \frac{1}{(1-x)^n} \end{aligned}$$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Solution. $(1 + x + x^2 + \dots)(1 + x + x^2 + \dots) \dots (1 + x + x^2 + \dots)$
 $= (1 + x + x^2 + \dots)^n$
 $= \frac{1}{(1-x)^n}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

Midsem question

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$
If it contained $2^r m$, where m is odd,

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$
Given any two $x, y \in A_i$, either $x|y$ or $y|x$.
So by pigeonhole principle, if we choose at least $k + 1$ numbers
then there will exist an x, y so that $x|y$
There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$
What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$
If it contained $2^r m$, where m is odd,
then 2^{r+1}

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$
If it contained $2^r m$, where m is odd, then 2^{r+1} would not divide it

$$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$$

$$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$
choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$
so that it has the highest power of 2

i.e. there exists m_1 and m_2

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$
If it contained $2^r m$, where m is odd, then 2^{r+1} would not divide it

$$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$$

$$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$
choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$
so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

such that $2^{k_1} m_1 \leq 60$ and $2^{k_2} m_2 \leq 60$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

such that $2^{k_1} m_1 \leq 60$ and $2^{k_2} m_2 \leq 60$

but $2^{k_1+1} m_1 > 60$ and $2^{k_2+1} m_2 > 60$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

such that $2^{k_1} m_1 \leq 60$ and $2^{k_2} m_2 \leq 60$

but $2^{k_1+1} m_1 > 60$ and $2^{k_2+1} m_2 > 60$

If $2^{k_1} m_1 | 2^{k_2} m_2$ (m_1 and m_2 are odd)

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

such that $2^{k_1} m_1 \leq 60$ and $2^{k_2} m_2 \leq 60$

but $2^{k_1+1} m_1 > 60$ and $2^{k_2+1} m_2 > 60$

If $2^{k_1} m_1 | 2^{k_2} m_2$ (m_1 and m_2 are odd)

$\implies m_1 | m_2 \implies m_2 \geq 2m_1$

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

such that $2^{k_1} m_1 \leq 60$ and $2^{k_2} m_2 \leq 60$

but $2^{k_1+1} m_1 > 60$ and $2^{k_2+1} m_2 > 60$

If $2^{k_1} m_1 | 2^{k_2} m_2$ (m_1 and m_2 are odd)

$\implies m_1 | m_2 \implies m_2 \geq 2m_1$

If $k_2 \geq k_1$, then $2^{k_2} m_2 \geq 2^{k_1+1} m_1 > 60$.

Midsem question

What is the minimum number of numbers one should select from $\{1, 2, 3, \dots, 60\}$ to guarantee that there is at least one number which divides another?

Solution. $\{1, 2, 3, \dots, 60\} = \sqcup_{i=1}^k A_i$

Given any two $x, y \in A_i$, either $x|y$ or $y|x$.

So by pigeonhole principle, if we choose at least $k + 1$ numbers

then there will exist an x, y so that $x|y$

There exists $a_i \in A_i$ so that $\{a_1, \dots, a_k\}$ so that a_i does not divide a_j for any $i \neq j$

What is the right choice of A_i s?

$A_1 := \{1, 2, 4, 8, \dots, 2^s\}$, where s is such that $2^{s+1} > 60$

If it contained $2^r m$, where m is odd,

then 2^{r+1} would not divide it

$A_3 := \{3, 2 \times 3, 2^2 \times 3, \dots, 2^? \times 3\}$

$A_m := \{m, 2 \times m, 2^2 \times m, \dots, 2^? \times m\}$

where m is odd

Given the set A_{m_1} and A_{m_2} , $m_1 \neq m_2$

choose $a_{m_1} \in A_{m_1}$ and $a_{m_2} \in A_{m_2}$

so that it has the highest power of 2

i.e. there exists m_1 and m_2 odd and k_1 and k_2

such that $2^{k_1} m_1 \leq 60$ and $2^{k_2} m_2 \leq 60$

but $2^{k_1+1} m_1 > 60$ and $2^{k_2+1} m_2 > 60$

If $2^{k_1} m_1 | 2^{k_2} m_2$ (m_1 and m_2 are odd)

$\implies m_1 | m_2 \implies m_2 \geq 2m_1$

If $k_2 \geq k_1$, then $2^{k_2} m_2 \geq 2^{k_1+1} m_1 > 60$.

which contradicts the choice of k_2

□