

Theorem.

**Theorem.** *The number of partitions of  $m$*

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers*

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice*

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* □

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)$  □



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)$   $\square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$= \frac{1-x^3}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1-x^3} \dots$$



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

□

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

□

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$\begin{aligned}
 & (1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots \\
 &= \frac{\cancel{1-x^3} 1-x^6 1-x^9}{1-x \cancel{1-x^2} \cancel{1-x^3}} \dots \\
 &= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots \\
 &= (1 + x + x^2 + \dots)(1 + x^2 + x^4 + \dots)(1 + x^4 + x^8 + \dots) \dots \quad \square
 \end{aligned}$$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

$$= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$$

$$= \text{number of} \quad \square$$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$

$$\begin{aligned}
 &= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots \\
 &= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots \\
 &= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots \\
 &= \text{number of partitions with sizes } \textit{not} \text{ multiples of } 3 \quad \square
 \end{aligned}$$



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes  $m$  not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

$$= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$$

$$= \text{number of partitions with sizes } not \text{ multiples of } 3 \quad \square$$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler).

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

$$= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$$

= number of partitions with sizes *not* multiples of 3     $\square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$*

*Proof.* Number of partitions is the  $m$ th coefficient of
 
$$\begin{aligned}
 &(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots \\
 &= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots \\
 &= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots \\
 &= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots \\
 &= \text{number of partitions with sizes } \textit{not} \text{ multiples of } 3 \quad \square
 \end{aligned}$$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $= \text{number of partitions with sizes } not \text{ multiples of } 3 \quad \square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes so that each term occurs at most once not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$$

$$= \frac{1-x^3}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1-x^3} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

$$= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$$

$$= \text{number of partitions with sizes } not \text{ multiples of } 3 \quad \square$$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$ .

$$\begin{aligned}
 &= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots \\
 &= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots \\
 &= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots \\
 &= \text{number of partitions with sizes } \textit{not} \text{ multiples of } 3 \quad \square
 \end{aligned}$$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

*Proof.*  $\square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1-x^3}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $\square$



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

$$= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$$

= number of partitions with sizes *not* multiples of 3     $\square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)$   $\square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1-x^3}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2) \dots$      $\square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$ 

$\square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of
 
$$(1+x+x^2)(1+x^2+x^4)(1+x^3+x^6)\dots$$

$$= \frac{\cancel{1-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{\cancel{1-x^3}} \dots$$

$$= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^4} \dots$$

$$= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots)\dots$$

$$= \text{number of partitions with sizes } \textit{not} \text{ multiples of } 3 \quad \square$$

*Proof.* Number of partitions is the  $m$ th coefficient of
 
$$(1+x)(1+x^2)(1+x^3)\dots$$

$$= \frac{1-x^2}{1-x} \frac{1-x^4}{1-x^2} \frac{1-x^6}{1-x^3} \dots$$

□

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1\cancel{-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1\cancel{-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1\cancel{-x^4}} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $= \text{number of partitions with sizes } not \text{ multiples of } 3 \quad \square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1\cancel{-x^2}}{1-x} \frac{1-x^4}{1\cancel{-x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$

□

.

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1\cancel{-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1\cancel{-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1\cancel{-x^4}} \dots$   
 $= (1 + x + x^2 + \dots)(1 + x^2 + x^4 + \dots)(1 + x^4 + x^8 + \dots) \dots$   
 $= \text{number of partitions with sizes } not \text{ multiples of } 3 \quad \square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1\cancel{-x^2}}{1-x} \frac{1-x^4}{1\cancel{-x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1 + x + x^2 + \dots)(1 + x^3 + x^6 + \dots)(1 + x^5 + x^{10} + \dots) \dots \quad \square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1\cancel{-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1\cancel{-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1\cancel{-x^4}} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $= \text{number of partitions with sizes } not \text{ multiples of } 3 \quad \square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of...*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1\cancel{-x^2}}{1-x} \frac{1-x^4}{1\cancel{-x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of} \quad \square$

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1\cancel{-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1\cancel{-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1\cancel{-x^4}} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1\cancel{-x^2}}{1-x} \frac{1-x^4}{1\cancel{-x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes .     $\square$



**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1\cancel{-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1\cancel{-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1\cancel{-x^4}} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1\cancel{-x^2}}{1-x} \frac{1-x^4}{1\cancel{-x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes .     $\square$

**Exercise.**

.

**Theorem.** *The number of partitions of  $m$  into a sum of non-zero integers so that each term occurs at most twice is equal to the number of partitions with sizes not multiples of 3.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x + x^2)(1 + x^2 + x^4)(1 + x^3 + x^6) \dots$   
 $= \frac{1\cancel{-x^3}}{1-x} \frac{1-x^6}{1-x^2} \frac{1-x^9}{1\cancel{-x^3}} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1\cancel{-x^4}} \dots$   
 $= (1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^4+x^8+\dots) \dots$   
 $=$  number of partitions with sizes *not* multiples of 3     $\square$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1\cancel{-x^2}}{1-x} \frac{1-x^4}{1\cancel{-x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes .     $\square$

**Exercise.** Generalize the above theorems to paritions of sizes at most  $k$ .

.

**Theorem.**

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1+x)(1+x^2)(1+x^3)\dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots)\dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} . \quad \square$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler).

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1+x)(1+x^2)(1+x^3)\dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots)\dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} . \quad \square$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} . \quad \square$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1 \cancel{x^2}}{1-x} \frac{1-x^4}{1 \cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with odd sizes} . \quad \square$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.*

□

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} .$  □

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

□ *Proof.* Number of partitions is the  $m$ th coefficient of

$$\begin{aligned}
 & (1+x)(1+x^2)(1+x^3)\dots \\
 &= \frac{\cancel{1-x^2}}{1-x} \frac{\cancel{1-x^4}}{\cancel{1-x^2}} \frac{1-x^6}{1-x^3} \dots \\
 &= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots \\
 &= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots)\dots \\
 &= \text{number of partitions with } \textit{odd} \text{ sizes} . \qquad \square
 \end{aligned}$$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1 :$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

□ *Proof.* Number of partitions is the  $m$ th coefficient of

$$\begin{aligned}
 &(1+x)(1+x^2)(1+x^3)\dots \\
 &= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots \\
 &= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots \\
 &= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots)\dots \\
 &= \text{number of partitions with } \textit{odd} \text{ sizes} .
 \end{aligned}$$

□

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

□ *Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$\begin{aligned} \square & (1+x)(1+x^2)(1+x^3)\dots \\ &= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots \\ &= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots \\ &= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots)\dots \\ &= \text{number of partitions with } \textit{odd} \text{ sizes} . \qquad \square \end{aligned}$$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  :

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of

$$\begin{aligned} \square & (1+x)(1+x^2)(1+x^3)\dots \\ &= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots \\ &= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots \\ &= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots)\dots \\ &= \text{number of partitions with } \textit{odd} \text{ sizes} . \quad \square \end{aligned}$$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $\square (1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} . \quad \square$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   

$$\begin{aligned} &= \frac{1-x^2}{1-x} \frac{1-x^4}{1-x^2} \frac{1-x^6}{1-x^3} \dots \\ &= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots \\ &= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots \\ &= \text{number of partitions with } \textit{odd} \text{ sizes} . \end{aligned}$$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$   
 We will define a bijection,

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $\square = \frac{1-x^2}{1-x} \frac{1-x^4}{1-x^2} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with odd sizes} . \quad \square$

**Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 =$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

□ **Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$   
 We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 +$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

□ **Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 +$$

$$\begin{matrix} * & * & * & * & * \end{matrix}$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

□ **Exercise.** Generalize the above theorems to paritions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 + 3 +$$

\* \* \* \* \*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-\cancel{x^4}}{1-x^2} \frac{1-\cancel{x^6}}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

□ **Exercise.** Generalize the above theorems to partitions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$\begin{array}{l} 12 = 5 + 3 + \\ * \quad * \quad * \quad * \quad * \\ * \quad * \quad * \end{array}$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

□ **Exercise.** Generalize the above theorems to paritions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 + 3 + 2 +$$

$$\begin{array}{ccccc} * & * & * & * & * \\ * & * & * & & \end{array}$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} .$  □

□ **Exercise.** Generalize the above theorems to paritions of sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$   
 We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$\begin{array}{l} 12 = 5 + 3 + 2 + \\ * \quad * \quad * \quad * \quad * \\ * \quad * \quad * \\ * \quad * \end{array}$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-x^4}{1-\cancel{x^2}} \frac{1-x^6}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $= \text{number of partitions with } \textit{odd} \text{ sizes} . \quad \square$

**Exercise.** Generalize the above theorems to paritions of  
 □ sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$   
 We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$\begin{array}{l} 12 = 5 + 3 + 2 + 2 \\ * \quad * \quad * \quad * \quad * \\ * \quad * \quad * \\ * \quad * \end{array}$$

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-\cancel{x^4}}{1-x^2} \frac{1-\cancel{x^6}}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

**Exercise.** Generalize the above theorems to paritions of □  
 sizes at most  $k$ .

.

**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$   
 We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 + 3 + 2 + 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*

**Theorem** (Euler). *The number of partitions of  $m$  into a sum of non-zero integers of distinct sizes, i.e. so that each term occurs at most once is equal to the number of partitions with odd sizes.*

*Proof.* Number of partitions is the  $m$ th coefficient of  
 $(1 + x)(1 + x^2)(1 + x^3) \dots$   
 $= \frac{1-\cancel{x^2}}{1-x} \frac{1-\cancel{x^4}}{1-x^2} \frac{1-\cancel{x^6}}{1-x^3} \dots$   
 $= \frac{1}{1-x} \frac{1}{1-x^3} \frac{1}{1-x^5} \dots$   
 $= (1+x+x^2+\dots)(1+x^3+x^6+\dots)(1+x^5+x^{10}+\dots) \dots$   
 $=$  number of partitions with *odd* sizes . □

**Exercise.** Generalize the above theorems to paritions of sizes at most  $k$ .

□



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,

$\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts

$\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 + 3 + 2 + 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

\* \* \* \*  
 \* \* \* \*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

12 = 5+ 3+ 2+ 2  
 \* \* \* \*  
 \* \* \*  
 \* \*  
 \* \*



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

\* \* \* \*  
 \* \* \* \*  
 \* \*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

12 = 5+ 3+ 2+ 2  
 \* \* \* \*  
 \* \* \*  
 \* \*  
 \* \*



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

\* \* \* \*  
 \* \* \* \*  
 \* \* .  
 \*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 + 3 + 2 + 2$$

\* \* \* \*  
 \* \* \*  
 \* \*  
 \* \*



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5+ 3+ 2+ 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

.



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,

$\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts

$\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5+ 3+ 2+ 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

$$12 =4 + .$$



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,

$\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts

$\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5+ 3+ 2+ 2$$

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + .



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,

$\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts

$\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5 + 3 + 2 + 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

$$12 = 4 + 4 + 2 + .$$





**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5+ 3+ 2+ 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

12 =4 + 4 + 2 + 1 + .



**Theorem** (Euler). *The number of partitions of  $n$  into  $k$  parts is equal to the number of partitions of  $n$  into parts of at most size  $k$ .*

*Proof.* Denote,  
 $\mathcal{P}_1$  : # of partitions of  $n$  into  $k$  parts  
 $\mathcal{P}_2$  : # of partitions of  $n$  into parts of at most size  $k$

We will define a bijection,

$$f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$$

$$12 = 5+ 3+ 2+ 2$$

\* \* \* \* \*

\* \* \*

\* \*

\* \*

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.



# Exponential generating function

\* \* \* \*  
\* \* \* \*  
\* \*  
\*  
\*

$12 = 4 + 4 + 2 + 1 + 1$

.

# Exponential generating function

Definition.

$$\begin{aligned} & * \quad * \quad * \quad * \\ & * \quad * \quad * \quad * \\ & * \quad * \\ & * \\ & * \\ & 12 = 4 + 4 + 2 + 1 + 1 \\ & . \end{aligned}$$

# Exponential generating function

**Definition** (Exponential generating function).

$$\begin{aligned} & * \quad * \quad * \quad * \\ & * \quad * \quad * \quad * \\ & * \quad * \\ & * \\ & * \end{aligned}$$

$$12 = 4 + 4 + 2 + 1 + 1$$

.

Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

\* \* \* \*  
\* \* \* \*  
\* \*  
\*  
\*

$$12 = 4 + 4 + 2 + 1 + 1$$

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example.

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.



Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )  
Generating function:

\* \* \* \*

\* \* \* \*

\* \*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots, )$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots$$

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )  
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )  
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example.

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots)$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence:  $(1, 3, 3^2, \dots)$

\* \* \* \*  
\* \* \* \*  
\* \*  
\*  
\*

$$12 = 4 + 4 + 2 + 1 + 1$$

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots, )$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence:  $(1, 3, 3^2, \dots, )$   
Generating function:

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots, )$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence:  $(1, 3, 3^2, \dots, )$   
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots$$

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots, )$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence:  $(1, 3, 3^2, \dots, )$   
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.



Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots, )$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence:  $(1, 3, 3^2, \dots, )$   
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

Example.

\* \* \* \*  
\* \* \* \*  
\* \*  
\*  
\*

$$12 = 4 + 4 + 2 + 1 + 1$$

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence:  $(1, 1, 1, \dots, )$   
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence:  $(1, 3, 3^2, \dots, )$   
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

Example. Sequence:  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

\* \* \* \*  
\* \* \* \*  
\* \*  
\*  
\*

$$12 = 4 + 4 + 2 + 1 + 1$$

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )  
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence: (1, 3, 3^2, ..., )  
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

Example. Sequence: (P\_0^n, P\_1^n, P\_2^n, ..., P\_n^n)  
Generating function:

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 =4 + 4 + 2 + 1 + 1

.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )  
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence: (1, 3, 3^2, ..., )  
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

Example. Sequence: (P\_0^n, P\_1^n, P\_2^n, ..., P\_n^n)  
Generating function:

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!}$$

\* \* \* \*  
\* \* \* \*  
\* \*  
\*  
\*  
  
12 = 4 + 4 + 2 + 1 + 1  
  
.

Exponential generating function

Definition (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

Example. Sequence: (1, 1, 1, ..., )  
Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

Example. Sequence: (1, 3, 3^2, ..., )  
Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

Example. Sequence: (P\_0^n, P\_1^n, P\_2^n, ..., P\_n^n)  
Generating function:

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

\*

12 = 4 + 4 + 2 + 1 + 1

.

# Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2x^2 + \cdots = a_0 + (a_11!)\frac{x}{1!} + (a_22!)\frac{x^2}{2!} + \cdots$$

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots, )$   
*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots, )$   
*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$   
*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

# Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots, )$   
*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots, )$   
*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$   
*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

.

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots,)$

*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots,)$

*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.* .



## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots,)$

*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots,)$

*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$\left(1 + \frac{x}{1!} + \frac{x^2}{2!}\right)$$

.

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** Sequence:  $(1, 1, 1, \dots,)$

Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** Sequence:  $(1, 3, 3^2, \dots,)$

Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** Sequence:  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

Generating function:

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

.

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** Sequence:  $(1, 1, 1, \dots,)$

Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** Sequence:  $(1, 3, 3^2, \dots,)$

Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** Sequence:  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

Generating function:

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

$k!$  times the  $k$ th coefficient

.

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots,)$

*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots,)$

*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

$k!$  times the  $k$ th coefficient

**Example.** In how many ways can  $k$  balls chosen out of infinitely many red and yellow ones be arranged?

.

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots,)$

*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots,)$

*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

$k!$  times the  $k$ th coefficient

**Example.** In how many ways can  $k$  balls chosen out of infinitely many red and yellow ones be arranged?

*Solution.* .

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** Sequence:  $(1, 1, 1, \dots)$

Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** Sequence:  $(1, 3, 3^2, \dots)$

Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** Sequence:  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

Generating function:

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

$k!$  times the  $k$ th coefficient

**Example.** In how many ways can  $k$  balls chosen out of infinitely many red and yellow ones be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!} + \cdots)$$

.

## Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** Sequence:  $(1, 1, 1, \dots)$

Generating function:

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** Sequence:  $(1, 3, 3^2, \dots)$

Generating function:

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** Sequence:  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$

Generating function:

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

$k!$  times the  $k$ th coefficient

**Example.** In how many ways can  $k$  balls chosen out of infinitely many red and yellow ones be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!} + \cdots)^2$$

.

# Exponential generating function

**Definition** (Exponential generating function).

$$a_0 + a_1x + a_2\frac{x^2}{2!} + a_3\frac{x^3}{3!} + \cdots$$

**Example.** *Sequence:*  $(1, 1, 1, \dots, )$   
*Generating function:*

$$1 + 1x + 1\frac{x^2}{2!} + 1\frac{x^3}{3!} + \cdots = e^x$$

**Example.** *Sequence:*  $(1, 3, 3^2, \dots, )$   
*Generating function:*

$$1 + 3x + 3^2\frac{x^2}{2!} + 3^3\frac{x^3}{3!} + \cdots = e^{3x}$$

**Example.** *Sequence:*  $(P_0^n, P_1^n, P_2^n, \dots, P_n^n)$   
*Generating function:*

$$P_0^n + P_1^n x + P_2^n \frac{x^2}{2!} + P_3^n \frac{x^3}{3!} + \cdots + P_n^n \frac{x^n}{n!} = (1 + x)^n$$

**Example.** In how many ways can  $k$  balls chosen out of 2 red and 3 yellow be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!})(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!})$$

$k!$  times the  $k$ th coefficient

**Example.** In how many ways can  $k$  balls chosen out of infinitely many red and yellow ones be arranged?

*Solution.*

$$(1 + \frac{x}{1!} + \frac{x^2}{2!} + \cdots)^2$$

$k!$  times the  $k$ th coefficient

.