

Example. Count the number of ways

Example. Count the number of ways to form n pairs

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution.

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n :

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{ \hspace{10cm} \}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \quad \}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{ (x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_1, x_2 \, , \, x_3, x_4 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\ x_1, x_2 \ , \ x_4, x_3 \ , \dots, \ x_{2n-1}, x_{2n} \] \mid x_i \in X\}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_1, x_2 \, , \, x_4, x_3 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_1, x_2 \, , \, x_4, x_3 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

$$|\mathcal{P}_n| \times 2!$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_4, x_3 \, , \, x_1, x_2 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

$$|\mathcal{P}_n| \times 2!$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_4, x_3 \, , \, x_1, x_2 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

$$|\mathcal{P}_n| \times 2! \times n!$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\ x_4, x_3 \ , \ x_1, x_2 \ , \dots, \ x_{2n-1}, x_{2n} \] \mid x_i \in X\}$$

$$(2n)! = |\mathcal{P}_n| \times 2! \times n!$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_4, x_3 \, , \, x_1, x_2 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

$$(2n)! = |\mathcal{P}_n| \times 2! \times n!$$

$$|\mathcal{P}_n|$$

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[x_4, x_3, x_1, x_2, \dots, x_{2n-1}, x_{2n}] \mid x_i \in X\}$$

$$(2n)! = |\mathcal{P}_n| \times 2! \times n!$$

$$|\mathcal{P}_n| = \frac{(2n)!}{n! \times 2!}$$

Example. Count the number of subsets

Example. Count the number of subsets of a set X of n elements.

Example. Count the number of subsets of a set X of n elements.

Solution.

Example. Count the number of subsets of a set X of n elements.

Solution. $X =$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\{A \subset X\}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$$\begin{aligned}\mathcal{P}(X) &:= \{A \subset X\} \\ &\rightarrow \mathcal{P}(X)\end{aligned}$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$\{0, 1\}^n \rightarrow \mathcal{P}(X)$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n)$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{ \hspace{10em} \}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$$\mathcal{P}(X) := \{A \subset X\}$$

$$f : \{0, 1\}^n \rightarrow \mathcal{P}(X), \text{ bijection}$$

$$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$$

$$|\mathcal{P}(X)|$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n|$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of subsets of a set X of n elements. **Example.** Count the number of n -tuples of $0, 1$ with exactly m 0s and $n - m$ 1s.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution.

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

Example. Count the number of subsets of a set X of n elements.

Example. Count the number of n -tuples of $0, 1$ with exactly m 0s and $n - m$ 1s.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Solution. X^n_m : set of such n -tuples
 $\rightarrow X^n_m$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$$\mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

where $\mathcal{C}_m(\{1, \dots, n\})$

Example. Count the number of subsets of a set X of n elements.

Example. Count the number of n -tuples of $0, 1$ with exactly m 0s and $n - m$ 1s.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Solution. X^n_m : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X^n_m$
where $\mathcal{C}_m(\{1, \dots, n\}) := \{ \hspace{10cm} \}$

Example. Count the number of subsets of a set X of n elements.

Example. Count the number of n -tuples of $0, 1$ with exactly m 0s and $n - m$ 1s.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Solution. X^n_m : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X^n_m$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A$

$\}$

Example. Count the number of subsets of a set X of n elements.

Example. Count the number of n -tuples of $0, 1$ with exactly m 0s and $n - m$ 1s.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Solution. X^n_m : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X^n_m$
where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

Example. Count the number of subsets of a set X of n elements.

Example. Count the number of n -tuples of $0, 1$ with exactly m 0s and $n - m$ 1s.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Solution. X^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X^n$
where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
and $f(A)$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$
 $\mathcal{P}(X) := \{A \subset X\}$
 $f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection
 $f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n|$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})|$$

Example. Count the number of subsets of a set X of n elements.

Solution. $X = \{x_1, x_2, \dots, x_n\}$

$\mathcal{P}(X) := \{A \subset X\}$

$f : \{0, 1\}^n \rightarrow \mathcal{P}(X)$, bijection

$f(i_1, i_2, \dots, i_n) = \{x_m \in X \mid i_m = 1\}$

$$|\mathcal{P}(X)| = |\{0, 1\}^n| = 2^n$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$


Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.



Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

X				

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

X	X			

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

X	X	X		

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

		X		
X	X	X		

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

		X	X	
X	X	X		

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	
		X	X	
X	X	X		

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y :

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $\rightarrow Y$

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $X_{m-1}^{n-1} \rightarrow Y$

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $X_{m-1}^{n-1} \rightarrow Y$
 where X

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, 1, 0, 1, 0) = \begin{array}{|c|c|c|c|c|} \hline & & & \text{X} & \text{X} \\ \hline & & \text{X} & \text{X} & \\ \hline \text{X} & \text{X} & \text{X} & & \\ \hline \end{array}$$

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(\textcolor{red}{0}, 0, 1, 0, 1, 0) =$$

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, \textcolor{red}{0}, 1, 0, 1, 0) =$$

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, \textcolor{red}{1}, 0, 1, 0) = \begin{array}{|c|c|c|c|c|} \hline & & & \text{X} & \text{X} \\ \hline & & \text{X} & \text{X} & \\ \hline \text{X} & \text{X} & \text{X} & & \\ \hline \end{array}$$

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, 1, \textcolor{red}{0}, 1, 0) =$$

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, 1, 0, \textcolor{red}{1}, 0) =$$

			X	X
		X	X	
X	X	X		

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, 1, 0, 1, \textcolor{red}{0}) = \begin{array}{|c|c|c|c|c|} \hline & & & \text{X} & \text{X} \\ \hline & & \text{X} & \text{X} & \\ \hline \text{X} & \text{X} & \text{X} & & \\ \hline \end{array}$$

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0,0,1,0,1,0) = \begin{array}{|c|c|c|c|c|} \hline & & & X & X \\ \hline & & X & X & \\ \hline X & X & X & & \\ \hline \end{array}$$

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

$$|Y|$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, 1, 0, 1, 0) = \begin{array}{|c|c|c|c|c|} \hline & & & X & X \\ \hline & & X & X & \\ \hline X & X & X & & \\ \hline \end{array}$$

Solution. Y : set of such routes

$$f : X_{m-1}^{n-1} \rightarrow Y$$

where X : is defined as before

and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where, i th step is right if $x_i = 0$ and up if $x_i = 1$.

$$|Y| = |X_{m-1}^{n-1}|$$

Example. Count the number of n -tuples of 0, 1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples

$$f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$$

where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$

and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,

such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

Example. Count the number of ways to go from the bottom left corner of an $m \times n$ -grid to the top right corner in the shortest way.

$$f(0, 0, 1, 0, 1, 0) = \begin{array}{|c|c|c|c|c|} \hline & & & \text{X} & \text{X} \\ \hline & & \text{X} & \text{X} & \\ \hline \text{X} & \text{X} & \text{X} & & \\ \hline \end{array}$$

Solution. Y : set of such routes
 $f : X_{m-1}^{n-1} \rightarrow Y$
 where X : is defined as before
 and $f(x_1, \dots, x_i, \dots, x_{m+n-2}) =$ a route, where,
 i th step is right if $x_i = 0$ and up if $x_i = 1$.

Example. Count the number of n -tuples of 0,1 with exactly m 0s and $n - m$ 1s.

Solution. X_m^n : set of such n -tuples
 $f : \mathcal{C}_m(\{1, \dots, n\}) \rightarrow X_m^n$
 where $\mathcal{C}_m(\{1, \dots, n\}) := \{A \subset \{1, \dots, n\} \mid |A| = m\}$
 and $f(A) = (x_1, \dots, x_i, \dots, x_n)$,
 such that $x_i = 0$ if and only if $i \in A$

$$|X_m^n| = |\mathcal{C}_m(\{1, \dots, n\})| = \binom{n}{m}$$

$$|Y| = |X_{m-1}^{n-1}| = \binom{m+n-2}{m-1}$$