



Example. Solve the recurrence relation $a_n = 2a_{n-1} + 3n^2$,

Example. Solve the recurrence relation $a_n = 2a_{n-1} + 3n^2$, $a_0 = 1$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

$$a_n = 2a_{n-1} + 3n^2$$

$$a_n = 2a_{n-1} + 3n^2, a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only $a_n = 2a_{n-1} + 3n^2$ (not $a_0 = 1$) i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If
$$f(n)$$
, $g(n)$ are solutions of only $a_n = 2a_{n-1} + 3n^2$ (not $a_0 = 1$) i.e.

$$f(n) - 2f(n-1) = 3n^{2}$$

$$g(n) - 2g(n-1) = 3n^{2}$$

$$(f-q)(n) - 2(f-q)(n-1) = 0$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

$$a_n = 2a_{n-1} + 3n^2, a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n) - g(n) by the usual method,

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

$$a_n = 2a_{n-1} + 3n^2, a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

E.g.
$$g(n) = A + Bn + Cn^2$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

 $a_n - 2a_{n-1} = 3n^2$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

Example.

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

Example. Solve the recurrence relation $a_n = 2a_{n-1} + 3^n$,

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

Example. Solve the recurrence relation $a_n = 2a_{n-1} + 3^n$, $a_0 = 1$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

Example. Solve the recurrence relation $a_n = 2a_{n-1} + 3^n$, $a_0 = 1$

$$a_n - 2a_{n-1} = 3^n$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g. $g(n) = A + Bn + Cn^2$ find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

Example. Solve the recurrence relation $a_n = 2a_{n-1} + 3^n$, $a_0 = 1$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$ E.g. $g(n) = A + Bn + Cn^2$ find A, B, Cf(n) = (f - g)(n) + g(n)

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only $a_n = 2a_{n-1} + 3n^2$ (not $a_0 = 1$) i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$ E.g. $g(n) = A + Bn + Cn^2$ find A, B, Cf(n) = (f - g)(n) + g(n)

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only $a_n = 2a_{n-1} + 3n^2$ (not $a_0 = 1$) i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$ E.g. $g(n) = A + Bn + Cn^2$ find A, B, Cf(n) = (f - g)(n) + g(n)

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

 $a_n - 2a_{n-1} = 3n^2$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g. $g(n) = A + Bn + Cn^2$ find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C
 $f(n) = (f - q)(n) + g(n)$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If
$$f(n)$$
, $g(n)$ are solutions of only $a_n = 2a_{n-1} + 3n^2$ (not $a_0 = 1$) i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$ E.g. $q(n) = A + Bn + Cn^2$ find A, B, Cf(n) = (f - q)(n) + q(n)

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$
$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$a_n = 2a_{n-1} + 3n^2, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3n^2$$

Observe:

If f(n), g(n) are solutions of only

$$a_n = 2a_{n-1} + 3n^2 \ (not \ a_0 = 1)$$
 i.e.

$$f(n) - 2f(n-1) = 3n^2$$

$$g(n) - 2g(n-1) = 3n^2$$

$$(f-g)(n) - 2(f-g)(n-1) = 0$$

then, f(n) - g(n) is a solution to $a_n - 2a_{n-1} = 0$

Solve, f(n)-g(n) by the usual method, satisfying $a_0=1$

$$f(n) - g(n) = 2^n$$

Let g(n) be a solution of only $a_n = 2a_{n-1} + 3n^2$

E.g.
$$g(n) = A + Bn + Cn^2$$
 find A, B, C

$$f(n) = (f - g)(n) + g(n)$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B=3,$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer:
$$3^{n+1} - 2^{n+1}$$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Subsitute

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n + 2a_{n-1} - 6a_{n-2} = 0$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n + 2a_{n-1} - 6a_{n-2} = 0$$

$$a_n = 2a_{n-1} + 3^n, \ a_0 = 1$$

$$a_n - 2a_{n-1} = 3^n$$

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

$$2xf(x) = 2a_0x + 2a_1x^2 + 2a_2x^2 + 2a_2x^4 + \cdots$$

$$f(x) - 2xf(x) = a_0 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$

$$f(x)(1-2x) = 1 + 3x + 3^2x^2 + 3^3x^3 + 3^4x^4 + \cdots$$
$$= \frac{1}{1-3x}$$

$$f(x) = \frac{1}{(1 - 2x)(1 - 3x)}$$

$$= \frac{A}{1 - 2x} + \frac{B}{1 - 3x}$$

$$= A \sum_{n=0}^{\infty} (2x)^n + B \sum_{n=0}^{\infty} (3x)^n$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n + 2a_{n-1} - 6a_{n-2} = 0$$

. . .

Exercise. Find a solution for the system of recurrence relations,

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n + 2a_{n-1} - 6a_{n-2} = 0$$

. . .

Exercise. Find a solution for the system of recurrence relations,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n + 2a_{n-1} - 6a_{n-2} = 0$$

. . .

Exercise. Find a solution for the system of recurrence relations,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + a_{n-1} + b_{n-1} = 0$$

Solution.

$$1 = a_0 = A + B$$

$$5 = a_1 = 2A + 3B$$

$$B = 3, A = -2$$

Answer: $3^{n+1} - 2^{n+1}$

Example.

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + 2a_{n-1} = 0$$

Substitute $b_{n-1} = -2a_{n-2}$ in,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

to get,

$$a_n + 2a_{n-1} - 6a_{n-2} = 0$$

. . .

Exercise. Find a solution for the system of recurrence relations,

$$a_n + 2a_{n-1} + 3b_{n-1} = 0$$

$$b_n + a_{n-1} + b_{n-1} = 0$$

Solution. Use generating functions for f(x) and g(x) for a_n and b_n respectively . . .