

$s(n, m)$  denote the number of ways to arrange

$s(n, m)$  denote the number of ways to arrange  $\{1, 2, \dots, n\}$  around  $m$  identical circles so that each circle has at least one number. Note the difference with  $S(n, m)$  that was done during a lecture ( $s(n, m)$  are called Stirling numbers of the first kind, and  $S(n, m)$  are called Stirling numbers of the second kind).

$s(n, m)$  denote the number of ways to arrange  $\{1, 2, \dots, n\}$  around  $m$  identical circles so that each circle has at least one number. Note the difference with  $S(n, m)$  that was done during a lecture ( $s(n, m)$  are called Stirling numbers of the first kind, and  $S(n, m)$  are called Stirling numbers of the second kind).

1. Prove that  $s(n, m) = s(n-1, m-1) + (n-1)s(n-1, m)$ .

*Case 1:* 1 is the only number in its circle

$s(n, m)$  denote the number of ways to arrange  $\{1, 2, \dots, n\}$  around  $m$  identical circles so that each circle has at least one number. Note the difference with  $S(n, m)$  that was done during a lecture ( $s(n, m)$  are called Stirling numbers of the first kind, and  $S(n, m)$  are called Stirling numbers of the second kind).

1. Prove that  $s(n, m) = s(n-1, m-1) + (n-1)s(n-1, m)$ .

*Case 1:* 1 is the only number in its circle  
answer:  $s(n-1, m-1)$

$s(n, m)$  denote the number of ways to arrange  $\{1, 2, \dots, n\}$  around  $m$  identical circles so that each circle has at least one number. Note the difference with  $S(n, m)$  that was done during a lecture ( $s(n, m)$  are called Stirling numbers of the first kind, and  $S(n, m)$  are called Stirling numbers of the second kind).

1. Prove that  $s(n, m) = s(n-1, m-1) + (n-1)s(n-1, m)$ .

*Case 1:* 1 is the only number in its circle

answer:  $s(n-1, m-1)$

*Case 2:* 1 is not the only number in its circle

$s(n, m)$  denote the number of ways to arrange  $\{1, 2, \dots, n\}$  around  $m$  identical circles so that each circle has at least one number. Note the difference with  $S(n, m)$  that was done during a lecture ( $s(n, m)$  are called Stirling numbers of the first kind, and  $S(n, m)$  are called Stirling numbers of the second kind).

1. Prove that  $s(n, m) = s(n-1, m-1) + (n-1)s(n-1, m)$ .

*Case 1:* 1 is the only number in its circle

answer:  $s(n-1, m-1)$

*Case 2:* 1 is not the only number in its circle

First place  $\{2, \dots, n\}$  around the circles.  $s(n-1, m)$ .

$s(n, m)$  denote the number of ways to arrange  $\{1, 2, \dots, n\}$  around  $m$  identical circles so that each circle has at least one number. Note the difference with  $S(n, m)$  that was done during a lecture ( $s(n, m)$  are called Stirling numbers of the first kind, and  $S(n, m)$  are called Stirling numbers of the second kind).

Example: 245 376 89

$$\begin{aligned} s(n, n) &= 1 \\ s(n, m) &= 0 \text{ if } n < m \\ s(n, 1) &= (n-1)! \\ s(n, n-1) &= \binom{n}{2} \end{aligned}$$

1. Prove that  $s(n, m) = s(n-1, m-1) + (n-1)s(n-1, m)$ .

*Case 1:* 1 is the only number in its circle  
 answer:  $s(n-1, m-1)$

*Case 2:* 1 is not the only number in its circle  
 First place  $\{2, \dots, n\}$  around the circles.  $s(n-1, m)$ .

Place 1 next to any of the  $n-1$  numbers that are already placed.  
 To the right the number.

$$s(n,m) = s(n-1,m-1) + (n-1)s(n-1,m).$$

$$\begin{aligned} s(n,n-1) &= s(n-1,n-2) + (n-1)s(n-1,n-1) \\ &= s(n-1,n-2) + (n-1) \\ &= s(n-2,n-3) + (n-2) + (n-1) \\ &= \dots \\ &= 1 + 2 + \dots + n-1 \\ &= \frac{n(n-1)}{2} = \binom{n}{2} \end{aligned}$$

$$\{A \subset B \subset X \mid |B| = k\} \leftrightarrow \{C \subset X \setminus A \mid |C| = k - |A|\}$$