

Example. Count the number of ways

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to permute $AAABBB$.

Solution.

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:
number of ways

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$

Example. Count the number of ways to permute $AAABB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$

$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Example. Count the number of ways to permute $AAABB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$

$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class:

Example. Count the number of ways to permute $AAABB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$

$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3!$

Example. Count the number of ways to permute $AAABB$.

Solution. More familiar modification:

number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$

$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Example. Count the number of ways to permute $AAABBB$. **Example.** Count the number of ways

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Solution.

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent, if the permutations are only among the A_i s and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Solution. If $m = 5$,

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Solution. If $m = 5$, examples of selections:

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Solution. If $m = 5$, examples of selections:

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.
 Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
 number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$
Solution. If $m = 5$, examples of selections:
 $RRRGG$,

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent, if the permutations are only among the A_i s and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$

$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
 number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRRGB, RRGGB,$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
 if the permutations are only among the A_i s
 and / or only among the B_i s.

$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct}, x_i \in \{A_1, A_2, A_3\}\}$

$B := \{(y_1, y_2) \mid y_i \text{ are distinct}, y_i \in \{B_1, B_2\}\}$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
 number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$
 Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
 if the permutations are only among the A_i s
 and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$

$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
 number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$
 Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
 if the permutations are only among the A_i s
 and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$

$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Solution. If $m = 5$, examples of selections:
 $\textcolor{red}{RRRGB}, RRGGB, RGGGG$
 Alternatively,
 $XXX|X|X,$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Solution. If $m = 5$, examples of selections:
 $RRRGG, \textcolor{red}{RRGGG}, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X,$

Example. Count the number of ways to permute $AAABBB$.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. More familiar modification:
 number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent, if the permutations are only among the A_i s and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_1 \in \{A_1, A_2, A_3\}\}$$

$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_1 \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer:

Example. Count the number of ways to permute $AAABBB$.

Solution. More familiar modification:
number of ways to permute $A_1A_2A_3B_1B_2$. *Answer:* $5!$

Two rearrangements of $A_1A_2A_3B_1B_2$ are equivalent,
if the permutations are only among the A_i s
and / or only among the B_i s.

$$A := \{(x_1, x_2, x_3) \mid x_i \text{ are distinct, } x_i \in \{A_1, A_2, A_3\}\}$$
$$B := \{(y_1, y_2) \mid y_i \text{ are distinct, } y_i \in \{B_1, B_2\}\}$$

$$f : A \times B \rightarrow X$$

$$f((x_1, x_2, x_3), (y_1, y_2)) = x_1x_2x_3y_1y_2$$

Size of each equivalence class: $3! \times 2!$

Final answer: $\frac{5!}{3! \times 2!}$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:

$RRRGB, RRGGB, RGGGG$

Alternatively,

$XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGB, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution.

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$
Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set,

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$
Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2, \dots, x_m\}$$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m-1)\}$$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 1, \dots, x_m - 1\}$$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 1, \dots, x_m - 1\}$$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 1, \dots, x_m - 1\}$$

where, $x_{i+1} - x_i \geq 2$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 1, \dots, x_m - 1\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 2$,

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 1, \dots, x_m - 1\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 2$, otherwise, $x_{i+1} - x_i \geq 1$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 1, \dots, x_m - 1\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 2$, otherwise, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - 1$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - 1\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 2$, otherwise, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - 1$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - 2\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 2$, otherwise, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - 1$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - 2\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 3$, otherwise, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - 1$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - 2\}$$

where, $x_{i+1} - x_i \geq 2$ for $i \geq 3$, otherwise, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - 2$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - 2$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - (m - 1)$

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - (m - 1)$

Rest, exercise!

Example. Count the number of ways to select m balls from an infinite supply of red, green, and blue ones.

Solution. If $m = 5$, examples of selections:
 $RRRGG, RRGGB, RGGGG$

Alternatively,
 $XXX|X|X, XX|XX|X, X|XXXX|$

Final answer: $\binom{m+2}{m}$

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:

Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$

and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$

and $x_m \leq n - (m - 1)$

Rest, exercise!

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - (m - 1)$

Rest, exercise!

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution.

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - (m - 1)$

Rest, exercise!

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea:

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:
Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$
and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$
and $x_m \leq n - (m - 1)$

Rest, exercise!

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1x_2 \dots x_n$ be an element that is not in the set

Example. Count the number of ways to select m integers out of $\{1, \dots, n\}$ so that they contain no consecutive integers.

Solution. Consider such a set, $A = \{x_1, \dots, x_m\}$. Main idea:

Order the elements

$$A := \{x_1, x_2, x_3, \dots, x_m\}$$

where, $x_{i+1} - x_i \geq 2$

and $x_m \leq n$

$$B := \{x_1, x_2 - 1, x_3 - 2, \dots, x_m - (m - 1)\}$$

where, $x_{i+1} - x_i \geq 1$

and $x_m \leq n - (m - 1)$

Rest, exercise!

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object:

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object:

$$\frac{n!}{(n-m)!}$$

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object:
$$\frac{n!}{(n-m)!}$$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects:

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object: $\frac{n!}{(n-m)!}$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects: n^m

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object:
$$\frac{n!}{(n-m)!}$$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects: n^m
3. Number of ways to *arrange* m distinct objects in n distinct boxes:

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object: $\frac{n!}{(n-m)!}$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects: n^m
3. Number of ways to *arrange* m distinct objects in n distinct boxes: $\frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute m identical objects in n distinct boxes if each box can contain only 1 object:

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object: $\frac{n!}{(n-m)!}$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects: n^m
3. Number of ways to *arrange* m distinct objects in n distinct boxes: $\frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute m identical objects in n distinct boxes if each box can contain only 1 object: $\binom{n}{m}$

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object: $\frac{n!}{(n-m)!}$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects: n^m
3. Number of ways to *arrange* m distinct objects in n distinct boxes: $\frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute m identical objects in n distinct boxes if each box can contain only 1 object: $\binom{n}{m}$
5. Number of ways to distribute m identical objects in n distinct boxes if each box can contain any number of objects:

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.

Example.

1. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only 1 object: $\frac{n!}{(n-m)!}$
2. Number of ways to distribute m distinct objects in n distinct boxes if each box can contain only any number of objects: n^m
3. Number of ways to *arrange* m distinct objects in n distinct boxes: $\frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute m identical objects in n distinct boxes if each box can contain only 1 object: $\binom{n}{m}$
5. Number of ways to distribute m identical objects in n distinct boxes if each box can contain any number of objects: n^m

Example. Show that the number of permutations of $1, 2, 3, \dots, 2n$ with consecutive terms that differ by n are more than permutations which do not.

Solution. Main idea: Let $x_1 x_2 \dots x_n$ be an element that is not in the set

We can create an element which is in the set:

If $|x_j - x_1| = n$, consider the element $x_2 \dots x_1 x_j \dots x_n$.