

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\mathcal{C}_r^n = \{ \hspace{10em} \}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\mathcal{C}_r^n = \{ \{x_1, x_2, \dots, x_r\} \}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\mathcal{C}_r^n = \{ \{x_1, x_2, \dots, x_r\} \mid x_i \in X \}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{ \end{aligned}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)]\}\end{aligned}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X\}\end{aligned}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\}\end{aligned}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned} \mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{ \end{aligned}$$

5

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

[illegible]

1

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if □

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class:



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!}$ □

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes: $|\mathcal{C}_r^n|$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

number of arrangements =

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes: $|\mathcal{C}_r^n|$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

$$\begin{aligned} \mathcal{C}_r^n &= \{ \{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct} \} \\ &= \{ [(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct} \} \\ &= \{ [a] \mid a \in r - \text{arrangements} \} \end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes: $|\mathcal{C}_r^n|$



Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes: $|\mathcal{C}_r^n|$

□

$$\frac{n!}{(n-r)!} = \text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes: $|\mathcal{C}_r^n|$

$$\frac{n!}{(n-r)!} = \text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

$$|\mathcal{C}_r^n| = \frac{n!}{(n-r)!r!}$$

□

Example. Number of ways to choose r distinct objects from a set of X of n objects.

Solution.

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$ is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if (x_1, \dots, x_r) is a rearrangement of (y_1, \dots, y_r) .

Size of each equivalence class: $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes: $|\mathcal{C}_r^n|$

□

$$\frac{n!}{(n-r)!} = \text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

$$\binom{n}{r} := |\mathcal{C}_r^n| = \frac{n!}{(n-r)!r!}$$

Example. Find the number of positive divisors of 50.

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10$$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2$$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j\}$ □

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, \}$ □

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
[[n]] □

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$f :$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]]$$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection



Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| \qquad \qquad \qquad \square$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| \quad \square$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 \qquad \square$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution. □

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\} \qquad \qquad \qquad \square$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \qquad \qquad \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{ \qquad \qquad \qquad \}$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\} \tag*{\square}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \tag*{\square}$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \hspace{10em}\}$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \qquad \}$$

\square

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

$$S_i = \{ \qquad \qquad \qquad \}$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

$$S_i = \{(a, i) \qquad \qquad \qquad \}$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, \qquad \}$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

\square

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=?}^{10} S_i \qquad \square$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i \qquad \square$$

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6$$

□

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

$$|S|$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[[2]]| \times |[[1]]| = 3 \times 2 = 6$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

$$|S| = 1 + 2 + \dots + 9$$

□

Example. Find the number of positive divisors of 50.

Solution.

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors: $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

f is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6$$

Example. Count the number of pairs (a, b) so that $a, b \in \{1, 2, \dots, 10\}$ and a is smaller.

Solution.

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

$$|S| = 1 + 2 + \dots + 9 = \dots$$

□

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. □

Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair



Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:



Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{ \hspace{10cm} \}$$



Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \}$$



Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$



Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{ (x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$



Example. Count the number of ways to form n pairs out of a set X of $2n$ elements.

Solution. \mathcal{P}_n : Number of distinct ways to pair
Each element of \mathcal{P}_n looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[\, x_1, x_2 \, , \, x_3, x_4 \, , \dots , \, x_{2n-1}, x_{2n} \,] \mid x_i \in X\}$$

