

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.*



**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: □



**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1 x_2 \dots x_{2n}$  be an element that is not in the set  $\square$

**Example.** Show that the number of permutations of  $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1 x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1 x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

**Example.** Show that the number of permutations of **Example.**  
 $1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  ,  
are more than permutations which do not.

For instance, consider permutations of 123456  
 $A := \{125346, 253416, \dots, 231456, 234156, \dots\}$   
 $\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1 x_2 \dots x_{2n}$  be an element  
that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1 x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:

For instance, consider permutations of 123456  
 $A := \{125346, 253416, \dots, 231456, 234156, \dots\}$   
 $\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$

For instance, consider permutations of 123456

$$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$$

$$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$

For instance, consider permutations of 123456

$$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$$

$$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$

For instance, consider permutations of 123456

$$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$$

$$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  
are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1 x_2 \dots x_{2n}$  be an element  
that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1 x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

1. Number of ways to distribute  $m$  distinct objects in  
 $n$  distinct boxes if each box can contain at most 1  
object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  
 $n$  distinct boxes if each box can contain any number  
of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$   
distinct boxes:



**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$  are more than permutations which do not.

For instance, consider permutations of 123456

$$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$$

$$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

- 2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
- 3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$  are more than permutations which do not.

For instance, consider permutations of 123456

$$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$$

$$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$  , 1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$

For instance, consider permutations of 123456  
 $A := \{125346, 253416, \dots, 231456, 234156, \dots\}$   
 $\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$ , are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1 x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1 x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:

**Example.** Show that the number of permutations of **Example.**

$1, 2, 3, \dots, 2n$  with consecutive terms that differ by  $n$ , are more than permutations which do not.

For instance, consider permutations of 123456

$A := \{125346, 253416, \dots, 231456, 234156, \dots\}$

$\{1, \dots, n\} \setminus A := \{123456, 423156, \dots\}$

*Solution.* Main idea: Let  $x_1 x_2 \dots x_{2n}$  be an element that is not in the set

We can create an element which is in the set:

If  $|x_j - x_1| = n$ , consider the element  $x_2 \dots x_1 x_j \dots x_{2n}$ .

*Exercise:* Show that this map is injective □

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*


After placing the blue balls,

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*



After placing the blue balls, there are  $m + 1$  “boxes”

- Example.**
1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
  2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
  3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
  4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
  5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*



After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*



After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*




After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”


$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*     

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*     

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges  
 Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.*     

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges  
 Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges  
 Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges  
 Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 


After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues  
 2 red balls must be placed at the edges  
 Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

- Example.**
1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
  2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
  3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
  4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
  5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n$$

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box.

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”  
 $m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box.

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*

**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.**

1. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\frac{n!}{(n-m)!}$
2. Number of ways to distribute  $m$  distinct objects in  $n$  distinct boxes if each box can contain any number of objects:  $n^m$
3. Number of ways to *arrange*  $m$  distinct objects in  $n$  distinct boxes:  $\frac{(n-1+m)!}{(n-1)!m!} \times m! = \frac{(n-1+m)!}{(n-1)!}$
4. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain at most 1 object:  $\binom{n}{m}$
5. Number of ways to distribute  $m$  identical objects in  $n$  distinct boxes if each box can contain any number of objects:  $\binom{m+n-1}{m}$



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

**Example.** Number of ways to partition  $\{1, \dots, n\}$

*Solution.*


After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$ 
□


**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \dots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$ 
□

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □


**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \dots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □


**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \dots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
 Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes


*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \dots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).


*Solution. Idea:* Split the partitions into two disjoint types.

1.

2.



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).


*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

2.



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset
- 1 is always accompanied



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).


*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset  
# of such partitions:
- 1 is always accompanied





**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
- 1 is always accompanied



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
- 1 is always accompanied  
# of such partitions:  $S(n - 1, m)$ .



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions: .

First place  $\{2, \dots, n\}$  into the  $m$  subsets!

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
- 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times 2$ .



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times$  .

then place 1 into one of the  $m$  subsets!

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into  $m$  subsets.  
 Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset  
 # of such partitions:  $S(n - 1, m - 1)$ .
- 1 is always accompanied  
 # of such partitions:  $S(n - 1, m) \times m$ .



**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red balls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
- 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*

**Example.** Count the number of ways to arrange  $m$  blue balls, and  $n$  red balls so that none of the blue balls are adjacent and the ends are red balls?

*Solution.* 

After placing the blue balls, there are  $m + 1$  “boxes”

$m - 1$  red balls must be placed to separate the blues

2 red balls must be placed at the edges

Remaining  $n - m - 1$  red calls must be placed in  $m + 1$  boxes

*Answer:*  $\binom{n-1}{n-m-1}$  □

**Example.** Count the number of non-negative integer solutions to

$$x_1 + x_2 + \cdots + x_n = m$$

*Solution.*  $n$  boxes.  $m$  1s.  $x_i$ s are the number of 1s in each box. *Answer:*  $\binom{n+m-1}{m}$  □

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

- 1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
- 2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$

□



$$S(n, m) =$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = ?$$


**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$S(n, m) = ?$  if  $n < m$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$S(n, m) = 0$  if  $n < m$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$
$$S(n, 1) =$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$
$$S(n, 1) = 1$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) =$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$





$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2)$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) +$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2) = S(2, 1) + S(2, 2)$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2) = S(2, 1) + S(2, 2) \times 2$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$

2.  $S(4, 2)$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) +$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$





$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2)$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

1.  $S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$

2.  $S(4, 2) = S(3, 1) + S(3, 2) \times 2$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3)$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) +$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3)$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3) \times 3$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$





$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$S(n, n - 1)$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n - 1, m - 1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$S(n, n-1) = S(n-1, n-2) +$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$S(n, n-1) = S(n-1, n-2) + S(n-1, n-1)$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$S(n, n-1) = S(n-1, n-2) + S(n-1, n-1) \times n$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$

$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$\begin{aligned} S(n, n-1) &= S(n-1, n-2) + S(n-1, n-1) \times n \\ &= S(n-1, n-2) + (n-1) \end{aligned}$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$\begin{aligned} S(n, n-1) &= S(n-1, n-2) + S(n-1, n-1) \times n - 1 \\ &= S(n-1, n-2) + (n-1) \\ &= S(n-2, n-3) + (n-2) + (n-1) \end{aligned}$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$





$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. \ S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. \ S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. \ S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$\begin{aligned} S(n, n-1) &= S(n-1, n-2) + S(n-1, n-1) \times n \\ &= S(n-1, n-2) + (n-1) \\ &= S(n-2, n-3) + (n-2) + (n-1) \\ &\dots \\ &= S(2, 1) + \dots + (n-2) + (n-1) \end{aligned}$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied  
# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$



$$\begin{aligned}
 S(n, m) &= 0 \text{ if } n < m \\
 S(n, 1) &= 1 \\
 S(n, n) &= 1
 \end{aligned}$$

**Examples.**

1.  $S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$
2.  $S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$
3.  $S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$
- 4.

$$\begin{aligned}
 S(n, n - 1) &= S(n - 1, n - 2) + S(n - 1, n - 1) \times n - 1 \\
 &= S(n - 1, n - 2) + (n - 1) \\
 &= S(n - 2, n - 3) + (n - 2) + (n - 1) \\
 &\dots \\
 &= \underbrace{S(2, 1)}_1 + \dots + (n - 2) + (n - 1)
 \end{aligned}$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.  
 Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset  
 # of such partitions:  $S(n - 1, m - 1)$ .
2. 1 is always accompanied  
 # of such partitions:  $S(n - 1, m) \times m$ .

*Answer:*  $S(n, m) = S(n - 1, m - 1) + S(n - 1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$S(n, n-1) = S(n-1, n-2) + S(n-1, n-1) \times n-1$$

$$= S(n-1, n-2) + (n-1)$$

$$= S(n-2, n-3) + (n-2) + (n-1)$$

...

$$= \underbrace{S(2, 1)}_1 + \dots + (n-2) + (n-1)$$

$$= \frac{(n-1)n}{2}$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$



$$S(n, m) = 0 \text{ if } n < m$$

$$S(n, 1) = 1$$

$$S(n, n) = 1$$

### Examples.

$$1. S(3, 2) = S(2, 1) + S(2, 2) \times 2 = 1 + 1 \times 2 = 3$$

$$2. S(4, 2) = S(3, 1) + S(3, 2) \times 2 = 1 + 3 \times 2 = 7$$

$$3. S(4, 3) = S(3, 2) + S(3, 3) \times 3 = 3 + 1 \times 3 = 6$$

4.

$$S(n, n-1) = S(n-1, n-2) + S(n-1, n-1) \times n-1$$

$$= S(n-1, n-2) + (n-1)$$

$$= S(n-2, n-3) + (n-2) + (n-1)$$

...

$$= \underbrace{S(2, 1)}_1 + \dots + (n-2) + (n-1)$$

$$= \frac{(n-1)n}{2} = \binom{n}{2}$$

**Example.** Number of ways to partition  $\{1, \dots, n\}$  into a  $m$  subsets.

Denoted  $S(n, m)$  (Sterling number of the second kind).

*Solution. Idea:* Split the partitions into two disjoint types.

1. 1 forms a singletons subset

# of such partitions:  $S(n-1, m-1)$ .

2. 1 is always accompanied

# of such partitions:  $S(n-1, m) \times m$ .

*Answer:*  $S(n, m) = S(n-1, m-1) + S(n-1, m) \times m$

