

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$f(x) = \frac{x}{1 - x - x^2}$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \end{aligned}$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$.

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$f(x) = \frac{x}{1 - x - x^2}$$

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \end{aligned}$$

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1 - x - x^2$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1 - x - x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

Rest: easy

Example. Using generating functions to find a closed form formula of the Fibonacci sequence $(0, 1, 1, 2, 3, 5, 8, \dots)$. Alternatively, leave A and B unknown for now...

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

$$f(x) = 0 + 1x + 1x^2 + 2x^3 + 3x^4 + \dots$$

$$xf(x) = 0x + 1x^2 + 1x^3 + 2x^4 + 3x^5 + \dots$$

$$x^2f(x) = 0x^2 + 1x^3 + 1x^4 + 2x^5 + 3x^6 + \dots$$

$$f(x) - xf(x) - x^2f(x) = x$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= x \left(\frac{A}{\alpha - x} + \frac{B}{\beta - x} \right) \\ &= \frac{x}{\beta - \alpha} \left(\frac{1}{\alpha - x} - \frac{1}{\beta - x} \right) \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{x}{1 - x - x^2} \\ &= \frac{A}{\alpha - x} + \frac{B}{\beta - x} \\ &= \frac{A/\alpha}{1 - x/\alpha} + \frac{B/\beta}{1 - x/\beta} \\ &= \frac{A'}{1 - x/\alpha} + \frac{B'}{1 - x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

$$\alpha, \beta \text{ are roots of } 1 - x - x^2$$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$$\alpha' := 1/\alpha, \beta' := 1/\beta \text{ are roots of } x^2 - x - 1$$

Rest: easy

$$\alpha' = \frac{1+\sqrt{5}}{2}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned} f(x) &= \frac{x}{1-x-x^2} \\ &= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\ &= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\ &= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\alpha' = \frac{1+\sqrt{5}}{2}$$

$$\beta' = \frac{1-\sqrt{5}}{2}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned} f(x) &= \frac{x}{1-x-x^2} \\ &= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\ &= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\ &= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i} \end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\begin{aligned}\alpha' &= \frac{1+\sqrt{5}}{2} \\ \beta' &= \frac{1-\sqrt{5}}{2} \\ a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n\end{aligned}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned}f(x) &= \frac{x}{1-x-x^2} \\ &= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\ &= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\ &= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i}\end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\begin{aligned}\alpha' &= \frac{1+\sqrt{5}}{2} \\ \beta' &= \frac{1-\sqrt{5}}{2} \\ a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\ a_0 &= A' + B' = 0\end{aligned}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned}f(x) &= \frac{x}{1-x-x^2} \\ &= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\ &= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\ &= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i}\end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\begin{aligned}\alpha' &= \frac{1+\sqrt{5}}{2} \\ \beta' &= \frac{1-\sqrt{5}}{2} \\ a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\ a_0 &= A' + B' = 0, \text{ so } A' = -B'\end{aligned}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned}f(x) &= \frac{x}{1-x-x^2} \\ &= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\ &= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\ &= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i}\end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\begin{aligned}\alpha' &= \frac{1+\sqrt{5}}{2} \\ \beta' &= \frac{1-\sqrt{5}}{2} \\ a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\ a_0 &= A' + B' = 0, \text{ so } A' = -B' \\ a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1\end{aligned}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned}f(x) &= \frac{x}{1-x-x^2} \\ &= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\ &= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\ &= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\ &= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i}\end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}}
\end{aligned}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned}
f(x) &= \frac{x}{1-x-x^2} \\
&= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\
&= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\
&= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\
&= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i}
\end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

Alternatively, leave A and B unknown for now...

$$\begin{aligned}
f(x) &= \frac{x}{1-x-x^2} \\
&= \frac{A}{\alpha-x} + \frac{B}{\beta-x} \\
&= \frac{A/\alpha}{1-x/\alpha} + \frac{B/\beta}{1-x/\beta} \\
&= \frac{A'}{1-x/\alpha} + \frac{B'}{1-x/\beta} \\
&= A' \sum_{i=0}^{\infty} \frac{x^i}{\alpha^i} + B' \sum_{i=0}^{\infty} \frac{x^i}{\beta^i}
\end{aligned}$$

$$a_n = A' \frac{1}{\alpha^n} + B' \frac{1}{\beta^n}$$

α, β are roots of $1-x-x^2$

$$a_n = A' \alpha'^n + B' \beta'^n$$

$\alpha' := 1/\alpha, \beta' := 1/\beta$ are roots of $x^2 - x - 1$

$$\alpha' = \frac{1+\sqrt{5}}{2}$$

$$\beta' = \frac{1-\sqrt{5}}{2}$$

$$a_n = A'(\frac{1+\sqrt{5}}{2})^n + B'(\frac{1-\sqrt{5}}{2})^n$$

$$a_0 = A' + B' = 0, \text{ so } A' = -B'$$

$$a_1 = A'(\frac{1+\sqrt{5}}{2}) - A'(\frac{1-\sqrt{5}}{2}) = 1$$

$$A' = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

In general:

$$\alpha' = \frac{1+\sqrt{5}}{2}$$

$$\beta' = \frac{1-\sqrt{5}}{2}$$

$$a_n = A'(\frac{1+\sqrt{5}}{2})^n + B'(\frac{1-\sqrt{5}}{2})^n$$

$$a_0 = A' + B' = 0, \text{ so } A' = -B'$$

$$a_1 = A'(\frac{1+\sqrt{5}}{2}) - A'(\frac{1-\sqrt{5}}{2}) = 1$$

$$A' = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

In general:

Theorem.

$$\alpha' = \frac{1+\sqrt{5}}{2}$$

$$\beta' = \frac{1-\sqrt{5}}{2}$$

$$a_n = A'(\frac{1+\sqrt{5}}{2})^n + B'(\frac{1-\sqrt{5}}{2})^n$$

$$a_0 = A' + B' = 0, \text{ so } A' = -B'$$

$$a_1 = A'(\frac{1+\sqrt{5}}{2}) - A'(\frac{1-\sqrt{5}}{2}) = 1$$

$$A' = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^n - \left(\frac{1-\sqrt{5}}{2} \right)^n \right]$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence,*

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$\begin{aligned}\alpha' &= \frac{1+\sqrt{5}}{2} \\ \beta' &= \frac{1-\sqrt{5}}{2} \\ a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\ a_0 &= A' + B' = 0, \text{ so } A' = -B' \\ a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\ A' &= \frac{1}{\sqrt{5}} \\ a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \cdots + c_r a_{n-r} = 0$$

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots.

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

$$\begin{aligned}
\alpha' &= \frac{1+\sqrt{5}}{2} \\
\beta' &= \frac{1-\sqrt{5}}{2} \\
a_n &= A' \left(\frac{1+\sqrt{5}}{2}\right)^n + B' \left(\frac{1-\sqrt{5}}{2}\right)^n \\
a_0 &= A' + B' = 0, \text{ so } A' = -B' \\
a_1 &= A' \left(\frac{1+\sqrt{5}}{2}\right) - A' \left(\frac{1-\sqrt{5}}{2}\right) = 1 \\
A' &= \frac{1}{\sqrt{5}} \\
a_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]
\end{aligned}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example.

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \cdots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \cdots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \cdots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \cdots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \cdots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \cdots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

In general:

Theorem. *Let $(a_0, a_1, a_2, a_3, \dots)$ be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its roots. If they are distinct,

$$a_n = A_1 \alpha_1^n + \dots + A_r \alpha_r^n$$

for some fixed A_1, \dots, A_r

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + A_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1 \left(\frac{1+\sqrt{5}}{2}\right) - A_1 \left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$$\alpha_1 = 1$$

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$$\alpha_2 = 2$$

$$a_n = (A_1 + A_2n)1^n + A_32^n$$

Example (Returning to Fibonacci).

$$a_n = a_{n-1} + a_{n-2}$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Characteristic polynomial:

$$x^2 - x - 1 = 0$$

Its roots,

$$\alpha_1 = \frac{1+\sqrt{5}}{2}$$

$$\alpha_2 = \frac{1-\sqrt{5}}{2}$$

$$a_n = A_1\left(\frac{1+\sqrt{5}}{2}\right)^n + A_2\left(\frac{1-\sqrt{5}}{2}\right)^n$$

$$a_0 = A_1 + A_2 = 0, \text{ so } A_1 = -A_2$$

$$a_1 = A_1\left(\frac{1+\sqrt{5}}{2}\right) - A_1\left(\frac{1-\sqrt{5}}{2}\right) = 1$$

$$A_1 = \frac{1}{\sqrt{5}}$$

$$a_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right]$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$$\alpha_2 = 2$$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!

In general:

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2n)1^n + A_32^n$$

Rest: exercise!

In general:

Theorem.

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2n)1^n + A_32^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_2, a_3, \dots) be a sequence,*

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_2, a_3, \dots) be a sequence, satisfying a recurrence relation of the form*

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2n)1^n + A_32^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_2, a_3, \dots) be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_2, a_3, \dots) be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_2, a_3, \dots) be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_2, a_3, \dots) be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its root with multiplicity m_i .

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!

In general:

Theorem. *Let (a_0, a_1, a_2, \dots) be a sequence, satisfying a recurrence relation of the form*

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its root with multiplicity m_i .

$$a_n = (A_{11} + A_{12}n + \dots + A_{1m_1} n^{m_1-1}) \alpha_1^n + \dots + (A_{r1} + A_{r2}n + \dots + A_{rm_r} n^{m_r-1}) \alpha_r^n$$

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n) 1^n + A_3 2^n$$

Rest: exercise!

In general:

Theorem. Let (a_0, a_2, a_3, \dots) be a sequence, satisfying a recurrence relation of the form

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0$$

Define the associated characteristic polynomial,

$$c_0 x^r + c_1 x^{r-1} + \dots + c_r = 0$$

Let α_i be its root with multiplicity m_i .

$$a_n = (A_{11} + A_{12}n + \dots + A_{1m_1}n^{m_1-1})\alpha_1^n + \dots + (A_{r1} + A_{r2}n + \dots + A_{rm_r}n^{m_r-1})\alpha_r^n$$

for some fixed A_{ij} s

Example. Find a formula for the recurrence relation

$$a_n = 4a_{n-1} - 5a_{n-2} + 2a_{n-3}$$

Characteristic polynomial:

$$x^3 - 4x^2 + 5x - 2$$

Its roots,

$\alpha_1 = 1$ with multiplicity 2

$\alpha_2 = 2$

$$a_n = (A_1 + A_2 n)1^n + A_3 2^n$$

Rest: exercise!