

**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\mathcal{C}_r^n = \{ \hspace{10em} \}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\mathcal{C}_r^n = \{ \{x_1, x_2, \dots, x_r\} \}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\mathcal{C}_r^n = \{ \{x_1, x_2, \dots, x_r\} \mid x_i \in X \}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{ \end{aligned}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)]\}\end{aligned}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X\}\end{aligned}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\}\end{aligned}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{ \end{aligned}$$

1



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned} \mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \end{aligned}$$

1

**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if □

**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!}$  □

**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$





**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:  $|\mathcal{C}_r^n|$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

number of arrangements =

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:  $|\mathcal{C}_r^n|$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

$$\begin{aligned} \mathcal{C}_r^n &= \{ \{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct} \} \\ &= \{ [(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct} \} \\ &= \{ [a] \mid a \in r - \text{arrangements} \} \end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:  $|\mathcal{C}_r^n|$



**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:  $|\mathcal{C}_r^n|$

□

$$\frac{n!}{(n-r)!} = \text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:  $|\mathcal{C}_r^n|$

□

$$\frac{n!}{(n-r)!} = \text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

$$|\mathcal{C}_r^n| = \frac{n!}{(n-r)!r!}$$

**Example.** Number of ways to choose  $r$  distinct objects from a set of  $X$  of  $n$  objects.

*Solution.*

$$\begin{aligned}\mathcal{C}_r^n &= \{\{x_1, x_2, \dots, x_r\} \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[(x_1, x_2, \dots, x_r)] \mid x_i \in X, x_i \text{ distinct}\} \\ &= \{[a] \mid a \in r - \text{arrangements}\}\end{aligned}$$

$[(x_1, \dots, x_r)]$  is the equivalence class, for the relation,

$$(x_1, \dots, x_r) \sim (y_1, \dots, y_r)$$

if  $(x_1, \dots, x_r)$  is a rearrangement of  $(y_1, \dots, y_r)$ .

Size of each equivalence class:  $\frac{r!}{(r-r)!} = r!$

Number of equivalence classes:  $|\mathcal{C}_r^n|$

□

$$\frac{n!}{(n-r)!} = \text{number of arrangements} = |\mathcal{C}_r^n| \times r!$$

$$\binom{n}{r} := |\mathcal{C}_r^n| = \frac{n!}{(n-r)!r!}$$

**Example.** Find the number of positive divisors of 50.



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10$$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2$$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j\}$  □

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, \}$  □

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
□

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
[[n]] □



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$f :$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]]$$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| \qquad \qquad \qquad \square$$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| \quad \square$$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 \quad \square$$



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.* □

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\} \qquad \qquad \qquad \square$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \qquad \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{ \qquad \qquad \qquad \}$$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\} \qquad \qquad \qquad \square$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \qquad \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \qquad \qquad \qquad \}$$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \qquad \}$$

$\square$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

□

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

$$S_i = \{ \qquad \qquad \qquad \}$$

□

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

$$S_i = \{(a, i) \qquad \qquad \qquad \}$$

□



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, \ a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, \quad \quad \quad \}$$

□

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$

$$[[n]] := \{0, 1, \dots, n\}$$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$\square$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=?}^{10} S_i \qquad \square$$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i \qquad \square$$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6 \qquad \square$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$\square$

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

□

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[2]| \times |[1]| = 3 \times 2 = 6$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

$$|S|$$

□

**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

$$|S| = 1 + 2 + \dots + 9$$

□



**Example.** Find the number of positive divisors of 50.

*Solution.*

$$50 = 5 \times 10 = 5 \times 5 \times 2 = 5^2 \times 2$$

Set of divisors:  $\mathcal{D}_{50} := \{5^i \times 2^j \mid 0 \leq i \leq 2, 0 \leq j \leq 1\}$   
 $[[n]] := \{0, 1, \dots, n\}$

$$f : [[2]] \times [[1]] \rightarrow \mathcal{D}_{50}$$

$f$  is a bijection

$$|\mathcal{D}_{50}| = |[ [2] ]| \times |[ [1] ]| = 3 \times 2 = 6$$

**Example.** Count the number of pairs  $(a, b)$  so that  $a, b \in \{1, 2, \dots, 10\}$  and  $a$  is smaller.

*Solution.*

$$S := \{(a, b) \mid a, b \in \{1, 2, \dots, 10\}, a < b\}$$

$$S_i = \{(a, i) \mid a \in \{1, 2, \dots, 10\}, a < i\}$$

$$S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

$$S = \sqcup_{i=2}^{10} S_i$$

$$|S| = \sum_{i=2}^{10} |S_i|$$

$$\square \quad |S_i| = i - 1$$

$$|S| = 1 + 2 + \dots + 9 = \dots$$

□

**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.* □

**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair



**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair  
Each element of  $\mathcal{P}_n$  looks like:



**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair  
Each element of  $\mathcal{P}_n$  looks like:

$$\{ \hspace{10cm} \}$$



**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair  
Each element of  $\mathcal{P}_n$  looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \}$$



**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair  
Each element of  $\mathcal{P}_n$  looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$





**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair  
Each element of  $\mathcal{P}_n$  looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{ (x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$



**Example.** Count the number of ways to form  $n$  pairs out of a set  $X$  of  $2n$  elements.

*Solution.*  $\mathcal{P}_n$  : Number of distinct ways to pair  
Each element of  $\mathcal{P}_n$  looks like:

$$\{(x_1, x_2), (x_3, x_4), \dots, (x_{2n-1}, x_{2n}) \mid x_i \in X\}$$

$$\{[ \, x_1, x_2 \, , \, x_3, x_4 \, , \dots , \, x_{2n-1}, x_{2n} \, ] \mid x_i \in X\}$$

