

Example.

**Example.** Prove that any subset of cardinality  $n + 1$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A :=$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{ \qquad \qquad \qquad \}$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{ \hspace{10em} \}$

$f$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{ \hspace{10em} \}$

$f : I \rightarrow$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{ \hspace{10em} \}$

$f : I \rightarrow B,$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\} \hspace{10em}\}$

$f : I \rightarrow B,$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \quad \quad \quad \}$

$f : I \rightarrow B,$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B,$



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j)$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i]$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\}$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a$ ,  $b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j]$



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

□

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ . **Example.**

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

□

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

□

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$



**Example.** In a group of 6 people, there is either a subset

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .  
**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = [a_i, a_j] = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$



**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

□

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$



*Solution.*

Denote the set,

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

□

*Solution.*

Denote the set,  $A :=$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

□

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

□

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

□

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\}$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

□

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

$\square$

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*  
Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

*Solution.*  
Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$

□

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

$\square$

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)

Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$

If any  $b_i, b_j$

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)

Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$

If any  $b_i, b_j \in \{b_1, b_2, b_3\}$

□

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

□

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.

□

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
then  $\{a_1, b_i, b_j\}$  form the triple.  
□ Otherwise,

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
then  $\{a_1, b_i, b_j\}$  form the triple.  
□ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
then  $\{a_1, b_i, b_j\}$  form the triple.  
□ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
and form the triple □

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = [a_i, a_j] = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)

Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$

If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,

then  $\{a_1, b_i, b_j\}$  form the triple.

□ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other

and form the triple

□

**Example.**

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = [a_i, a_j] = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.  
 □ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
 and form the triple □

**Example.** Consider a graph with 6 vertices,

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)

Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$

If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other, then  $\{a_1, b_i, b_j\}$  form the triple.

□ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.  
 □ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
 and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue.

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.  
 □ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
 and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = [a_i, a_j] = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.  
 □ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
 and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.  
 □ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
 and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.



**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$

$I := \{1, 2, \dots, n, n + 1\}$

$B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$

$f : I \rightarrow B$ , defined,  $f(i) = [a_i]$

$f$  cannot be injective,

so  $f(i) = f(j) \iff [a_i] = \{a_i, a_j\} = [a_j] \in B$

i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$

$a_1$  either knows or does not know

$\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)

Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$

If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other, then  $\{a_1, b_i, b_j\}$  form the triple.

□ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

*Solution.*

**Example.** Prove that any subset of cardinality  $n + 1$  of the set  $\{1, 2, \dots, 2n\}$  has a pair,  $a, b$ , so that  $a$  divides  $b$ .

*Solution.*

Denote the subset,  $A := \{a_1, a_2, \dots, a_{n+1}\}$   
 $I := \{1, 2, \dots, n, n + 1\}$   
 $B := \{\{1, 2\}, \{2, 4\}, \dots, \{n, 2n\}\}$   
 $f : I \rightarrow B$ , defined,  $f(i) = [a_i]$   
 $f$  cannot be injective,  
 so  $f(i) = f(j) \iff [a_i] = [a_i, a_j] = [a_j] \in B$   
 i.e.  $a_i$  divides  $a_j$

**Example.** In a group of 6 people, there is either a subset of 3 who know each other, or a subset of 3 who do not know each other.

*Solution.*

Denote the set,  $A := \{a_1, a_2, \dots, a_6\}$   
 $a_1$  either knows or does not know  
 $\{b_1, b_2, b_3\} \subset \{a_2, \dots, a_6\}$  (pigeonhole principle)  
 Assume  $a_1$  knows  $\{b_1, b_2, b_3\}$   
 If any  $b_i, b_j \in \{b_1, b_2, b_3\}$  know each other,  
 then  $\{a_1, b_i, b_j\}$  form the triple.  
 □ Otherwise, none among  $\{b_1, b_2, b_3\}$  know each other  
 and form the triple □

**Example.** Consider a graph with 6 vertices, such that every pair vertices is joined by exactly one edge which can be coloured either red, or blue. Show that there are three vertices so that the edges joining each pair of them are of the same colour.

*Solution.* Same as above □