

Example.

Example. Find the number of ways to distribute

Example. Find the number of ways to distribute m identical objects

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

$$2 + 3 + 0$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial:

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial:

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial:

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \cdots (1 + x + x^2 + \cdots)}_{n \text{ times}}$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 2 + 3 + 0 Monomial: $x^2x^3x^0$
 0 + 2 + 3 Monomial: $x^0x^2x^3$
 1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \dots (1 + x + x^2 + \cdots)}$$

(because $a_mx^m = x^{r_1}x^{r_2} \overset{n \text{ times}}{\dots} x^{r_n} + \dots,$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 $2 + 3 + 0$ Monomial: $x^2x^3x^0$
 $0 + 2 + 3$ Monomial: $x^0x^2x^3$
 $1 + 1 + 3$ Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \cdots (1 + x + x^2 + \cdots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \cdots x^{r_n} + \cdots$,
such that $r_1 + r_2 + \cdots + r_n = m$)

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1+x+x^2+\cdots)(1+x+x^2+\cdots)(1+x+x^2+\cdots)$$

Solution.

$$\underbrace{(1+x+x^2+\cdots)(1+x+x^2+\cdots)\cdots(1+x+x^2+\cdots)}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$, n times

such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 $2 + 3 + 0$ Monomial: $x^2x^3x^0$
 $0 + 2 + 3$ Monomial: $x^0x^2x^3$
 $1 + 1 + 3$ Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \dots (1 + x + x^2 + \cdots)}$$

$$\begin{aligned} &\hspace{10em} n \text{ times} \\ \text{(because } a_mx^m &= x^{r_1}x^{r_2} \dots x^{r_n} + \dots, \end{aligned}$$

$$\text{such that } r_1 + r_2 + \cdots + r_n = m)$$

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 $2 + 3 + 0$ Monomial: $x^2x^3x^0$
 $0 + 2 + 3$ Monomial: $x^0x^2x^3$
 $1 + 1 + 3$ Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \dots (1 + x + x^2 + \cdots)}$$

$\qquad\qquad\qquad n \text{ times}$
 (because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \cdots$,
 such that $r_1 + r_2 + \cdots + r_n = m$)

$$\begin{aligned} &= (1 + x + x^2 + \cdots)^n \\ &= \frac{1}{(1-x)^n} \\ &= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots \end{aligned}$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 $2 + 3 + 0$ Monomial: $x^2x^3x^0$
 $0 + 2 + 3$ Monomial: $x^0x^2x^3$
 $1 + 1 + 3$ Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \dots (1 + x + x^2 + \cdots)}$$

$\qquad\qquad\qquad n \text{ times}$
 (because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
 such that $r_1 + r_2 + \cdots + r_n = m$)

$$\begin{aligned} &= (1 + x + x^2 + \cdots)^n \\ &= \frac{1}{(1-x)^n} \\ &= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots \end{aligned}$$

Answer: Coefficient of x^m ,

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 $2 + 3 + 0$ Monomial: $x^2x^3x^0$
 $0 + 2 + 3$ Monomial: $x^0x^2x^3$
 $1 + 1 + 3$ Monomial: $x^1x^1x^3$

$$(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots)$$

Solution.

$$\underbrace{(1 + x + x^2 + \cdots)(1 + x + x^2 + \cdots) \dots (1 + x + x^2 + \cdots)}$$

$\qquad\qquad\qquad n \text{ times}$
 (because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
 such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes
 $2 + 3 + 0$ Monomial: $x^2x^3x^0$
 $0 + 2 + 3$ Monomial: $x^0x^2x^3$
 $1 + 1 + 3$ Monomial: $x^1x^1x^3$

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \dots (1 + x + \cdots)}$$

$\qquad\qquad\qquad n \text{ times}$
 (because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
 such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes. **Example.** Find the number of ways

5 objects in 3 boxes
 2 + 3 + 0 Monomial: $x^2x^3x^0$
 0 + 2 + 3 Monomial: $x^0x^2x^3$
 1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \dots (1 + x + \cdots)}$$

$\qquad\qquad\qquad n \text{ times}$
 (because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
 such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Example. Find the number of ways to distribute m identical objects

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \dots (1 + x + \cdots)}$$

n times

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Example. Find the number of ways to distribute m identical objects in n distinct boxes

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \dots (1 + x + \cdots)}$$

n times

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \dots (1 + x + \cdots)}$$

n times

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

5 objects in 3 boxes
2 + 3 + 0 Monomial: $x^2x^3x^0$
0 + 2 + 3 Monomial: $x^0x^2x^3$
1 + 1 + 3 Monomial: $x^1x^1x^3$

Solution.

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \dots (1 + x + \cdots)}$$

$$\begin{aligned} & \text{\scriptsize n times} \\ \text{(because } a_mx^m &= x^{r_1}x^{r_2} \dots x^{r_n} + \dots, \\ \text{such that } r_1 + r_2 + \dots + r_n &= m) \end{aligned}$$

$$\begin{aligned} &= (1 + x + x^2 + \cdots)^n \\ &= \frac{1}{(1-x)^n} \\ &= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots \end{aligned}$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \cdots)(1 + x + \cdots)(1 + x + \cdots)$$

Solution.

$$\underbrace{(1 + x + \cdots)(1 + x + \cdots) \cdots (1 + x + \cdots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \cdots x^{r_n} + \cdots$,

such that $r_1 + r_2 + \cdots + r_n = m$)

$$= (1 + x + x^2 + \cdots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \cdots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (1 + x + x^2 + \dots)^n \\ &= \frac{1}{(1-x)^n} \\ &= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots \end{aligned}$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes.

5 objects in 3 boxes

2 + 3 + 0 Monomial: $x^2x^3x^0$

0 + 2 + 3 Monomial: $x^0x^2x^3$

1 + 1 + 3 Monomial: $x^1x^1x^3$

$$(1 + x + \dots)(1 + x + \dots)(1 + x + \dots)$$

Solution.

$$\underbrace{(1 + x + \dots)(1 + x + \dots) \dots (1 + x + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,

such that $r_1 + r_2 + \dots + r_n = m$)

$$= (1 + x + x^2 + \dots)^n$$

$$= \frac{1}{(1-x)^n}$$

$$= 1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots$$

Answer: Coefficient of x^m , i.e. $\binom{n+m-1}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_m x^m = x^{r_1} x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n (1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n \left(1 + nx + \binom{n+1}{2} x^2 + \binom{n+2}{3} x^3 + \dots \right)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2} x^{n+2} + \binom{n+2}{3} x^{n+3} + \dots$$

$$\text{General term: } \binom{n+k-1}{k} x^{n+k}$$

$$\text{Answer: } \binom{m-1}{m-n}$$

Example. Number of ways to choose

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_m x^m = x^{r_1} x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n (1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n \left(1 + nx + \binom{n+1}{2} x^2 + \binom{n+2}{3} x^3 + \dots \right)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2} x^{n+2} + \binom{n+2}{3} x^{n+3} + \dots$$

$$\text{General term: } \binom{n+k-1}{k} x^{n+k}$$

$$\text{Answer: } \binom{m-1}{m-n}$$

Example. Number of ways to choose m elements out of n elements

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_m x^m = x^{r_1} x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n (1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n \left(1 + nx + \binom{n+1}{2} x^2 + \binom{n+2}{3} x^3 + \dots \right)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2} x^{n+2} + \binom{n+2}{3} x^{n+3} + \dots$$

$$\text{General term: } \binom{n+k-1}{k} x^{n+k}$$

$$\text{Answer: } \binom{m-1}{m-n}$$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1+x) \dots (1+x)$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x+x^2+\dots)(x+x^2+\dots)\dots(x+x^2+\dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2}\dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x+x^2+x^3+\dots)^n$$

$$= x^n(1+x+x^2+x^3+\dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1+nx+\binom{n+1}{2}x^2+\binom{n+2}{3}x^3+\dots)$$

$$= x^n+nx^{n+1}+\binom{n+1}{2}x^{n+2}+\binom{n+2}{3}x^{n+3}+\dots)$$

$$\text{General term: } \binom{n+k-1}{k}x^{n+k}$$

$$\text{Answer: } \binom{m-1}{m-n}$$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1+x) \dots (1+x) = (1+x)^n$$

Answer: $\binom{n}{m}$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x+x^2+\dots)(x+x^2+\dots)\dots(x+x^2+\dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2}\dots x^{r_n} + \dots$,
such that $r_1+r_2+\dots+r_n=m$)

$$= (x+x^2+x^3+\dots)^n$$

$$= x^n(1+x+x^2+x^3+\dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1+nx+\binom{n+1}{2}x^2+\binom{n+2}{3}x^3+\dots)$$

$$= x^n+nx^{n+1}+\binom{n+1}{2}x^{n+2}+\binom{n+2}{3}x^{n+3}+\dots$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

3

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$\begin{aligned} &= (x + x^2 + x^3 \dots)^n \\ &= x^n(1 + x + x^2 + x^3 \dots)^n \\ &= \frac{x^n}{(1-x)^n} \\ &= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots) \\ &= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots \end{aligned}$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1 = 2+2$$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1 = 2+2$$

Example.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1 = 2+2$$

Example. Find the number of partitions of m into integers

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1 = 2+2$$

Example. Find the number of partitions of m into integers with allowed sizes 1, 2, 3.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1 = 2+2$$

Example. Find the number of partitions of m into integers with allowed sizes 1, 2, 3.

Solution.

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$

Example. Number of ways to choose m elements out of n elements

Solution. Generating function:

$$(1 + x) \dots (1 + x) = (1 + x)^n$$

Answer: $\binom{n}{m}$

Partitions of integers

Examples

$$3 = 1 + 1 + 1 = 2 + 1$$

$$4 = 1 + 1 + 1 + 1 = 2 + 1 + 1 = 2+2$$

Example. Find the number of partitions of m into integers with allowed sizes 1, 2, 3.

Solution. $(1+x+\dots)(1+x^2+x^4+\dots)(1+x^3+x^6+\dots)$

Example. Find the number of ways to distribute m identical objects in n distinct boxes so that each box contains at least one object.

Solution.

$$\underbrace{(x + x^2 + \dots)(x + x^2 + \dots) \dots (x + x^2 + \dots)}_{n \text{ times}}$$

(because $a_mx^m = x^{r_1}x^{r_2} \dots x^{r_n} + \dots$,
such that $r_1 + r_2 + \dots + r_n = m$)

$$= (x + x^2 + x^3 \dots)^n$$

$$= x^n(1 + x + x^2 + x^3 \dots)^n$$

$$= \frac{x^n}{(1-x)^n}$$

$$= x^n(1 + nx + \binom{n+1}{2}x^2 + \binom{n+2}{3}x^3 + \dots)$$

$$= x^n + nx^{n+1} + \binom{n+1}{2}x^{n+2} + \binom{n+2}{3}x^{n+3} + \dots)$$

General term: $\binom{n+k-1}{k}x^{n+k}$

Answer: $\binom{m-1}{m-n}$