



CW-complex

-


CW-complex




CW-complex




CW-complex





CW-complex





CW-complex




CW-complex




CW-complex





CW-complex




CW-complex





CW-complex



CW-complex




CW-complex



X_0 : set of points


CW-complex



X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$


CW-complex



X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

CW-complex




X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$$X = \cup X_i$$

CW-complex




X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

CW-complex



X_0 : set of points


$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

CW-complex

$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$




X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

CW-complex



$$e_{\alpha}^n = D_{\alpha}^n \setminus \partial D_{\alpha}^n$$
$$X_n \setminus X_{n-1} = \sqcup e_{\alpha}^n$$


X_0 : set of points

$$\phi_{\alpha} : S^{n-1} = \partial D_{\alpha}^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_{\alpha}^n / \sim, \text{ where } y \sim \phi_{\alpha}(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

CW-complex



$$e_{\alpha}^n = D_{\alpha}^n \setminus \partial D_{\alpha}^n$$
$$X_n \setminus X_{n-1} = \sqcup e_{\alpha}^n$$
$$\Phi_{\alpha} : D_{\alpha}^n \rightarrow X$$


X_0 : set of points

$$\phi_{\alpha} : S^{n-1} = \partial D_{\alpha}^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_{\alpha}^n / \sim, \text{ where } y \sim \phi_{\alpha}(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

CW-complex



$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$
$$X_n \setminus X_{n-1} = \sqcup e_\alpha^n$$
$$\Phi_\alpha : D_\alpha^n \rightarrow X \text{ (characteristic map)}$$


X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

CW-complex



X_0 : set of points


$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$
$$X_n \setminus X_{n-1} = \sqcup e_\alpha^n$$
$$\Phi_\alpha : D_\alpha^n \rightarrow X \text{ (characteristic map)}$$
$$\Phi_\alpha|_{\partial D_\alpha^n} = \phi$$

CW-complex



X_0 : set of points


$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$
$$X_n \setminus X_{n-1} = \sqcup e_\alpha^n$$
$$\Phi_\alpha : D_\alpha^n \rightarrow X \text{ (characteristic map)}$$
$$\Phi_\alpha|_{\partial D_\alpha^n} = \phi$$
$$\Phi_\alpha|_{Int D_\alpha^n} \text{ is a homeomorphism onto } e_\alpha^n$$

CW-complex



X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$


$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$
$$X_n \setminus X_{n-1} = \sqcup e_\alpha^n$$
$$\Phi_\alpha : D_\alpha^n \rightarrow X \text{ (characteristic map)}$$
$$\Phi_\alpha|_{\partial D_\alpha^n} = \phi$$
$$\Phi_\alpha|_{Int D_\alpha^n} \text{ is a homeomorphism onto } e_\alpha^n$$

(X_n, X_{n-1}) is a good pair?

CW-complex



X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$


$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$
$$X_n \setminus X_{n-1} = \sqcup e_\alpha^n$$
$$\Phi_\alpha : D_\alpha^n \rightarrow X \text{ (characteristic map)}$$
$$\Phi_\alpha|_{\partial D_\alpha^n} = \phi$$
$$\Phi_\alpha|_{Int D_\alpha^n} \text{ is a homeomorphism onto } e_\alpha^n$$

(X_n, X_{n-1}) is a good pair.

CW-complex



X_0 : set of points

$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$
 $X_n = X_{n-1} \sqcup D_\alpha^n / \sim$, where $y \sim \phi_\alpha(x)$


$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$
 $X_n \setminus X_{n-1} = \sqcup e_\alpha^n$
 $\Phi_\alpha : D_\alpha^n \rightarrow X$ (characteristic map)
 $\Phi_\alpha|_{\partial D_\alpha^n} = \phi$
 $\Phi_\alpha|_{Int D_\alpha^n}$ is a homeomorphism onto e_α^n

(X_n, X_{n-1}) is a good pair.
 $H_k(X_n, X_{n-1})$

CW-complex



X_0 : set of points

$$\phi_\alpha : S^{n-1} = \partial D_\alpha^n \rightarrow X_{n-1}$$
$$X_n = X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x)$$


$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes

$$e_\alpha^n = D_\alpha^n \setminus \partial D_\alpha^n$$
$$X_n \setminus X_{n-1} = \sqcup e_\alpha^n$$
$$\Phi_\alpha : D_\alpha^n \rightarrow X \text{ (characteristic map)}$$
$$\Phi_\alpha|_{\partial D_\alpha^n} = \phi$$
$$\Phi_\alpha|_{Int D_\alpha^n} \text{ is a homeomorphism onto } e_\alpha^n$$

$$(X_n, X_{n-1}) \text{ is a good pair.}$$
$$H_k(X_n, X_{n-1}) \cong \tilde{H}_k(X_n/X_{n-1})$$

CW-complex



X_0 : set of points

$$\begin{aligned} \phi_\alpha : S^{n-1} = \partial D_\alpha^n &\rightarrow X_{n-1} \\ X_n &= X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x) \end{aligned}$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. *Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes*


$$\begin{aligned} e_\alpha^n &= D_\alpha^n \setminus \partial D_\alpha^n \\ X_n \setminus X_{n-1} &= \sqcup e_\alpha^n \\ \Phi_\alpha : D_\alpha^n &\rightarrow X \text{ (characteristic map)} \\ \Phi_\alpha|_{\partial D_\alpha^n} &= \phi \\ \Phi_\alpha|_{Int D_\alpha^n} &\text{ is a homeomorphism onto } e_\alpha^n \end{aligned}$$

(X_n, X_{n-1}) is a good pair.

$$H_k(X_n, X_{n-1}) \cong \tilde{H}_k(X_n/X_{n-1}) \cong \tilde{H}_k(\vee S^n)$$

$$H_k(X_n, X_{n-1}) =$$

CW-complex



X_0 : set of points

$$\begin{aligned} \phi_\alpha : S^{n-1} = \partial D_\alpha^n &\rightarrow X_{n-1} \\ X_n &= X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x) \end{aligned}$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes


$$\begin{aligned} e_\alpha^n &= D_\alpha^n \setminus \partial D_\alpha^n \\ X_n \setminus X_{n-1} &= \sqcup e_\alpha^n \\ \Phi_\alpha : D_\alpha^n &\rightarrow X \text{ (characteristic map)} \\ \Phi_\alpha|_{\partial D_\alpha^n} &= \phi \\ \Phi_\alpha|_{Int D_\alpha^n} &\text{ is a homeomorphism onto } e_\alpha^n \end{aligned}$$

(X_n, X_{n-1}) is a good pair.

$$H_k(X_n, X_{n-1}) \cong \tilde{H}_k(X_n/X_{n-1}) \cong \tilde{H}_k(\vee S^n)$$

$$H_k(X_n, X_{n-1}) = \begin{cases} \oplus \mathbb{Z} & k = n \end{cases}$$

CW-complex



X_0 : set of points

$$\begin{aligned} \phi_\alpha : S^{n-1} = \partial D_\alpha^n &\rightarrow X_{n-1} \\ X_n &= X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x) \end{aligned}$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes


$$\begin{aligned} e_\alpha^n &= D_\alpha^n \setminus \partial D_\alpha^n \\ X_n \setminus X_{n-1} &= \sqcup e_\alpha^n \\ \Phi_\alpha : D_\alpha^n &\rightarrow X \text{ (characteristic map)} \\ \Phi_\alpha|_{\partial D_\alpha^n} &= \phi \\ \Phi_\alpha|_{Int D_\alpha^n} &\text{ is a homeomorphism onto } e_\alpha^n \end{aligned}$$

(X_n, X_{n-1}) is a good pair.

$$H_k(X_n, X_{n-1}) \cong \tilde{H}_k(X_n/X_{n-1}) \cong \tilde{H}_k(\vee S^n)$$

$$H_k(X_n, X_{n-1}) = \begin{cases} \oplus \mathbb{Z} & k = n \\ 0 & k \neq n \end{cases}$$

CW-complex



X_0 : set of points

$$\begin{aligned} \phi_\alpha : S^{n-1} = \partial D_\alpha^n &\rightarrow X_{n-1} \\ X_n &= X_{n-1} \sqcup D_\alpha^n / \sim, \text{ where } y \sim \phi_\alpha(x) \end{aligned}$$

$X = \cup X_i$, and U open in X if and only if $U \cap X_i$ open in each X_i .

Exercise. *Prove that the spheres (any dimension), torus, projective plane, Klein bottle, and any compact surface are all CW-complexes*

$$\begin{aligned} e_\alpha^n &= D_\alpha^n \setminus \partial D_\alpha^n \\ X_n \setminus X_{n-1} &= \sqcup e_\alpha^n \\ \Phi_\alpha : D_\alpha^n &\rightarrow X \text{ (characteristic map)} \\ \Phi_\alpha|_{\partial D_\alpha^n} &= \phi \\ \Phi_\alpha|_{Int D_\alpha^n} &\text{ is a homeomorphism onto } e_\alpha^n \end{aligned}$$

$$\begin{aligned} (X_n, X_{n-1}) &\text{ is a good pair.} \\ H_k(X_n, X_{n-1}) &\cong \tilde{H}_k(X_n/X_{n-1}) \cong \tilde{H}_k(\vee S^n) \end{aligned}$$

$$H_k(X_n, X_{n-1}) = \begin{cases} \oplus \mathbb{Z} & k = n \\ 0 & k \neq n \end{cases}$$

$$H_{n+1}(X_k, X_{k-1}) \rightarrow H_n(X_{k-1}) \rightarrow H_n(X_k) \rightarrow H_n(X_k, X_{k-1})$$

$$\underbrace{H_{n+1}(X_{n+2}, X_{n+1})}_{=0} \rightarrow H_n(X_{n+1}) \xrightarrow{\sim} H_n(X_{n+2}) \rightarrow \underbrace{H_n(X_{n+2}, X_{n+1})}_{=0}$$

$$\underbrace{H_{n+1}(X_{n+2}, X_{n+1})}_{=0} \rightarrow H_n(X_{n+1}) \overset{\sim}{\rightarrow} H_n(X_{n+2}) \rightarrow \underbrace{H_n(X_{n+2}, X_{n+1})}_{=0}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2})$$

$$\underbrace{H_{n+1}(X_{n+3}, X_{n+2})}_{=0} \rightarrow H_n(X_{n+2}) \xrightarrow{\sim} H_n(X_{n+3}) \rightarrow \underbrace{H_n(X_{n+3}, X_{n+2})}_{=0}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3})$$

$$\underbrace{H_{n+1}(X_{n+3}, X_{n+2})}_{=0} \rightarrow H_n(X_{n+2}) \xrightarrow{\sim} H_n(X_{n+3}) \rightarrow \underbrace{H_n(X_{n+3}, X_{n+2})}_{=0}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$$

$$\underbrace{H_{n+1}(X_{n+3}, X_{n+2})}_{=0} \rightarrow H_n(X_{n+2}) \overset{\sim}{\rightarrow} H_n(X_{n+3}) \rightarrow \underbrace{H_n(X_{n+3}, X_{n+2})}_{=0}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$$

$$\underbrace{H_{n+1}(X_{n-1}, X_{n-2})}_{=0} \rightarrow H_n(X_{n-2}) \overset{\sim}{\rightarrow} H_n(X_{n-1}) \rightarrow \underbrace{H_n(X_{n-1}, X_{n-2})}_{=0}$$

$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$

$$\underbrace{H_{n+1}(X_{n-1}, X_{n-2})}_{=0} \rightarrow H_n(X_{n-2}) \overset{\sim}{\rightarrow} H_n(X_{n-1}) \rightarrow \underbrace{H_n(X_{n-1}, X_{n-2})}_{=0}$$

$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$

$$H_n(X_{n-1}) \cong H_n(X_{n-2})$$

$$\underbrace{H_{n+1}(X_{n-2}, X_{n-3})}_{=0} \rightarrow H_n(X_{n-3}) \overset{\sim}{\rightarrow} H_n(X_{n-2}) \rightarrow \underbrace{H_n(X_{n-2}, X_{n-3})}_{=0}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3})$$

$$\underbrace{H_{n+1}(X_{n-2}, X_{n-3})}_{=0} \rightarrow H_n(X_{n-3}) \overset{\sim}{\rightarrow} H_n(X_{n-2}) \rightarrow \underbrace{H_n(X_{n-2}, X_{n-3})}_{=0}$$

$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots$$

$$\underbrace{H_{n+1}(X_{n-2}, X_{n-3})}_{=0} \rightarrow H_n(X_{n-3}) \xrightarrow{\sim} H_n(X_{n-2}) \rightarrow \underbrace{H_n(X_{n-2}, X_{n-3})}_{=0}$$

$$\boxed{H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0$$

(if $n > 0$)

$$\underbrace{H_{n+1}(X_{n-2}, X_{n-3}) \rightarrow H_n(X_{n-3})}_{=0} \overset{\sim}{\rightarrow} H_n(X_{n-2}) \rightarrow \underbrace{H_n(X_{n-2}, X_{n-3})}_{=0}$$

$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$

$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0 \text{)}$

$$H_{n+1}(X_{n+1}, X_n) \rightarrow H_n(X_n) \rightarrow H_n(X_{n+1}) \rightarrow \underbrace{H_n(X_{n+1}, X_n)}_{=0}$$

$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$

$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0\text{)}$

$$H_{n+1}(X_{n+1}, X_n) \rightarrow H_n(X_n) \rightarrow H_n(X_{n+1}) \rightarrow 0$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0 \text{)}$$

$$H_{n+1}(X_{n+1}, X_n) \rightarrow H_n(X_n) \rightarrow H_n(X) \rightarrow 0$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0\text{)}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$H_{n+1}(X_{n+1}, X_n) \xrightarrow{\partial_{n+1}} H_n(X_n) \longrightarrow H_n(X_{n+1}) \cong H_n(X) \rightarrow 0$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$H_{n+1}(X_{n+1}, X_n) \xrightarrow{\partial_{n+1}} H_n(X_n) \longrightarrow H_n(X_{n+1}) \cong H_n(X) \rightarrow 0$$

$$H_n(X) \cong H_n(X_n)/Im \partial_{n+1}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc} H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\ & & \downarrow j_n & & & & \\ & & H_n(X_n, X_{n-1}) & & & & \end{array}$$

$$H_n(X) \cong H_n(X_n)/Im \partial_{n+1}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & &
 \end{array}$$

$$H_n(X) \cong H_n(X_n)/Im \partial_{n+1}$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & &
 \end{array}$$

$$H_n(X) \cong H_n(X_n)/Im \partial_{n+1} \cong j_n(H_n(X_n))/Im (j_n \circ \partial_{n+1})$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & & \\
 & & \downarrow \partial_n & & & & \\
 & & H_{n-1}(X_{n-1}) & & & &
 \end{array}$$

$$H_n(X) \cong H_n(X_n)/\text{Im } \partial_{n+1} \cong j_n(H_n(X_n))/\text{Im } (j_n \circ \partial_{n+1})$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & & \\
 & & \downarrow \partial_n & & & & \\
 & & H_{n-1}(X_{n-1}) & & & &
 \end{array}$$

$$H_n(X) \cong H_n(X_n)/\text{Im } \partial_{n+1} \cong j_n(H_n(X_n))/\text{Im } (j_n \circ \partial_{n+1}) \cong \ker \partial_n / \text{Im } (j_n \circ \partial_{n+1})$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & & \\
 & & \downarrow \partial_n & & & & \\
 H_{n-1}(X_{n-2}) \cong 0 & \longrightarrow & H_{n-1}(X_{n-1}) & \xrightarrow{j_{n-1}} & H_{n-1}(X_{n-1}, X_{n-2}) & &
 \end{array}$$

$$H_n(X) \cong H_n(X_n)/Im \partial_{n+1} \cong j_n(H_n(X_n))/Im (j_n \circ \partial_{n+1}) \cong ker \partial_n/Im (j_n \circ \partial_{n+1})$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0)$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & \searrow d_{n+1} & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & & \\
 & & \downarrow \partial_n & \searrow d_n & & & \\
 H_{n-1}(X_{n-2}) \cong 0 & \longrightarrow & H_{n-1}(X_{n-1}) & \xrightarrow{j_{n-1}} & H_{n-1}(X_{n-1}, X_{n-2}) & &
 \end{array}$$

$$H_n(X) \cong H_n(X_n)/Im \partial_{n+1} \cong j_n(H_n(X_n))/Im (j_n \circ \partial_{n+1}) \cong ker \partial_n/Im (j_n \circ \partial_{n+1}) \cong ker (j_{n-1} \circ \partial_n)/Im (j_n \circ \partial_{n+1})$$

$$H_n(X_{n+1}) \cong H_n(X_{n+2}) \cong H_n(X_{n+3}) \cong \dots \cong H_n(X) \text{ exercise!}$$

$$H_n(X_{n-1}) \cong H_n(X_{n-2}) \cong H_n(X_{n-3}) \cong \dots \cong H_n(X_0) \cong 0 \text{ (if } n > 0 \text{)}$$

$$\begin{array}{ccccccc}
 & & H_n(X_{n-1}) = 0 & & & & \\
 & & \downarrow & & & & \\
 H_{n+1}(X_{n+1}, X_n) & \xrightarrow{\partial_{n+1}} & H_n(X_n) & \longrightarrow & H_n(X_{n+1}) \cong H_n(X) & \rightarrow & 0 \\
 & \searrow^{d_{n+1}} & \downarrow j_n & & & & \\
 & & H_n(X_n, X_{n-1}) & & & & \\
 & & \downarrow \partial_n & & & & \\
 H_{n-1}(X_{n-2}) \cong 0 & \longrightarrow & H_{n-1}(X_{n-1}) & \xrightarrow{j_{n-1}} & H_{n-1}(X_{n-1}, X_{n-2}) & &
 \end{array}$$

$$\begin{aligned}
 H_n(X) &\cong H_n(X_n)/\text{Im } \partial_{n+1} \cong j_n(H_n(X_n))/\text{Im } (j_n \circ \partial_{n+1}) \cong \ker \partial_n / \text{Im } (j_n \circ \partial_{n+1}) \\
 &\cong \ker (j_{n-1} \circ \partial_n) / \text{Im } (j_n \circ \partial_{n+1}) \cong \ker d_n / \text{Im } d_{n+1}
 \end{aligned}$$

$$\begin{array}{ccc} H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) \\ & \searrow d_n & \downarrow j \\ & & H_{n-1}(X_{n-1}, X_{n-2}) \end{array}$$

$$H_n(D^n, \partial D^n)$$

$$\downarrow \Phi_*$$

$$H_n(X_n, X_{n-1}) \xrightarrow{\partial_n} H_{n-1}(X_{n-1})$$

$$\searrow d_n$$

$$\downarrow j$$

$$H_{n-1}(X_{n-1}, X_{n-2})$$

$$d_n(\Phi_*(1))$$

$$H_n(D^n, \partial D^n) \xrightarrow{\partial'_n} H_{n-1}(\partial D^n)$$

$$\downarrow \Phi_*$$

$$H_n(X_n, X_{n-1}) \xrightarrow{\partial_n} H_{n-1}(X_{n-1})$$

$$\searrow d_n$$

$$\downarrow j$$

$$H_{n-1}(X_{n-1}, X_{n-2})$$

$$d_n(\Phi_*(\partial_n'^{-1}(1)))$$

$$\begin{array}{ccc}
 H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) \\
 \downarrow \Phi_* & & \\
 H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) \\
 & \searrow d_n & \downarrow j \\
 & & H_{n-1}(X_{n-1}, X_{n-2})
 \end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1))))$$

$$\begin{array}{ccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) \\
\downarrow \Phi_* & & \downarrow \phi_* \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) \\
& \searrow d_n & \downarrow j \\
& & H_{n-1}(X_{n-1}, X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$\begin{array}{ccc}
 H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) \\
 \downarrow \Phi_* & & \downarrow \phi_* \\
 H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) \\
 & \searrow d_n & \downarrow j \\
 & & H_{n-1}(X_{n-1}, X_{n-2})
 \end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \Sigma_i n_i g_i$$

$$\begin{array}{ccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) \\
\downarrow \Phi_* & & \downarrow \phi_* \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) \\
& \searrow d_n & \downarrow j \\
& & H_{n-1}(X_{n-1}, X_{n-2}) \xrightarrow{q_*} H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & \\
\downarrow \Phi_* & & \downarrow \phi_* & & \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & \\
\downarrow \Phi_* & & \downarrow \phi_* & & \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & \\
\downarrow \Phi_* & & \downarrow \phi_* & & \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \Sigma_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \Sigma_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \Sigma_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \Sigma_i n_i j_*'^{-1}(q_*(g_i))$$

$$\textbf{Lemma. } H_{n-1}(X_{n-1}/X_{n-2}) \cong \vee S^{n-1}_\alpha$$

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & \\
\downarrow \Phi_* & & \downarrow \phi_* & & \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$\textbf{Lemma. } H_{n-1}(X_{n-1}/X_{n-2}) \cong \vee S^{n-1}_\alpha \xrightarrow{q''} S^{n-1}_\beta$$

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & \\
\downarrow \Phi_* & & \downarrow \phi_* & & \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

Lemma. $H_{n-1}(X_{n-1}/X_{n-2}) \cong \vee S_\alpha^{n-1} \xrightarrow{q''} S_\beta^{n-1}$ induces $q''_* : \oplus_\alpha \mathbb{Z} \rightarrow \mathbb{Z}$,

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & \\
\downarrow \Phi_* & & \downarrow \phi_* & & \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

Lemma. $H_{n-1}(X_{n-1}/X_{n-2}) \cong \vee S_{\alpha}^{n-1} \xrightarrow{q''} S_{\beta}^{n-1}$ induces $q''_* : \oplus_{\alpha} \mathbb{Z} \rightarrow \mathbb{Z}$, which is a projection onto the β th term.

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & & S_\beta^{n-1} \\
\downarrow \Phi_* & & \downarrow \phi_* & & \uparrow q'' \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

Lemma. $H_{n-1}(X_{n-1}/X_{n-2}) \cong \vee S_\alpha^{n-1} \xrightarrow{q''} S_\beta^{n-1}$ induces $q''_* : \oplus_\alpha \mathbb{Z} \rightarrow \mathbb{Z}$, which is a projection onto the β th term.

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & \xrightarrow{\quad\quad\quad} & S_\beta^{n-1} \\
\downarrow \Phi_* & & \downarrow \phi_* & & \uparrow q'' \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$d_n(\Phi_*(\partial_n'^{-1}(1))) = j(\partial_n(\Phi_*(\partial_n'^{-1}(1)))) = j(\phi_*(1))$$

$$j(\phi_*(1)) = \sum_i n_i g_i$$

$$q_*(j(\phi_*(1))) = \sum_i n_i q_*(g_i)$$

$$j_*'^{-1}(q_*(j(\phi_*(1)))) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

$$q'_*(\phi_*(1)) = \sum_i n_i j_*'^{-1}(q_*(g_i))$$

Lemma. $H_{n-1}(X_{n-1}/X_{n-2}) \cong \vee S_\alpha^{n-1} \xrightarrow{q''} S_\beta^{n-1}$ induces $q''_* : \oplus_\alpha \mathbb{Z} \rightarrow \mathbb{Z}$, which is a projection onto the β th term.

$$\begin{array}{ccccc}
H_n(D^n, \partial D^n) & \xrightarrow{\partial'_n} & H_{n-1}(\partial D^n) & \longrightarrow & S^{n-1}_\beta \\
\downarrow \Phi_* & & \downarrow \phi_* & & \uparrow q'' \\
H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X_{n-1}) & \xrightarrow{q'_*} & H_{n-1}(X_{n-1}/X_{n-2}) \\
& \searrow d_n & \downarrow j & & \downarrow j' \\
& & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow{q_*} & H_{n-1}(X_{n-1}/X_{n-2}, X_{n-2}/X_{n-2})
\end{array}$$

$$\begin{array}{ccc}
\partial D^n & & S^{n-1}_\beta \\
\downarrow \phi & & \uparrow q'' \\
X_{n-1} & \xrightarrow{q'} & X_{n-1}/X_{n-2}
\end{array}$$

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Generators g_{α}^n correspond to n -cells e_{α}

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Genarators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Genarators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Genarators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

$$d_n(g_{\alpha}^n) = \sum n_{\alpha, \beta} g_{\beta}^{n-1}$$

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Genarators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

$$d_n(g_{\alpha}^n) = \sum n_{\alpha, \beta} g_{\beta}^{n-1}$$

To find $n_{\alpha, \beta}$,

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Genarators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

$$d_n(g_{\alpha}^n) = \sum n_{\alpha, \beta} g_{\beta}^{n-1}$$

To find $n_{\alpha, \beta}$,

Let $\phi_{\alpha} : \partial D_{\alpha}^n \rightarrow X_{n-1}$ be the gluing map.

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Genarators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

$$d_n(g_{\alpha}^n) = \sum n_{\alpha, \beta} g_{\beta}^{n-1}$$

To find $n_{\alpha, \beta}$,

Let $\phi_{\alpha} : \partial D_{\alpha}^n \rightarrow X_{n-1}$ be the gluing map.

Compose with $q : X_{n-1} \rightarrow X_{n-1} / (X_{n-1} \setminus e_{\beta}) \cong S^{n-1}$,

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Generators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

$$d_n(g_{\alpha}^n) = \sum n_{\alpha, \beta} g_{\beta}^{n-1}$$

To find $n_{\alpha, \beta}$,

Let $\phi_{\alpha} : \partial D_{\alpha}^n \rightarrow X_{n-1}$ be the gluing map.

Compose with $q : X_{n-1} \rightarrow X_{n-1}/(X_{n-1} \setminus e_{\beta}) \cong S^{n-1}$,

$$q \circ \phi_{\alpha} : S^{n-1} \cong \partial D^{n-1} \rightarrow X_{n-1}/(X_{n-1} \setminus e_{\beta}) \cong S^{n-1}$$

Summary of cellular homology

$$H_n(X_n, X_{n-1}) = \oplus_{\alpha} \mathbb{Z}$$

Generators g_{α}^n correspond to n -cells e_{α}

$$\rightarrow H_{n+1}(X_{n+1}, X_n) \xrightarrow{d_{n+1}} H_n(X_n, X_{n-1}) \xrightarrow{d_n} H_n(X_n, X_{n-1}) \rightarrow$$

$$d_n \circ d_{n+1} = 0$$

$$H_n(X) = \ker d_n / \operatorname{Im} d_{n+1}$$

$$d_n(g_{\alpha}^n) = \sum n_{\alpha, \beta} g_{\beta}^{n-1}$$

To find $n_{\alpha, \beta}$,

Let $\phi_{\alpha} : \partial D_{\alpha}^n \rightarrow X_{n-1}$ be the gluing map.

Compose with $q : X_{n-1} \rightarrow X_{n-1}/(X_{n-1} \setminus e_{\beta}) \cong S^{n-1}$,

$$q \circ \phi_{\alpha} : S^{n-1} \cong \partial D^{n-1} \rightarrow X_{n-1}/(X_{n-1} \setminus e_{\beta}) \cong S^{n-1}$$

$$n_{\alpha, \beta} = \text{degree}(q \circ \phi_{\alpha})$$

Recall:

$$\phi : S^n \rightarrow S^n$$

Recall:

$$\phi : S^n \rightarrow S^n$$

$$\text{For } y \in S^n, \phi^{-1}(y) = \{x_1, \dots, x_k\}$$

Recall:

$$\phi : S^n \rightarrow S^n$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

Recall:

$$\phi : S^n \rightarrow S^n$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

Recall:

$$\phi : S^n \rightarrow S^n$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

$(\phi_{U_i})_*$ is multiplication by $\text{degree } \phi|_{x_i}$

Recall:


$$\phi : S^n \rightarrow S^n$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

$(\phi|_{U_i})_*$ is multiplication by $\text{degree } \phi|_{x_i}$



Recall:

$$\theta : S^n \rightarrow S^n,$$

$$\phi : S^n \rightarrow S^n$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

$(\phi|_{U_i})_*$ is multiplication by $\text{degree } \phi|_{x_i}$

$H_n(S^n)$

$H_n(S^n, S^n \setminus x_i) \xleftarrow{\sim} H_n(U_i, U_i \setminus x_i)$

$H_n(S^n, S^n \setminus x_j) \xleftarrow{\sim} H_n(U_j, U_j \setminus x_j)$

Recall:

$$\phi : S^n \rightarrow S^n$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

$(\phi|_{U_i})_*$ is multiplication by $\text{degree } \phi|_{x_i}$

$H_n(S^n)$

$H_n(S^n, S^n \setminus x_i) \longleftarrow_{\sim} H_n(U_i, U_i \setminus x_i)$

$H_n(S^n, S^n \setminus x_j) \longleftarrow_{\sim} H_n(U_j, U_j \setminus x_j)$

$$\theta : S^n \rightarrow S^n, \text{ such that } \theta(x_i) = x_j,$$

Recall:
 $\phi : S^n \rightarrow S^n$


$\theta : S^n \rightarrow S^n$, such that $\theta(x_i) = x_j$, $\theta(U_i) = U_j$ and $\theta \sim Id$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$

$(\phi_{U_i})_*$ is multiplication by $\text{degree } \phi|_{x_i}$



Recall:

$\phi : S^n \rightarrow S^n$
 $\theta : S^n \rightarrow S^n$, such that $\theta(x_i) = x_j$, $\theta(U_i) = U_j$ and $\theta \sim Id$


For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

$\theta_* = Id$

degree $\phi = \sum_i \text{degree } \phi|_{x_i}$

$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$

$(\phi_{U_i})_*$ is multiplication by $\text{degree } \phi|_{x_i}$



Recall:
$$\phi : S^n \rightarrow S^n$$

$$\theta : S^n \rightarrow S^n, \text{ such that } \theta(x_i) = x_j, \theta(U_i) = U_j \text{ and } \theta \sim Id$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

$$\text{degree } \phi = \sum_i \text{degree } \phi|_{x_i}$$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

$$(\phi_{U_i})_* \text{ is multiplication by } \text{degree } \phi|_{x_i}$$

$$\begin{array}{ccccc} H_n(S^n) & \longrightarrow & H_n(S^n, S^n \setminus x_i) & \xleftarrow{\sim} & H_n(U_i, U_i \setminus x_i) \\ \downarrow Id & & \downarrow \theta & & \\ H_n(S^n) & \longrightarrow & H_n(S^n, S^n \setminus x_j) & \xleftarrow{\sim} & H_n(U_j, U_j \setminus x_j) \end{array}$$

Recall:
$$\phi : S^n \rightarrow S^n$$

$$\theta : S^n \rightarrow S^n, \text{ such that } \theta(x_i) = x_j, \theta(U_i) = U_j \text{ and } \theta \sim Id$$

For $y \in S^n$, $\phi^{-1}(y) = \{x_1, \dots, x_k\}$

$$\text{degree } \phi = \sum_i \text{degree } \phi|_{x_i}$$

$$\theta_* = Id$$

$$\phi|_{U_i} : H_n(U_i, U_i \setminus x_i) \rightarrow H_n(V, V \setminus y)$$

$$(\phi_{U_i})_* \text{ is multiplication by } \text{degree } \phi|_{x_i}$$

$$\begin{array}{ccccc} H_n(S^n) & \longrightarrow & H_n(S^n, S^n \setminus x_i) & \xleftarrow{\sim} & H_n(U_i, U_i \setminus x_i) \\ \downarrow Id & & \downarrow \theta & & \downarrow \theta|_{U_i} \\ H_n(S^n) & \longrightarrow & H_n(S^n, S^n \setminus x_j) & \xleftarrow{\sim} & H_n(U_j, U_j \setminus x_j) \end{array}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$V:=-1,1)\subset \mathbb{R}$$

$$\tau(t)=-t$$

$$\tau_*:\widetilde{H}_1(V,V\setminus 0)\rightarrow \widetilde{H}_1(V,V\setminus 0)$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \widetilde{H}_1(V, V \setminus 0) \rightarrow \widetilde{H}_1(V, V \setminus 0)$$

What is $\tau_*=$?

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_*=$?

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_*=?$

Solution:

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\downarrow \tau'_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$V := (-1,1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_*=?$

Solution:

$$I := [-1/2,1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I,\partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V,V\setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I,\partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V,V\setminus 0) \end{array}$$

$$\tau'_*(g) = ng$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_*=?$

Solution:

$$I := [-1/2, 1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \end{array}$$

$$\tau'_*(g) = ng \implies i_*(\tau'_*(g)) = ni_*(g)$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \end{array}$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \end{array}$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \end{array}$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_*=?$

Solution:

$$I := [-1/2, 1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \end{array}$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* =?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \overset{i_*}{\longrightarrow} \widetilde{H}_1(V, V \setminus 0)$$

$$\downarrow \tau'_*$$

$$\downarrow \tau_*$$

$$\widetilde{H}_1(I, \partial I) \overset{i_*}{\longrightarrow} \widetilde{H}_1(V, V \setminus 0)$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$\tau'_* = ?$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\downarrow \tau'_*$$

$$\downarrow \tau_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$\tau'_* = ?$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* =?$

Solution:

$$I := [-1/2, 1/2]$$

$$\begin{array}{ccc} \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \\ \downarrow \tau'_* & & \downarrow \tau_* \\ \widetilde{H}_1(I, \partial I) & \xrightarrow{i_*} & \widetilde{H}_1(V, V \setminus 0) \end{array}$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$\tau'_* =?$$

$$\widetilde{H}_1(I, \partial I) \overset{\partial}{\longrightarrow} \widetilde{H}_0(\partial I)$$

$$\widetilde{H}_1(I, \partial I) \overset{\partial}{\longrightarrow} \widetilde{H}_0(\partial I)$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* =?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\downarrow \tau'_* \qquad \qquad \qquad \downarrow \tau_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$\tau'_* =?$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\downarrow \tau'_* \qquad \qquad \qquad \downarrow \tau'_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* = ?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\downarrow \tau'_*$$

$$\downarrow \tau_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$\tau'_* = ?$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\downarrow \tau'_*$$

$$\downarrow \tau'_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\tau'_* = -Id$$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

$$V := (-1, 1) \subset \mathbb{R}$$

$$\tau(t) = -t$$

$$\tau_* : \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}} \rightarrow \underbrace{\widetilde{H}_1(V, V \setminus 0)}_{\mathbb{Z}}$$

What is $\tau_* =?$

Solution:

$$I := [-1/2, 1/2]$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\downarrow \tau'_*$$

$$\downarrow \tau_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{i_*} \widetilde{H}_1(V, V \setminus 0)$$

$$\tau' : (I, \partial I) \rightarrow (I, \partial I)$$

$$\tau'(t) = -t$$

$$\tau' : \widetilde{H}_1(I, \partial I) \rightarrow \widetilde{H}_1(I, \partial I)$$

$$\tau'_* = ?$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\downarrow \tau'_*$$

$$\downarrow \tau'_*$$

$$\widetilde{H}_1(I, \partial I) \xrightarrow{\partial} \widetilde{H}_0(\partial I)$$

$$\tau'_* = -Id$$

Therefore, $\tau_* = -Id$

$$\begin{aligned} \tau'_*(g) = ng &\implies i_*(\tau'_*(g)) = ni_*(g) \\ &\implies \tau_*(i_*(g)) = n(i_*(g)) \end{aligned}$$

Example: S^n

Example: S^n

0 cells 1

Example: S^n

0 cells $\rightarrow H_0(X_0) = \mathbb{Z}$

Example: S^n

0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

n cells 1

Example: S^n

0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

n cells 1 ($H_n(X_n, X_{n-1}) = \mathbb{Z}$)

Example: S^n

0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

n cells 1 ($H_n(X_n, X_{n-1}) = \mathbb{Z}$)

$$H_n(X_{n+1}, X_n) \longrightarrow H_n(X_n, X_{n-1}) \longrightarrow H_n(X_{n-1}, X_{n-2}) \longrightarrow \dots \longrightarrow H_0(X_0)$$

Example: S^n

0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

n cells 1 ($H_n(X_n, X_{n-1}) = \mathbb{Z}$)

$$H_n(X_{n+1}, X_n) \longrightarrow H_n(X_n, X_{n-1}) \longrightarrow H_n(X_{n-1}, X_{n-2}) \longrightarrow \dots \longrightarrow H_0(X_0)$$

$$0 \longrightarrow \mathbb{Z} \longrightarrow 0 \longrightarrow \dots \longrightarrow \mathbb{Z}$$

Example: S^n

0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

n cells 1 ($H_n(X_n, X_{n-1}) = \mathbb{Z}$)

$$H_n(X_{n+1}, X_n) \longrightarrow H_n(X_n, X_{n-1}) \longrightarrow H_n(X_{n-1}, X_{n-2}) \longrightarrow \dots \longrightarrow H_0(X_0)$$

$$0 \longrightarrow \mathbb{Z} \longrightarrow 0 \longrightarrow \dots \longrightarrow \mathbb{Z}$$

$$H_k(S^n) = \begin{cases} \mathbb{Z} & k = 0, n \end{cases}$$

Example: S^n

0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

n cells 1 ($H_n(X_n, X_{n-1}) = \mathbb{Z}$)

$$H_n(X_{n+1}, X_n) \longrightarrow H_n(X_n, X_{n-1}) \longrightarrow H_n(X_{n-1}, X_{n-2}) \longrightarrow \dots \longrightarrow H_0(X_0)$$

$$0 \longrightarrow \mathbb{Z} \longrightarrow 0 \longrightarrow \dots \longrightarrow \mathbb{Z}$$


$$H_k(S^n) = \begin{cases} \mathbb{Z} & k = 0, n \\ 0 & \text{otherwise} \end{cases}$$

Example: \mathbb{RP}^2


Example: \mathbb{RP}^2

•


Example: \mathbb{RP}^2




Example: \mathbb{RP}^2



Example: \mathbb{RP}^2




Example: \mathbb{RP}^2




0 cells 1

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1


Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

Example: \mathbb{RP}^2




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1

Example: \mathbb{RP}^2




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

Example: \mathbb{RP}^2




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \end{cases}$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \end{cases}$$

Example: \mathbb{RP}^2



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 1 ($H_1(X_1, X_0) = \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & \text{otherwise} \end{cases}$$

Example: $S^1 \times S^1$


Example: $S^1 \times S^1$

-


Example: $S^1 \times S^1$




Example: $S^1 \times S^1$




Example: $S^1 \times S^1$




Example: $S^1 \times S^1$




Example: $S^1 \times S^1$




Example: $S^1 \times S^1$



Example: $S^1 \times S^1$




Example: $S^1 \times S^1$




0 cells **1**

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2


Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

Example: $S^1 \times S^1$




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1

Example: $S^1 \times S^1$




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

Example: $S^1 \times S^1$




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \longrightarrow \mathbb{Z}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \longrightarrow \mathbb{Z}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{0} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{0} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \end{cases}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{0} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z} \oplus \mathbb{Z} & k = 1 \end{cases}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{0} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z} \oplus \mathbb{Z} & k = 1 \\ \mathbb{Z} & k = 2 \end{cases}$$

Example: $S^1 \times S^1$



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{0} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z} \oplus \mathbb{Z} & k = 1 \\ \mathbb{Z} & k = 2 \\ 0 & \text{otherwise} \end{cases}$$

Example: Klein bottle


Example: Klein bottle

-


Example: Klein bottle




Example: Klein bottle




Example: Klein bottle




Example: Klein bottle



Example: Klein bottle




Example: Klein bottle




0 cells 1

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2


Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

Example: Klein bottle




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1

Example: Klein bottle




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

Example: Klein bottle




0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \longrightarrow \mathbb{Z}$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \longrightarrow \mathbb{Z}$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)


1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{1 \rightarrow (0,2)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{1 \rightarrow (0,2)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \end{cases}$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)


2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{1 \rightarrow (0,2)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z} \oplus \mathbb{Z}/2 & k = 1 \end{cases}$$

Example: Klein bottle



0 cells 1 ($H_0(X_0) = \mathbb{Z}$)

1 cells 2 ($H_1(X_1, X_0) = \mathbb{Z} \oplus \mathbb{Z}$)

2 cells 1 ($H_2(X_2, X_1) = \mathbb{Z}$)

$$H_2(X_2, X_1) \longrightarrow H_1(X_1, X_0) \longrightarrow H_0(X_0)$$

$$\mathbb{Z} \xrightarrow{1 \rightarrow (0,2)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{0} \mathbb{Z}$$

$$H_k(S^1 \times S^1) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z} \oplus \mathbb{Z}/2 & k = 1 \\ 0 & \text{otherwise} \end{cases}$$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
rank $G = k$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , *finitely generated, abelian*

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
rank $G = k$

Theorem. A, B, C , *finitely generated, abelian*
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ *short exact*

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
rank $G = k$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, rank $B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$ □

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
rank $G = k$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, rank $B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free □

G abelian, $G \cong \bigoplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \bigoplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \bigoplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator
maps to a different copy of \mathbb{Z} . \square

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , *finitely generated, abelian*
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ *short exact*
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , *finitely generated, abelian*
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ *short exact*
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$C_n := H_n(X_n, X_{n-1})$ (number of cells of dimension n)

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$C_n := H_n(X_n, X_{n-1})$ (number of cells of dimension n)

$\text{rank } H_n(X) = \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1}$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , *finitely generated, abelian*
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ *short exact*
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$C_n := H_n(X_n, X_{n-1})$ (number of cells of dimension n)

$\text{rank } H_n(X) = \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1}$
 $\text{rank } \ker d_n = \text{rank } C_n - \text{rank } \text{Im } d_n$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , *finitely generated, abelian*
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ *short exact*
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$C_n := H_n(X_n, X_{n-1})$ (number of cells of dimension n)

$\text{rank } H_n(X) = \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1}$
 $\text{rank } \ker d_n = \text{rank } C_n - \text{rank } \text{Im } d_n$
 $\text{rank } H_n(X) = \text{rank } C_n - \text{rank } \text{Im } d_n - \text{rank } \text{Im } d_{n+1}$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

$$\text{rank } H_0(X) = \text{rank } C_0 - \text{rank } Im \, d_0 - \text{rank } Im \, d_1$$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \overset{i}{\rightarrow} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$$C_n := H_n(X_n, X_{n-1}) \text{ (number of cells of dimension } n\text{)}$$

$$\begin{aligned} \text{rank } H_n(X) &= \text{rank } \ker d_n - \text{rank } Im \, d_{n+1} \\ \text{rank } \ker d_n &= \text{rank } C_n - \text{rank } Im \, d_n \\ \text{rank } H_n(X) &= \text{rank } C_n - \text{rank } Im \, d_n - \text{rank } Im \, d_{n+1} \end{aligned}$$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

$$\begin{aligned} \text{rank } H_0(X) &= \text{rank } C_0 - \text{rank } \text{Im } d_0 - \text{rank } \text{Im } d_1 \\ \text{rank } H_1(X) &= \text{rank } C_1 - \text{rank } \text{Im } d_1 - \text{rank } \text{Im } d_2 \end{aligned}$$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$$C_n := H_n(X_n, X_{n-1}) \text{ (number of cells of dimension } n\text{)}$$

$$\begin{aligned} \text{rank } H_n(X) &= \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1} \\ \text{rank } \ker d_n &= \text{rank } C_n - \text{rank } \text{Im } d_n \\ \text{rank } H_n(X) &= \text{rank } C_n - \text{rank } \text{Im } d_n - \text{rank } \text{Im } d_{n+1} \end{aligned}$$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

$$\begin{aligned} \text{rank } H_0(X) &= \text{rank } C_0 - \text{rank } \text{Im } d_0 - \text{rank } \text{Im } d_1 \\ \text{rank } H_1(X) &= \text{rank } C_1 - \text{rank } \text{Im } d_1 - \text{rank } \text{Im } d_2 \\ \text{rank } H_2(X) &= \text{rank } C_2 - \text{rank } \text{Im } d_2 - \text{rank } \text{Im } d_3 \end{aligned}$$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ □

$$C_n := H_n(X_n, X_{n-1}) \text{ (number of cells of dimension } n)$$

$$\begin{aligned} \text{rank } H_n(X) &= \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1} \\ \text{rank } \ker d_n &= \text{rank } C_n - \text{rank } \text{Im } d_n \\ \text{rank } H_n(X) &= \text{rank } C_n - \text{rank } \text{Im } d_n - \text{rank } \text{Im } d_{n+1} \end{aligned}$$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$C_n := H_n(X_n, X_{n-1})$ (number of cells of dimension n)

$\text{rank } H_n(X) = \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1}$
 $\text{rank } \ker d_n = \text{rank } C_n - \text{rank } \text{Im } d_n$
 $\text{rank } H_n(X) = \text{rank } C_n - \text{rank } \text{Im } d_n - \text{rank } \text{Im } d_{n+1}$

$\text{rank } H_0(X) = \text{rank } C_0 - \text{rank } \text{Im } d_0 - \text{rank } \text{Im } d_1$
 $\text{rank } H_1(X) = \text{rank } C_1 - \text{rank } \text{Im } d_1 - \text{rank } \text{Im } d_2$
 $\text{rank } H_2(X) = \text{rank } C_2 - \text{rank } \text{Im } d_2 - \text{rank } \text{Im } d_3$
 \dots
 $\Sigma(-1)^n \text{rank } H_n(X) = \Sigma(-1)^n \text{rank } C_n$

G abelian, $G \cong \oplus_{i=1}^k \mathbb{Z} \oplus \text{torsion part}$
 $\text{rank } G = k$

Theorem. A, B, C , finitely generated, abelian
 $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ short exact
then, $\text{rank } B = \text{rank } A + \text{rank } C$

Proof. $0 \rightarrow \oplus_{i=1}^m \mathbb{Z} \oplus T \xrightarrow{i} \oplus_{i=1}^n \mathbb{Z} \oplus T' \rightarrow F \oplus T'' \rightarrow 0$

Injectivity of $i \implies$ free maps to free, each generator maps to a different copy of \mathbb{Z} .

Those copies will become torsion in $B/i(A)$ \square

$C_n := H_n(X_n, X_{n-1})$ (number of cells of dimension n)

$\text{rank } H_n(X) = \text{rank } \ker d_n - \text{rank } \text{Im } d_{n+1}$
 $\text{rank } \ker d_n = \text{rank } C_n - \text{rank } \text{Im } d_n$
 $\text{rank } H_n(X) = \text{rank } C_n - \text{rank } \text{Im } d_n - \text{rank } \text{Im } d_{n+1}$

$\text{rank } H_0(X) = \text{rank } C_0 - \text{rank } \text{Im } d_0 - \text{rank } \text{Im } d_1$
 $\text{rank } H_1(X) = \text{rank } C_1 - \text{rank } \text{Im } d_1 - \text{rank } \text{Im } d_2$
 $\text{rank } H_2(X) = \text{rank } C_2 - \text{rank } \text{Im } d_2 - \text{rank } \text{Im } d_3$
 \dots

$\chi(X) = \sum (-1)^n \text{rank } H_n(X) = \sum (-1)^n \text{rank } C_n$
(Euler characteristic)

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$H_2(x_0 \times S^1) \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$\underbrace{H_2(x_0 \times S^1)}_0 \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$\underbrace{H_2(x_0 \times S^1)}_0 \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

$$r : S^1 \times S^1 \rightarrow x_0 \times S^1, r(x, y) = (x_0, y)$$

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$\underbrace{H_2(x_0 \times S^1)}_0 \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

$$r : S^1 \times S^1 \rightarrow x_0 \times S^1, r(x, y) = (x_0, y)$$

$$r \circ i = Id$$

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$\underbrace{H_2(x_0 \times S^1)}_0 \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

$$r : S^1 \times S^1 \rightarrow x_0 \times S^1, r(x, y) = (x_0, y)$$

$$r \circ i = Id$$

$$r_* \circ i_* = Id_*$$

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$\underbrace{H_2(x_0 \times S^1)}_0 \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

$$r : S^1 \times S^1 \rightarrow x_0 \times S^1, r(x, y) = (x_0, y)$$

$$r \circ i = Id$$

$$r_* \circ i_* = Id_*$$

$$i_* \text{ injective}$$

Prove that the quotient map $q : S^1 \times S^1 \rightarrow (S^1 \times S^1)/(x_0 \times S^1)$, where $x_0 \in S^1$, induces an isomorphism $q_* : H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1))$.

$$\underbrace{H_2(x_0 \times S^1)}_0 \rightarrow H_2(S^1 \times S^1) \rightarrow H_2((S^1 \times S^1)/(x_0 \times S^1)) \rightarrow H_1(x_0 \times S^1) \xrightarrow{i_*} H_1(S^1 \times S^1)$$

$$r : S^1 \times S^1 \rightarrow x_0 \times S^1, r(x, y) = (x_0, y)$$

$$r \circ i = Id$$

$$r_* \circ i_* = Id_*$$

$$i_* \text{ injective}$$

$$q_* \text{ is an isomorphism}$$

Let p_1 , p_2 , and p_3 denote 3 distinct points chosen on S^2 .

1. Compute $H_i(S^2 \setminus \{p_1, p_2, p_3\})$ for all $i \geq 0$.

Let p_1 , p_2 , and p_3 denote 3 distinct points chosen on S^2 .

1. Compute $H_i(S^2 \setminus \{p_1, p_2, p_3\})$ for all $i \geq 0$.
2. Compute $H_i(S^2 / \{p_1, p_2, p_3\})$ for all $i \geq 0$.

Let p_1 , p_2 , and p_3 denote 3 distinct points chosen on S^2 .

1. Compute $H_i(S^2 \setminus \{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $S^2 \setminus \{p_1, p_2, p_3\} \cong \mathbb{R}^2 \setminus \text{two points}$, which deformation retracts to the wedge of two circles...

2. Compute $H_i(S^2/\{p_1, p_2, p_3\})$ for all $i \geq 0$.

Let p_1 , p_2 , and p_3 denote 3 distinct points chosen on S^2 .

1. Compute $H_i(S^2 \setminus \{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $S^2 \setminus \{p_1, p_2, p_3\} \cong \mathbb{R}^2 \setminus \text{two points}$, which deformation retracts to the wedge of two circles...

2. Compute $H_i(S^2/\{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $\tilde{H}_i(S^2) \xrightarrow{\sim} \tilde{H}_i(S^2/\{p_1, p_2, p_3\})$

Let p_1 , p_2 , and p_3 denote 3 distinct points chosen on S^2 .

1. Compute $H_i(S^2 \setminus \{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $S^2 \setminus \{p_1, p_2, p_3\} \cong \mathbb{R}^2 \setminus \text{two points}$, which deformation retracts to the wedge of two circles...

2. Compute $H_i(S^2/\{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $\tilde{H}_i(S^2) \xrightarrow{\sim} \tilde{H}_i(S^2/\{p_1, p_2, p_3\})$, if $i > 1$

$$0 \rightarrow \tilde{H}_1(S^2) \rightarrow H_1(S^2, \{p_1, p_2, p_3\}) \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow 0$$

Let p_1 , p_2 , and p_3 denote 3 distinct points chosen on S^2 .

1. Compute $H_i(S^2 \setminus \{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $S^2 \setminus \{p_1, p_2, p_3\} \cong \mathbb{R}^2 \setminus \text{two points}$, which deformation retracts to the wedge of two circles...

2. Compute $H_i(S^2/\{p_1, p_2, p_3\})$ for all $i \geq 0$.

Solution: $\tilde{H}_i(S^2) \xrightarrow{\sim} \tilde{H}_i(S^2/\{p_1, p_2, p_3\})$, if $i > 1$

$$0 \rightarrow \underbrace{\tilde{H}_1(S^2)}_0 \rightarrow H_1(S^2, \{p_1, p_2, p_3\}) \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow 0$$

Let $r : S^1 \times S^1 \rightarrow S^1$ denote the map $r(s, t) = s$. Prove that $r_* : H_1(S^1 \times S^1) \cong \mathbb{Z} \oplus \mathbb{Z} \rightarrow H_1(S^1) \cong \mathbb{Z}$ is the projection onto the first factor.