

Mayer-Vietoris sequence

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n^{\{A,B\}}(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Observation:

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Observation:

If $A \cap B$ is path connected,

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Observation:

If $A \cap B$ is path connected,

$$\cdots \rightarrow \tilde{H}_1(A \cap B) \xrightarrow{i_*} \tilde{H}_1(A) \oplus \tilde{H}_1(B) \xrightarrow{j_*} \tilde{H}_1(X) \xrightarrow{\partial} 0$$

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Observation:

If $A \cap B$ is path connected,

$$\cdots \rightarrow \tilde{H}_1(A \cap B) \xrightarrow{i_*} \tilde{H}_1(A) \oplus \tilde{H}_1(B) \xrightarrow{j_*} \tilde{H}_1(X) \xrightarrow{\partial} 0$$

$$H_1(X) \cong (H_1(A) \oplus H_1(B)) / \text{Im } i_*$$

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Observation:

If $A \cap B$ is path connected,

$$\cdots \rightarrow \tilde{H}_1(A \cap B) \xrightarrow{i_*} \tilde{H}_1(A) \oplus \tilde{H}_1(B) \xrightarrow{j_*} \tilde{H}_1(X) \xrightarrow{\partial} 0$$

$$H_1(X) \cong (H_1(A) \oplus H_1(B)) / \text{Im } i_*$$

$$c \xrightarrow{i} (c, -c)$$

Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Reduced Mayer-Vietoris sequence

If $X = \text{Int } A \cup \text{Int } B$,

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact (Exercise!)

Observation:

If $A \cap B$ is path connected,

$$\cdots \rightarrow \tilde{H}_1(A \cap B) \xrightarrow{i_*} \tilde{H}_1(A) \oplus \tilde{H}_1(B) \xrightarrow{j_*} \tilde{H}_1(X) \xrightarrow{\partial} 0$$

$$H_1(X) \cong (H_1(A) \oplus H_1(B)) / \text{Im } i_*$$

$$c \xrightarrow{i} (c, -c)$$

Recall Van-Kampen!

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$), □

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0)$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) \quad \square$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\tilde{H}_0(S^2 \setminus f(S^1))$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\tilde{H}_0(U \cap V)$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, □

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

□

Jordan curve theorem



Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V)$$

Jordan curve theorem



Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V)$$

Jordan curve theorem



Theorem. $f : S^1 \rightarrow S^2$ continuous and homeomorphism onto its image, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$ ($I_1 \cong I_2 \cong [0, 1]$),
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p)$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2,$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism}$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$): for all j

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-**
set $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset \dots$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$

<i>Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):</i>	non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$
$I = I_1 \cup I_2,$	for all j
$I_1 \cap I_2 = \{p\}$	
$U = S^2 \setminus f(I_1)$	But $\tilde{H}_i(S^2 \setminus f(q)) = 0$
$V = S^2 \setminus f(I_2)$	Therefore, $x = \partial y,$
$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2,$ so $(\tilde{H}_i(U \cup V) = 0)$	
$U \cap V = S^2 \setminus f(I)$	
$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism	
$[x] \rightarrow ([i_U(x)], [-i_V(x)])$	
$[x] = [0]$ if and only if $[i_U(x)] = 0$ AND $[i_V(x)] = 0$	
$[x] \neq [0] \implies [i_U(x)] \neq 0$ OR $[i_V(x)] \neq 0$	
i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in super-	
set $S^2 \setminus f(I_i)$ for $i = 1$ or 2	
Applying repeatedly:	
$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$	
$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$	
$\dots \supset S^2 \setminus f(q)$	

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-set** $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$ for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

Therefore, $x = \partial y, y \in C_n(S^2 \setminus q)$

<i>Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):</i>	non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$
$I = I_1 \cup I_2,$	for all j
$I_1 \cap I_2 = \{p\}$	
$U = S^2 \setminus f(I_1)$	But $\tilde{H}_i(S^2 \setminus f(q)) = 0$
$V = S^2 \setminus f(I_2)$	Therefore, $x = \partial y, y \in C_n(S^2 \setminus q)$
$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2,$ so $(\tilde{H}_i(U \cup V) = 0)$	
$U \cap V = S^2 \setminus f(I)$	$y = \sum_k \sigma_k$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism
 $[x] \rightarrow ([i_U(x)], [-i_V(x)])$
 $[x] = [0]$ if and only if $[i_U(x)] = 0$ AND $[i_V(x)] = 0$
 $[x] \neq [0] \implies [i_U(x)] \neq 0$ OR $[i_V(x)] \neq 0$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-**
set $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$\begin{aligned}
 I_i &= I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\} \\
 S^2 \setminus f(I_i) &= S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset \\
 &\dots \supset S^2 \setminus f(q)
 \end{aligned}$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-set** $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$
for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\sigma_k(\Delta_{i+1}) \subset S^2 \setminus q$$

<i>Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):</i>	non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$
$I = I_1 \cup I_2,$	for all j
$I_1 \cap I_2 = \{p\}$	But $\tilde{H}_i(S^2 \setminus f(q)) = 0$
$U = S^2 \setminus f(I_1)$	Therefore, $x = \partial y, y \in C_n(S^2 \setminus q)$
$V = S^2 \setminus f(I_2)$	$y = \sum_k \sigma_k$
$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2,$ so $(\tilde{H}_i(U \cup V) = 0)$	$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q$
$U \cap V = S^2 \setminus f(I)$	

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism
 $[x] \rightarrow ([i_U(x)], [-i_V(x)])$
 $[x] = [0]$ if and only if $[i_U(x)] = 0$ AND $[i_V(x)] = 0$
 $[x] \neq [0] \implies [i_U(x)] \neq 0$ OR $[i_V(x)] \neq 0$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-**
set $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$\begin{aligned}
 I_i &= I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\} \\
 S^2 \setminus f(I_i) &= S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset \\
 &\dots \supset S^2 \setminus f(q)
 \end{aligned}$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-set** $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$ for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q \cup S^2 \setminus f(I_{i_j})$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-**
set $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$
for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q \cup S^2 \setminus f(I_{i_j})$$

$$\sigma_k(\Delta_{i+1}) \text{ is compact}$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-set** $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$ for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q \cup S^2 \setminus f(I_{i_j})$$

$$\sigma_k(\Delta_{i+1}) \text{ is compact}$$

$$y \in C_n(S^2 \setminus f(I_{i_j})) \text{ for some } j.$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism}$$

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-**
set $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$
for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q \cup S^2 \setminus f(I_{i_j})$$

$$\sigma_k(\Delta_{i+1}) \text{ is compact}$$

$$y \in C_n(S^2 \setminus f(I_{i_j})) \text{ for some } j. \text{ Therefore, } [x] = [0] \in H_n(S^2 \setminus I_{i_j}),$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism}$$

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-set** $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$ for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q \cup S^2 \setminus f(I_{i_j})$$

$$\sigma_k(\Delta_{i+1}) \text{ is compact}$$

$$y \in C_n(S^2 \setminus f(I_{i_j})) \text{ for some } j. \text{ Therefore, } [x] = [0] \in H_n(S^2 \setminus I_{i_j}), \text{ contradicting } [x] \neq [0]$$

Proof of claim (i.e. $H_i(S^2 \setminus f([0, 1])) = 0$):

$$I = I_1 \cup I_2,$$

$$I_1 \cap I_2 = \{p\}$$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0)$$

$$U \cap V = S^2 \setminus f(I)$$

$$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism}$$

$$[x] \rightarrow ([i_U(x)], [-i_V(x)])$$

$$[x] = [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0$$

$$[x] \neq [0] \implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0$$

i.e. non-trivial in $S^2 \setminus f(I)$ implies non-trivial in **super-set** $S^2 \setminus f(I_i)$ for $i = 1$ or 2

Applying repeatedly:

$$I_i = I_{i_1} \supset I_{i_2} \supset I_{i_3} \supset \dots \subset \{q\}$$

$$S^2 \setminus f(I_i) = S^2 \setminus f(I_{i_1}) \subset S^2 \setminus f(I_{i_2}) \subset S^2 \setminus f(I_{i_3}) \subset$$

$$\dots \supset S^2 \setminus f(q)$$

non-trivial in $S^2 \setminus f(I)$ implies non-trivial in $S^2 \setminus f(I_{i_j})$ for all j

$$\text{But } \tilde{H}_i(S^2 \setminus f(q)) = 0$$

$$\text{Therefore, } x = \partial y, y \in C_n(S^2 \setminus q)$$

$$y = \sum_k \sigma_k$$

$$\cup \sigma_k(\Delta_{i+1}) \subset S^2 \setminus q \cup S^2 \setminus f(I_{i_j})$$

$$\sigma_k(\Delta_{i+1}) \text{ is compact}$$

$$y \in C_n(S^2 \setminus f(I_{i_j}) \text{ for some } j. \text{ Therefore, } [x] = [0] \in H_n(S^2 \setminus I_{i_j}), \text{ contradicting } [x] \neq [0]$$

Lemma (Five lemma). $A \longrightarrow B \longrightarrow C \longrightarrow D \longrightarrow E$

$$A' \longrightarrow B' \longrightarrow C' \longrightarrow D' \longrightarrow E'$$

rows exact

Lemma (Five lemma).
$$\begin{array}{ccccccccc} A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & D & \longrightarrow & E \\ \downarrow f & & \downarrow g & & \downarrow h & & \downarrow i & & \downarrow j \\ A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & D' & \longrightarrow & E' \end{array}$$

rows exact

Lemma (Five lemma).
$$\begin{array}{ccccccccc} A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & D & \longrightarrow & E \\ \downarrow f & & \downarrow g & & \downarrow h & & \downarrow i & & \downarrow j \\ A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & D' & \longrightarrow & E' \end{array}$$

rows exact and f, g, i, j isomorphisms, diagram commutes $\implies h$ isomorphism

Proof. Exercise!



Lemma. $0 \rightarrow A_* \rightarrow B_* \rightarrow C_*$ an exact sequence of chain complexes,

Lemma. $0 \rightarrow A_* \rightarrow B_* \rightarrow C_*$ an exact sequence of chain complexes,
 $0 \rightarrow A'_* \rightarrow B'_* \rightarrow C'_*$ an exact sequence of chain complexes,

Lemma. $0 \rightarrow A_* \rightarrow B_* \rightarrow C_*$ an exact sequence of chain complexes,

$0 \rightarrow A'_* \rightarrow B'_* \rightarrow C'_*$ an exact sequence of chain complexes, following diagram commutes,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & A_i & \longrightarrow & B_i & \longrightarrow & C_i \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & A'_i & \longrightarrow & B'_i & \longrightarrow & C'_i \longrightarrow 0
 \end{array}
 \quad \text{then}$$

Lemma. $0 \rightarrow A_* \rightarrow B_* \rightarrow C_*$ an exact sequence of chain complexes,

$0 \rightarrow A'_* \rightarrow B'_* \rightarrow C'_*$ an exact sequence of chain complexes, following diagram commutes,

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_i & \longrightarrow & B_i & \longrightarrow & C_i \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A'_i & \longrightarrow & B'_i & \longrightarrow & C'_i \longrightarrow 0 \end{array} \quad \text{then}$$

$$\begin{array}{cccccccc} \cdots & \longrightarrow & H_i(A) & \longrightarrow & H_i(B) & \longrightarrow & H_i(C) & \longrightarrow & H_{i-1}(A) & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & H_i(A') & \longrightarrow & H_i(B') & \longrightarrow & H_i(C') & \longrightarrow & H_{i-1}(A) & \longrightarrow & \cdots \end{array}$$

also commutes

Proof. Exercise!

□

If $X = A \cup B$,

If $X = A \cup B$, such that there exist open $U, V \subset X$ with $A \subset U$ and $B \subset V$

If $X = A \cup B$, such that there exist open $U, V \subset X$ with $A \subset U$ and $B \subset V$, and U deformation retracts onto A , V deformation retracts onto B ,

If $X = A \cup B$, such that there exist open $U, V \subset X$ with $A \subset U$ and $B \subset V$,
and U deformation retracts onto A , V deformation retracts onto B ,
and $U \cap V$ deformation retracts onto $A \cap B$

If $X = A \cup B$, such that there exist open $U, V \subset X$ with $A \subset U$ and $B \subset V$,
and U deformation retracts onto A , V deformation retracts onto B ,
and $U \cap V$ deformation retracts onto $A \cap B$, then

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

If $X = A \cup B$, such that there exist open $U, V \subset X$ with $A \subset U$ and $B \subset V$,
and U deformation retracts onto A , V deformation retracts onto B ,
and $U \cap V$ deformation retracts onto $A \cap B$, then

$$\cdots \rightarrow H_{n+1}(X) \xrightarrow{\partial} H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

and

$$\cdots \rightarrow \tilde{H}_{n+1}(X) \xrightarrow{\partial} \tilde{H}_n(A \cap B) \xrightarrow{i_*} \tilde{H}_n(A) \oplus \tilde{H}_n(B) \xrightarrow{j_*} \tilde{H}_n(X) \xrightarrow{\partial} \tilde{H}_{n-1}(A \cap B) \rightarrow \cdots$$

is exact

Proof. Exercise!

□