

$$0 \rightarrow A \rightarrow 0 \implies A \cong 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$0 \rightarrow A \rightarrow B \rightarrow ?? \rightarrow 0 \implies ?? \cong B/i(A)$$

$$0 \rightarrow ?? \rightarrow B \xrightarrow{f} C \rightarrow 0 \implies ?? \cong \operatorname{Ker} f$$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C)$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

b



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b)$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$s(q(b))$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$b - s(q(b))$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b)))$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b))$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b)) = i(a) + s(c), \text{ where } c := q(b)$$

□

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$



Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b)) = i(a) + s(c), \text{ where } c := q(b)$$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

$$i(a) = s(c)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b)) = i(a) + s(c), \text{ where } c := q(b)$$



$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} i(a) &= s(c) \\ \implies q(i(a)) &= q(s(c)) \end{aligned}$$

□

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

$$\text{Then, } B \cong i(A) \oplus s(C) \cong A \oplus C$$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b)) = i(a) + s(c), \text{ where } c := q(b)$$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} & i(a) = s(c) \\ \implies & q(i(a)) = q(s(c)) \\ \implies & 0 = c \end{aligned}$$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$
If there exists, $s : C \rightarrow B$, such that
 $q \circ s = Id$,
Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

□

Proof. Consider $b \in B$
 $q(b - s(q(b))) = 0$

Therefore,
 $b - s(q(b)) = i(a)$ for some $a \in A$
 $b = i(a) + s(q(b)) = i(a) + s(c)$, where $c := q(b)$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} & i(a) = s(c) \\ \implies & q(i(a)) = q(s(c)) \\ \implies & 0 = c \\ \implies & 0 = s(c) \end{aligned}$$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b)) = i(a) + s(c), \text{ where } c := q(b)$$

□

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} & i(a) = s(c) \\ \implies & q(i(a)) = q(s(c)) \\ \implies & 0 = c \\ \implies & 0 = s(c) = i(a) \end{aligned}$$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that

$$q \circ s = Id,$$

$$\text{Then, } B \cong i(A) \oplus s(C) \cong A \oplus C$$

Proof. Consider $b \in B$

$$q(b - s(q(b))) = 0$$

Therefore,

$$b - s(q(b)) = i(a) \text{ for some } a \in A$$

$$b = i(a) + s(q(b)) = i(a) + s(c), \text{ where } c := q(b)$$

□

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} & i(a) = s(c) \\ \implies & q(i(a)) = q(s(c)) \\ \implies & 0 = c \\ \implies & 0 = s(c) = i(a) \end{aligned}$$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$

If there exists, $s : C \rightarrow B$, such that
 $q \circ s = Id$,

Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Therefore,
 $i(A) \cap s(C) = 0$

□

Proof. Consider $b \in B$
 $q(b - s(q(b))) = 0$

Therefore,
 $b - s(q(b)) = i(a)$ for some $a \in A$
 $b = i(a) + s(q(b)) = i(a) + s(c)$, where $c := q(b)$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} & i(a) = s(c) \\ \implies & q(i(a)) = q(s(c)) \\ \implies & 0 = c \\ \implies & 0 = s(c) = i(a) \end{aligned}$$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$
If there exists, $s : C \rightarrow B$, such that
 $q \circ s = Id$ (the exact sequence splits),
Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Therefore,
 $i(A) \cap s(C) = 0$

□

Proof. Consider $b \in B$
 $q(b - s(q(b))) = 0$

Therefore,
 $b - s(q(b)) = i(a)$ for some $a \in A$
 $b = i(a) + s(q(b)) = i(a) + s(c)$, where $c := q(b)$

$$0 \rightarrow A \rightarrow ?? \rightarrow C \rightarrow 0$$

$$B = i(A) + s(B)$$

Example. $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$
 $\mathbb{Z} \not\cong \mathbb{Z} \oplus \mathbb{Z}/2$

$$\begin{aligned} & i(a) = s(c) \\ \implies & q(i(a)) = q(s(c)) \\ \implies & 0 = c \\ \implies & 0 = s(c) = i(a) \end{aligned}$$

Lemma. $0 \rightarrow A \xrightarrow{i} B \xrightarrow{q} C \rightarrow 0$
If there exists, $s : C \rightarrow B$, such that
 $q \circ s = Id$ *(the exact sequence splits),*
Then, $B \cong i(A) \oplus s(C) \cong A \oplus C$

Therefore,
 $i(A) \cap s(C) = 0$

□

Proof. Consider $b \in B$
 $q(b - s(q(b))) = 0$

Therefore,
 $b - s(q(b)) = i(a)$ for some $a \in A$
 $b = i(a) + s(q(b)) = i(a) + s(c)$, where $c := q(b)$

Exercise. *If $r : X \rightarrow A$ is a retract,*
 $H_i(X) \cong H_i(A) \oplus H_i(X, A)$

$$X = S^1 \times S^1$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\})$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$U := S^1 \times (S^1 \setminus x_0)$ deformation retracts onto A

$V := S^1 \times (S^1 \setminus x_1)$ deformation retracts onto B

$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\})$ deformation retracts onto

$A \cap B$

$i_p : S^1 \hookrightarrow S^1 \times I_1$, where $i_p(s) = (s, p)$

$i_q : S^1 \hookrightarrow S^1 \times I_1$, where $i_q(s) = (s, q)$

$i_{p*}(1) = i_{q*}(1)$ because i_p is homotopic to i_q

$y_A := i_{p*}(1)(= i_{q*}(1))$ generates $H_1(A) = \mathbb{Z}$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}$$

$$j_p : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}$$

$$j_p : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p)$$

$$j_q : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q)$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}$$

$$j_p : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p)$$

$$j_q : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q)$$

$$x_p = j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}$$

$$j_p : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p)$$

$$j_q : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q)$$

$$x_p = j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A$$

$$k_A \circ j_p = i_p \text{ and } k_A \circ j_q = i_q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}$$

$$j_p : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p)$$

$$j_q : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q)$$

$$x_p = j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}$$

$$X = S^1 \times S^1$$

$$A = S^1 \times I_1$$

$$B = S^1 \times I_2$$

$$A \cup B = X$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q$$

$$U := S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A$$

$$V := S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B$$

$$V \cap V = S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto}$$

$$A \cap B$$

$$i_p : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p)$$

$$i_q : S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q)$$

$$i_{p*}(1) = i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q$$

$$y_A := i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}$$

$$j_p : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p)$$

$$j_q : S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q)$$

$$x_p = j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}$$

$$A \cap B = S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A$$

$$k_A \circ j_p = i_p \text{ and } k_A \circ j_q = i_q$$

$$k_{A*} \circ j_{p*} = i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p = j_{p*}(1) &\text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p &= j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$k_{A*}(x_p) = y_A = k_{A*}(x_q)$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p &= j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
&\text{Similarly,}
\end{aligned}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p &= j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p &= j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$H_1(A \cap B) \xrightarrow{i_*} H_1(A) \oplus H_1(B)$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p = j_{p*}(1) \text{ and } x_q = j_{q*}(1) &\text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\underbrace{H_1(A \cap B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle x_p, x_q \rangle} \xrightarrow{i_*} H_1(A) \oplus H_1(B)$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q &: S^1 \hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q &: S^1 \hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p &= j_{p*}(1) \text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\underbrace{H_1(A \cap B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle x_p, x_q \rangle} \xrightarrow{i_*} \underbrace{H_1(A) \oplus H_1(B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle y_A, y_B \rangle}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p = j_{p*}(1) &\text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\underbrace{H_1(A \cap B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle x_p, x_q \rangle} \xrightarrow[i_*]{x_p \rightarrow (y_A, y_B)} \underbrace{H_1(A) \oplus H_1(B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle y_A, y_B \rangle}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p = j_{p*}(1) &\text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\underbrace{H_1(A \cap B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle x_p, x_q \rangle} \xrightarrow[i_*]{x_p \rightarrow (y_A, y_B), x_q \rightarrow (y_A, y_B)} \underbrace{H_1(A) \oplus H_1(B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle y_A, y_B \rangle}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1)(= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p = j_{p*}(1) &\text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
\underbrace{H_1(A \cap B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle x_p, x_q \rangle} &\xrightarrow[i_*]{x_p \rightarrow (y_A, y_B), x_q \rightarrow (y_A, y_B)} \underbrace{H_1(A) \oplus H_1(B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle y_A, y_B \rangle} \\
Im \ i_* &= \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\begin{aligned}
U &:= S^1 \times (S^1 \setminus x_0) \text{ deformation retracts onto } A \\
V &:= S^1 \times (S^1 \setminus x_1) \text{ deformation retracts onto } B \\
V \cap V &= S^1 \times (S^1 \setminus \{x_0, x_1\}) \text{ deformation retracts onto } \\
&A \cap B
\end{aligned}$$

$$\begin{aligned}
i_p : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_p(s) = (s, p) \\
i_q : S^1 &\hookrightarrow S^1 \times I_1, \text{ where } i_q(s) = (s, q) \\
i_{p*}(1) &= i_{q*}(1) \text{ because } i_p \text{ is homotopic to } i_q \\
y_A &:= i_{p*}(1) (= i_{q*}(1)) \text{ generates } H_1(A) = \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
j_p : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_p(s) = (s, p) \\
j_q : S^1 &\hookrightarrow S^1 \times p \sqcup S^1 \times q, \text{ where } j_q(s) = (s, q) \\
x_p = j_{p*}(1) &\text{ and } x_q = j_{q*}(1) \text{ generate } H_1(A \cap B) = \mathbb{Z} \oplus \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
A \cap B &= S^1 \times p \sqcup S^1 \times q \xrightarrow{k_A} S^1 \times I_p = A \\
k_A \circ j_p &= i_p \text{ and } k_A \circ j_q = i_q \\
k_{A*} \circ j_{p*} &= i_{p*} \text{ and } k_{A*} \circ j_{q*} = i_{q*} \\
k_{A*}(j_{p*}(1)) &= i_{p*}(1) = i_{q*}(1) = k_{A*}(j_{q*}(1))
\end{aligned}$$

$$\begin{aligned}
k_{A*}(x_p) &= y_A = k_{A*}(x_q) \\
\text{Similarly,} \\
k_{B*}(x_p) &= y_B = k_{B*}(x_q), \text{ where } y_B \text{ generates } H_1(B) = \mathbb{Z}
\end{aligned}$$

$$\underbrace{H_1(A \cap B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle x_p, x_q \rangle} \xrightarrow[i_*]{x_p \rightarrow (y_A, y_B), x_q \rightarrow (y_A, y_B)} \underbrace{H_1(A) \oplus H_1(B)}_{\mathbb{Z} \oplus \mathbb{Z} \cong \langle y_A, y_B \rangle}$$

$$\begin{aligned}
Im \ i_* &= \mathbb{Z} \\
Ker \ i_* &= \mathbb{Z}
\end{aligned}$$

$$\begin{aligned}
X &= S^1 \times S^1 \\
A &= S^1 \times I_1 \\
B &= S^1 \times I_2 \\
A \cup B &= X \\
A \cap B &= S^1 \times p \sqcup S^1 \times q
\end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$0$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow{i_*} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \mapsto (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$X=S^1\times S^1$$

$$A=S^1\times I_1$$

$$B=S^1\times I_2$$

$$A\cup B=X$$

$$A\cap B=S^1\times p\sqcup S^1\times q$$

$$\widetilde{H}_1(A\cap B)\longrightarrow \widetilde{H}_1(A)\oplus \widetilde{H}_1(B)\longrightarrow \widetilde{H}_1(X)\longrightarrow \widetilde{H}_0(A\cap B)\longrightarrow \widetilde{H}_0(A)\oplus \widetilde{H}_0(B)\qquad \cdots$$

$$\mathbb{Z}\oplus\mathbb{Z}\overset{x_p,x_q\rightarrow(y_A,y_B)}{\underset{i_*}{\longrightarrow}}\mathbb{Z}\oplus\mathbb{Z}\overset{\quad\quad\quad}{\underset{j_*}{\longrightarrow}}??\overset{\quad\quad\quad}{\longrightarrow}\mathbb{Z}\overset{\quad\quad\quad}{\longrightarrow}0$$

$$Im\; i_* = \mathbb{Z}$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \overset{x_p, x_q \rightarrow (y_A, y_B)}{\underset{i_*}{\longrightarrow}} \mathbb{Z} \oplus \mathbb{Z} \overset{\hspace{1cm}}{\underset{j_*}{\longrightarrow}} ?? \overset{\hspace{1cm}}{\longrightarrow} \mathbb{Z} \overset{\hspace{1cm}}{\longrightarrow} 0$$

$$Im\ i_* = \mathbb{Z} = ker\ j_*$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \overset{x_p, x_q \rightarrow (y_A, y_B)}{\underset{i_*}{\longrightarrow}} \mathbb{Z} \oplus \mathbb{Z} \overset{\quad \quad \quad}{\underset{j_*}{\longrightarrow}} ?? \overset{\quad \quad \quad}{\longrightarrow} \mathbb{Z} \overset{\quad \quad \quad}{\longrightarrow} 0$$

$$\begin{aligned} Im\ i_* &= \mathbb{Z} = ker\ j_* \\ 0 \rightarrow \mathbb{Z} &\rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 \end{aligned}$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \overset{x_p, x_q \rightarrow (y_A, y_B)}{\underset{i_*}{\longrightarrow}} \mathbb{Z} \oplus \mathbb{Z} \overset{\quad \quad \quad}{\underset{j_*}{\longrightarrow}} ?? \overset{\quad \quad \quad}{\longrightarrow} \mathbb{Z} \overset{\quad \quad \quad}{\longrightarrow} 0$$

$$\begin{aligned} Im\ i_* &= \mathbb{Z} = ker\ j_* \\ 0 \rightarrow \mathbb{Z} \rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 &\text{ splits.} \end{aligned}$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$


$$\begin{aligned} Im\ i_* &= \mathbb{Z} = ker\ j_* \\ 0 \rightarrow \mathbb{Z} \rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 &\text{ splits.} \\ \text{Therefore,} \\ H_1(X) &= \mathbb{Z} \oplus \mathbb{Z} \end{aligned}$$

$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$

$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \mapsto (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} Im\ i_* &= \mathbb{Z} = ker\ j_* \\ 0 \rightarrow \mathbb{Z} \rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 &\text{ splits.} \\ \text{Therefore,} \\ H_1(X) &= \mathbb{Z} \oplus \mathbb{Z} \end{aligned}$$



$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$


$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \mapsto (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} Im \ i_* &= \mathbb{Z} = ker \ j_* \\ 0 \rightarrow \mathbb{Z} \rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 &\text{ splits.} \end{aligned}$$

Therefore,

$$H_1(X) = \mathbb{Z} \oplus \mathbb{Z}$$



$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$


$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} Im\ i_* &= \mathbb{Z} = ker\ j_* \\ 0 \rightarrow \mathbb{Z} \rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 &\text{ splits.} \end{aligned}$$

Therefore,

$$H_1(X) = \mathbb{Z} \oplus \mathbb{Z}$$



$$\begin{aligned} X &= S^1 \times S^1 \\ A &= S^1 \times I_1 \\ B &= S^1 \times I_2 \\ A \cup B &= X \\ A \cap B &= S^1 \times p \sqcup S^1 \times q \end{aligned}$$


$$\widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \widetilde{H}_1(X) \longrightarrow \widetilde{H}_0(A \cap B) \longrightarrow \widetilde{H}_0(A) \oplus \widetilde{H}_0(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{j_*} ?? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$\begin{aligned} Im \ i_* &= \mathbb{Z} = ker \ j_* \\ 0 \rightarrow \mathbb{Z} \rightarrow H_1(X) \rightarrow \mathbb{Z} \rightarrow 0 &\text{ splits.} \end{aligned}$$

Therefore,

$$H_1(X) = \mathbb{Z} \oplus \mathbb{Z}$$



$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \cdots$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \cdots$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$??$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \cdots$$

$$0$$

$$??$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \longrightarrow \cdots$$

$$0 \qquad \qquad \qquad 0 \qquad \qquad \qquad ?? \qquad \qquad \qquad \mathbb{Z} \oplus \mathbb{Z} \qquad \qquad \qquad \mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$0 \qquad \qquad \qquad 0 \longrightarrow ?? \xrightarrow{\partial} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{i_*} \mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$0 \qquad \qquad \qquad 0 \longrightarrow ?? \xrightarrow[\partial]{} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$0 \qquad \qquad \qquad 0 \longrightarrow ?? \xrightarrow[\partial]{} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z}$$

$$Im \, \partial$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$0 \qquad \qquad \qquad 0 \longrightarrow ?? \xrightarrow[\partial]{} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z}$$

$$Im \, \partial = Ker \, i_*$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$0 \qquad \qquad \qquad 0 \longrightarrow ?? \xrightarrow[\partial]{} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z}$$

$$Im \, \partial = Ker \, i_* = \mathbb{Z}$$

$$\widetilde{H}_2(A \cap B) \longrightarrow \widetilde{H}_2(A) \oplus \widetilde{H}_2(B) \longrightarrow \widetilde{H}_2(X) \longrightarrow \widetilde{H}_1(A \cap B) \longrightarrow \widetilde{H}_1(A) \oplus \widetilde{H}_1(B) \qquad \dots$$

$$0 \qquad \qquad \qquad 0 \longrightarrow ?? \xrightarrow[\partial]{} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow[i_*]{x_p, x_q \rightarrow (y_A, y_B)} \mathbb{Z} \oplus \mathbb{Z}$$

$$Im \, \partial = Ker \, i_* = \mathbb{Z}$$

$$0 \rightarrow H_2(X) \rightarrow \mathbb{Z} \rightarrow 0$$

$$H_0(S^1 \times S^1) = \mathbb{Z}$$

$$H_0(S^1 \times S^1) = \mathbb{Z}$$

$$H_1(S^1 \times S^1) = \mathbb{Z} \oplus \mathbb{Z}$$

$$H_0(S^1 \times S^1) = \mathbb{Z}$$

$$H_1(S^1 \times S^1) = \mathbb{Z} \oplus \mathbb{Z}$$

$$H_2(S^1 \times S^1) = \mathbb{Z}$$

$$H_0(S^1 \times S^1) = \mathbb{Z}$$

$$H_1(S^1 \times S^1) = \mathbb{Z} \oplus \mathbb{Z}$$

$$H_2(S^1 \times S^1) = \mathbb{Z}$$

$$H_n(S^1 \times S^1) = 0 \text{ if } n \geq 3$$

$$H_0(S^1 \times S^1) = \mathbb{Z}$$

$$H_1(S^1 \times S^1) = \mathbb{Z} \oplus \mathbb{Z}$$

$$H_2(S^1 \times S^1) = \mathbb{Z}$$

$$H_n(S^1 \times S^1) = 0 \text{ if } n \geq 3 \text{ (exercise!)}$$

$$H_0(S^1 \times S^1) = \mathbb{Z}$$

$$H_1(S^1 \times S^1) = \mathbb{Z} \oplus \mathbb{Z}$$

$$H_2(S^1 \times S^1) = \mathbb{Z}$$

$$H_n(S^1 \times S^1) = 0 \text{ if } n \geq 3 \text{ (exercise!)}$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$$\sigma(s) = (\cos s, \sin s)$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p)$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial x + a$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial x + a$$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$\partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$
$$\sigma(s) = (\cos s, \sin s) \text{ is a cycle}$$
$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$
$$\sigma = q_{\#}(Id_{[0,1]})$$
$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$
$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$
$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

$$\text{But } [c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \widetilde{H}_0(\{0, 1\})$

$H_1([0, 1], \{0, 1\})$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

$$\text{But } [c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$$

$$H_1([0, 1], \{0, 1\}) \xrightarrow{\partial} \tilde{H}_0(\{0, 1\})$$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)$?

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0$?
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \widetilde{H}_0(\{0, 1\})$

$H_1([0, 1], \{0, 1\}) \xrightarrow{\partial} \widetilde{H}_0(\{0, 1\}) \rightarrow \widetilde{H}_0([0, 1])$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

$$\text{But } [c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$$

$$H_1([0, 1], \{0, 1\}) \xrightarrow{\partial} \tilde{H}_0(\{0, 1\}) \rightarrow 0$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

$$\text{But } [c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$$

$$H_1([0, 1]) \rightarrow H_1([0, 1], \{0, 1\}) \xrightarrow{\partial} \tilde{H}_0(\{0, 1\}) \rightarrow 0$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

$$\text{But } [c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$$

$$0 \rightarrow H_1([0, 1], \{0, 1\}) \xrightarrow{\partial} \tilde{H}_0(\{0, 1\}) \rightarrow 0$$

Representative of a homology class

$$\sigma : \Delta_1 \rightarrow S^1$$

$\sigma(s) = (\cos s, \sin s)$ is a cycle

$$[\sigma] = [0] \in H_1(S^1)?$$

$$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$$

$$\sigma = q_{\#}(Id_{[0,1]})$$

$$q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$$

$$Id_{[0,1]} = 0?$$

$$Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$$

$$c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$$

$$\text{But } [c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$$

$$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \tilde{H}_0(\{0, 1\}) \rightarrow 0$$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0]>} \rightarrow 0$

Representative of a homology class

$$[Id_{\Delta_2}] \in H_2(\Delta_2, \partial \Delta_2)$$

$$\begin{aligned} \sigma : \Delta_1 &\rightarrow S^1 \\ \sigma(s) &= (\cos s, \sin s) \text{ is a cycle} \\ [\sigma] &= [0] \in H_1(S^1)? \end{aligned}$$

$$\begin{aligned} q : ([0, 1], \{0, 1\}) &\rightarrow (S^1, p) \\ \sigma &= q_{\#}(Id_{[0,1]}) \\ q_* : H_1([0, 1], \{0, 1\}) &\xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1) \end{aligned}$$

$$\begin{aligned} Id_{[0,1]} &= 0? \\ Id_{[0,1]} &= \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}} \\ c_1 - c_0 &= \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a \\ \text{But } [c_1 - c_0] &\neq 0 \in \tilde{H}_0(\{0, 1\}) \end{aligned}$$

$$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$$

Representative of a homology class

$$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial\Delta_2)?$$

$$\begin{aligned} \sigma : \Delta_1 &\rightarrow S^1 \\ \sigma(s) &= (\cos s, \sin s) \text{ is a cycle} \\ [\sigma] &= [0] \in H_1(S^1)? \end{aligned}$$

$$\begin{aligned} q : ([0, 1], \{0, 1\}) &\rightarrow (S^1, p) \\ \sigma &= q_{\#}(Id_{[0,1]}) \\ q_* : H_1([0, 1], \{0, 1\}) &\xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1) \end{aligned}$$

$$\begin{aligned} Id_{[0,1]} &= 0? \\ Id_{[0,1]} &= \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}} \\ c_1 - c_0 &= \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a \\ \text{But } [c_1 - c_0] &\neq 0 \in \tilde{H}_0(\{0, 1\}) \end{aligned}$$

$$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$

$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial \Delta_2)?$
 $H_2(\Delta_2, \partial \Delta_2) \xrightarrow{\partial} H_1(\underbrace{\partial \Delta_2}_{S^1})$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$

$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial \Delta_2)?$
 $H_2(\Delta_2, \partial \Delta_2) \xrightarrow{\partial} H_1(\Delta_1, \partial \Delta_1)??$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$

$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial \Delta_2)?$
 $H_2(\Delta_2, \partial \Delta_2) \xrightarrow{\partial} H_1(\partial \Delta_2, ??) \xleftarrow{\sim} H_1(\Delta_1, \partial \Delta_1)$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$

$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial \Delta_2)?$
 $H_2(\Delta_2, \partial \Delta_2) \xrightarrow{\partial} H_1(\partial \Delta_2, \wedge) \xleftarrow{\sim} H_1(\Delta_1, \partial \Delta_1)$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$

$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial \Delta_2)?$
 $H_2(\Delta_2, \partial \Delta_2) \xrightarrow{\partial} H_1(\partial \Delta_2, \wedge) \xrightarrow{\sim} \tilde{H}(\partial \Delta_1 / \partial \Delta_1) \xleftarrow{\sim} H_1(\Delta_1, \partial \Delta_1)$

Representative of a homology class

$\sigma : \Delta_1 \rightarrow S^1$
 $\sigma(s) = (\cos s, \sin s)$ is a cycle
 $[\sigma] = [0] \in H_1(S^1)?$

$q : ([0, 1], \{0, 1\}) \rightarrow (S^1, p)$
 $\sigma = q_{\#}(Id_{[0,1]})$
 $q_* : H_1([0, 1], \{0, 1\}) \xrightarrow{\sim} H_1(S^1, p) \cong H_1(S^1)$

$Id_{[0,1]} = 0?$
 $Id_{[0,1]} = \partial \underbrace{x}_{\in [0,1]} + \underbrace{a}_{\in \{0,1\}}$
 $c_1 - c_0 = \partial Id_{[0,1]} = \partial \partial x + \partial a = \partial a$
But $[c_1 - c_0] \neq 0 \in \tilde{H}_0(\{0, 1\})$

$0 \rightarrow \underbrace{H_1([0, 1], \{0, 1\})}_{<Id_{[0,1]}>} \xrightarrow{\partial} \underbrace{\tilde{H}_0(\{0, 1\})}_{<[c_1 - c_0] = \partial Id_{[0,1]}>} \rightarrow 0$

$[Id_{\Delta_2}] \neq 0 \in H_2(\Delta_2, \partial \Delta_2)?$
 $H_2(\Delta_2, \partial \Delta_2) \xrightarrow{\partial} H_1(\partial \Delta_2, \wedge) \xrightarrow{\sim} \tilde{H}(\partial \Delta_1 / \partial \Delta_1) \xleftarrow{\sim} H_1(\Delta_1, \partial \Delta_1)$

$\wedge \subset \partial \Delta_2 \subset \Delta_2$

Degree

$$f : S^n \rightarrow S^n$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : H_n(S^n) \rightarrow H_n(S^n)$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

$$f_*(1) = \deg f$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

$$f_*(1) = \deg f$$

Local degree

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

$$f_*(1) = \deg f$$

Local degree

$$f : U \rightarrow V \text{ homeomorphism}$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

$$f_*(1) = \deg f$$

Local degree

$$f : U \rightarrow V \text{ homeomorphism}$$

$$f : H_n(U, U \setminus x) \rightarrow H_n(V, V \setminus f(x))$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

$$f_*(1) = \deg f$$

Local degree

$$f : U \rightarrow V \text{ homeomorphism}$$

$$f : \mathbb{Z} \cong H_n(U, U \setminus x) \rightarrow H_n(V, V \setminus f(x)) \cong \mathbb{Z}$$

Degree

$$f : S^n \rightarrow S^n$$

$$f_* : \mathbb{Z} \cong H_n(S^n) \rightarrow H_n(S^n) \cong \mathbb{Z}$$

$$f_*(1) = \deg f$$

Local degree

$$f : U \rightarrow V \text{ homeomorphism}$$

$$f : \mathbb{Z} \cong H_n(U, U \setminus x) \rightarrow H_n(V, V \setminus f(x)) \cong \mathbb{Z}$$

$$f_*(1) = \deg f|_x$$

Theorem. $f : S^n \rightarrow S^n$

Theorem. $f : S^n \rightarrow S^n$

$f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y

Theorem. $f : S^n \rightarrow S^n$

$f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y

$\deg f = \sum_i \deg f|_{x_i}$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ *for some* y
 $\deg f = \sum_i \deg f|_{x_i}$

$$H_n(S^n) \xrightarrow{f_*} H_n(S^n)$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$H_n(S^n, S^n \setminus y)$$

$$j_* \uparrow$$

$$H_n(S^n) \xrightarrow{f_*} H_n(S^n)$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y))$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i))$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y)$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y)$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:

$$H_n(U_i, U_i \setminus x_i)$$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \overset{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \overset{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $H_n(U_i, U_i \setminus x_i) \rightarrow H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \overset{\sim}{\rightarrow} H_n(S^n, S^n \setminus \{x_1, \dots, x_k\})$ □

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \overset{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \overset{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $H_n(U_i, U_i \setminus x_i) \rightarrow H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \overset{\sim}{\rightarrow} H_n(S^n, S^n \setminus \{x_1, \dots, x_k\}) \rightarrow H_n(S^n, S^n \setminus x_j)$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & \uparrow j_* \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:

$$\begin{array}{ccc} H_n(U_i, U_i \setminus x_i) & \rightarrow & H_n(S^n, S^n \setminus \{x_1, \dots, x_k\}) \\ \downarrow & & \downarrow \\ H_n(U_i, U_i \setminus x_i) & \rightarrow & H_n(S^n, S^n \setminus x_j) \end{array}$$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $\underbrace{H_n(U_i, U_i \setminus x_i) \rightarrow H_n(S^n, S^n \setminus \{x_1, \dots, x_k\})}_{\text{picking out the } i\text{th generator}} \rightarrow H_n(S^n,$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $\underbrace{H_n(U_i, U_i \setminus x_i) \rightarrow H_n(S^n, S^n \setminus \{x_1, \dots, x_k\})}_{\text{picking out the } i\text{th generator}} \rightarrow H_n(S^n,$

Is equal to:
 $H_n(U_i, U_i \setminus x_i) \rightarrow H_n(U_i, U_i \setminus x_i) \rightarrow H_n(S^n, S^n \setminus x_j) \quad \square$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $\underbrace{H_n(U_i, U_i \setminus x_i) \rightarrow H_n(S^n, S^n \setminus \{x_1, \dots, x_k\})}_{\text{picking out the } i\text{th generator}} \rightarrow H_n(S^n,$

Is equal to:
 $H_n(U_i, U_i \setminus x_i) \rightarrow \underbrace{H_n(U_i, U_i \setminus x_i)}_0 \rightarrow H_n(S^n, S^n \setminus x_j)$

$$H_n(S^n, S^n \setminus \{x_1, \dots, x_k\}) \xrightarrow{i_*} H_n(S^n, S^n \setminus x_i) \quad \square$$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $\underbrace{H_n(U_i, U_i \setminus x_i) \rightarrow H_n(S^n, S^n \setminus \{x_1, \dots, x_k\})}_{\text{picking out the } i\text{th generator}} \rightarrow H_n(S^n,$

Is equal to:
 $H_n(U_i, U_i \setminus x_i) \rightarrow \underbrace{H_n(U_i, U_i \setminus x_i)}_0 \rightarrow H_n(S^n, S^n \setminus x_j)$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus \{x_1, \dots, x_k\}) & \xrightarrow{i_*} & H_n(S^n, S^n \setminus x_i) \\ \alpha_* \uparrow & & \\ H_n(U_i, U_i \setminus x_i) & & \end{array}$$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Proof. The composition:
 $\underbrace{H_n(U_i, U_i \setminus x_i) \rightarrow H_n(S^n, S^n \setminus \{x_1, \dots, x_k\})}_{\text{picking out the } i\text{th generator}} \rightarrow H_n(S^n,$

Is equal to:
 $H_n(U_i, U_i \setminus x_i) \rightarrow \underbrace{H_n(U_i, U_i \setminus x_i)}_0 \rightarrow H_n(S^n, S^n \setminus x_j)$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus \{x_1, \dots, x_k\}) & \xrightarrow{i_*} & H_n(S^n, S^n \setminus x_i) \\ \alpha_* \uparrow & \nearrow \beta_* & \\ H_n(U_i, U_i \setminus x_i) & & \end{array}$$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Lemma. $H_n(S^n) \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\})$

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Lemma. $H_n(S^n) \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\})$ maps the generator

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Lemma. $H_n(S^n) \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\})$ maps the generator to the sum of the generators.

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Lemma. $H_n(S^n) \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\})$ maps the generator to the sum of the generators.

$$\begin{array}{ccc} & H_n(S^n) & \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \\ \text{Proof.} & \downarrow & \\ & H_n(S^n, S^n \setminus x_i) & \end{array}$$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Lemma. $H_n(S^n) \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\})$ maps the generator to the sum of the generators.

Proof.

$$\begin{array}{ccc} H_n(S^n) & \xrightarrow{j'_*} & H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \\ \downarrow & & \swarrow p_* \\ H_n(S^n, S^n \setminus x_i) & & \end{array}$$

□

Theorem. $f : S^n \rightarrow S^n$
 $f^{-1}(y) = \{x_1, x_2, \dots, x_n\}$ for some y
 $\deg f = \sum_i \deg f|_{x_i}$

$$\begin{array}{ccc} H_n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ j'_* \uparrow & & j'_* \uparrow \\ H_n(S^n) & \xrightarrow{f_*} & H_n(S^n) \end{array}$$

$$H_n(S^n, S^n \setminus f^{-1}(y)) \stackrel{\sim}{\leftarrow} H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) \cong \oplus \mathbb{Z}$$

$$H_n(S^n, S^n \setminus y) \stackrel{\sim}{\leftarrow} H_n(V, V \setminus y) \cong \mathbb{Z}$$

Lemma. $\deg f = f'_*(j'_*(1))$

Lemma. $H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\}) \xrightarrow{p_*} H_n(S^n, S^n \setminus x_i)$ is a projection onto the i th factor.

Lemma. $H_n(S^n) \xrightarrow{j'_*} H_n(S^n, S^n \setminus \{x_1, x_2, \dots, x_k\})$ maps the generator to the sum of the generators.

$$\begin{array}{ccc} H^n(S^n, S^n \setminus f^{-1}(y)) & \xrightarrow{f'_*} & H_n(S^n, S^n \setminus y) \\ \uparrow & & \uparrow \\ H_n(\sqcup U_i, \sqcup (U_i \setminus x_i)) & \xrightarrow{f'_*} & H_n(V, V \setminus y) \end{array}$$