

M n-dimensional manifold

M n-dimensional manifold

Local orientation at p :

M n-dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p)$$

M n-dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p),$$

(U homeomorphic to \mathbb{R}^n)

M n-dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p) \cong \mathbb{Z},$$

(U homeomorphic to \mathbb{R}^n)

$$p \in B \subset U \subset M,$$

B homeomorphic to an open ball in \mathbb{R}^n .

M n -dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p) \cong \mathbb{Z},$$

(U homeomorphic to \mathbb{R}^n)

$$p \in B \subset U \subset M,$$

B homeomorphic to an open ball in \mathbb{R}^n .

The map, $i_{p*} : H_n(M, M \setminus B) \rightarrow H_n(M, M \setminus p)$
induces an isomorphism

M n-dimensional manifold

An orientation on M :

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p) \cong \mathbb{Z},$$

(U homeomorphic to \mathbb{R}^n)

$$p \in B \subset U \subset M,$$

B homeomorphic to an open ball in \mathbb{R}^n .

The map, $i_{p*} : H_n(M, M \setminus B) \rightarrow H_n(M, M \setminus p)$
induces an isomorphism

M n -dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p) \cong \mathbb{Z},$$

(U homeomorphic to \mathbb{R}^n)

$$p \in B \subset U \subset M,$$

B homeomorphic to an open ball in \mathbb{R}^n .

The map, $i_{p*} : H_n(M, M \setminus B) \rightarrow H_n(M, M \setminus p)$
induces an isomorphism

An orientation on M :

choice of assignments

$$p \rightarrow \mu_p \in H_n(M, M \setminus p) \text{ such that}$$

M n-dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p) \cong \mathbb{Z},$$

(U homeomorphic to \mathbb{R}^n)

$$p \in B \subset U \subset M,$$

B homeomorphic to an open ball in \mathbb{R}^n .

The map, $i_{p*} : H_n(M, M \setminus B) \rightarrow H_n(M, M \setminus p)$
induces an isomorphism

An orientation on M :

choice of assignments

$$p \rightarrow \mu_p \in H_n(M, M \setminus p) \text{ such that}$$

for each p , there exists a B (homeomorphic to an open ball in \mathbb{R}^n) such that

M n -dimensional manifold

Local orientation at p :

Choice of generator,

$$\mu_p \in H_n(M, M \setminus p) \cong H_n(U, U \setminus p) \cong \mathbb{Z},$$

(U homeomorphic to \mathbb{R}^n)

$$p \in B \subset U \subset M,$$

B homeomorphic to an open ball in \mathbb{R}^n .

The map, $i_{p*} : H_n(M, M \setminus B) \rightarrow H_n(M, M \setminus p)$
induces an isomorphism

An orientation on M :

choice of assignments

$$p \rightarrow \mu_p \in H_n(M, M \setminus p) \text{ such that}$$

for each p , there exists a B (homeomorphic to an open ball in \mathbb{R}^n) such that

if $\mu_p = i_{p*}(\mu_B)$ for some $\mu_B \in H_n(M, M \setminus B)$ for some $\mu_B \in B$, then for any $q \in B$, $\mu_q = i_{q*}(\mu_B)$

Theorem. *There is a 2 sheeted covering*
 $\pi : \widetilde{M} \rightarrow M$ such that

Theorem. *There is a 2 sheeted covering*
 $\pi : \widetilde{M} \rightarrow M$ *such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- 1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$
- 2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$

□

Theorem. *There is a 2 sheeted covering*

$\pi : \widetilde{M} \rightarrow M$ *such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} *is disconnected if and only if* M *is orientable.*

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$

$\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$
2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n
and B homeomorphic to an open ball in \mathbb{R}^n ,
and $\mu_B \in H_n(M, M \setminus B)$, define
 $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$ □

Theorem. *There is a 2 sheeted covering*

$\pi : \widetilde{M} \rightarrow M$ *such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} *is disconnected if and only if* M *is orientable.*

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$

$\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n
and B homeomorphic to an open ball in \mathbb{R}^n ,

and $\mu_B \in H_n(M, M \setminus B)$, define

$$U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B_1 \cap B_2$. □

Theorem. *There is a 2 sheeted covering*

$\pi : \widetilde{M} \rightarrow M$ *such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$
2. \widetilde{M} *is disconnected if and only if* M *is orientable.*

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$

$\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n
and B homeomorphic to an open ball in \mathbb{R}^n ,

and $\mu_B \in H_n(M, M \setminus B)$, define

$$U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- 1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$
- 2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} H_n(M, M \setminus B) & \longrightarrow & H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ H_n(M, M \setminus B_i) & & \end{array}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

- \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ H_n(M, M \setminus B_i) & & \end{array}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- 1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$
- 2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ H_n(M, M \setminus B_i) & & \end{array}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- 1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$
- 2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.
 If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc}
 \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\
 \uparrow & \nearrow & \\
 \mu_{B_1} \in H_n(M, M \setminus B_i) & &
 \end{array}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

- \widetilde{M} is disconnected if and only if M is orientable.

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_i) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. □

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_i) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.

□

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_i) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_i) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis. □

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

- $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

- \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_i) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis.

Each B is evenly covered:

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_1) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis.

Each B is evenly covered:

$$\pi^{-1}(B) = \{\mu_p \mid p \in B\}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_1) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis.

Each B is evenly covered:

$$\begin{aligned} \pi^{-1}(B) &= \{\mu_p \mid p \in B\} \\ &= \{i_{p*}(\mu_B) \mid p \in B\} \sqcup \{-i_{p*}(\mu_B) \mid p \in B\} \end{aligned}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_1) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis.

Each B is evenly covered:

$$\begin{aligned} \pi^{-1}(B) &= \{\mu_p \mid p \in B\} \\ &= \{i_{p*}(\mu_B) \mid p \in B\} \sqcup \{-i_{p*}(\mu_B) \mid p \in B\} \\ &= U_{\mu_B} \sqcup U_{-\mu_B} \end{aligned}$$

□

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_1) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis.

Each B is evenly covered:

$$\begin{aligned} \pi^{-1}(B) &= \{\mu_p \mid p \in B\} \\ &= \{i_{p*}(\mu_B) \mid p \in B\} \sqcup \{-i_{p*}(\mu_B) \mid p \in B\} \\ &= U_{\mu_B} \sqcup U_{-\mu_B} \end{aligned}$$

$\pi|_{U_{\mu_B}}$ bijection onto U_{μ_B} . □

Theorem. *There is a 2 sheeted covering $\pi : \widetilde{M} \rightarrow M$ such that*

1. $\pi^{-1}(p) = \{\mu_p, -\mu_p\}$

2. \widetilde{M} is disconnected if and only if M is orientable.

Proof. $\widetilde{M} := \{\mu_p \mid \mu_p \in H_n(M, M \setminus p)\}$
 $\pi : \widetilde{M} \rightarrow M$, defined by $\pi(\mu_p) := p$.

Given $B \subset U \subset M$, where U is homeomorphic to \mathbb{R}^n and B homeomorphic to an open ball in \mathbb{R}^n , and $\mu_B \in H_n(M, M \setminus B)$, define $U_{\mu_B} := \{i_{p*}(\mu_B) \mid p \in B\}$

If $\mu_x \in U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$, then $x \in B \subset B_1 \cap B_2$.

If $\mu_y \in U_{\mu_B}$, then $\mu_y = i_{y*}(\mu_B)$

$$\begin{array}{ccc} \mu_B \in H_n(M, M \setminus B) & \longrightarrow & \mu_y \in H_n(M, M \setminus y) \\ \uparrow & \nearrow & \\ \mu_{B_1} \in H_n(M, M \setminus B_1) & & \end{array}$$

Therefore, $\mu_y \in U_{\mu_{B_1}}$. Similarly, $\mu_y \in U_{\mu_{B_2}}$.
 $U_{\mu_B} \subset U_{\mu_{B_1}} \cap U_{\mu_{B_2}}$

So U_{μ_B} form a basis.

Each B is evenly covered:

$$\begin{aligned} \pi^{-1}(B) &= \{\mu_p \mid p \in B\} \\ &= \{i_{p*}(\mu_B) \mid p \in B\} \sqcup \{-i_{p*}(\mu_B) \mid p \in B\} \\ &= U_{\mu_B} \sqcup U_{-\mu_B} \end{aligned}$$

$\pi|_{U_{\mu_B}}$ bijection onto U_{μ_B} .

Maps basic open sets (i.e. U_{μ_B}) to basic open sets (i.e. B). □

Lemma. \widetilde{M} is orientable.

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism □

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$ □

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p



Lemma. \widetilde{M} is orientable.

$$H_n(\widetilde{M}, \widetilde{M} \setminus U_{\mu_B}) \xrightarrow{\pi_*} H_n(M, M \setminus B) \quad \square$$

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

$$H_n(\widetilde{M}, \widetilde{M} \setminus U_{\mu_B}) \xrightarrow{\pi_*} H_n(M, M \setminus B)$$

$$H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \xrightarrow{\pi_*} H_n(M, M \setminus p)$$

□

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

$$\begin{array}{ccc} H_n(\widetilde{M}, \widetilde{M} \setminus U_{\mu_B}) & \xrightarrow{\pi_*} & H_n(M, M \setminus B) \\ \downarrow & & \downarrow \\ H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) & \xrightarrow{\pi_*} & H_n(M, M \setminus p) \end{array}$$

□

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

$$\begin{array}{ccc} H_n(\widetilde{M}, \widetilde{M} \setminus U_{\mu_B}) & \xrightarrow{\pi_*} & H_n(M, M \setminus B) \\ \downarrow & & \downarrow \\ H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) & \xrightarrow{\pi_*} & H_n(M, M \setminus p) \end{array}$$

□

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

$$\begin{array}{ccc} H_n(\widetilde{M}, \widetilde{M} \setminus U_{\mu_B}) & \xrightarrow{\pi_*} & H_n(M, M \setminus B) \\ \downarrow & & \downarrow \\ H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) & \xrightarrow{\pi_*} & H_n(M, M \setminus p) \end{array}$$

□

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$
Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

$$\begin{array}{ccc} H_n(\widetilde{M}, \widetilde{M} \setminus U_{\mu_B}) & \xrightarrow{\pi_*} & H_n(M, M \setminus B)^{\mu_B} \\ \downarrow & & \downarrow \\ H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)^{\pi^{-1}(\mu_p)} & \xrightarrow{\pi_*} & H_n(M, M \setminus p)^{\mu_p} \end{array}$$

□

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$
Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

$$\begin{array}{ccc} H_n(\widetilde{M}, \overset{\mu_{\mu_B}}{\widetilde{M}} \setminus U_{\mu_B}) & \xrightarrow{\pi_*} & H_n(\overset{\mu_B}{M}, M \setminus B) \\ \downarrow & & \downarrow \\ \overset{\pi^{-1}(\mu_p)}{H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)} & \xrightarrow{\pi_*} & \overset{\mu_p}{H_n(M, M \setminus p)} \end{array}$$

□

Lemma. \widetilde{M} is orientable.

Proof. $\pi_* : H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p) \rightarrow H_n(M, M \setminus p)$ is an isomorphism

$\mu_p \in \widetilde{M}$ also determines $\mu_p \in H_n(M, M \setminus p)$

Assign, $\pi_*^{-1}(\mu_p) \in H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)$ as local orientation at μ_p .

Note: Given $\mu_p \in \widetilde{M}$, there is a B containing p therefore, there is a U_{μ_B} containing μ_p .

The assignment follows the conpatibility condition.

$$\begin{array}{ccc} H_n(\widetilde{M}, \overset{\mu_{\mu_B}}{\widetilde{M}} \setminus U_{\mu_B}) & \xrightarrow{\pi_*} & H_n(\overset{\mu_B}{M}, M \setminus B) \\ \downarrow & & \downarrow \\ \overset{\pi^{-1}(\mu_p)}{H_n(\widetilde{M}, \widetilde{M} \setminus \mu_p)} & \xrightarrow{\pi_*} & \overset{\mu_p}{H_n(M, M \setminus p)} \end{array}$$

□

Lemma. *A 2-sheeted cover of a connected space can have no more than two components.*

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.



Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

□

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. □

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path □

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path that joins \tilde{x} with an element of the fibre of x_0 , □

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path that joins \tilde{x} with an element of the fibre of x_0 , so the elements of the fibre represent the path homotopy classes. \square

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path that joins \tilde{x} with an element of the fibre of x_0 , so the elements of the fibre represent the path homotopy classes.

\tilde{X} is connected if and only if the two points in the same fibre can be connected by a path. \square

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path that joins \tilde{x} with an element of the fibre of x_0 , so the elements of the fibre represent the path homotopy classes.

\tilde{X} is connected if and only if the two points in the same fibre can be connected by a path.

Each component contains exactly one point of each fibre.

□

Lemma. *A 2-sheeted cover of a connected space can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \tilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \tilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path that joins \tilde{x} with an element of the fibre of x_0 , so the elements of the fibre represent the path homotopy classes.

\tilde{X} is connected if and only if the two points in the same fibre can be connected by a path.

Each component contains exactly one point of each fibre. π restricted to each component is a bijection. Also continuous and open. \square

Lemma. *A 2-sheeted cover of a connected space \widetilde{M} disconnected $\implies M$ orientable can have no more than two components. If it has two components, then each is homeomorphic to the ground space.*

Proof. $\pi : \widetilde{X} \rightarrow X$ is a 2-sheeted cover.

Fix $x_0 \in X$

Consider any $\tilde{x} \in \widetilde{X}$. Any path joining $\pi(\tilde{x})$ with x_0 can lift to a path that joins \tilde{x} with an element of the fibre of x_0 , so the elements of the fibre represent the path homotopy classes.

\widetilde{X} is connected if and only if the two points in the same fibre can be connected by a path.

Each component contains exactly one point of each fibre. π restricted to each component is a bijection. Also continuous and open. \square

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Conversely, if s is continuous,

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Conversely, if s is continuous, then if V is connected

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Conversely, if s is continuous, then if V is connected then
 $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Conversely, if s is continuous, then if V is connected then
 $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous
 $s : M \rightarrow \widetilde{M}$

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that If \widetilde{M} is connected,
 $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Conversely, if s is continuous, then if V is connected then
 $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous
 $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that If \widetilde{M} is connected, connect two points of a fibre by γ , $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.

$$\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so
continuous.

Conversely, if s is continuous, then if V is connected then
 $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous
 $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$. If \widetilde{M} is connected, connect two points of a fibre by γ , so $\pi(\gamma)$ is a loop.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so continuous.

Conversely, if s is continuous, then if V is connected then $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$. If \widetilde{M} is connected, connect two points of a fibre by γ , so $\pi(\gamma)$ is a loop. Then $s(\pi(\gamma))$ is a lift of a loop

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so continuous.

Conversely, if s is continuous, then if V is connected then $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$. If \widetilde{M} is connected, connect two points of a fibre by γ , so $\pi(\gamma)$ is a loop. Then $s(\pi(\gamma))$ is a lift of a loop contradicting uniqueness of a lift.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so continuous.

Conversely, if s is continuous, then if V is connected then $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Assignment $p \rightarrow \mu_p$ equivalent to $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.

Restrict to an evenly covered neighbourhood $V \subset M$.
 $\pi^{-1}(V) = \widetilde{V}_1 \sqcup \widetilde{V}_2$

If \widetilde{M} is connected, connect two points of a fibre by γ , so $\pi(\gamma)$ is a loop. Then $s(\pi(\gamma))$ is a lift of a loop contradicting uniqueness of a lift.

So, \widetilde{M} connected $\implies M$ not orientable.

If s is such that $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 ,
then s is an inverse of the homeomorphism $\pi|_{\widetilde{V}_1}$ so continuous.

Conversely, if s is continuous, then if V is connected then $s(V) = \widetilde{V}_i$ for $i = 0$ or 1 .

So M orientable is equivalent to finding continuous $s : M \rightarrow \widetilde{M}$ such that $\pi \circ s = Id$.