

$$C_n(X) \xrightarrow{j} C_n(X)/C_n(A)$$

$$C_n(X) \xrightarrow{j} C_n(X)/C_n(A)$$

$$H_n(X) \xrightarrow{j_*} H_n(X, A)$$

$$C_n(A) \xrightarrow{i^\#} C_n(X) \xrightarrow{j} C_n(X)/C_n(A)$$

$$H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A)$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A)$$

$$H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A)$$

$$0 \rightarrow C_n(A) \xrightarrow{i^\#} C_n(X) \xrightarrow{j} C_n(X)/C_n(A)$$

$$H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A)$$

$$0 \rightarrow C_n(A) \xrightarrow{i^\#} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A)$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A)$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

$$0 \rightarrow C_n(A) \xrightarrow{i^\#} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Exact sequence

$$\cdots \rightarrow A_{i+i} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \rightarrow \cdots$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X,A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X,A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Exact sequence

$$\cdots \rightarrow A_{i+i} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \rightarrow \cdots$$

$$Ker\ f_i = Im\ f_{i+1}$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Exact sequence

$$\cdots \rightarrow A_{i+i} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \rightarrow \cdots$$

$$\text{Ker } f_i = \text{Im } f_{i+1}$$

Short exact sequence

$$0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Exact sequence

$$\cdots \rightarrow A_{i+i} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \rightarrow \cdots$$

$$\text{Ker } f_i = \text{Im } f_{i+1}$$

Short exact sequence

$$0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$$

$$\text{Ker } g = \text{Im } f$$

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Exact sequence

$$\cdots \rightarrow A_{i+i} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \rightarrow \cdots$$

$$\text{Ker } f_i = \text{Im } f_{i+1}$$

Short exact sequence

$$0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$$

$$\text{Ker } g = \text{Im } f$$

f injective

$$0 \rightarrow C_n(A) \xrightarrow{i_{\#}} C_n(X) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Exact sequence

$$\cdots \rightarrow A_{i+i} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \rightarrow \cdots$$

$$\text{Ker } f_i = \text{Im } f_{i+1}$$

Short exact sequence

$$0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$$

$$\text{Ker } g = \text{Im } f$$

f injective

g surjective

Chain complexes

Chain complexes

$$\dots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \dots$$

Chain complexes

$$\cdots \xrightarrow{\partial_{n+2}} C_{n+1}(X) \xrightarrow{\partial_{n+1}} C_n(X) \xrightarrow{\partial_n} C_{n-1}(X) \xrightarrow{\partial_{n-1}} \cdots$$

Chain complexes

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & C_{n+1}(X, A) & \xrightarrow{\partial_{n+1}} & C_n(X, A) & \xrightarrow{\partial_n} & \\ C_{n-1}(X, A) & \xrightarrow{\partial_{n-1}} & \cdots & & & & \end{array}$$

Chain complexes

$$\cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

Chain complexes

$$\cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \rightarrow \cdots$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \rightarrow \cdots$$

$$\cdots \xrightarrow{\partial'_{n+2}} B_{n+1} \xrightarrow{\partial'_{n+1}} B_n \xrightarrow{\partial'_n} B_{n-1} \rightarrow \cdots$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & A_{n+1} & \xrightarrow{\partial_{n+1}} & A_n & \xrightarrow{\partial_n} & A_{n-1} \rightarrow \cdots \\ & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} \\ \cdots & \xrightarrow{\partial'_{n+2}} & B_{n+1} & \xrightarrow{\partial'_{n+1}} & B_n & \xrightarrow{\partial'_n} & B_{n-1} \rightarrow \cdots \end{array}$$

Chain complexes

$$0 \rightarrow A_* \rightarrow B_* \rightarrow C_* \rightarrow 0$$

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & A_{n+1} & \xrightarrow{\partial_{n+1}} & A_n & \xrightarrow{\partial_n} & A_{n-1} \rightarrow \cdots \\ & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} \\ \cdots & \xrightarrow{\partial'_{n+2}} & B_{n+1} & \xrightarrow{\partial'_{n+1}} & B_n & \xrightarrow{\partial'_n} & B_{n-1} \rightarrow \cdots \end{array}$$

$$\partial'_n \circ f_n = f_{n-1} \circ \partial_n$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & A_{n+1} & \xrightarrow{\partial_{n+1}} & A_n & \xrightarrow{\partial_n} & A_{n-1} \rightarrow \cdots \\ & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} \\ \cdots & \xrightarrow{\partial'_{n+2}} & B_{n+1} & \xrightarrow{\partial'_{n+1}} & B_n & \xrightarrow{\partial'_n} & B_{n-1} \rightarrow \cdots \end{array}$$

$$\partial'_n \circ f_n = f_{n-1} \circ \partial_n$$

$$0 \rightarrow A_* \rightarrow B_* \rightarrow C_* \rightarrow 0 \text{ implies } H_n(A) \rightarrow H_n(B) \rightarrow H_n(C)$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & A_{n+1} & \xrightarrow{\partial_{n+1}} & A_n & \xrightarrow{\partial_n} & A_{n-1} \rightarrow \cdots \\ & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} \\ \cdots & \xrightarrow{\partial'_{n+2}} & B_{n+1} & \xrightarrow{\partial'_{n+1}} & B_n & \xrightarrow{\partial'_n} & B_{n-1} \rightarrow \cdots \end{array}$$

$$\partial'_n \circ f_n = f_{n-1} \circ \partial_n$$

$$0 \rightarrow A_* \rightarrow B_* \rightarrow C_* \rightarrow 0 \text{ implies } \rightarrow H_{n+1}(A) \rightarrow H_n(A) \rightarrow H_n(B) \rightarrow H_n(C)$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & A_{n+1} & \xrightarrow{\partial_{n+1}} & A_n & \xrightarrow{\partial_n} & A_{n-1} \rightarrow \cdots \\ & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} \\ \cdots & \xrightarrow{\partial'_{n+2}} & B_{n+1} & \xrightarrow{\partial'_{n+1}} & B_n & \xrightarrow{\partial'_n} & B_{n-1} \rightarrow \cdots \end{array}$$

$$\partial'_n \circ f_n = f_{n-1} \circ \partial_n$$

$$\begin{aligned} 0 \rightarrow A_* \rightarrow B_* \rightarrow C_* \rightarrow 0 \text{ implies} \\ \rightarrow H_{n+1}(A) \rightarrow H_n(A) \rightarrow H_n(B) \rightarrow H_n(C) \rightarrow \\ H_{n-1}(A) \rightarrow \end{aligned}$$

Chain complexes

$$A_* := \cdots \xrightarrow{\partial_{n+2}} A_{n+1} \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} \cdots$$

$$\partial_n \circ \partial_{n+1} = 0$$

$$H_n(A) := \frac{\ker \partial_n}{\operatorname{Im} \partial_{n+1}}$$

Chain map

$$f : A_* \rightarrow B_*$$

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{n+2}} & A_{n+1} & \xrightarrow{\partial_{n+1}} & A_n & \xrightarrow{\partial_n} & A_{n-1} \rightarrow \cdots \\ & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} \\ \cdots & \xrightarrow{\partial'_{n+2}} & B_{n+1} & \xrightarrow{\partial'_{n+1}} & B_n & \xrightarrow{\partial'_n} & B_{n-1} \rightarrow \cdots \end{array}$$

$$\partial'_n \circ f_n = f_{n-1} \circ \partial_n$$

$$\begin{aligned} 0 \rightarrow A_* \rightarrow B_* \rightarrow C_* \rightarrow 0 \text{ implies} \\ \rightarrow H_{n+1}(A) \rightarrow H_n(A) \rightarrow H_n(B) \rightarrow H_n(C) \rightarrow \\ H_{n-1}(A) \rightarrow H_{n-1}(B) \rightarrow H_{n-1}(C) \rightarrow \cdots \end{aligned}$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(\overset{c_n}{X}, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow c_n \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \\
 & & & & & & \downarrow \partial c_n
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n^{c_n}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}^{\partial c_n=0}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\text{red}} & C_n(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \text{red} \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X,A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X,A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X,A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\text{red}} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\text{red}} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \xrightarrow{\text{red}} & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Defining ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\text{red}} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\text{red}} & C_{n-1}(X) & \xrightarrow{\text{red}} & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of preimage of j

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) & \longrightarrow & 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of preimage of j

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(\overset{0}{X}, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of preimage of j

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of preimage of j

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of preimage of j

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \\
 & & & & \downarrow & & \\
 & & & & \partial i(a_n) & &
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of preimage of j

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \\
& & & & \downarrow & & \downarrow \\
& & & & \partial i(a_n) & & 0
\end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of preimage of j

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of preimage of j

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\quad a_{n-1} \quad} & C_{n-1}(X) & \xrightarrow{\quad \partial i(a_n) \quad} & C_{n-1}(X, A) \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of homology class

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) & \longrightarrow & 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of homology class

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}^c(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n^{\partial c}(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of homology class

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\text{red}} & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \text{red} \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of homology class

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} \\
0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{\partial b}(X) & \longrightarrow & C_n^{\partial(j(b))}(X, A) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness

Independence of homology class

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{\partial b}(X) & \xrightarrow{\text{red}} & C_n^{\partial(j(b))}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of homology class

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{\partial b}(X) & \xrightarrow{\text{red}} & C_n^{\partial(j(b))}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}^0(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking well definedness
Independence of homology class

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{\partial b}(X) & \xrightarrow{\text{red}} & C_n^{\partial(j(b))}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}^0(A) & \xrightarrow{\text{red}} & C_{n-1}^0(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \xrightarrow{\quad j(b) \quad} 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{b} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{j(b_n)} & C_n(X, A) \xrightarrow{j(b_n)} 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \xrightarrow{\quad j(b) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \xrightarrow{\quad j(b_n) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \xrightarrow{\quad j(b) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \xrightarrow{\quad j(b_n) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \xrightarrow{\quad \partial b_n \quad} & C_{n-1}(X, A) \xrightarrow{\quad 0 \quad} 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{b} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{b_n} & C_n(X, A) \xrightarrow{j(b_n)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \xrightarrow{\partial b_n} & C_{n-1}(X, A) \xrightarrow{0} 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{b} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{b_n} & C_n(X, A) \xrightarrow{j(b_n)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{a_{n-1}} & C_{n-1}(X) & \xrightarrow{\partial b_n = i(a_{n-1})} & C_{n-1}(X, A) \xrightarrow{0} 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \xrightarrow{\quad j(b) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \xrightarrow{\quad j(b_n) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\quad a_{n-1}=\partial a_n \quad} & C_{n-1}(X) & \xrightarrow{\quad \partial b_n=i(a_{n-1}) \quad} & C_{n-1}(X, A) \xrightarrow{\quad 0 \quad} 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Ker } \partial \subset \text{Im } j_*$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad b_n, b_n - i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\quad a_{n-1} = \partial a_n \quad} & C_{n-1}(X) & \xrightarrow{\quad \partial b_n = i(a_{n-1}) \quad} & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$Im\ j_* \subset Ker\ \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \longrightarrow & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Im } j_* \subset \text{Ker } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{b} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{b_n} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Im } j_* \subset \text{Ker } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \xrightarrow{\quad j(b) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{ (red) } & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \xrightarrow{\quad \partial b_n = 0 \quad} & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\operatorname{Im} j_* \subset \operatorname{Ker} \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{b} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{b_n} & C_n(X, A) \xrightarrow{j(b_n)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \xrightarrow{\partial b_n=0} & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X, A)$

$$\text{Im } j_* \subset \text{Ker } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X, A) \xrightarrow{\quad j(b) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\quad b_n \quad} & C_n(X, A) \xrightarrow{\quad j(b_n) \quad} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\quad 0 \quad} & C_{n-1}(X) & \xrightarrow{\quad \partial b_n = 0 \quad} & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$\text{Ker } i \subset \text{Im } \partial$$

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \longrightarrow & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$\text{Ker } i \subset \text{Im } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^b(X) & \longrightarrow & C_{n+1}^{j(b)}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n^{a_n}(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$\text{Ker } i \subset \text{Im } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{b} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{a_n} & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\partial a_n=0} & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$
Ker i \subset *Im* ∂

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad b \quad} & C_{n+1}(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X,A) \longrightarrow 0 \\
 & & \downarrow & & & & \\
 & & \partial a_n = 0 & & & &
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X,A) \xrightarrow{\quad \partial \quad} H_n(A) \xrightarrow{\quad i_* \quad} H_n(X) \xrightarrow{\quad j_* \quad} H_n(X,A) \xrightarrow{\quad \partial \quad} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$\text{Ker } i \subset \text{Im } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{bb_{n+1}} & C_{n+1}(X, A) \xrightarrow{j(b)} 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{a_n} & C_n(X) & \xrightarrow{i(a_n)=\partial(b_{n+1})} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \xrightarrow{\partial a_n=0} & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$\text{Ker } i \subset \text{Im } \partial$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{\quad j(b)j(b_{n+1}) \quad} & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad i(a_n)=\partial(b_{n+1}) \quad} & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$Im \partial \subset Ker i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$Im \partial \subset Ker i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{j(b_{n+1})} & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$Im \partial \subset Ker i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \xrightarrow{j(b_{n+1})} & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$Im \partial \subset Ker i$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^{b_{n+1}}(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b_{n+1})}(X, A) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \text{red} & & \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n^0(X, A) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) & \longrightarrow & 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$Im \partial \subset Ker i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^{b_{n+1}}(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b_{n+1})}(X, A) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} & & \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{\partial b_{n+1}}(X) & \longrightarrow & C_n^0(X, A) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) & \longrightarrow & 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(A)$

$$\operatorname{Im} \partial \subset \operatorname{Ker} i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^{b_{n+1}}(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b_{n+1})}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} \\
 0 & \longrightarrow & C_n^{a_n}(A) & \xrightarrow{\text{red}} & C_n^{\partial b_{n+1}}(X) & \longrightarrow & C_n^0(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\text{Ker } j \subset \text{Im } i$$

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) & \longrightarrow & 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\text{Ker } j \subset \text{Im } i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$
Ker j \subset *Im i*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X,A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X,A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X,A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$
Ker j \subset *Im i*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \xrightarrow{\text{red}} & C_n(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X,A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X,A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X,A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\text{Ker } j \subset \text{Im } i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}^{c_{n+1}}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{b_n}(X) & \xrightarrow{\text{red}} & C_n^{j(b_n)=\partial c_{n+1}}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}^0(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\text{Ker } j \subset \text{Im } i$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^{b_{n+1}}(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b_{n+1})}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \text{red} \\
 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n^{b_n}(X) & \xrightarrow{\text{red}} & C_n^{j(b_n)=\partial j(b_{n+1})}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}^0(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$\text{Ker } j \subset \text{Im } i$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^{b_{n+1}}(X) & \xrightarrow{\text{red}} & C_{n+1}^{j(b_{n+1})}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \text{red} \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad b_n \text{ homologous } b_n - \partial b_{n+1} \quad} & C_n(X) & \xrightarrow{\text{red}} & C_n^{j(b_n) = \partial j(b_{n+1})}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow \text{red} & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}^0(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$
 $Ker\ j \subset Im\ i$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}^{b_{n+1}}(X) & \xrightarrow{\hspace{1cm}} & C_{n+1}^{j(b_{n+1})}(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n^{a_n}(A) & \xrightarrow{\hspace{1cm}} & C_n^{b_n \text{ homologous } b_n - \partial b_{n+1} = i(a_n)}(X) & \xrightarrow{\hspace{1cm}} & C_n^{j(b_n) = \partial j(b_{n+1})}(X,A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}^0(X) & \longrightarrow & C_{n-1}(X,A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X,A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X,A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\operatorname{Im} i \subset \operatorname{Ker} j$$

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_n(A) & \longrightarrow & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$Im\ i \subset Ker\ j$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{a_n} & C_n(X) & \longrightarrow & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\operatorname{Im} i \subset \operatorname{Ker} j$$

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \longrightarrow & C_n(X, A) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) & \longrightarrow & 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$Im\ i \subset Ker\ j$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

Checking exactness at $H_n(X)$

$$\textit{Im } i \subset \textit{Ker } j$$

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_{n+1}(A) & \longrightarrow & C_{n+1}(X) & \longrightarrow & C_{n+1}(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_n(A) & \xrightarrow{\quad a_n \quad} & C_n(X) & \xrightarrow{\quad i(a_n) \quad} & C_n(X, A) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C_{n-1}(A) & \longrightarrow & C_{n-1}(X) & \longrightarrow & C_{n-1}(X, A) \longrightarrow 0
 \end{array}$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

$$B \subset A \subset X$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A)/C_n(B) \overset{i}{\hookrightarrow} C_n(X)/C_n(B)$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A)/C_n(B) \xrightarrow{i} C_n(X)/C_n(B) \xrightarrow{j} C_n(X)/C_n(A)$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A)/C_n(B) \xrightarrow{i} C_n(X)/C_n(B) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xrightarrow{i} C_n(X)/C_n(B) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X)/C_n(A) \rightarrow 0$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$H_n(A, B) \xrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A)$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$H_n(A, B) \xrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$C_n(A) \oplus C_n(B) \xrightarrow[i_A + i_B]{j} C_n(X)$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$C_n(A) \oplus C_n(B) \xrightarrow[i_A+i_B]{j} C_n^{A,B}(X)$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$C_n(A) \oplus C_n(B) \xrightarrow[i_A + i_B]{j} C_n^{A, B}(X) \rightarrow 0$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$0 \rightarrow C_n(A \cap B) \xrightarrow[c \mapsto (i(c), -i(c))]{i} C_n(A) \oplus C_n(B) \xrightarrow[i_A + i_B]{j} C_n^{A, B}(X) \rightarrow 0$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A,B) \overset{i}{\hookrightarrow} C_n(X,B) \overset{j}{\rightarrow} C_n(X,A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X,A) \overset{\partial}{\rightarrow} H_n(A,B) \overset{i_*}{\hookrightarrow} H_n(X,B) \overset{j_*}{\rightarrow} H_n(X,A) \overset{\partial}{\rightarrow} H_{n-1}(A,B) \rightarrow \cdots$$

$$0 \rightarrow C_n(A \cap B) \overset{i}{\underset{c \rightarrow (i(c), -i(c))}{\longrightarrow}} C_n(A) \oplus C_n(B) \overset{j}{\underset{i_A + i_B}{\longrightarrow}} C_n^{A,B}(X) \rightarrow 0$$

$$H_n(A \cap B) \overset{i_*}{\rightarrow} H_n(A) \oplus H_n(B) \overset{j_*}{\rightarrow} H_n^{A,B}(X)$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$0 \rightarrow C_n(A \cap B) \xrightarrow[c \rightarrow (i(c), -i(c))]{i} C_n(A) \oplus C_n(B) \xrightarrow[i_A + i_B]{j} C_n^{A,B}(X) \rightarrow 0$$

$$H_n(A \cap B) \xrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n^{A,B}(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

$$0 \rightarrow C_n(A \cap B) \xrightarrow[c \mapsto (i(c), -i(c))]{i} C_n(A) \oplus C_n(B) \xrightarrow[i_A + i_B]{j} C_n^{A, B}(X) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(A \cap B) \xrightarrow{\partial} H_n(A \cap B) \xhookrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n^{A, B}(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

Long exact sequence of triple

$$B \subset A \subset X$$

$$0 \rightarrow C_n(A, B) \xhookrightarrow{i} C_n(X, B) \xrightarrow{j} C_n(X, A) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(X, A) \xrightarrow{\partial} H_n(A, B) \xhookrightarrow{i_*} H_n(X, B) \xrightarrow{j_*} H_n(X, A) \xrightarrow{\partial} H_{n-1}(A, B) \rightarrow \cdots$$

Mayer-Vietoris sequence

$$0 \rightarrow C_n(A \cap B) \xrightarrow[c \mapsto (i(c), -i(c))]{i} C_n(A) \oplus C_n(B) \xrightarrow[i_A + i_B]{j} C_n^{A, B}(X) \rightarrow 0$$

$$\cdots \rightarrow H_{n+1}(A \cap B) \xrightarrow{\partial} H_n(A \cap B) \xhookrightarrow{i_*} H_n(A) \oplus H_n(B) \xrightarrow{j_*} H_n^{A, B}(X) \xrightarrow{\partial} H_{n-1}(A \cap B) \rightarrow \cdots$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$*

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A ,*

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

Reduced homology

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E$, and f, g , isomorphisms, then $C \cong 0$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$\tilde{H}_0(X) := \frac{\ker \epsilon}{\operatorname{Im} \partial_1}$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$\tilde{H}_0(X) := \frac{\ker \epsilon}{\operatorname{Im} \partial_1}$$

$$H_0(X) = \tilde{H}_0(X) \oplus \mathbb{Z} \text{ (Exercise!)}$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$\tilde{H}_0(X) := \frac{\ker \epsilon}{\operatorname{Im} \partial_1}$$

$$H_0(X) = \tilde{H}_0(X) \oplus \mathbb{Z} \text{ (Exercise!)}$$

$$\tilde{H}_i(X) = H_i(X) \text{ for } i \geq 1$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$\tilde{H}_0(X) := \frac{\ker \epsilon}{\operatorname{Im} \partial_1}$$

$$H_0(X) = \tilde{H}_0(X) \oplus \mathbb{Z} \text{ (Exercise!)}$$

$$\tilde{H}_i(X) = H_i(X) \text{ for } i \geq 1$$

$$\cdots \rightarrow \tilde{H}_n(A) \rightarrow \tilde{H}_n(X) \rightarrow H_n(X, A) \rightarrow \tilde{H}_{n-1}(A) \rightarrow \cdots$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$\tilde{H}_0(X) := \frac{\ker \epsilon}{\operatorname{Im} \partial_1}$$

$$H_0(X) = \tilde{H}_0(X) \oplus \mathbb{Z} \text{ (Exercise!)}$$

$$\tilde{H}_i(X) = H_i(X) \text{ for } i \geq 1$$

$$\cdots \rightarrow \tilde{H}_n(A) \rightarrow \tilde{H}_n(X) \rightarrow H_n(X, A) \rightarrow \tilde{H}_{n-1}(A) \rightarrow \cdots$$

$$\textbf{Exercise. } H_n(X, x_0) = \tilde{H}_n(X)$$

$$0 \rightarrow A \rightarrow 0 \implies A = 0$$

$$0 \rightarrow A \rightarrow B \rightarrow 0 \implies A \cong B$$

$$A \xrightarrow{f} B \rightarrow C \rightarrow D \xrightarrow{g} E, \text{ and } f, g, \text{ isomorphisms, then } C \cong 0$$

Exercise. If $A \subset V \subset X$ and V deformation retracts onto A , then $i_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.

Reduced homology

$$\cdots \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z}$$

$$\tilde{H}_0(X) := \frac{\ker \epsilon}{\operatorname{Im} \partial_1}$$

$$H_0(X) = \tilde{H}_0(X) \oplus \mathbb{Z} \text{ (Exercise!)}$$

$$\tilde{H}_i(X) = H_i(X) \text{ for } i \geq 1$$

$$\cdots \rightarrow \tilde{H}_n(A) \rightarrow \tilde{H}_n(X) \rightarrow H_n(X, A) \rightarrow \tilde{H}_{n-1}(A) \rightarrow \cdots$$

$$\textbf{Exercise. } H_n(X, x_0) = \tilde{H}_n(X)$$

Exercise. Prove that $\tilde{H}_n(A) \xrightarrow{i_*} \tilde{H}_n(X)$ is an isomorphism for all n if and only if $H_n(X, A) = 0$ for all n

$$A \subset X$$

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$.

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map)

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map), $q(x) = [x]$

$q^{-1}(U)$ open if and only if U is open.

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map), $q(x) = [x]$

$q^{-1}(U)$ open if and only if U is open. “ X/A is a quotient of X ”.

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map), $q(x) = [x]$

$q^{-1}(U)$ open if and only if U is open. “ X/A is a quotient of X ”.

Examples:

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map), $q(x) = [x]$

$q^{-1}(U)$ open if and only if U is open. “ X/A is a quotient of X ”.

Examples:

1. $D^n/S^{n-1} \simeq S^n$

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map), $q(x) = [x]$

$q^{-1}(U)$ open if and only if U is open. “ X/A is a quotient of X ”.

Examples:

1. $D^n/S^{n-1} \simeq S^n$

2. $[0, 1]/\{0, 1\} \simeq S^1$

$A \subset X$ $x \sim y$ iff $x = y$ or $x, y \in A$. X/A denotes set of equivalence classes.

$q : X \rightarrow X/A$ (quotient map), $q(x) = [x]$

$q^{-1}(U)$ open if and only if U is open. “ X/A is a quotient of X ”.

Examples:

1. $D^n/S^{n-1} \simeq S^n$

2. $[0, 1]/\{0, 1\} \simeq S^1$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$H_n(X \setminus A, V \setminus A) \xrightarrow{i_* \text{ excision}} H_n(X, V)$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & & \end{array}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) \end{aligned}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) \end{aligned}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} \end{aligned}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) \end{aligned}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \end{aligned}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_*$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \end{aligned}$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism Recall,

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \end{aligned}$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

Recall,

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

Exercise. *If $A \subset V \subset X$*

$$\begin{array}{ccc}
 H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\
 \downarrow q_* & & \downarrow q'_* \\
 H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A)
 \end{array}$$

$$\begin{aligned}
 i' \circ q &= q' \circ i \\
 (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x))
 \end{aligned}$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

Recall,

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A ,*

$$\begin{array}{ccc}
 H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\
 \downarrow q_* & & \downarrow q'_* \\
 H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A)
 \end{array}$$

$$\begin{aligned}
 i' \circ q &= q' \circ i \\
 (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x))
 \end{aligned}$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$H_n(X, A) \xrightarrow{j_*} H_n(X, V)$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \\ i'_* \circ q_* &= q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1} \end{aligned}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ & & \downarrow q'_* \\ & & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \\ i'_* \circ q_* &= q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1} \end{aligned}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \\ i'_* \circ q_* &= q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1} \end{aligned}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \\ i'_* \circ q_* &= q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1} \end{aligned}$$

q_* is an isomorphism

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \end{aligned}$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} i' \circ q &= q' \circ i \\ (i'(q(x))) &= i'(\{x\}) = \{x\} = q'(x) = q'(i(x)) \\ i'_* \circ q_* &= q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1} \end{aligned}$$

q_* is an isomorphism

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$\begin{aligned} j' \circ q'' &= q' \circ j \\ (j'(q''(x))) & \end{aligned}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\})$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\}) = \{x\}$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\}) = \{x\} = q'(x)$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. *If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.*

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\}) = \{x\} = q'(x) = q'(j(x))$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\}) = \{x\} = q'(x) = q'(j(x))$$

$$j'_* \circ q''_* = q'_* \circ j_*$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\}) = \{x\} = q'(x) = q'(j(x))$$

$$j'_* \circ q''_* = q'_* \circ j_* \implies q''_* = j'^{-1}_* \circ q'_* \circ j_*$$

$q : X \setminus A \rightarrow X/A \setminus A/A$ is an isomorphism

If there exists $V \subset X$, such that $A \subset V$ and V deformation retracts onto A , then

$$\begin{array}{ccc} H_n(X \setminus A, V \setminus A) & \xrightarrow{i_* \text{ excision}} & H_n(X, V) \\ \downarrow q_* & & \downarrow q'_* \\ H_n(X/A \setminus A/A, V/A \setminus A/A) & \xrightarrow{i'_*(excision)} & H_n(X/A, V/A) \end{array}$$

$$i' \circ q = q' \circ i$$

$$(i'(q(x))) = i'(\{x\}) = \{x\} = q'(x) = q'(i(x))$$

$$i'_* \circ q_* = q'_* \circ i_* \implies q'_* = i'_* \circ q_* \circ i_*^{-1}$$

q_* is an isomorphism

Recall,

Exercise. If $A \subset V \subset X$ and V deformation retracts onto A , then $j_* : H_n(X, A) \rightarrow H_n(X, V)$ is an isomorphism.

$$\begin{array}{ccc} H_n(X, A) & \xrightarrow{j_*} & H_n(X, V) \\ \downarrow q''_* & & \downarrow q'_* \\ H_n(X/A, A/A) & \xrightarrow{j'_*} & H_n(X/A, V/A) \end{array}$$

$$j' \circ q'' = q' \circ j$$

$$(j'(q''(x))) = j'(\{x\}) = \{x\} = q'(x) = q'(j(x))$$

$$j'_* \circ q''_* = q'_* \circ j_* \implies q''_* = j'^{-1}_* \circ q'_* \circ j_*$$

q''_* is an isomorphism

Cone over A , CA

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (excision)

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (excision)

$H_n(X, A) \rightarrow H_n(X \cup CA \setminus p, CA \setminus p)$ is an isomorphism

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (excision)

$H_n(X, A) \rightarrow H_n(X \cup CA \setminus p, CA \setminus p)$ is an isomorphism (exercise!)

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (excision)

$H_n(X, A) \rightarrow H_n(X \cup CA \setminus p, CA \setminus p)$ is an isomorphism (exercise!)

$\tilde{H}_n(X \cup CA) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (excision)

$H_n(X, A) \rightarrow H_n(X \cup CA \setminus p, CA \setminus p)$ is an isomorphism (exercise!)

$\tilde{H}_n(X \cup CA) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (exercise!!)

Cone over A , $CA := A \times [0, 1]/(A \times \{0\})$

$$p := A \times \{0\}/A \times \{0\}$$

1. $A \hookrightarrow CA$ (identified with $A \times 1$)
2. A deformation retracts to p .
3. $A \setminus p$ deformation retracts to A .

$H_n(X \cup CA \setminus p, CA \setminus p) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (excision)

$H_n(X, A) \rightarrow H_n(X \cup CA \setminus p, CA \setminus p)$ is an isomorphism (exercise!)

$\tilde{H}_n(X \cup CA) \rightarrow H_n(X \cup CA, CA)$ is an isomorphism (exercise!!)

$$\boxed{H_n(X, A) \text{ is isomorphic to } \tilde{H}_n(X \cup CA)}$$

$$f : X \rightarrow Y$$

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism}$$

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{array}{l} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} \implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} \end{array}$$

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\begin{aligned} \cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \\ \cdots \end{aligned}$$

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Not in the syllabus

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

$$f : X \rightarrow Y$$

Assume X is also a CW complex

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

$$f : X \rightarrow Y$$

Assume X is also a CW complex and $\pi_1(X) = 0$, then

$$X \xrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Assume X is also a CW complex and $\pi_1(X) = 0$, then

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism for all } n &\implies \\ H_n(M_f, X) = 0 \text{ for all } n \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Assume X is also a CW complex and $\pi_1(X) = 0$, then

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism for all } n &\implies \\ H_n(M_f, X) = 0 \text{ for all } n &\implies \pi_n(M_f, X) = 0 \text{ for all } n \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

$$f : X \rightarrow Y$$

$$X \xhookrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

Assume X is also a CW complex and $\pi_1(X) = 0$, then

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism for all } n &\implies \\ H_n(M_f, X) = 0 \text{ for all } n &\implies \pi_n(M_f, X) = 0 \text{ for} \\ \text{all } n &\implies \pi_n(X) = \pi_n(M_f) \end{aligned}$$

$$f : X \rightarrow Y$$

$$X \xrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

Assume X is also a CW complex and $\pi_1(X) = 0$, then

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism for all } n &\implies \\ H_n(M_f, X) = 0 \text{ for all } n &\implies \pi_n(M_f, X) = 0 \text{ for} \\ \text{all } n &\implies \pi_n(X) = \pi_n(M_f) = \pi_n(Y) \end{aligned}$$

$$f : X \rightarrow Y$$

$$X \xrightarrow{i} M_f \xrightarrow{r} Y$$

$$f = r \circ i$$

r_* an isomorphism (r deformation retract)

$$\begin{aligned} H_n(X) \xrightarrow{f_*} H_n(Y) \text{ isomorphism} &\implies H_n(X) \xrightarrow{i_*} \\ H_n(M_f) \text{ isomorphism} &\implies H_n(M_f, X) = 0 \end{aligned}$$

Fact 1:

$$\cdots \rightarrow \pi_n(A) \rightarrow \pi_n(X) \rightarrow \pi_n(X, A) \rightarrow \pi_{n-1}(A) \rightarrow \cdots$$

Fact 2:

If $\pi_1(A) = 0$, then first non-zero $H_n(X, A)$ is the first non-zero $\pi_n(X, A)$ and they are isomorphic.

Fact 3:

$f : X \rightarrow Y$, X and Y "CW complexes", and $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ isomorphisms, then X is homotopically equivalent to Y .

Assume X is also a CW complex and $\pi_1(X) = 0$, then

$H_n(X) \xrightarrow{f_*} H_n(Y)$ isomorphism for all $n \implies H_n(M_f, X) = 0$ for all $n \implies \pi_n(M_f, X) = 0$ for all $n \implies \pi_n(X) = \pi_n(M_f) = \pi_n(Y) \implies X$ is homotopically equivalent to Y

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .



Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .



Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,
 $H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$

□

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1}) \quad \square$$

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

□

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

□

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $U \subset \mathbb{R}^n$ open and homeomorphic to $V \subset \mathbb{R}^n$ open, then $m = n$.

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $U \subset \mathbb{R}^n$ open and homeomorphic to $V \subset \mathbb{R}^n$ open, then $m = n$.

Proof. $H_k(U, U \setminus \{x\}) \xrightarrow{i_*} H_k(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\})$ □

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $U \subset \mathbb{R}^n$ open and homeomorphic to $V \subset \mathbb{R}^n$ open, then $m = n$.

Proof. $H_k(U, U \setminus \{x\}) \xrightarrow{i_*} H_k(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(\mathbb{R}^n \setminus \{x\})$ □

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $U \subset \mathbb{R}^n$ open and homeomorphic to $V \subset \mathbb{R}^n$ open, then $m = n$.

Proof. $H_k(U, U \setminus \{x\}) \xrightarrow{i_*} H_k(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(\mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(S^{n-1})$

□

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $U \subset \mathbb{R}^n$ open and homeomorphic to $V \subset \mathbb{R}^n$ open, then $m = n$.

Proof. $H_k(U, U \setminus \{x\}) \xrightarrow{i_*} H_k(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(\mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(S^{n-1})$

$f : U \rightarrow V$ homeomorphism induces, □

Theorem. $\tilde{H}_i(S^n)$ is \mathbb{Z} if $i = n$, otherwise, trivial

Proof. $\tilde{H}_i(D^n) = 0$ for all i .

$\tilde{H}_i(D^k) = 0$ for all i .

By the long exact sequence,

$$\tilde{H}_i(S^n) = \tilde{H}_i(D^n/S^{n-1}) = H_i(D^n, S^{n-1}) = \tilde{H}_{i-1}(S^{n-1})$$

$\tilde{H}_i(S^0) = \mathbb{Z}$ if $i = 0$, otherwise, trivial

Induction! □

Theorem. $U \subset \mathbb{R}^n$ open and homeomorphic to $V \subset \mathbb{R}^n$ open, then $m = n$.

Proof. $H_k(U, U \setminus \{x\}) \xrightarrow{i_*} H_k(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(\mathbb{R}^n \setminus \{x\}) \simeq H_{k-1}(S^{n-1})$

$f : U \rightarrow V$ homeomorphism induces,

$$f_* : H_n(U, U \setminus \{x\}) \rightarrow H_n(U, U \setminus \{f(x)\}) \quad \square$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$, □

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,

$I_1 \cap I_2 = S^0$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$U = S^2 \setminus f(I_1)$
 $V = S^2 \setminus f(I_2)$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0)$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) \quad \square$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\tilde{H}_0(S^2 \setminus f(S^1))$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\tilde{H}_0(U \cap V)$$



Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, □

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V)$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V)$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

□

Jordan curve theorem

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$

□

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$U = S^2 \setminus f(I_1)$$

$$V = S^2 \setminus f(I_2)$$

$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$

$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$$U = S^2 \setminus f(I_1^{1/2})$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$$U = S^2 \setminus f(I_1^{1/2})$$
$$V = S^2 \setminus f(I_2^{1/2})$$

□

$$U = S^2 \setminus f(I_1)$$
$$V = S^2 \setminus f(I_2)$$
$$U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$$
$$U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$U = S^2 \setminus f(I_1^{1/2})$
 $V = S^2 \setminus f(I_2^{1/2})$
 $U \cup V = S^2 \setminus f(p)$

□

$U = S^2 \setminus f(I_1)$
 $V = S^2 \setminus f(I_2)$
 $U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$
 $U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$

$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$U = S^2 \setminus f(I_1^{1/2})$
 $V = S^2 \setminus f(I_2^{1/2})$
 $U \cup V = S^2 \setminus f(p) = \mathbb{R}^2,$ □

$U = S^2 \setminus f(I_1)$
 $V = S^2 \setminus f(I_2)$
 $U \cup V = S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1$
 $U \cap V = S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1)$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$
 $U = S^2 \setminus f(I_1^{1/2})$
 $V = S^2 \setminus f(I_2^{1/2})$
 $U \cup V = S^2 \setminus f(p) = \mathbb{R}^2$, so $(\tilde{H}_i(U \cup V) = 0)$

□

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

$$\begin{aligned} \text{Proof of claim: } I &= I_1^{1/2} \cup I_2^{1/2}, \\ I_1^{1/2} \cap I_2^{1/2} &= \underbrace{\{1/2\}}_p \\ U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$$\begin{aligned} U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism}$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

$$\begin{aligned} \text{Proof of claim: } I &= I_1^{1/2} \cup I_2^{1/2}, \\ I_1^{1/2} \cap I_2^{1/2} &= \underbrace{\{1/2\}}_p \\ U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$\begin{aligned} H_i(U \cap V) &\xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism} \\ [x] &\rightarrow ([i_U(x)], [-i_V(x)]) \end{aligned}$$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$$\begin{aligned} U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$H_i(U \cap V) \xrightarrow{i_*} H_i(U) \oplus H_i(V)$ is an isomorphism
 $[x] \rightarrow ([i_U(x)], [-i_V(x)])$
 $[x] = [0]$ if and only if $[i_U(x)] = 0$ AND $[i_V(x)] = 0$

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\begin{aligned} \cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) &\rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \\ \tilde{H}_0(U) \oplus \tilde{H}_0(V) &\rightarrow \cdots \end{aligned}$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

Proof of claim: $I = I_1^{1/2} \cup I_2^{1/2}$,
 $I_1^{1/2} \cap I_2^{1/2} = \underbrace{\{1/2\}}_p$

$$\begin{aligned} U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$\begin{aligned} H_i(U \cap V) &\xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism} \\ [x] &\rightarrow ([i_U(x)], [-i_V(x)]) \\ [x] &= [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0 \\ [x] \neq [0] &\implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0 \end{aligned}$$

Repeatedly zooming in and subdividing, $x \in C_i(S^2 \setminus f(q))$
 (Exercise!)

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\begin{aligned} \cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) &\rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \\ \tilde{H}_0(U) \oplus \tilde{H}_0(V) &\rightarrow \cdots \end{aligned}$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

$$\begin{aligned} \textit{Proof of claim: } I &= I_1^{1/2} \cup I_2^{1/2}, \\ I_1^{1/2} \cap I_2^{1/2} &= \underbrace{\{1/2\}}_p \end{aligned}$$

$$\begin{aligned} U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$\begin{aligned} H_i(U \cap V) &\xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism} \\ [x] &\rightarrow ([i_U(x)], [-i_V(x)]) \\ [x] = [0] &\text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0 \\ [x] \neq [0] &\implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0 \end{aligned}$$

Repeatedly zooming in and subdividing, $x \in C_i(S^2 \setminus f(q))$
 (Exercise!)

Therefore, $x = \partial y$.

□

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\begin{aligned} \cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \\ \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots \end{aligned}$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

$$\begin{aligned} \textit{Proof of claim: } I &= I_1^{1/2} \cup I_2^{1/2}, \\ I_1^{1/2} \cap I_2^{1/2} &= \underbrace{\{1/2\}}_p \end{aligned}$$

$$\begin{aligned} U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$\begin{aligned} H_i(U \cap V) &\xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism} \\ [x] &\rightarrow ([i_U(x)], [-i_V(x)]) \\ [x] &= [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0 \\ [x] \neq [0] &\implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0 \end{aligned}$$

Repeatedly zooming in and subdividing, $x \in C_i(S^2 \setminus f(q))$
 (Exercise!)

Therefore, $x = \partial y$.
 But $y \in C_n(S^2 \setminus I_k^{1/2^m})$ for some m (Exercise) □

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\begin{aligned} \cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \\ \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots \end{aligned}$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

$$\begin{aligned} \textit{Proof of claim: } I &= I_1^{1/2} \cup I_2^{1/2}, \\ I_1^{1/2} \cap I_2^{1/2} &= \underbrace{\{1/2\}}_p \\ U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$\begin{aligned} H_i(U \cap V) &\xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism} \\ [x] &\rightarrow ([i_U(x)], [-i_V(x)]) \\ [x] = [0] &\text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0 \\ [x] \neq [0] &\implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0 \end{aligned}$$

Repeatedly zooming in and subdividing, $x \in C_i(S^2 \setminus f(q))$
 (Exercise!)

Therefore, $x = \partial y$.
 But $y \in C_n(S^2 \setminus I_k^{1/2^m})$ for some m (Exercise)
 Therefore, $[x] = [0] \in H_n(S^2 \setminus I_k^{1/2^m})$, □

Jordan curve theorem

Theorem. $f : S^1 \rightarrow S^2$ homeomorphism, then $S^2 \setminus f(S^1)$ has two components, i.e. $H_0(S^2 \setminus f(S^1)) = \mathbb{Z}$.

Proof. $S^1 = I_1 \cup I_2$,
 $I_1 \cap I_2 = S^0$

$$\begin{aligned} U &= S^2 \setminus f(I_1) \\ V &= S^2 \setminus f(I_2) \\ U \cup V &= S^2 \setminus f(S^0) = S^2 \setminus \{f(-1), f(1)\} \simeq S^1 \\ U \cap V &= S^2 \setminus (f(I_1) \cup f(I_2)) = S^2 \setminus f(S^1) \end{aligned}$$

$$\begin{aligned} \cdots \rightarrow \tilde{H}_1(U) \oplus \tilde{H}_1(V) \rightarrow \tilde{H}_1(U \cup V) \rightarrow \tilde{H}_0(U \cap V) \rightarrow \\ \tilde{H}_0(U) \oplus \tilde{H}_0(V) \rightarrow \cdots \end{aligned}$$

Claim: $H_i(S^2 \setminus f(I)) = 0$, i.e. $\tilde{H}_i(U) = \tilde{H}_i(V) = 0$

$$\mathbb{Z} = \tilde{H}_1(S^1) = \tilde{H}_1(U \cup V) = \tilde{H}_0(U \cap V) = \tilde{H}_0(S^2 \setminus f(S^1))$$

$$\begin{aligned} \textit{Proof of claim: } I &= I_1^{1/2} \cup I_2^{1/2}, \\ I_1^{1/2} \cap I_2^{1/2} &= \underbrace{\{1/2\}}_p \end{aligned}$$

$$\begin{aligned} U &= S^2 \setminus f(I_1^{1/2}) \\ V &= S^2 \setminus f(I_2^{1/2}) \\ U \cup V &= S^2 \setminus f(p) = \mathbb{R}^2, \text{ so } (\tilde{H}_i(U \cup V) = 0) \\ U \cap V &= S^2 \setminus f(I) \end{aligned}$$

$$\begin{aligned} H_i(U \cap V) &\xrightarrow{i_*} H_i(U) \oplus H_i(V) \text{ is an isomorphism} \\ [x] &\rightarrow ([i_U(x)], [-i_V(x)]) \\ [x] &= [0] \text{ if and only if } [i_U(x)] = 0 \text{ AND } [i_V(x)] = 0 \\ [x] \neq [0] &\implies [i_U(x)] \neq 0 \text{ OR } [i_V(x)] \neq 0 \end{aligned}$$

Repeatedly zooming in and subdividing, $x \in C_i(S^2 \setminus f(q))$
 (Exercise!)

Therefore, $x = \partial y$.
 But $y \in C_n(S^2 \setminus I_k^{1/2^m})$ for some m (Exercise)
 Therefore, $[x] = [0] \in H_n(S^2 \setminus I_k^{1/2^m})$, contradicting $[x] \neq [0]$ □