

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\mathbb{Z} \otimes \mathbb{Z}/2 \rightarrow \mathbb{Z} \otimes \mathbb{Z}/2 \rightarrow \mathbb{Z}/2 \otimes \mathbb{Z}/2 \rightarrow 0$$

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\mathbb{Z} \otimes \mathbb{Z}/2 \rightarrow \mathbb{Z} \otimes \mathbb{Z}/2 \rightarrow \mathbb{Z}/2 \otimes \mathbb{Z}/2 \rightarrow 0$$

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow \mathbb{Z} \otimes \mathbb{Z}/2 \rightarrow \mathbb{Z}/2 \otimes \mathbb{Z}/2 \rightarrow 0$$

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow \underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow \mathbb{Z}/2 \otimes \mathbb{Z}/2 \rightarrow 0$$

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow \underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow \underbrace{\mathbb{Z}/2 \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow 0$$

**Example.**  $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$

$$\underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \xrightarrow{0} \underbrace{\mathbb{Z} \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow \underbrace{\mathbb{Z}/2 \otimes \mathbb{Z}/2}_{\mathbb{Z}/2} \rightarrow 0$$

**Exercise.** *If*  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$



**Exercise.** *If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$*

**Exercise.** *If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups*

**Exercise.** *If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups*

**Definition.** *A free resolution of an abelian group,  $H$ ,*

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:

$$\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0,$$

where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,

$$\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$$

is a chain complex  
(exercise!)

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:

$$\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0,$$

where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,

$$\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$$

is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$



**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:

$$\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0,$$

where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,

$$\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$$

is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** Prove:

$$1. Tor_1(H_1 \oplus H_2, G) = Tor_1(H_2, G) \oplus Tor_1(H_1, G)$$

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** Prove:

1.  $Tor_1(H_1 \oplus H_2, G) = Tor_1(H_2, G) \oplus Tor_1(H_1, G)$
2.  $Tor_1(\mathbb{Z}, G) = 0$

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** Prove:

1.  $Tor_1(H_1 \oplus H_2, G) = Tor_1(H_2, G) \oplus Tor_1(H_1, G)$
2.  $Tor_1(\mathbb{Z}, G) = 0$
3.  $Tor_1(\mathbb{Z}/n, G) = \ker (G \xrightarrow{\times n} G)$

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** Prove:

1.  $Tor_1(H_1 \oplus H_2, G) = Tor_1(H_2, G) \oplus Tor_1(H_1, G)$
2.  $Tor_1(\mathbb{Z}, G) = 0$
3.  $Tor_1(\mathbb{Z}/n, G) = \ker (G \xrightarrow{\times n} G)$
4. all the lemmas and corollary on this page!



**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** Prove:

1.  $Tor_1(H_1 \oplus H_2, G) = Tor_1(H_2, G) \oplus Tor_1(H_1, G)$
2.  $Tor_1(\mathbb{Z}, G) = 0$
3.  $Tor_1(\mathbb{Z}/n, G) = \ker (G \xrightarrow{\times n} G)$
4. all the lemmas and corollary on this page!

**Exercise.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$   
then,  $A \otimes G \rightarrow B \otimes G \rightarrow C \otimes G \rightarrow 0$   
 $A, B, C, G$  are abelian groups

**Definition.** A free resolution of an abelian group,  $H$ ,  
is an exact sequence:  
 $\cdots \rightarrow F_i \rightarrow F_{i-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$ ,  
where  $F_i$  are free

**Lemma.** If,  $\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0$   
is a free resolution, then,  
 $\cdots \rightarrow F_i \otimes G \rightarrow \cdots \rightarrow F_1 \otimes G \rightarrow F_0 \otimes G \rightarrow H \otimes G \rightarrow 0$   
is a chain complex  
(exercise!)

**Lemma.** Any map  $H \rightarrow H'$  extends to a chain map

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & F'_1 & \longrightarrow & F'_0 & \longrightarrow & H' \longrightarrow 0 \end{array}$$

unique upto chain homotopy

**Lemma.** Any two free resolutions of the same group  
are chain homotopic

**Corollary.** Given a free resolution,  
 $\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \cdots \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H$ , consider the  
chain complex,  
 $\cdots \rightarrow F_i \otimes G \xrightarrow{f_i \otimes Id} \cdots \rightarrow F_1 \otimes G \xrightarrow{f_1 \otimes Id} F_0 \otimes G \xrightarrow{f_0 \otimes Id} H \otimes G$   
 $Tor_i(H, G) := \ker f_i \otimes Id / \text{Im } f_{i+1} \otimes Id$  depends only  
on  $H$  and  $G$ .

**Exercise.** Prove:

1.  $Tor_1(H_1 \oplus H_2, G) = Tor_1(H_2, G) \oplus Tor_1(H_1, G)$
2.  $Tor_1(\mathbb{Z}, G) = 0$
3.  $Tor_1(\mathbb{Z}/n, G) = \ker (G \xrightarrow{\times n} G)$
4. all the lemmas and corollary on this page!

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
 & & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
 \cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
 \end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

$$\begin{array}{ccccc}
 \cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
 & & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
 \cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
 \end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

$$\begin{array}{ccccc}
 \cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
 & & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
 \cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
 \end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,  
 $\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))$ ?

Assume,  $D_k$  defined for  $k < n$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,  
 $\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))$ ?

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x)))$$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,  
 $\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))$ ?

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
= & \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x))
\end{aligned}$$



$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
&= \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - \partial'_n D_{n-1}(\partial_n(x))
\end{aligned}$$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
= & \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x)) \\
= & \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - \partial'_n D_{n-1}(\partial_n(x)) \\
= & \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - (\alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - D_{n-2}(\partial_{n-1} \partial_n(x)))
\end{aligned}$$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
&= \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - (\alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - D_{n-2}(\partial_{n-1} \partial_n(x))) \\
&= 0
\end{aligned}$$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
&= \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - (\alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - D_{n-2}(\partial_{n-1} \partial_n(x))) \\
&= 0
\end{aligned}$$

Therefore, if  $H_n(C'_*) = 0$ , then  $\partial'_{n+1}(x') = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))$  for some  $x'$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
&= \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - (\alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - D_{n-2}(\partial_{n-1} \partial_n(x))) \\
&= 0
\end{aligned}$$

Therefore, if  $H_n(C'_*) = 0$ , then  $\partial'_{n+1}(x') = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))$  for some  $x'$ . Define,  $D_n(x) = x'$

$$\begin{array}{ccccc}
\cdots & \longrightarrow & C_2 & \longrightarrow & C_1 & \longrightarrow & C_0 \\
& & \downarrow \alpha_2, \beta_2 & & \downarrow \alpha_1, \beta_1 & & \downarrow \alpha_0, \beta_0 \\
\cdots & \longrightarrow & C'_2 & \longrightarrow & C'_1 & \longrightarrow & C'_0
\end{array}$$

$$\alpha_n - \beta_n = \partial'_{n+1} \circ D_n + D_{n-1} \circ \partial_n$$

Can we define  $D_n$  so that,

$$\partial'_{n+1}(D_n(x)) = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))?$$

Assume,  $D_k$  defined for  $k < n$ . Observe,

$$\begin{aligned}
& \partial'_n(\alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))) \\
&= \partial'_n \alpha_n(x) - \partial'_n \beta_n(x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - \partial'_n D_{n-1}(\partial_n(x)) \\
&= \alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - (\alpha_{n-1}(\partial_n x) - \beta_{n-1}(\partial_n x) - D_{n-2}(\partial_{n-1} \partial_n(x))) \\
&= 0
\end{aligned}$$

Therefore, if  $H_n(C'_*) = 0$ , then  $\partial'_{n+1}(x') = \alpha_n(x) - \beta_n(x) - D_{n-1}(\partial_n(x))$  for some  $x'$ . Define,  $D_n(x) = x'$

**Lemma.** *If  $C_*$  and  $C'_*$  are chain complexes and  $H_k(C'_*) = 0$  for all  $k$ , then any chain maps  $\alpha, \beta : C_* \rightarrow C'_*$  are chain homotopic.*

$$\cdots \rightarrow C_n(X) \xrightarrow{\partial_n} C_{n-1}(X) \rightarrow \cdots \rightarrow C_1(X) \xrightarrow{\partial_1} C_0(X)$$

$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian



$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian

$$(\partial_n \otimes Id) \circ (\partial_{n-1} \otimes Id) = 0$$

$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian

$$(\partial_n \otimes Id) \circ (\partial_{n-1} \otimes Id) = 0$$

$$Z_n(X; G) := \ker (\partial_n \otimes Id)$$

$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian

$$(\partial_n \otimes Id) \circ (\partial_{n-1} \otimes Id) = 0$$

$$Z_n(X; G) := \ker (\partial_n \otimes Id)$$

$$B_n(X; G) := \operatorname{Im} (\partial_{n+1} \otimes Id)$$

$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian

$$(\partial_n \otimes Id) \circ (\partial_{n-1} \otimes Id) = 0$$

$$Z_n(X; G) := \ker (\partial_n \otimes Id)$$

$$B_n(X; G) := \operatorname{Im} (\partial_{n+1} \otimes Id)$$

$$H_n(X; G) := Z_n(X; G) / B_n(X; G)$$

$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian

$$(\partial_n \otimes Id) \circ (\partial_{n-1} \otimes Id) = 0$$

$$Z_n(X; G) := \ker (\partial_n \otimes Id)$$

$$B_n(X; G) := \operatorname{Im} (\partial_{n+1} \otimes Id)$$

$$H_n(X; G) := Z_n(X; G) / B_n(X; G)$$

$$H_n(X) \otimes G \rightarrow H_n(X; G)$$

$$\cdots \rightarrow C_n(X) \otimes G \xrightarrow{\partial_n \otimes Id} C_{n-1}(X) \otimes G \rightarrow \cdots \rightarrow C_1(X) \otimes G \xrightarrow{\partial_1 \otimes Id} C_0(X) \otimes G$$

$G$  abelian

$$(\partial_n \otimes Id) \circ (\partial_{n-1} \otimes Id) = 0$$

$$Z_n(X;G) := ker \; (\partial_n \otimes Id)$$

$$B_n(X;G) := Im \; (\partial_{n+1} \otimes Id)$$

$$H_n(X;G) := Z_n(X;G)/B_n(X;G)$$

$$H_n(X) \otimes G \rightarrow H_n(X;G) \; [c] \otimes g \rightarrow [c \otimes g]$$

$$0 \longrightarrow Z_{n+1} \longrightarrow C_{n+1} \longrightarrow B_n \longrightarrow 0$$

$$0 \longrightarrow Z_{n+1} \longrightarrow C_{n+1} \longrightarrow B_n \longrightarrow 0$$

$$0 \longrightarrow Z_n \longrightarrow C_n \longrightarrow B_{n-1} \longrightarrow 0$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & & & \downarrow \partial & & \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial & & \downarrow \partial & & \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \rightarrow Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$b_n = \partial c_{n+1}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \rightarrow Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \rightarrow Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \overset{i}{\rightarrow} Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \overset{i}{\rightarrow} Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$

$$Z_n \overset{j_*}{\rightarrow} H_n(C_*) \rightarrow \ker i \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \overset{i}{\rightarrow} Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$

$$Z_n \overset{j_*}{\rightarrow} H_n(C_*) \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \overset{i}{\rightarrow} Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$

$$0 \rightarrow Z_n / \ker j \overset{j_*}{\rightarrow} H_n(C_*) \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \overset{i}{\rightarrow} Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
&b_n = \partial c_{n+1} \\
&b_n \in B_n \rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$

$$0 \rightarrow Z_n/i(B_n) \overset{j_*}{\rightarrow} H_n(C_*) \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \longrightarrow & C_{n+1} & \longrightarrow & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0
\end{array}$$

$$\cdots \rightarrow B_n \overset{i}{\rightarrow} Z_n \overset{j}{\rightarrow} H_n(C_*) \rightarrow B_{n-1} \rightarrow \cdots$$

$$\begin{aligned}
b_n &= \partial c_{n+1} \\
b_n \in B_n &\rightarrow \partial c_{n+1} = b_n \in Z_n
\end{aligned}$$

$$\text{Therefore, } b_n \in B_n \rightarrow b_n \in Z_n$$

$$0 \rightarrow H_n(C_*) \overset{j_*}{\rightarrow} H_n(C_*) \rightarrow 0$$

$$0 \longrightarrow Z_{n+1} \xrightarrow{j} C_{n+1} \xrightarrow{\partial} B_n \longrightarrow 0$$

$$0 \longrightarrow Z_{n+1} \xrightarrow{j} C_{n+1} \xrightarrow{\partial} B_n \longrightarrow 0$$

$$0 \longrightarrow Z_n \xrightarrow{j} C_n \xrightarrow{\partial} B_{n-1} \longrightarrow 0$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \xrightarrow{j} & C_{n+1} & \xrightarrow{\partial} & B_n \longrightarrow 0 \\
& & & & \downarrow \partial & & \\
0 & \longrightarrow & Z_n & \xrightarrow{j} & C_n & \xrightarrow{\partial} & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \xrightarrow{j} & C_{n+1} & \xrightarrow{\partial} & B_n \longrightarrow 0 \\
& & \downarrow \partial & & \downarrow \partial & & \\
0 & \longrightarrow & Z_n & \xrightarrow{j} & C_n & \xrightarrow{\partial} & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \xrightarrow{j} & C_{n+1} & \xrightarrow{\partial} & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \\
0 & \longrightarrow & Z_n & \xrightarrow{j} & C_n & \xrightarrow{\partial} & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \xrightarrow{j} & C_{n+1} & \xrightarrow{\partial} & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial \\
0 & \longrightarrow & Z_n & \xrightarrow{j} & C_n & \xrightarrow{\partial} & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} & \xrightarrow{j} & C_{n+1} & \xrightarrow{\partial} & B_n \longrightarrow 0 \\
& & \downarrow \partial=0 & & \downarrow \partial & & \downarrow \partial=0 \\
0 & \longrightarrow & Z_n & \xrightarrow{j} & C_n & \xrightarrow{\partial} & B_{n-1} \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots H_n(Z_* \otimes G)$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots H_n(Z_* \otimes G) \rightarrow H_n(C_* \otimes G)$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots H_n(Z_* \otimes G) \rightarrow H_n(C_* \otimes G) \rightarrow H_n(B_{*-1} \otimes G)$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots H_n(Z_* \otimes G) \rightarrow H_n(C_* \otimes G) \rightarrow H_n(B_{*-1} \otimes G) \rightarrow H_{n-1}(Z_* \otimes G) \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C_* \otimes G) \rightarrow H_n(B_{*-1} \otimes G) \rightarrow H_{n-1}(Z_* \otimes G) \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow H_n(B_{*-1} \otimes G) \rightarrow H_{n-1}(Z_* \otimes G) \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \rightarrow H_{n-1}(Z_* \otimes G) \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \rightarrow Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \overset{??}{\rightarrow} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b} \otimes g} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \overset{??}{\rightarrow} Z_{n-1} \otimes G \rightarrow \cdots$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \overset{??}{\rightarrow} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \overset{??}{\rightarrow} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \overset{??}{\rightarrow} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \overset{??}{\rightarrow} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1}^{\textcolor{blue}{c \otimes g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b \otimes g = \partial c \otimes g}} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n^{\textcolor{blue}{b \otimes g}} \otimes G & \xrightarrow{j \otimes Id} & C_n^{\textcolor{blue}{\partial c \otimes g = b \otimes g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow Ker \ (\partial \otimes Id)_* \rightarrow H_n(C; G) \rightarrow Im \ (\partial \otimes Id)_* \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1}^{\textcolor{blue}{c} \otimes g} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b} \otimes g = \partial c \otimes g} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n^{\textcolor{blue}{b} \otimes g} \otimes G & \xrightarrow{j \otimes Id} & C_n^{\textcolor{blue}{\partial c} \otimes g = \textcolor{blue}{b} \otimes g} \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow Ker \ (\partial \otimes Id)_* \rightarrow H_n(C; G) \rightarrow Ker \ (i \otimes Id) \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow Im \, (j \otimes Id)_* \rightarrow H_n(C; G) \rightarrow Ker \, (i \otimes Id) \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow Z_n \otimes G / Im \ (i \otimes Id) \rightarrow H_n(C; G) \rightarrow Ker \ (i \otimes Id) \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow (Z_n / i(B_n)) \otimes G \rightarrow H_n(C; G) \rightarrow Ker (i \otimes Id) \rightarrow 0$$

because,

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow (Z_n / i(B_n)) \otimes G \rightarrow H_n(C; G) \rightarrow Ker (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n / B_n \rightarrow 0$$



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$\text{Labels: } C_{n+1} \text{ has } c \otimes g, B_n \text{ has } b \otimes g = \partial c \otimes g, C_n \text{ has } \partial c \otimes g = b \otimes g$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow (Z_n / i(B_n)) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker } (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n / B_n \rightarrow 0$$

$$\implies B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow (Z_n / B_n) \otimes G \rightarrow 0$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$\text{Labels: } C_{n+1} \otimes G \text{ has } c \otimes g; B_n \otimes G \text{ has } b \otimes g = \partial c \otimes g; C_n \otimes G \text{ has } \partial c \otimes g = b \otimes g$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow (Z_n / i(B_n)) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker } (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n / B_n \rightarrow 0$$

$$\implies B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow (Z_n / B_n) \otimes G \rightarrow 0$$

$$\implies Z_n \otimes G / \text{Im } (i \otimes Id) \cong (Z_n / B_n) \otimes G$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1}^{\textcolor{blue}{c} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b} \otimes \textcolor{blue}{g} = \partial \textcolor{blue}{c} \otimes \textcolor{blue}{g}} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n^{\textcolor{blue}{b} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{j \otimes Id} & C_n^{\textcolor{blue}{\partial c} \otimes \textcolor{blue}{g} = \textcolor{blue}{b} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow (Z_n / i(B_n)) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n / B_n \rightarrow 0$$

$$\implies B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow (Z_n / B_n) \otimes G \rightarrow 0$$

$$\implies Z_n \otimes G / \text{Im} (i \otimes Id) \cong (Z_n / B_n) \otimes G \cong H_n(C) \otimes G$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1}^{\textcolor{blue}{c} \otimes g} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b} \otimes g = \partial c \otimes g} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n^{\textcolor{blue}{b} \otimes g} \otimes G & \xrightarrow{j \otimes Id} & C_n^{\textcolor{blue}{\partial c} \otimes g = \textcolor{blue}{b} \otimes g} \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow Z_n \otimes G / \text{Im} (i \otimes Id) \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n / B_n \rightarrow 0$$

$$\implies B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow (Z_n / B_n) \otimes G \rightarrow 0$$

$$\implies Z_n \otimes G / \text{Im} (i \otimes Id) \cong (Z_n / B_n) \otimes G \cong H_n(C) \otimes G$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1}^{\textcolor{blue}{c} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b} \otimes \textcolor{blue}{g} = \partial \textcolor{blue}{c} \otimes \textcolor{blue}{g}} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n^{\textcolor{blue}{b} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{j \otimes Id} & C_n^{\textcolor{blue}{\partial c} \otimes \textcolor{blue}{g} = \textcolor{blue}{b} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow (Z_n / i(B_n)) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n / B_n \rightarrow 0$$

$$\implies B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow (Z_n / B_n) \otimes G \rightarrow 0$$

$$\implies Z_n \otimes G / \text{Im} (i \otimes Id) \cong (Z_n / B_n) \otimes G \cong H_n(C) \otimes G$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1}^{\textcolor{blue}{c} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_n^{\textcolor{blue}{b} \otimes \textcolor{blue}{g} = \partial \textcolor{blue}{c} \otimes \textcolor{blue}{g}} \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n^{\textcolor{blue}{b} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{j \otimes Id} & C_n^{\textcolor{blue}{\partial c} \otimes \textcolor{blue}{g} = \textcolor{blue}{b} \otimes \textcolor{blue}{g}} \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

because,

$$0 \rightarrow B_n \xrightarrow{i} Z_n \rightarrow Z_n/B_n \rightarrow 0$$

$$\implies B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow (Z_n/B_n) \otimes G \rightarrow 0$$

$$\implies Z_n \otimes G / \text{Im} (i \otimes Id) \cong (Z_n/B_n) \otimes G \cong H_n(C) \otimes G$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

Note,  $0 \rightarrow B_{n-1} \rightarrow Z_{n-1} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

Note,  $0 \rightarrow B_{n-1} \rightarrow Z_{n-1} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )



$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \otimes G & \xrightarrow{\partial \otimes Id} & B_n \otimes G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \otimes G & \xrightarrow{j \otimes Id} & C_n \otimes G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$\text{Labels: } C_{n+1} \text{ has } c \otimes g, B_n \text{ has } b \otimes g = \partial c \otimes g, Z_n \text{ has } b \otimes g, C_n \text{ has } \partial c \otimes g = b \otimes g$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

Note,  $0 \rightarrow \underbrace{B_{n-1}}_{F_1} \rightarrow \underbrace{Z_{n-1}}_{F_0} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

Note,  $0 \rightarrow \underbrace{B_{n-1}}_{F_1} \rightarrow \underbrace{Z_{n-1}}_{F_0} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )

$0 \rightarrow B_{n-1} \otimes G \xrightarrow{i_{n-1} \otimes Id} Z_{n-1} \otimes G \rightarrow H_{n-1}(C) \otimes G \rightarrow 0$  (only a complex. **not** exact)

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow \text{Ker} (i \otimes Id) \rightarrow 0$$

Note,  $0 \rightarrow \underbrace{B_{n-1}}_{F_1} \rightarrow \underbrace{Z_{n-1}}_{F_0} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )

$$0 \rightarrow \underbrace{B_{n-1} \otimes G}_{F_1 \otimes G} \xrightarrow{i_{n-1} \otimes Id} \underbrace{Z_{n-1} \otimes G}_{F_0 \otimes G} \rightarrow H_{n-1}(C) \otimes G \rightarrow 0 \text{ (only a complex. **not** exact)}$$

$$\text{ker} (i_{n-1} \otimes Id) = \text{Tor}_1(H_{n-1}(C), G)$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

Note,  $0 \rightarrow \underbrace{B_{n-1}}_{F_1} \rightarrow \underbrace{Z_{n-1}}_{F_0} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )

$$0 \rightarrow \underbrace{B_{n-1} \otimes G}_{F_1 \otimes G} \xrightarrow{i_{n-1} \otimes Id} \underbrace{Z_{n-1} \otimes G}_{F_0 \otimes G} \rightarrow H_{n-1}(C) \otimes G \rightarrow 0 \text{ (only a complex. **not** exact)}$$

$$\ker (i_{n-1} \otimes Id) = Tor_1(H_{n-1}(C), G)$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

Note,  $0 \rightarrow \underbrace{B_{n-1}}_{F_1} \rightarrow \underbrace{Z_{n-1}}_{F_0} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )

$$0 \rightarrow \underbrace{B_{n-1} \otimes G}_{F_1 \otimes G} \xrightarrow{i_{n-1} \otimes Id} \underbrace{Z_{n-1} \otimes G}_{F_0 \otimes G} \rightarrow H_{n-1}(C) \otimes G \rightarrow 0 \text{ (only a complex. **not** exact)}$$

$$\ker (i_{n-1} \otimes Id) = Tor_1(H_{n-1}(C), G)$$

$$\begin{array}{ccccccc}
0 & \longrightarrow & Z_{n+1} \otimes G & \xrightarrow{j \otimes Id} & C_{n+1} \overset{c \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_n \overset{b \otimes g = \partial c \otimes g}{\otimes} G \longrightarrow 0 \\
& & \downarrow \partial \otimes Id = 0 & & \downarrow \partial \otimes Id & & \downarrow \partial \otimes Id = 0 \\
0 & \longrightarrow & Z_n \overset{b \otimes g}{\otimes} G & \xrightarrow{j \otimes Id} & C_n \overset{\partial c \otimes g = b \otimes g}{\otimes} G & \xrightarrow{\partial \otimes Id} & B_{n-1} \otimes G \longrightarrow 0
\end{array}$$

$$\cdots B_n \otimes G \xrightarrow{i \otimes Id} Z_n \otimes G \rightarrow H_n(C; G) \rightarrow B_{n-1} \otimes G \xrightarrow{i \otimes Id} Z_{n-1} \otimes G \rightarrow \cdots$$

$$0 \rightarrow H_n(C) \otimes G \rightarrow H_n(C; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

Note,  $0 \rightarrow \underbrace{B_{n-1}}_{F_1} \rightarrow \underbrace{Z_{n-1}}_{F_0} \rightarrow H_{n-1}(C) \rightarrow 0$  (exact and a free resolution of  $H_{n-1}(C)$ )

$$0 \rightarrow \underbrace{B_{n-1} \otimes G}_{F_1 \otimes G} \xrightarrow{i_{n-1} \otimes Id} \underbrace{Z_{n-1} \otimes G}_{F_0 \otimes G} \rightarrow H_{n-1}(C) \otimes G \rightarrow 0 \text{ (only a complex. **not** exact)}$$

$$\ker (i_{n-1} \otimes Id) = Tor_1(H_{n-1}(C), G)$$

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$0 \longrightarrow H_n(C_*) \otimes G \longrightarrow H_n(C_*; G) \longrightarrow \operatorname{Tor}_1(H_{n-1}(C), G) \longrightarrow 0$$



**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$0 \longrightarrow H_n(C_*) \otimes G \longrightarrow H_n(C_*; G) \longrightarrow \text{Tor}_1(H_{n-1}(C), G) \longrightarrow 0$$

$$0 \longrightarrow H_n(C'_*) \otimes G \longrightarrow H_n(C'_*; G) \longrightarrow \text{Tor}_1(H_{n-1}(C'), G) \longrightarrow 0$$

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

*commutes*

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

commutes

*Proof.*  $0 \longrightarrow Z_n \longrightarrow C_n \longrightarrow B_{n-1} \longrightarrow 0$

□

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

commutes

$$0 \longrightarrow Z_n \longrightarrow C_n \longrightarrow B_{n-1} \longrightarrow 0$$

*Proof.*

$$0 \longrightarrow Z'_n \longrightarrow C'_n \longrightarrow B'_{n-1} \longrightarrow 0$$

□

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

commutes

$$0 \longrightarrow Z_n \longrightarrow C_n \longrightarrow B_{n-1} \longrightarrow 0$$

*Proof.*

$$0 \longrightarrow Z'_n \longrightarrow C'_n \longrightarrow B'_{n-1} \longrightarrow 0$$

Naturality of long exact sequence... □

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

commutes

$$0 \longrightarrow Z_n \longrightarrow C_n \longrightarrow B_{n-1} \longrightarrow 0$$

*Proof.*

$$0 \longrightarrow Z'_n \longrightarrow C'_n \longrightarrow B'_{n-1} \longrightarrow 0$$

Naturality of long exact sequence...(exercise!) □

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

commutes

$$\begin{array}{ccccccc}
 0 & \longrightarrow & Z_n & \longrightarrow & C_n & \longrightarrow & B_{n-1} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 \textit{Proof.} & & 0 & \longrightarrow & Z'_n & \longrightarrow & C'_n \longrightarrow B'_{n-1} \longrightarrow 0
 \end{array}$$

Naturality of long exact sequence...(exercise!) □

**Theorem.**  $f : C \rightarrow C'$  a chain map, then,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_n(C_*) \otimes G & \longrightarrow & H_n(C_*; G) & \longrightarrow & Tor_1(H_{n-1}(C), G) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H_n(C'_*) \otimes G & \longrightarrow & H_n(C'_*; G) & \longrightarrow & Tor_1(H_{n-1}(C'), G) \longrightarrow 0
 \end{array}$$

commutes

$$\begin{array}{ccccccc}
 0 & \longrightarrow & Z_n \otimes G & \longrightarrow & C_n \otimes G & \longrightarrow & B_{n-1} \otimes G \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 \textit{Proof.} & & 0 & \longrightarrow & Z'_n \otimes G & \longrightarrow & C'_n \otimes G \longrightarrow B'_{n-1} \otimes G \longrightarrow 0
 \end{array}$$

Naturality of long exact sequence...(exercise!) □



**Theorem.** *The short exact sequence,*  
 $0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$  *splits.*

**Theorem.** *The short exact sequence,*  
 $0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow \operatorname{Tor}_1(H_{n-1}(C), G) \rightarrow 0$  *splits.*

*Proof.*  $0 \rightarrow Z_n \rightarrow C_n \rightarrow B_{n-1} \rightarrow 0$  splits.



**Theorem.** *The short exact sequence,*  
 $0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$  *splits.*

*Proof.*  $0 \rightarrow Z_n \rightarrow C_n \rightarrow B_{n-1} \rightarrow 0$  splits.  
There exists,  $a : C_n \rightarrow Z_n$ , such that  $a \circ i = Id$



**Theorem.** *The short exact sequence,*  
 $0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$  *splits.*

*Proof.*  $0 \rightarrow Z_n \rightarrow C_n \rightarrow B_{n-1} \rightarrow 0$  splits.  
 There exists,  $a : C_n \rightarrow Z_n$ , such that  $a \circ i = Id$   
 ... □

**Theorem.** *The short exact sequence,*

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow \operatorname{Tor}_1(H_{n-1}(C), G) \rightarrow 0 \text{ splits.}$$

*Proof.*  $0 \rightarrow Z_n \rightarrow C_n \rightarrow B_{n-1} \rightarrow 0$  splits.

There exists,  $a : C_n \rightarrow Z_n$ , such that  $a \circ i = Id$

... (exercise!)



The splitting is not natural

The splitting is not natural

$$f : D^2 \rightarrow S^2 \text{ (collapse the boundary),}$$

The splitting is not natural

$$f : D^2 \rightarrow S^2 \text{ (collapse the boundary),}$$

$$\text{induces } f : \mathbb{RP}^2 \rightarrow S^2$$

$$H_2(\mathbb{RP}^2; \mathbb{Z}/2) \longrightarrow Tor_1(H_1(\mathbb{RP}^2), \mathbb{Z}/2) \oplus (H_2(\mathbb{RP}^2) \otimes \mathbb{Z}/2)$$

$$\downarrow f_*$$

$$\downarrow$$

$$H_2(S^2; \mathbb{Z}/2) \longrightarrow Tor_1(H_1(S^2), \mathbb{Z}/2) \oplus (H_2(S^2) \otimes \mathbb{Z}/2)$$



The splitting is not natural

$f : D^2 \rightarrow S^2$  (collapse the boundary),

induces  $f : \mathbb{RP}^2 \rightarrow S^2$

$$H_2(\mathbb{RP}^2; \mathbb{Z}/2) \longrightarrow Tor_1(H_1(\mathbb{RP}^2), \mathbb{Z}/2) \oplus (H_2(\mathbb{RP}^2) \otimes \mathbb{Z}/2)$$

$$\downarrow f_*$$

$$\downarrow 0$$

$$H_2(S^2; \mathbb{Z}/2) \longrightarrow Tor_1(H_1(S^2), \mathbb{Z}/2) \oplus (H_2(S^2) \otimes \mathbb{Z}/2)$$

The splitting is not natural

$f : D^2 \rightarrow S^2$  (collapse the boundary),

induces  $f : \mathbb{RP}^2 \rightarrow S^2$

$$H_2(\mathbb{RP}^2; \mathbb{Z}/2) \longrightarrow \text{Tor}_1(H_1(\mathbb{RP}^2), \mathbb{Z}/2) \oplus (H_2(\mathbb{RP}^2) \otimes \mathbb{Z}/2)$$

$$\downarrow f_*$$

$$\downarrow 0$$

$$H_2(S^2; \mathbb{Z}/2) \longrightarrow \text{Tor}_1(H_1(S^2), \mathbb{Z}/2) \oplus (H_2(S^2) \otimes \mathbb{Z}/2)$$

To show  $f_*$  is an isomorphism:

$$H_2(\mathbb{RP}^2; \mathbb{Z}/2) \longrightarrow H_2(\mathbb{RP}^2, \mathbb{RP}^2 \setminus p; \mathbb{Z}/2) \xrightarrow{\text{excision}} H_2(U, U \setminus p)$$

$$\downarrow f_*$$

$$\downarrow$$

$$\downarrow$$

$$H_2(S^2; \mathbb{Z}/2) \longrightarrow H_2(S^2, \mathbb{RP}^2 \setminus p; \mathbb{Z}/2) \xrightarrow{\text{excision}} H_2(U, U \setminus p)$$

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

**Example.**

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & otherwise \end{cases}$$

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

**Example.**

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & otherwise \end{cases}$$

$$H_k(\mathbb{RP}^2; \mathbb{Z}/2) = \begin{cases} \mathbb{Z}/2 & k = 0 \end{cases}$$

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

**Example.**

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & otherwise \end{cases}$$

$$H_k(\mathbb{RP}^2; \mathbb{Z}/2) = \begin{cases} \mathbb{Z}/2 & k = 0 \\ \mathbb{Z}/2 & k = 1 \end{cases}$$

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

**Example.**

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & otherwise \end{cases}$$

$$H_k(\mathbb{RP}^2; \mathbb{Z}/2) = \begin{cases} \mathbb{Z}/2 & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ \mathbb{Z}/2 & k = 2 \end{cases}$$

$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

**Example.**

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & otherwise \end{cases}$$

$$H_k(\mathbb{RP}^2; \mathbb{Z}/2) = \begin{cases} \mathbb{Z}/2 & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ \mathbb{Z}/2 & k = 2 \\ 0 & otherwise \end{cases}$$



$$0 \rightarrow H_n(C_*) \otimes G \rightarrow H_n(C_*; G) \rightarrow Tor_1(H_{n-1}(C), G) \rightarrow 0$$

**Example.**

$$H_k(\mathbb{RP}^2) = \begin{cases} \mathbb{Z} & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ 0 & otherwise \end{cases}$$

$$H_k(\mathbb{RP}^2; \mathbb{Z}/2) = \begin{cases} \mathbb{Z}/2 & k = 0 \\ \mathbb{Z}/2 & k = 1 \\ \mathbb{Z}/2 & k = 2 \\ 0 & otherwise \end{cases}$$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p\#} C_k(X; \mathbb{Z}/2)$$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}$ ,

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p\#} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma,$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$



$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{aligned} & p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0 \\ \implies & p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0 \end{aligned}$$

$$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2) \text{ etc.}$$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{aligned} & p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0 \\ \implies & p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X};\mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X;\mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, \, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$

$$\implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{array}{ccccc} < \textit{injt} > & C_k(X;\mathbb{Z}/2) & \xrightarrow{\tau} & C_k(\tilde{X};\mathbb{Z}/2) & \xrightarrow{p_{\#}} \\ & C_k(X;\mathbb{Z}/2) & \rightarrow & 0 & \end{array}$$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X};\mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X;\mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma$ , i.e.  $p_{\#}(\tilde{\sigma}) = \sigma$

$$\begin{aligned} p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) &= 0 \\ \implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) &= 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} < \textit{injt} > \quad C_k(X;\mathbb{Z}/2) \quad \xrightarrow{\tau} \quad C_k(\tilde{X};\mathbb{Z}/2) \quad \xrightarrow{p_{\#}} \\ C_k(X;\mathbb{Z}/2) &\rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2, \tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma$ , i.e.  $p_{\#}(\tilde{\sigma}) = \sigma$

$$\begin{aligned} p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) &= 0 \\ \implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) &= 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} < \text{ injt } > \quad C_k(X; \mathbb{Z}/2) \quad \xrightarrow{\tau} \quad C_k(\tilde{X}; \mathbb{Z}/2) \quad \xrightarrow{p_{\#}} \\ C_k(X; \mathbb{Z}/2) &\rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2, \tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, \; f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, \; p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{aligned} p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) &= 0 \\ \implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) &= 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} < \text{ injt } > \quad C_k(X; \mathbb{Z}/2) \quad \xrightarrow{\tau} \quad C_k(\tilde{X}; \mathbb{Z}/2) \quad \xrightarrow{p_{\#}} \\ C_k(X; \mathbb{Z}/2) &\rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, \; f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, \; p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{aligned} p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) &= 0 \\ \implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) &= 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} < \text{ injt } > \quad C_k(X; \mathbb{Z}/2) \quad \xrightarrow{\tau} \quad C_k(\tilde{X}; \mathbb{Z}/2) \quad \xrightarrow{p_{\#}} \\ C_k(X; \mathbb{Z}/2) &\rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0 \qquad S^n \xrightarrow{p} \mathbb{RP}^n$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma$ , i.e.  $p_{\#}(\tilde{\sigma}) = \sigma$

$$\begin{aligned} p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) &= 0 \\ \implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) &= 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} < \text{ injt } > \quad C_k(X; \mathbb{Z}/2) \quad \xrightarrow{\tau} \quad C_k(\tilde{X}; \mathbb{Z}/2) \quad \xrightarrow{p_{\#}} \\ C_k(X; \mathbb{Z}/2) &\rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2, \tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$



$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0 \qquad S^n \xrightarrow{p} \mathbb{RP}^n$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  $S^n \xrightarrow{p} \mathbb{RP}^n$   
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma$ , i.e.  $p_{\#}(\tilde{\sigma}) = \sigma$

$$\begin{aligned} p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) &= 0 \\ \implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) &= 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} < \text{ injt } > \quad C_k(X; \mathbb{Z}/2) \xrightarrow{\tau} C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} \\ C_k(X; \mathbb{Z}/2) &\rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2, \tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, \, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma$ , i.e.  $p_{\#}(\tilde{\sigma}) = \sigma$

$$\begin{array}{ccc} S^n & \xrightarrow{p} & \mathbb{RP}^n \\ \downarrow f & & \\ S^n & \xrightarrow{p} & \mathbb{RP}^n \end{array}$$

$$\begin{aligned} & p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0 \\ \implies & p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0 \end{aligned}$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{aligned} & \langle \text{injt} \rangle \quad C_k(X; \mathbb{Z}/2) \xrightarrow{\tau} C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} \\ & C_k(X; \mathbb{Z}/2) \rightarrow 0 \end{aligned}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, \, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

$$\begin{array}{ccc} S^n & \xrightarrow{p} & \mathbb{RP}^n \\ \downarrow f & & \downarrow f' \\ S^n & \xrightarrow{p} & \mathbb{RP}^n \end{array}$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to  
 $\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma$ , i.e.  $p_{\#}(\tilde{\sigma}) = \sigma$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$

$$\implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{array}{ccccc} < \text{injt} > & C_k(X; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(\tilde{X}; \mathbb{Z}/2) & \xrightarrow{p_{\#}} \\ & C_k(X; \mathbb{Z}/2) & \rightarrow & 0 & \end{array}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2, \tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

$$C_k(S^n; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(\mathbb{RP}^n; \mathbb{Z}/2)$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{array}{ccc} \downarrow f_{\#} & & \downarrow f'_{\#} \\ C_k(S^n; \mathbb{Z}/2) & \xrightarrow{p_{\#}} & C_k(\mathbb{RP}^n; \mathbb{Z}/2) \end{array}$$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$

$$\implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{array}{ccccc} < \text{ injt } > & C_k(X; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(\tilde{X}; \mathbb{Z}/2) & \xrightarrow{p_{\#}} \\ & C_k(X; \mathbb{Z}/2) & \rightarrow & 0 & \end{array}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

$$C_k(\mathbb{RP}^n; \mathbb{Z}/2) \xrightarrow{\tau} C_k(S^n; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(\mathbb{RP}^n; \mathbb{Z}/2)$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{array}{ccccc} & & \downarrow f_{\#} & & \downarrow f'_{\#} \\ C_k(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(S^n; \mathbb{Z}/2) & \xrightarrow{p_{\#}} & C_k(\mathbb{RP}^n; \mathbb{Z}/2) \end{array}$$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$

$$\implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{array}{ccccccc} < \text{ injt } & & C_k(X; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(\tilde{X}; \mathbb{Z}/2) & \xrightarrow{p_{\#}} & \\ & & C_k(X; \mathbb{Z}/2) & \rightarrow & 0 & & \end{array}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

$$C_k(\mathbb{RP}^n; \mathbb{Z}/2) \xrightarrow{\tau} C_k(S^n; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(\mathbb{RP}^n; \mathbb{Z}/2)$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{array}{ccccc} \downarrow f'_{\#} & & \downarrow f_{\#} & & \downarrow f'_{\#} \\ C_k(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(S^n; \mathbb{Z}/2) & \xrightarrow{p_{\#}} & C_k(\mathbb{RP}^n; \mathbb{Z}/2) \end{array}$$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$

$$\implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{array}{ccccc} < \text{ injt } & & C_k(X; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(\tilde{X}; \mathbb{Z}/2) & \xrightarrow{p_{\#}} \\ & & C_k(X; \mathbb{Z}/2) & \rightarrow & 0 \end{array}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$\tilde{X} \xrightarrow{p} X$  two sheeted covering

$$f : S^n \rightarrow S^n, f(-x) = -f(x)$$

$$C_k(\tilde{X}; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(X; \mathbb{Z}/2) \rightarrow 0$$

$$0 \longrightarrow C_k(\mathbb{RP}^n; \mathbb{Z}/2) \xrightarrow{\tau} C_k(S^n; \mathbb{Z}/2) \xrightarrow{p_{\#}} C_k(\mathbb{RP}^n; \mathbb{Z}/2) \longrightarrow 0$$

because  $\sigma : \Delta_k \rightarrow X$  lifts to

$$\tilde{\sigma} : \Delta_k \rightarrow \tilde{X}, p \circ \tilde{\sigma} = \sigma, \text{ i.e. } p_{\#}(\tilde{\sigma}) = \sigma$$

$$\begin{array}{ccccccc} & & \downarrow f'_{\#} & & \downarrow f_{\#} & & \downarrow f'_{\#} \\ 0 & \longrightarrow & C_k(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(S^n; \mathbb{Z}/2) & \xrightarrow{p_{\#}} & C_k(\mathbb{RP}^n; \mathbb{Z}/2) \longrightarrow 0 \end{array}$$

$$p_{\#}(\tilde{\sigma}_1 + \tilde{\sigma}_2 + \cdots + \tilde{\sigma}_m) = 0$$

$$\implies p_{\#}(\tilde{\sigma}_1) + p_{\#}(\tilde{\sigma}_2) + \cdots + p_{\#}(\tilde{\sigma}_m) = 0$$

$p_{\#}(\tilde{\sigma}_1) = p_{\#}(\tilde{\sigma}_2)$  etc., i.e.  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_2$  are lifts of the same simplex

Implies exactness of,

$$\begin{array}{ccccc} < \text{ injt } > & C_k(X; \mathbb{Z}/2) & \xrightarrow{\tau} & C_k(\tilde{X}; \mathbb{Z}/2) & \xrightarrow{p_{\#}} \\ & C_k(X; \mathbb{Z}/2) & \rightarrow & 0 & \end{array}$$

where,  $\tau(\sigma) = \tilde{\sigma}_1 + \tilde{\sigma}_2$ ,  $\tilde{\sigma}_1$ , where  $\tilde{\sigma}_2$  are the lifts of  $\sigma$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_{n-1}(S^n; \mathbb{Z}/2) \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \downarrow f_* \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_{n-1}(S^n; \mathbb{Z}/2)
\end{array}$$



$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z]))$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)])$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)]$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)]$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)]$$



$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z]$$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective.

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{p_*} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$f$  has odd degree

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$f$  has odd degree

**Theorem** (Borsuk-Ulam).  $f : S^n \rightarrow \mathbb{R}^n$ , then  $f(-x) = f(x)$  for some  $x \in S^n$

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$f$  has odd degree

**Theorem** (Borsuk-Ulam).  $f : S^n \rightarrow \mathbb{R}^n$ , then  $f(-x) = f(x)$  for some  $x \in S^n$

*Proof.* Assume  $f(x) - f(-x) \neq 0$  for any  $x$

□



$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$f$  has odd degree

**Theorem** (Borsuk-Ulam).  $f : S^n \rightarrow \mathbb{R}^n$ , then  $f(-x) = f(x)$  for some  $x \in S^n$

*Proof.* Assume  $f(x) - f(-x) \neq 0$  for any  $x$

$$h(x) = \frac{f(x)-f(-x)}{|f(x)-f(-x)|} \text{ is well defined and } h(-x) = -h(x).$$

□

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$f$  has odd degree

**Theorem** (Borsuk-Ulam).  $f : S^n \rightarrow \mathbb{R}^n$ , then  $f(-x) = f(x)$  for some  $x \in S^n$

*Proof.* Assume  $f(x) - f(-x) \neq 0$  for any  $x$

$$h(x) = \frac{f(x)-f(-x)}{|f(x)-f(-x)|} \text{ is well defined and } h(-x) = -h(x).$$

$$S^n \overset{h}{\rightarrow} S^{n-1}$$

□

$$\begin{array}{ccccccccc}
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0 \\
& & \downarrow f'_* & & \downarrow f_* & & \downarrow f'_* & & \downarrow f'_* & & \\
0 & \longrightarrow & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\sim} & H_n(S^n; \mathbb{Z}/2) & \xrightarrow{0} & H_n(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{<iso>} & H_{n-1}(\mathbb{RP}^n; \mathbb{Z}/2) & \xrightarrow{\tau_*} & 0
\end{array}$$

$\theta$  is the (only) non-trivial deck transformation

$$p_*([z]) = [p_{\#}(z)]??$$

$$\tau_*(p_*([z])) = \tau_*([p_{\#}(z)]) = [z + \theta_{\#}(z)] = [z] + [\theta_{\#}(z)] = [z] + \theta_*[(z)] = 2[z] = 0$$

But  $\tau_*$  is injective. Therefore,  $p_* = 0$

$f$  has odd degree

**Theorem** (Borsuk-Ulam).  $f : S^n \rightarrow \mathbb{R}^n$ , then  $f(-x) = f(x)$  for some  $x \in S^n$

*Proof.* Assume  $f(x) - f(-x) \neq 0$  for any  $x$

$$h(x) = \frac{f(x)-f(-x)}{|f(x)-f(-x)|} \text{ is well defined and } h(-x) = -h(x).$$

$$S^{n-1} \xrightarrow{i} S^n \xrightarrow{h} S^{n-1} \text{ has odd degree}$$

