

ON CANONICITY FOR INTEGRAL MODELS OF SHIMURA VARIETIES WITH HYPERSPECIAL LEVEL

KEERTHI MADAPUSI AND ALEX YOUNG

ABSTRACT. We give a new definition—and in some cases, a new construction—of integral canonical models of Shimura varieties that uses the notion of an aperture appearing in work of the first author with Gardner on some conjectures of Drinfeld. This applies to Shimura varieties of pre-abelian type at primes of hyperspecial level, recovering and extending previous work of Kisin, Kim–Madapusi and Imai–Kato–Younis, but also to exceptional Shimura varieties for large enough primes. The characterization in the exceptional case is *a priori* different from the one recently shown by Bakker–Shankar–Tsimmerman, and recovers many of their results, such as the existence of prime-to- p Hecke operators, the non-emptiness of the μ -ordinary stratum and the theory of the canonical lift. In fact, we give a uniform proof of the non-emptiness of *all* possible Newton strata, and—in the pre-abelian case—of the non-emptiness of Ekedahl–Oort strata and central leaves as well. An important ingredient in the proofs is a generalization of Tate’s full faithfulness theorem for p -divisible groups to the context of apertures. This leads to a mapping property for the integral canonical model that characterizes maps into it from all normal, flat and excellent schemes over $\mathbb{Z}_{(p)}$.

CONTENTS

1. Introduction	1
2. Prismatic F -gauges	8
3. $(\mathcal{G}, \mu)$ -apertures	14
4. Tate’s full faithfulness for apertures and its consequences	26
5. Criteria for the existence of apertures	32
6. Integral canonical models: Definitions and first properties	40
7. Integral canonical models: Constructions and further properties	46
8. Applications and further results	55
References	62

1. INTRODUCTION

The purpose of this article is to lay the groundwork for applying the results of [26] to global questions. In doing so, we give a primarily p -adic characterization of integral canonical models of Shimura varieties at unramified primes, including for those of exceptional type considered by Bakker–Shankar–Tsimmerman [4]. This characterization is *a priori* different from the primarily ℓ -adic ones given in *op. cit.* and others, and is closely related to the ones introduced by Pappas [72] and Imai–Kato–Younis [39]. For Shimura varieties of abelian type, this method actually gives a new interpretation of Kisin’s twisting construction, and extends to give models for Shimura data of *pre*-abelian type as well.¹

1.1. Definition and characterization. The key observation animating this paper is that apertures, the mixed characteristic analogues of the local shtukas of Genestier–Lafforgue, introduced by Drinfeld [24] and studied in the work of the first author with Gardner [26], have very limpid connections with the integral theory of Shimura

¹Pre-abelian Shimura data have a simple type-theoretic characterization: the adjoint datum is of type $A, B, C, D^{\mathbb{R}}$, or $D^{\mathbb{H}}$. The first non-trivial example of one that is not of abelian type is a type $D^{\mathbb{H}}$ Shimura datum with simply connected derived subgroup—see Example 7.8.3.

varieties, and in fact characterize the ‘good’ models completely, reifying an expectation of Drinfeld [24]. Significant progress along these lines was made in [39], where it is shown that the prismatic realization on an integral model of a Shimura variety of abelian type is sufficient to pin down the model at *finite* level, without considering the whole prime-to- p Hecke tower.

Our primary aim here is to situate this result in what we believe to be the right context, and to substantially broaden the kinds of Shimura data it applies to by exploiting the geometry made available now by the theory of [26].

Setup 1.1.1. We will fix a prime p . An **unramified Shimura datum** is a triple $(G, \mathcal{G}, X)$ where (G, X) is a Shimura datum and $\mathcal{G}$ is a reductive model of G over $\mathbb{Z}_p$. We also set $K_p = \mathcal{G}(\mathbb{Z}_p)$, and only consider level subgroups $K \subset G(\mathbb{A}_f)$ of the form $K_p K^p$ for a *neat* compact open subgroup $K^p \subset G(\mathbb{A}_f^p)$. We call the tuple $(G, \mathcal{G}, X, K)$ an **unramified Shimura tuple**. We also fix a place $v|p$ of the reflex field E of (G, X) . Over the associated Shimura variety Sh_K , we have a canonical $\mathcal{G}^c(\mathbb{Z}_p)$ -local system $\mathbf{Et}_{K,p}$, where $\mathcal{G}^c$ is a reductive model over $\mathbb{Z}_p$ of the cuspidal quotient G^c (see §6.1).

Remark 1.1.2. The Shimura datum gives an E_v -rational conjugacy class of cocharacters $\{\mu_v\}$. We in fact have a representative $\mu_v: \mathbb{G}_{m, \mathcal{O}_{E_v}} \rightarrow \mathcal{G}_{\mathcal{O}_{E_v}}$ for $\{\mu_v\}$. We will view μ_v as a cocharacter of $\mathcal{G}_{\mathcal{O}_{E_v}}^c$. To this we can attach the smooth algebraic formal stacks $\mathrm{BT}_n^{\mathcal{G}^c, -\mu_v}$ over $\mathcal{O}_{E_v}$ constructed in [26]. We can also consider their inverse limit $\mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}$. These stacks generalize the formal stacks of (truncated) polarized p -divisible groups of height $2g$ in the case of the Siegel Shimura data, and give a group theoretic way of defining ‘ p -divisible groups with $\mathcal{G}^c$ -structure’ without actually requiring one to work with p -divisible groups. Moreover, they can be constructed from data that is not necessarily of ‘Hodge type’, where there might not be any connection with p -divisible groups at all.

Remark 1.1.3. For any p -adic formal algebraic space $\mathfrak{X}$, there is a canonical étale realization functor

$$T_{\mathrm{ét}}: \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}(\mathfrak{X}) \rightarrow \mathrm{Loc}_{\mathcal{G}^c(\mathbb{Z}_p)}(\mathfrak{X}_\eta),$$

where the right-hand side is the groupoid of $\mathcal{G}^c(\mathbb{Z}_p)$ -local systems over the generic fiber of $\mathfrak{X}$. When (G, X) is the Siegel Shimura datum, then what we have here is simply the functor taking a polarized p -divisible group to its Tate module viewed as a symplectic space.

Definition 1.1.4 (Crystalline integral canonical models). An integral model² $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_{E, (v)}$ with v -adic formal completion $\widehat{\mathcal{S}}_K$ is a **crystalline integral canonical model** (**crystalline ICM** for short) if the following conditions hold:

- (1) (Serre–Tate property) There exists a formally étale map of p -adic formal stacks over $\mathrm{Spf} \mathcal{O}_{E, v}$

$$\varpi: \widehat{\mathcal{S}}_K \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v},$$

such that the induced $\mathcal{G}^c(\mathbb{Z}_p)$ -local system on the generic fiber is isomorphic to $\mathbf{Et}_{K,p}$.

- (2) (Pointwise criterion) For any mixed characteristic $(0, p)$ complete discrete valuation field F over $\mathcal{O}_{E_v}$ with perfect residue field and $x \in \mathrm{Sh}_K(F)$ ³, the following are equivalent:
 - (a) $x \in \mathcal{S}_K(\mathcal{O}_F)$;
 - (b) $\mathbf{Et}_{K,p}|_x$ is a crystalline $\mathcal{G}^c(\mathbb{Z}_p)$ -local system;
 - (c) $\mathbf{Et}_{K,p}|_x$ is a potentially crystalline $\mathcal{G}^c(\mathbb{Z}_p)$ -local system;

With the definition in hand, we get the following quite limpid mapping property.

Theorem A (Mapping property of crystalline ICMs, Theorem 6.5.3). *Let $\mathcal{S}_K$ be a crystalline ICM over $\mathcal{O}_{E, (v)}$. Suppose that we have a normal flat excellent algebraic space $\mathcal{X}$ over $\mathcal{O}_{E, (v)}$. Then giving a map $\mathcal{X} \rightarrow \mathcal{S}_K$ of $\mathcal{O}_{E, (v)}$ -schemes is equivalent to giving the following data:*

- (1) A map $f: \mathcal{X}_\eta \rightarrow \mathrm{Sh}_K$;

²By ‘integral model’, we will mean a separated *algebraic space* of finite type over $\mathcal{O}_{E, (v)}$ whose generic fiber is equipped with an isomorphism to Sh_K as an E -scheme. In practice, the models we encounter will be quasi-projective schemes, but, even for Siegel modular varieties, knowing this requires a non-trivial result of Mumford.

³Here and elsewhere, we will write $\mathrm{Sh}_K(F)$ for what should properly be written as $(\mathrm{Sh}_K \otimes_E E_v)(F)$.

(2) A map $\eta: \widehat{\mathcal{X}} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}, \cdot \mu_v}$ such that the associated $\mathcal{G}^c(\mathbb{Z}_p)$ -local system on $\widehat{\mathcal{X}}_{\eta}$ is isomorphic to $\mathbf{Et}_{K,p}|_{\widehat{\mathcal{X}}_{\eta}}$.

In particular, the crystalline ICM $\mathcal{S}_K$ is unique up to unique isomorphism, and is functorial in the unramified tuple $(G, \mathcal{X}, X, K)$.

Remark 1.1.5. The proof of the theorem is based on an argument of Pappas [72, Theorem 7.1.7]. His use of Tate’s full faithfulness theorem for p -divisible groups is replaced with that of a version for apertures; see §1.6 below. A closely related argument also appears in [39, Theorem 3.13].

Remark 1.1.6 (Mapping property for Pappas–Rapoport models). The method of proof of Theorem A combined with a fully faithfulness result of Pappas–Rapoport [73] establishes a mapping property for integral models at parahoric levels satisfying the axioms from *loc. cit.*; see Theorem 8.6.17.

For $p > 3$, we have a simpler mapping property for crystalline ICMs with respect to smooth $\mathcal{O}_{E,(v)}$ -schemes.

Theorem B (Pointwise mapping property, Proposition 7.9.2). *Suppose that $p > 3$, that $\mathcal{S}_K$ is a crystalline ICM for Sh_K , and that $\mathcal{X}$ is a smooth $\mathcal{O}_{E,(v)}$ -scheme with generic fiber X . Then a map $f: X \rightarrow \mathrm{Sh}_K$ extends to a map $\mathcal{X} \rightarrow \mathcal{S}_K$ if and only if for all finite extensions F/E_v and all $x \in \mathcal{X}(\mathcal{O}_F)$ the local system $\mathbf{Et}_{K,p}|_x$ is crystalline. In particular, if $\mathcal{S}_K$ is proper over $\mathcal{O}_{E,(v)}$, then it is a Néron model for Sh_K in the sense of [11, Definition 1]: For any smooth $\mathcal{O}_{E,(v)}$ -scheme S , every map $S_{\eta} \rightarrow \mathrm{Sh}_K$ of E -schemes extends to a map $S \rightarrow \mathcal{S}_K$.*

In terms of existence of such models, we have:

Theorem C (Pre-abelian-type canonical models, Theorem 7.8.4). *Suppose that (G, X) is of pre-abelian type. Then Sh_K admits a crystalline ICM over $\mathcal{O}_{E,(v)}$. In fact, if (G, X) is of abelian type, then the models constructed by Kisin [47] (and Kim–Madapusi [45] for $p = 2$) are crystalline ICMs in the sense used here.*

See §1.4 below for a comparison with existing results.

1.2. Limpid ICMs and their properties.

Remark 1.2.1 (Yet another canonical model?). The experienced reader may be wondering at this point if the world really needs yet another notion of integral canonical models for Shimura varieties. The arguments we would proffer in favor of the definition here—which we believe to be the *correct* one—are as follows:

- (1) First, of course, one has the quite pleasing mapping property from Theorem A, which we should note is a *consequence* of the intrinsic definition of a crystalline ICM.
- (2) More convincingly, after one also accounts systematically for the prime-to- p realizations, one is able to deduce various non-trivial properties for such models using *only* their defining features, without invoking any moduli interpretation, as we will now see.
- (3) In forthcoming work of the first author with Si Ying Lee [57], one will moreover find that various properties of integral models of abelian type (Rapoport–Zink uniformization, existence of CM lifts, Igusa stacks, Langlands–Rapoport type point counting formulas, etc.), for which the use of abelian varieties has hitherto seemed essential, are in fact *properties* of all limpid ICMs as defined below.

Remark 1.2.2 (Prime-to- p étale realizations). For $\ell \neq p$, we also have K_{ℓ}^c -local systems $\mathbf{Et}_{K,\ell}$, with $K_{\ell}^c \subset G^c(\mathbb{Q}_{\ell})$ a compact open subgroup.

To present our further results, we will need two further ways of pinning down integral models.

Definition 1.2.3 (Limpid integral canonical models). A crystalline ICM $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_{E,(v)}$ with v -adic formal completion $\widehat{\mathcal{S}}_K$ is **limpid** if, for all $\ell \neq p$, the K_{ℓ}^c -local system $\mathbf{Et}_{K,\ell}$ extends over $\mathcal{S}_K$. Note that this is a *property* of the unique crystalline ICM when it exists.

Remark 1.2.4 (Pre-abelian ICMs are limpid). In the context of Theorem C, crystalline ICMs are in fact limpid. This follows from our proof, which uses the three theorems of this subsection—which do not use any sort of moduli interpretation—to reduce to the (essentially standard) fact that the moduli of principally polarized abelian schemes (with appropriate level structure) is a limpid ICM for the Siegel modular variety.

structure

Theorem D (Integral canonical models from reduction of structure group, §7.3). *Suppose that $(G, \mathcal{G}, X, K) \rightarrow (G^\sharp, \mathcal{G}^\sharp, X^\sharp, K^\sharp)$ is a map of unramified Shimura tuples with $\mathcal{G} \rightarrow \mathcal{G}^\sharp$ a closed immersion of reductive group schemes. If $\mathrm{Sh}_{K^\sharp}$ admits a crystalline (resp. limpid) ICM over $\mathcal{O}_{E^\sharp, (v^\sharp)}$ (for the place $v^\sharp \mid p$ of the reflex field $E^\sharp \subset E$ lying under v), then so does Sh_K .*

Remark 1.2.5. Theorem D is already known by work of Pappas–Rapoport and Daniels–van Hoften–Kim–Zhang in the context of Pappas–Rapoport canonical models, even for quasi-parahoric level structures: see [16, Theorem 4.1.8].

Next, we have two results concerning the behavior of limpid ICMs with respect to central modifications.

up:central

Setup 1.2.6. We will fix a map of unramified Shimura tuples

$$(G_1, \mathcal{G}_1, X_1, K_1) \rightarrow (G_2, \mathcal{G}_2, X_2, K_2)$$

with reflex fields $E_2 \subset E_1$ such that $G_1 \rightarrow G_2$ is surjective with central kernel. We will also fix a place $v_1 \mid p$ of E_1 and let $v_2 \mid p$ be the place of E_2 under it.

First, we have the following *ascent* statement, which implies that constructing ICMs for adjoint Shimura data is sufficient to construct them for all Shimura data:

thm:ascent

Theorem E (Ascent of ICMs along central isogenies). *If Sh_{K_2} admits a limpid ICM $\mathcal{S}_{K_2}$ over $\mathcal{O}_{E_2, (v_2)}$, then Sh_{K_1} admits a limpid ICM $\mathcal{S}_{K_1}$ over $\mathcal{O}_{E_1, (v_1)}$, and $\mathcal{S}_{K_1} \rightarrow \mathcal{S}_{K_2}$ is finite étale.*

Remark 1.2.7. There is an obvious way to construct an integral model $\mathcal{S}_{K_2}$ for Sh_{K_2} from an integral model $\mathcal{S}_{K_1}$ for Sh_{K_1} : One can take the normalization of $\mathcal{S}_{K_1}$ in Sh_{K_2} . The issue is to show that this preserves the property of being a crystalline ICM. The heart of the matter is to see that $\mathcal{S}_{K_1}$ is finite étale over $\mathcal{S}_{K_2}$. We prove this by showing it holds for the restriction to the μ -ordinary locus and using purity. Over the μ -ordinary locus, we need the theory of the canonical lift; see Theorem K below. The fact that the canonical lift is CM makes use of the prime-to- p realizations, and so requires us to have a limpid and not merely crystalline ICM.

is:enough

Remark 1.2.8 (Sufficiency of adjoint data). Theorem E tells us that, in order to construct limpid ICMs for tuples with underlying data $(G, \mathcal{G}, X)$, it suffices to do so for tuples with underlying *adjoint* Shimura data $(G^{\mathrm{ad}}, \mathcal{G}^{\mathrm{ad}}, X^{\mathrm{ad}})$.

Next, we have an abstract form of the *twisting* construction from [48], which uses the modified Weil restriction construction from Definition 7.6.1 below.

thm:descent

Theorem F (Descent of ICMs along central isogenies). *Suppose that one of the following holds:*

- (1) *we have $G_1 = G_2$ and Sh_{K_1} admits an ICM $\mathcal{S}_{K_1}$ over $\mathcal{O}_{E_1, (v_1)}$;*
- (2) *for every totally real number field F in which p is unramified, all unramified Shimura tuples with underlying unramified Shimura datum $\mathrm{Res}_{F/\mathbb{Q}}(G_1, \mathcal{G}_1, X_1)$ admit ICMs over $\mathcal{O}_{E_1, (v_1)}$.*

Then Sh_{K_2} admits an ICM $\mathcal{S}_{K_2}$ over $\mathcal{O}_{E_2, (v_2)}$, and the map $\mathcal{S}_{K_1} \rightarrow \mathcal{S}_{K_2}$ is finite étale.

Remark 1.2.9 (Role of the modified Weil restriction condition). The role of (2) is essentially to overcome the fact that in the generality that we work with here, we do not have a good *intrinsic* notion of ‘ F -linear isogenies’ between points on the Shimura variety. In the context of Shimura varieties of Hodge type, such isogenies, defined through the F -linearization of the isogeny category of abelian varieties, are an important actor in Kisin’s twisting construction. This is replaced here by a notion of isogeny obtained via restriction from the ICMs for the modified Weil restrictions. The condition of being of Hodge type is preserved under the modified Weil restriction construction, and so this gives another perspective on the construction of Shimura varieties of abelian type from those of Hodge type.

exceptional

1.3. Exceptional Shimura varieties. To show the existence of ICMs beyond the pre-abelian context, we need to change our definition to prioritize the prime-to- p realizations a bit more than we have so far.

Definition 1.3.1 (Étale integral canonical models). An integral model $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_{E, (v)}$ is an **étale integral canonical model** (**étale ICM** for short) if it satisfies the Serre–Tate property, and if the following conditions hold:

- (1) (Étale realizations) For all $\ell \neq p$, the K_ℓ^c -local system $\mathbf{Et}_{K,\ell}$ extends over $\mathcal{S}_K$.
- (2) (Pointwise criterion) For any mixed characteristic $(0, p)$ discrete valuation field F over $\mathcal{O}_{E_v}$ and $x \in \mathrm{Sh}_K(F)$, the following are equivalent:
 - (a) $x \in \mathcal{S}_K(\mathcal{O}_F)$;
 - (b) $\mathbf{Et}_{K,\ell}|_x$ is an unramified K_ℓ^c -local system for all $\ell \neq p$;
 - (c) $\mathbf{Et}_{K,\ell}|_x$ is a potentially unramified K_ℓ^c -local system for all $\ell \neq p$;

Remark 1.3.2 (Mapping property for étale ICMs). There is a variant of Theorem A for étale ICMs (see Theorem 6.5.3): One has to also require that, for all $x \in \mathcal{X}(\mathcal{O}_F) \subset \mathcal{X}_\eta(F)$, the local systems $\mathbf{Et}_{K,\ell}|_x$ over $\mathrm{Spec} F$ are unramified for all $\ell \neq p$. In particular, one finds that étale ICMs are unique up to isomorphism and functorial in unramified tuples.

limpid_etalé **Remark 1.3.3** (Limpid vs. étale for proper ICMs). If $\mathcal{S}_K$ is proper, then the pointwise criteria are irrelevant, and a limpid ICM is the same as an étale ICM.

hodge_type **Remark 1.3.4** (Pre-abelian ICMs are also étale ICMs). The limpid ICMs for pre-abelian Shimura variety are also étale ICMs. This comes down to the fact that, for an abelian variety A over a local field, both unramifiedness of the ℓ -adic Tate module (by the usual Néron–Ogg–Shafarevich criterion) and the crystallinity of the p -adic Tate module (by a result of Coleman–Iovita [14, Theorem 1]) are equivalent to saying that A has good reduction.

icm_enough **Remark 1.3.5.** In all the results of §1.2, one can replace ‘limpid ICM’ with ‘étale ICM’ without affecting the conclusions. The proofs of Theorems E and F depend mainly on the existence of the universal aperture and of prime-to- p Hecke correspondences, and the latter emerge from the unr

With this setup, we now have the following version of the theorem of Bakker–Shankar–Tsimmerman [4]:

exceptional **Theorem G** (Exceptional integral canonical models, §7.10). *For any unramified Shimura tuple $(G, \mathcal{G}, X, K)$, and all p large enough, an étale ICM $\mathcal{S}_K$ exists over $\mathcal{O}_{E,(v)}$ for any $v|p$. If Sh_K is proper (equivalently, if G^{ad} is anisotropic), then $\mathcal{S}_K$ is in fact a limpid ICM.*

Remark 1.3.6. Given Theorem C, the content of Theorem G is of course that it applies to the so-called exceptional Shimura data that are not of pre-abelian type and hence cannot be related to the moduli of abelian varieties in any reasonably direct way.

comparison **1.4. Comparison with existing results.**

canonicity **Remark 1.4.1** (Comparison with existing notions of canonicity). Our notion of canonicity is *different* from the ones appearing in [71], [47] or [4], which only use ℓ -adic realizations for $\ell \neq p$ to characterize the model. Also, the characterizations appearing in the first two of these, are not of models at any finite level, but of the whole prime-to- p Hecke tower. As such, they are slightly tricky to formulate and check. Also, our definition of ICMs is *intrinsic* to an integral model, with the mapping property arising as a consequence, while in the other three

The notion of a crystalline ICM here was introduced by Imai–Kato–Yucis [39], and this in turn has its antecedents in the work of Pappas [72] and Pappas–Rapoport [73]. The important observation that this should give a characterization at *finite level* is found in [39].

orem:kisin **Remark 1.4.2** (Comparison with the work of Kisin). The first general construction of integral canonical models of abelian type is due to Kisin [47].⁴ For Shimura varieties of Hodge type, what one will find in this paper is essentially a streamlined version of the arguments in *loc. cit.*, making use of the geometry of the stacks $\mathrm{BT}_n^{\mathcal{G}^c, \cdot \mu_v}$ to compress some of the p -adic Hodge theory used there.

To construct Shimura varieties of abelian type, Kisin introduces his *twisting* construction, which ultimately uses the global moduli interpretation of Siegel-type Shimura varieties in terms of abelian varieties. We actually give a *different* approach to twisting here. The main new observation here is that one does not need a moduli description:

⁴There is earlier work of Vasu [91]. Note, however, that despite the title of that paper, no actual construction of models of *pre*-abelian type appears within it.

One can formulate the construction in a way that is valid for all maps between Shimura data where the underlying map of groups is a central cover. We also make the additional new observation that the theory of the canonical lift for μ -ordinary points suffices to also *lift* crystalline ICMs along central covers. This leads to our construction of ICMs of pre-abelian typ.

ato_youcis

Remark 1.4.3 (Comparison with Imai–Kato–Youcis). The verification that the models constructed by Kisin and Kim–Madapusi are ICMs in the sense used here is originally due to Imai–Kato–Youcis [39]. The proof we give for Theorem C is different from that in [39] and works also when $p = 2$. But, we do make use of intermediate results from [37, 38, 39].

comparison

Remark 1.4.4 (Comparison with Bakker–Shankar–Tsimmerman). The idea that one should be able to *provably* characterize integral canonical models of *all* Shimura varieties, including those not of abelian type, first appears in the already mentioned recent work of Bakker–Shankar–Tsimmerman. Their results extend beyond the ambit of Shimura varieties, and apply to integral models of *non-minuscule* variations of Hodge structure. This is in some sense the main philosophical difference between these works: Here, we make full use of the minuscule (though in the sense of G -bundles, not just vector bundles) of the family of realizations over a Shimura variety, for instance, in the form of the full faithfulness result Theorem L. But these methods say nothing about what happens beyond the minuscule regime.

The étale ICMs here are also canonical in the sense of BST and our proof of canonicity is independent from theirs—though it uses essentially the same inputs. Moreover, the lower bound on the prime p is *a priori* smaller than the one in *loc. cit.*

1.5. Applications. In this subsection, the term ‘ICM’ without further qualification can mean any one of crystalline, limpid or étale ICM. Our first application is essentially an algebraization result for apertures.

surjective

Theorem H (Surjectivity of the syntomic realization, Theorem 8.1.8). *Suppose that Sh_K admits an ICM $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$ and that one of the following conditions holds:*

- (1) *the Shimura datum (G, X) is of pre-abelian type;*
- (2) *the ICM $\mathcal{S}_K$ is proper.*

Then, for all $n \geq 1$, the map of formal Artin stacks

$$\widehat{\mathcal{S}}_K \rightarrow \mathrm{BT}_n^{G^c, -\mu_v}$$

is smooth and surjective.

Remark 1.5.1. Proposition 8.1.7 shows that this surjectivity follows as soon as $(G, \mathcal{G}, X, K)$ satisfies the **de Jong extension property** as in Definition 8.1.4. One can see Theorem H as the claim that the extension property holds in these cases, and we expect that it holds for arbitrary ICMs.

Remark 1.5.2. The surjectivity assertion above is already known for Shimura varieties of PEL type by work of Viehmann–Wedhorn [93]. Their proof amounts to first establishing the non-emptiness of Newton strata, followed by an argument via isogenies. A similar argument works for Shimura varieties for Hodge and perhaps even abelian type, but one has to work quite a bit harder by appealing to the non-emptiness results of [56] or [49], as well as to some deeper results from [48] on p -isogenies between mod- p points of the Shimura variety.

The proof given here, on the other hand, is entirely geometric, works uniformly for all Shimura varieties of pre-abelian type, and requires no *a priori* knowledge of non-emptiness statements or of the existence of isogenies. In fact, it can be used to *deduce* such statements.

dahl--Oort

Theorem I (Non-emptiness of Ekedahl–Oort strata). *Suppose that $\mathcal{S}_K$ is an ICM. Then there exists a smooth map of stacks*

$$\mathcal{S}_K \otimes k(v) \rightarrow \mathcal{G}^c\text{-zip}_{-\mu_v}.$$

Here, the right-hand side is the stack of $\mathcal{G}^c$ -zips of type $-\mu_v$ defined in [78], and admits a canonical open and dense stratum known as the μ -ordinary stratum. Moreover:

- (1) *The μ -ordinary locus of $\mathcal{S}_K \otimes k(v)$ (which is the pre-image of the μ -ordinary stratum) is dense.*

- (2) If $\mathcal{S}_K$ satisfies the hypotheses of Theorem H (so in particular if $\mathcal{S}_K$ is pre-abelian), then this map is also surjective: That is, all Ekedahl–Oort strata of the special fiber are non-empty.

Remark 1.5.3. The non-emptiness of Ekedahl–Oort strata for Shimura varieties of PEL type is due to Viehmann–Wedhorn [93], and is also known in the abelian type case by work of Shen–Wang [86] (though also see earlier work of Wortmann [96] and [98]). The proof of Shen–Wang once again uses deep results of Kisin from [48]. On the other hand, our proof is essentially an immediate consequence of Theorem H, and the fact (shown in [26]) that there is a natural smooth surjective map $\mathrm{BT}_1^{\mathcal{G}^c, \tau^{\mu_v}} \otimes \mathbb{F}_p \rightarrow \mathcal{G}^c\text{-zip}_{-\mu_v}$.

There is another stratification where we can show non-emptiness for *all* ICMs: the *Newton stratification* using the Kottwitz set $B(G^c, \{-\mu_v\})$ defined in [53] and studied in [79].

ton_strata

Theorem J (Non-emptiness of Newton strata). *Let $\mathcal{S}_K$ be an ICM for Sh_K over $\mathcal{O}_{E,(v)}$. Then the natural map $\mathcal{S}_K \rightarrow B(G^c, \{-\mu_v\})$ is surjective.*

Next, we have applications to the μ -ordinary locus.

u_ordinary

Theorem K (Canonical CM lifts for μ -ordinary points, Theorem 7.4.2). *Suppose that $\mathcal{S}_K$ is an étale or limpid ICM for Sh_K and that $x \in \mathcal{S}_K(\kappa)$ is a closed point in the μ -ordinary locus where κ is a finite field. Let $\widehat{U}_x$ be the formal completion of $\mathcal{S}_K$ at x . Then:*

- (1) *There exists a canonical tower*

$$\widehat{U}_x = \widehat{U}_{x,n} \rightarrow \widehat{U}_{x,n-1} \rightarrow \cdots \rightarrow \widehat{U}_{x,0} = \mathrm{Spf} W(\kappa)$$

of formal schemes over $W(\kappa)$, where, for each $i > 0$, $\widehat{U}_{x,i}$ is a formal p -divisible group over $\widehat{U}_{x,i-1}$;

- (2) *The identity section⁵ $x^{\mathrm{can}} \in \widehat{U}_x(W(\kappa))$, the canonical lifting, is a CM point of $\mathcal{S}_K$.*

Remark 1.5.4. Bakker–Shankar–Tsimmerman also prove the existence of the CM canonical lift for their integral models [4, §8]. Unlike their proof, ours does not use *a priori* knowledge of the ℓ -independence of the conjugacy classes of Frobenius elements acting on the étale realizations (see Klevdal–Patrikis [52]): We only use the defining properties of ICMs, and deduce this ℓ -independence from the existence of the canonical CM lift.

Remark 1.5.5 (Integral canonical models for subhyperspecial levels). The results here can be combined with work of Takaya [88] to show the existence of integral canonical models in the sense of Pappas–Rapoport when the level at p is parahoric and *contained* in a hyperspecial subgroup of $G(\mathbb{Q}_p)$; see Proposition 8.6.13.

Remark 1.5.6 (Future applications and directions). In subsequent work, we expect the results here to be applied (among other questions) to:

- the problem of defining p -Hecke correspondences and Igusa stacks for limpid ICMs, as well as counting their mod- p points [57];
- an extension of the Langlands–Kottwitz–Scholze method to exceptional Shimura varieties by the second author;
- the understanding of central leaves [65];
- the study of special cycles on Shimura varieties.

In addition, in ongoing work, joint with Teruhisa Koshikawa and Kentaro Inoue, we will also extend these methods to the logarithmic context and define integral canonical models for toroidal and minimal compactifications of Shimura varieties.

rosub:tate

1.6. Tate full faithfulness. Finally, a key ingredient in our proofs of the canonicity results, and especially for establishing the mapping property in Theorem A, is a generalization of Tate’s full faithfulness theorem for the Tate module of p -divisible groups.

This result holds in the following setting: $\mathcal{G}$ will be a smooth affine group scheme over $\mathbb{Z}_p$ and μ will be a cocharacter of $\mathcal{G}$ defined over an unramified ring of integers $\hat{\mathcal{O}}$. We will assume that μ is 1-bounded in the sense of

⁵By this, we mean the successive composition of the identity sections of each map in the tower.

Lau [55, Definition 6.3.1] (when $\mathcal{G}$ is reductive, this is the same as being minuscule). Then we have the associated formal algebraic stacks $\mathrm{BT}_n^{\mathcal{G},\mu}$ constructed in [26, §9], and their inverse limit $\mathrm{BT}_\infty^{\mathcal{G},\mu}$. We also have the étale realization functor from $\mathrm{BT}_\infty^{\mathcal{G},\mu}$ to $\mathcal{G}(\mathbb{Z}_p)$ -local systems on the generic fiber, generalizing the Tate module of a p -divisible group.

Theorem L (Tate full faithfulness, Theorem 4.1.5). *If $\mathfrak{X}$ is a normal p -completely flat formal scheme over $\hat{\mathcal{O}}$, the étale realization functor from $\mathrm{BT}_\infty^{\mathcal{G},\mu}(\mathfrak{X})$ to $\mathcal{G}(\mathbb{Z}_p)$ -local systems over $\mathfrak{X}_\eta$ is fully faithful.*

Theorem M (Algebraization, Theorem 4.1.3). *For each $1 \leq n < \infty$, there is a unique smooth 0-dimensional algebraic stack $\mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}}$ over $\mathcal{O}$ with generic fiber $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}$ and with p -adic formal completion $\mathrm{BT}_n^{\mathcal{G},\mu}$. Moreover, the stack has affine diagonal.*

1.7. Notational and terminological conventions.

- Our notation and conventions concerning derived algebraic geometry agree with those in [26].
- We uniformly suppress notation indicating what ideal a completion (e.g., of a scheme or ring) is with respect to, as it will always be clear from context.
- A filtration (on an abelian group, ring, or module) is always assumed to be exhaustive and decreasing unless stated otherwise.
- For a ring A and ideal I , we write $\mathrm{Fil}_I^\bullet A$ for the I -adic filtration. The trivial filtration is $\mathrm{Fil}_{\mathrm{triv}} A \stackrel{\mathrm{defn}}{=} \mathrm{Fil}_0 A$. If I is invertible, we write $\mathrm{Fil}_{I,\pm}^\bullet A$ for the filtered ring with $\mathrm{Fil}_{I,\pm}^k A = I^k$.
- 1
- 2
- 3
- 4

2. PRISMATIC F -GAUGES

2.1. The stacks of Bhatt–Lurie and Drinfeld. Canonically associated with any derived p -adic formal scheme $\mathfrak{X}$ are three derived p -adic formal stacks⁶ $\mathfrak{X}^\Delta$, $\mathfrak{X}^\vee$, and $\mathfrak{X}^{\mathrm{syn}}$ (shortened to R^Δ , $R^\vee$, and R^{syn} when $\mathfrak{X} = \mathrm{Spf}(R)$), the **prismatization**, **filtered prismatization** and **syntomification**, respectively. We will review what we'll need to know of these cohomological stacks for this paper. The reader can find more details in [23], [5], [26, §6] and [67, §3]. The first two named sources explain how to construct these stacks as classical formal prestacks, and the third extends the constructions—following Bhatt–Lurie [6]—to the animated framework.

Remark 2.1.1 (Prisms and the prismatization). The prismatization $\mathfrak{X}^\Delta$ parameterizes *Cartier–Witt divisors* on $\mathfrak{X}$. These are related to the prismatic site for $\mathfrak{X}$. Indeed, suppose that we have an object $(A, I, \mathrm{Spf}(\bar{A}) \rightarrow \mathfrak{X})$ in the absolute prismatic site for $\mathfrak{X}$. Here A is endowed with the (p, I) -adic topology, and $\bar{A} = A/I$. Then, as in [6, Construction 3.10], we find a canonical map $\iota_{(A,I)}: \mathrm{Spf} A \rightarrow \mathfrak{X}^\Delta$ taking a map $\mathrm{Spec}(B) \rightarrow \mathrm{Spf}(A)$, where B is a p -nilpotent ring, to the Cartier–Witt divisor $I \otimes_A W(B) \rightarrow W(B)$ where $A \rightarrow W(B)$ is the unique δ -map lifting $A \rightarrow B$, and where we equip this Cartier–Witt divisor with structure map

$$\mathrm{Spec}(W(B)/{}^{\mathbb{L}}(I \otimes_A W(B))) \rightarrow \mathrm{Spf}(\bar{A}) \rightarrow \mathfrak{X}.$$

Remark 2.1.2 (Relationship between the stacks). The stack $\mathfrak{X}^\vee$ is a *filtered* formal stack, meaning that it is fibered naturally over the formal Artin stack $\mathbb{A}^1/\mathbb{G}_m$ parameterizing line bundles L over p -complete rings C equipped with a cosection $L \rightarrow C$. The pre-image of the open point

$$\mathrm{Spf} \mathbb{Z}_p \simeq \mathbb{G}_m/\mathbb{G}_m \subset \mathbb{A}^1/\mathbb{G}_m$$

is canonically isomorphic to $\mathfrak{X}^\Delta$, and is called the **de Rham locus** of $\mathfrak{X}^\vee$ and its inclusion is denoted $j_{\mathrm{dR}}: \mathfrak{X}^\Delta \rightarrow \mathfrak{X}^\vee$. There is also another open immersion $j_{\mathrm{HT}}: \mathfrak{X}^\Delta \rightarrow \mathfrak{X}^\vee$ called the **Hodge–Tate locus** which is physically disjoint from the de Rham locus. The stack $\mathfrak{X}^{\mathrm{syn}}$ is the coequalizer of these two open immersions and is therefore equipped with a canonical open immersion $j_\Delta: \mathfrak{X}^\Delta \rightarrow \mathfrak{X}^{\mathrm{syn}}$.

⁶This means that they are étale (and in fact fpqc) sheaves on p -nilpotent derived affine schemes, i.e. derived schemes of the form $\mathrm{Spec} A$ where A is a p -nilpotent animated commutative ring.

[1] **Alex:** Add something about o tions/convent B(G).

[2] **Alex:** Som about the sub and $\setminus X$.

[3] **Alex:** Som about the me $\Lambda^\otimes$.

[4] **Alex:** Giv references to C

cohomology

Remark 2.1.3 (Nygaard filtered absolute prismatic cohomology). Bhatt–Lurie show that the values of these stacks on quasiregular semiperfectoid (qrsp) inputs can be described in terms of Nygaard filtered absolute prismatic cohomology. For such rings R , their **absolute prismatic cohomology** is a classical p -complete ring Δ_R , which is in fact a δ -ring equipped with a canonical prism structure (Δ_R, I_R) : this is also the *initial* prism for R . In [9, §12], Bhatt–Scholze construct the **Nygaard filtration** $\mathrm{Fil}_{\mathcal{N}}^{\bullet} \Delta_R$ on absolute prismatic cohomology. When R is qrsp, we have

$$\mathrm{Fil}_{\mathcal{N}}^i \Delta_R = \{x \in \Delta_R : \varphi(x) \in \mathrm{Fil}_{I_R}^i \Delta_R\},$$

where $\mathrm{Fil}_{I_R}^{\bullet} \Delta_R$ is the I_R -adic filtration. The Nygaard filtered prismatization $R^{\mathcal{N}}$ is now canonically isomorphic to the *formal Rees stack* $\mathcal{R}(\mathrm{Fil}_{\mathcal{N}}^{\bullet} \Delta_R)$ associated with this filtration (e.g., see [39, §1.1.2]). The open immersion j_{dR} amounts to ‘forgetting’ the filtration (that is, restriction to the open substack $\mathbb{G}_m/\mathbb{G}_m \subset \mathbb{A}^1/\mathbb{G}_m$), while j_{HT} arises from the filtered Frobenius lift

$$\mathrm{Fil}_{\mathcal{N}}^{\bullet} \Delta_R \rightarrow \mathrm{Fil}_{I_R}^{\bullet} \Delta_R.$$

This description can be extended to all *semiperfectoid* rings; see [26, Theorem 6.11.7].

rham_point

Remark 2.1.4 (The (filtered) de Rham point). There are canonical maps

$$x_{\mathrm{dR}}^{\mathcal{N}} : \mathbb{A}^1/\mathbb{G}_m \times \mathfrak{X} \rightarrow \mathfrak{X}^{\mathcal{N}} ; x_{\mathrm{dR}} : \mathfrak{X} \rightarrow \mathfrak{X}^{\Delta}$$

with the second being the restriction of the first over $\mathbb{G}_m/\mathbb{G}_m$. In terms of the ‘affine’ description from Remark 2.1.3 for semiperfectoid rings, the first map corresponds to the map on formal Rees stacks associated to the canonical map of filtered rings from $\mathrm{Fil}_{\mathcal{N}}^{\bullet} \Delta_R$ to R equipped with its trivial descending filtration supported in graded degree 0.

Remark 2.1.5 (Functoriality and étale descent). Each of the assignments $\mathfrak{X} \mapsto \mathfrak{X}^?$, for $?$ equal to Δ , $\mathcal{N}$, or syn , is functorial in the derived formal scheme $\mathfrak{X}$, and preserves products: That is, we have canonical isomorphisms $(\mathfrak{X} \times \mathfrak{Y})^? \simeq \mathfrak{X}^? \times \mathfrak{Y}^?$. Furthermore, if $\mathfrak{Y} \rightarrow \mathfrak{X}$ is an étale and faithfully flat map⁷, then $\mathfrak{Y}^? \rightarrow \mathfrak{X}^?$ is an étale cover of derived formal stacks (see [26, Proposition 6.12.3]).

Definition 2.1.6. A map $\mathfrak{Y} \rightarrow \mathfrak{X}$ of derived p -adic formal schemes is **quasisyntomic** if it is p -completely flat (that is, $\mathfrak{Y} \otimes^{\mathbb{L}} \mathbb{F}_p \rightarrow \mathfrak{X} \otimes^{\mathbb{L}} \mathbb{F}_p$ is flat), and if $\mathbb{L}_{\mathfrak{Y}/\mathfrak{X}}$ has p -complete Tor amplitude $[-1, 0]$: that is, $\mathbb{L}_{\mathfrak{Y}/\mathfrak{X}}/\mathbb{L}^p$ has Tor amplitude $[-1, 0]$ as a quasi-coherent sheaf on $\mathfrak{Y} \otimes^{\mathbb{L}} \mathbb{F}_p$. The map $\mathfrak{Y} \rightarrow \mathfrak{X}$ is a **quasisyntomic cover** if it is quasisyntomic and $\mathfrak{Y} \otimes^{\mathbb{L}} \mathbb{F}_p \rightarrow \mathfrak{X} \otimes^{\mathbb{L}} \mathbb{F}_p$ is an fpqc cover.

nt_descent

Proposition 2.1.7. *If $\mathfrak{Y} \rightarrow \mathfrak{X}$ is a quasisyntomic cover, then $\mathfrak{Y}^{\mathcal{N}} \rightarrow \mathfrak{X}^{\mathcal{N}}$ is surjective in the p -completely flat topology. In fact, if $\mathfrak{Y} = \mathrm{Spf} S$ and $\mathfrak{X} = \mathrm{Spf} R$ with R and S semiperfectoid rings, then the map $\mathfrak{Y}^{\mathcal{N}} \rightarrow \mathfrak{X}^{\mathcal{N}}$ is faithfully flat.*

Proof. See [26, Proposition 6.12.3 and Corollary 6.12.8]. $\square$

tomic_qrsp

Remark 2.1.8. If $\mathfrak{X} = \mathrm{Spf} R$ is affine and the structure map $\mathfrak{X} \rightarrow \mathrm{Spf}(\mathbb{Z}_p)$ is quasisyntomic, there exists an affine quasisyntomic cover $\mathrm{Spf} R_{\infty} = \mathfrak{X}_{\infty} \rightarrow \mathfrak{X}$ such that $R_{\infty}^{\otimes_{Rm}}$ is qrsp and p -torsion free for all m (see [7, Lemmas 4.28 and 4.30]). Therefore, in many cases, Proposition 2.1.7 reduces questions about the stacks $\mathfrak{X}^?$ to ones about the corresponding stacks associated with (the formal spectra of) qrsp and p -torsion free rings.

d_f_gauges

2.2. F -gauges and prismatic F -crystals. Let $\mathfrak{X}$ be a derived p -adic formal scheme.

Definition 2.2.1 ((Prismatic) F -gauges). A **vector bundle in prismatic F -gauges** or simply **vector bundle F -gauge** over $\mathfrak{X}$ is a vector bundle over $\mathfrak{X}^{\mathrm{syn}}$. We will write $\mathrm{Vect}(\mathfrak{X}^{\mathrm{syn}})$ for the category of such objects.⁸

f-crystals

Remark 2.2.2 (Prismatic F -crystals). Recall from [8] the notion of *prismatic crystals* and *prismatic F -crystals* over $\mathfrak{X}$: These are objects over the absolute prismatic site $\mathfrak{X}_{\Delta}$ (shortened to R_{Δ} when $\mathfrak{X} = \mathrm{Spf}(R)$). To begin, we have the *structure sheaf* $\mathcal{O}_{\Delta}$ and a generalized Cartier divisor $\mathcal{I}_{\Delta} \rightarrow \mathcal{O}_{\Delta}$ given by the assignment

$$(\mathcal{I}_{\Delta} \rightarrow \mathcal{O}_{\Delta}) : (A, I, \mathrm{Spf}(\overline{A}) \rightarrow \mathfrak{X}) \mapsto (I \rightarrow A).$$

⁷Recall this means that the induced map of schemes $\mathfrak{Y} \otimes^{\mathbb{L}} \mathbb{F}_p \rightarrow \mathfrak{X} \otimes^{\mathbb{L}} \mathbb{F}_p$ is étale and faithfully flat.

⁸In general, $\mathrm{Vect}(\mathfrak{X}^{\mathrm{syn}})$ is actually an ∞ -category, but when $\mathfrak{X}$ is p -quasisyntomic and flat over $\mathbb{Z}_p$, Remark 2.1.8 combined with Remark 2.1.3 tells us that this is in fact a classical category.

A **prismatic crystal** (in vector bundles) over $\mathfrak{X}$ is a vector bundle $\mathcal{E}$ over $(\mathfrak{X}_\Delta, \mathcal{O}_\Delta)$, and a **prismatic F -crystal** is a pair $(\mathcal{E}, \varphi_\mathcal{E})$ where $\mathcal{E}$ is a prismatic crystal and

$$\varphi_\mathcal{E} : \varphi^* \mathcal{E}[1/I_\Delta] \xrightarrow{\sim} \mathcal{E}[1/I_\Delta]$$

is an $\mathcal{O}_\Delta[1/I_\Delta]$ -linear isomorphism of sheaves on $\mathfrak{X}_\Delta$. Write $\text{Vect}^\varphi(\mathfrak{X}_\Delta, \mathcal{O}_\Delta)$ for the category of prismatic F -crystals over $\mathfrak{X}$.

f-crystals

Remark 2.2.3. Concretely, a prismatic crystal is an assignment

$$(A, I, \text{Spf}(\overline{A}) \rightarrow \mathfrak{X}) \mapsto \mathcal{E}(A, I, \text{Spf}(\overline{A}) \rightarrow \mathfrak{X})$$

with the value being a finite locally free A -module, and satisfying the usual crystal property: For all maps $(A, I, \text{Spf}(\overline{A}) \rightarrow \mathfrak{X}) \rightarrow (B, J, \text{Spf}(\overline{B}) \rightarrow \mathfrak{X})$ in $\mathfrak{X}_\Delta$, we have isomorphisms

$$\mathcal{E}(A, I, \text{Spf}(\overline{A}) \rightarrow \mathfrak{X}) \otimes_A B \xrightarrow{\sim} \mathcal{E}(B, J, \text{Spf}(\overline{B}) \rightarrow \mathfrak{X})$$

satisfying the expected compatibility relations. Endowing this with the structure of a prismatic F -crystal now amounts to giving isomorphisms

$$\mathcal{E}(A, I, R \rightarrow \overline{A})[1/I] \otimes_{A, \varphi} A \xrightarrow{\sim} \mathcal{E}(A, I, R \rightarrow \overline{A})[1/I]$$

that are compatible with the isomorphisms coming from the crystal property.

-prismatic

Remark 2.2.4 (F -gauges and prismatic F -crystals). By Remark 2.1.1, any vector bundle on $\mathfrak{X}^\Delta$ yields a prismatic crystal over $\mathfrak{X}$. If the vector bundle arises from a vector bundle F -gauge via pullback along $j_\Delta : \mathfrak{X}^\Delta \rightarrow \mathfrak{X}^{\text{syn}}$, then $\mathcal{E}$ has a canonical structure of a prismatic F -crystal over $\mathfrak{X}$; see [5, Remark 6.3.4] and [39, Construction 1.21]. This gives rise to a natural functor $\text{Vect}(\mathfrak{X}^{\text{syn}}) \rightarrow \text{Vect}^\varphi(\mathfrak{X}_\Delta, \mathcal{O}_\Delta)$.

_prismatic

Proposition 2.2.5. Suppose that $\mathfrak{X}$ is p -quasisyntomic and flat over $\mathbb{Z}_p$. Then the natural functor

$$\text{Vect}(\mathfrak{X}^{\text{syn}}) \rightarrow \text{Vect}^\varphi(\mathfrak{X}_\Delta, \mathcal{O}_\Delta)$$

is fully faithful

Proof. See [33, Corollary 3.53]. □

realization

2.3. Analytic prismatic F -crystals and crystalline local systems. We now recall some generalizations due to Du–Liu–Moon–Shimizu [25] and Guo–Reinecke [34] of the classification of Bhatt–Scholze [8] of crystalline Galois representations. For more details, see the discussion in [37, §2.3]. In this subsection, K will denote a complete mixed characteristic $(0, p)$ discrete valuation field with perfect residue field.

Remark 2.3.1 (Analytic prismatic F -crystals). We will take $\text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_\Delta, \mathcal{O}_\Delta)$ to be the category of *analytic prismatic F -crystals* defined in [34, §3]. Concretely, this amounts to giving essentially the same data as that of a prismatic F -crystal, explained in Remark 2.2.3, except that $\mathcal{E}(A, I, \text{Spf}(\overline{A}) \rightarrow \mathfrak{X})$ is now a vector bundle over $\text{Spec}(A) \setminus V(p, I)$. Lemma 3.4 of [34] ensures that this is a sensible notion.

realization

Remark 2.3.2 (The étale realization). Via [5, Constructions 6.3.1 and 6.3.2] (see also [34, Construction 3.9]), we find a canonical functor

$$T_{\text{ét}} : \text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_\Delta, \mathcal{O}_\Delta) \rightarrow \text{Loc}_{\mathbb{Z}_p}(\mathfrak{X}_\eta)$$

where the right-hand side is the category of proétale $\mathbb{Z}_p$ -local systems on the generic fiber of $\mathfrak{X}$. Since both sides satisfy quasisyntomic descent (see [34, Lemma 3.6] for descent for analytic prismatic F -crystals), it suffices to specify the values for semiperfectoid $\text{Spf}(R) \rightarrow \mathfrak{X}$ with R semiperfectoid as:

$$\Gamma((\text{Spf } R)_\eta, T_{\text{ét}}(\mathcal{E})) = (\mathcal{E}(\Delta_R, I_R, \text{Spf}(\overline{\Delta}_R) \rightarrow \text{Spf}(R) \rightarrow \mathfrak{X})[1/I_R]^\wedge)^{\varphi_\mathcal{E} = \text{id}}.$$

Here, we are viewing $\mathcal{E}(\Delta_R, I_R, \text{Spf}(\overline{\Delta}_R) \rightarrow \text{Spf}(R) \rightarrow \mathfrak{X})[1/I_R]$ as a finite locally free module over $\Delta_R[1/I_R]$, and $(-)^\wedge$ denotes the p -adic completion.

Remark 2.3.3 (Relationship between various types of local systems). In the sequel we will frequently make use of the following natural equivalences:

- (1) Let X be a locally of finite type K -scheme, and $X^{\text{an}} \stackrel{\text{defn}}{=} X \times_{\text{Spec}(K)} \text{Spa}(K)$ its analytification. Then, there is a (bi-)exact monoidal equivalence $\text{Loc}_{\mathbb{Z}_p}(X) \xrightarrow{\sim} \text{Loc}_{\mathbb{Z}_p}(X^{\text{an}})$ (see [37, §2.1.3] and the references therein).
- (2) If (A, A^+) is a Huber pair, there is a (bi-)exact monoidal equivalence

$$\text{Loc}_{\mathbb{Z}_p}(\text{Spec}(A)) \xrightarrow{\sim} \text{Loc}_{\mathbb{Z}_p}(\text{Spa}(A, A^+)).$$

See [37, §2.1.4] and the references therein.

We will often glibly use the above identifications without further comment. Moreover, by (2) (and the independence of base points for connected spaces) there is no ambiguity in the notation $\text{Loc}_{\mathbb{Z}_p}(A)$ for a Huber pair (A, A^+) , and we use this notational shortcut often.

Proposition 2.3.4. *Suppose that $\mathfrak{X}$ is p -quasisyntomic and flat over $\mathbb{Z}_p$. Then the natural functor*

$$\text{Vect}^\varphi(\mathfrak{X}_\Delta, \mathcal{O}_\Delta) \rightarrow \text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_\Delta, \mathcal{O}_\Delta)$$

is fully faithful.

Proof. See [34, Proposition 3.7]. □

Remark 2.3.5 (Crystalline local systems). Suppose that $\mathfrak{X}$ is a *base formal $\mathcal{O}_K$ -scheme* as defined in [37, §1.1] and the references therein. This implies in particular that R is p -quasisyntomic and p -torsion free. Then within $\text{Loc}_{\mathbb{Z}_p}(\mathfrak{X}_\eta)$, we have the full subcategory $\text{Loc}_{\mathbb{Z}_p}^{\text{crys}}(\mathfrak{X})$ of *crystalline $\mathbb{Z}_p$ -local systems*; see [25, §2.2], [34, §2.4], or [37, §2.3.3]. When $\mathfrak{X} = \text{Spf}(\mathcal{O}_K)$, this recovers the classical notion of a Galois stable $\mathbb{Z}_p$ -lattice in a crystalline $\mathbb{Q}_p$ -representation of the absolute Galois group of K . Recent work of Guo–Yang [35, Theorem 1.1] shows that, when $\mathfrak{X}$ is smooth over $\mathcal{O}_K$, a $\mathbb{Z}_p$ -local system over $\mathfrak{X}$ is crystalline if and only if its restriction over every classical point of $\mathfrak{X}_\eta$ is crystalline.

Theorem 2.3.6. *Suppose that $\mathfrak{X}$ is p -quasisyntomic and flat over $\mathbb{Z}_p$. Then:*

- (1) *The étale realization functor $T_{\text{ét}}: \text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_\Delta, \mathcal{O}_\Delta) \rightarrow \text{Loc}_{\mathbb{Z}_p}(\mathfrak{X}_\eta)$ is faithful. In particular, its restriction to $\text{Vect}(\mathfrak{X}^{\text{syn}})$ is also faithful.*
- (2) *Suppose that $\mathfrak{X}$ is smooth over $\mathcal{O}_K$; or, more generally, that $\mathfrak{X}$ is a base formal $\mathcal{O}_K$ -scheme. Then the functor induces a bi-exact symmetric monoidal equivalence*

$$\text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_\Delta, \mathcal{O}_\Delta) \xrightarrow{\sim} \text{Loc}_{\mathbb{Z}_p}^{\text{crys}}(\mathfrak{X}_\eta).$$

In particular, its restriction to $\text{Vect}(\mathfrak{X}^{\text{syn}})$ is fully faithful.

Proof. This is essentially a translation of results of Guo–Reinecke [34] and Du–Liu–Moon–Shimizu [25]).

The first result follows from Propositions 2.2.5 and 2.3.4 combined with [34, Proposition 3.7] and the proof of [34, Lemma 4.1]. Note that the last result only claims to prove faithfulness for R p -completely smooth over $\mathcal{O}_K$. However, the proof actually shows that the functor $T_{\text{ét}}$ is faithful for any p -torsion free qrsp algebra⁹, and so implies what we need by quasisyntomic descent.

The second assertion follows from [34, Theorem 4.5], [25, Theorem 3.29], and [37, Proposition 2.22]. □

Definition 2.3.7 (Local systems for smooth affine group schemes). Suppose that $\mathcal{G}$ is a smooth affine group scheme over $\mathbb{Z}_p$ and that X is a locally Noetherian scheme or adic space. A $\mathcal{G}(\mathbb{Z}_p)$ -**local system** over X is a $\mathcal{G}(\mathbb{Z}_p)$ -torsor on the proétale site of X .¹⁰ We will write $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(X)$ for the groupoid of such local systems. If X is affine/affinoid with global sections R , we will also write $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(R)$ instead. We will also use analogous notation for the variants where $\mathcal{G}(\mathbb{Z}_p)$ is replaced by $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$. See [37, §2.1] for a more detailed discussion.

⁹The argument in this generality is actually needed to apply this lemma to the proof of Theorem 4.5 in *op. cit.*

¹⁰Concretely, this amounts to giving a compatible inverse system $\mathcal{P} = \{\mathcal{P}_n\}_{n \geq 1}$ where $\mathcal{P}_n \rightarrow X$ is a finite étale torsor for the finite group $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ (viewed as a locally constant sheaf), and the maps $\mathcal{P}_{n+1} \rightarrow \mathcal{P}_n$ are equivariant for the natural surjections $\mathcal{G}(\mathbb{Z}/p^{n+1}\mathbb{Z}) \rightarrow \mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$.

Remark 2.3.8 (Tannakian perspective on local systems). Giving a $\mathcal{G}(\mathbb{Z}_p)$ -local system $\mathcal{P}$ over X is equivalent to giving an exact symmetric monoidal functor $V \mapsto (V)_{\mathcal{P}}$ from the category $\text{Rep}_{\mathbb{Z}_p}(\mathcal{G})$ of algebraic representations $\mathcal{G} \rightarrow \text{GL}(V)$ defined over $\mathbb{Z}_p$ to the category $\text{Loc}_{\mathbb{Z}_p}(X)$. See [37, Propositions 2.3 and 2.8]. In fact, the cited result shows that giving a $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -local system $\mathcal{P}_n$ over X is the same as giving an exact symmetric monoidal functor $\text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \rightarrow \text{Loc}_{\mathbb{Z}/p^n\mathbb{Z}}(X)$.

Definition 2.3.9 (Crystalline $\mathcal{G}(\mathbb{Z}_p)$ -local systems). Let $\mathfrak{X}$ be a base formal $\mathcal{O}_K$ -scheme. A $\mathcal{G}(\mathbb{Z}_p)$ -local system $\mathcal{P}$ on $\mathfrak{X}_{\eta}$ is **crystalline** if, for every representation V in $\text{Rep}_{\mathbb{Z}_p}(\mathcal{G})$, the local system $(V)_{\mathcal{P}}$ is crystalline. Write $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}}(\mathfrak{X}_{\eta})$ for the full subcategory of $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(\mathfrak{X}_{\eta})$ of crystalline $\mathcal{G}(\mathbb{Z}_p)$ -local systems.

Remark 2.3.10. By [37, Proposition 2.20], to check that a $\mathcal{G}(\mathbb{Z}_p)$ -local system $\mathcal{P}$ is crystalline, it is sufficient to check that $(V)_{\mathcal{P}}$ is crystalline for a single faithful representation V .

Remark 2.3.11 ($\mathcal{G}$ -bundles in analytic prismatic F -crystals). Suppose that $\mathfrak{X}$ is a base formal $\mathcal{O}_K$ -scheme. Combining Remark 2.3.8 with Theorem 2.3.6, we see that the category $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}}(\mathfrak{X}_{\eta})$ is equivalent to the category of exact symmetric monoidal functors

$$\omega: \text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \rightarrow \text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_{\Delta}, \mathcal{O}_{\Delta}).$$

Corollary 2.3.12. Suppose that $\mathfrak{X}$ is p -quasisyntomic and flat over $\mathbb{Z}_p$.

(1) There is a natural faithful functor

$$T_{\text{ét}}: B\mathcal{G}(\mathfrak{X}^{\text{syn}}) \rightarrow \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(\mathfrak{X}_{\eta}).$$

(2) If $\mathfrak{X}$ is a base $\mathcal{O}_K$ -formal scheme, then the functor is also full and takes its values in $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}}(\mathfrak{X}_{\eta})$.

Proof. Under these hypotheses, $\mathfrak{X}^{\text{syn}}$ is a classical formal stack. Therefore, we find from [64, Corollary 9.3.7.3] that the groupoid $(B\mathcal{G})(\mathfrak{X}^{\text{syn}})$ is equivalent to the groupoid of exact symmetric monoidal functors

$$\text{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \rightarrow \text{Vect}(\mathfrak{X}^{\text{syn}}).$$

The corollary now follows from Theorem 2.3.6 and Remark 2.3.8. $\square$

2.4. F -gauges and p -divisible groups.

Definition 2.4.1 (Hodge–Tate weights). Let $\mathfrak{X}$ be a p -adic formal scheme. Every vector bundle F -gauge $\mathcal{V}$ over $\mathfrak{X}$ yields a graded finite locally free R -module via pullback along the composition

$$B\mathcal{G}_m \times \mathfrak{X} \rightarrow \mathbb{A}^1/\mathbb{G}_m \times \mathfrak{X} \xrightarrow{x_{\text{dR}}^{\mathcal{N}}} \mathfrak{X}^{\mathcal{N}} \rightarrow \mathfrak{X}^{\text{syn}}.$$

The **Hodge–Tate weights** of $\mathcal{V}$ are the graded degrees in which this graded module is non-zero.

The following result is essentially due to Anschütz–Le Bras [3] over quasisyntomic bases; the general statement here is shown in [26]. See Theorem 11.1.4 and Proposition 11.8.2 of *op. cit.*

Theorem 2.4.2 (Prismatic Dieudonné theory). Let $\mathfrak{X}$ be a derived p -adic formal scheme and let $B\mathcal{T}_p(\mathfrak{X})$ be the category of p -divisible groups over $\mathfrak{X}$, and let $\text{Vect}_{[0,1]}(\mathfrak{X}^{\text{syn}})$ be the category of vector bundle F -gauges over $\mathfrak{X}$ with Hodge–Tate weights in $\{0, 1\}$. Then:

(1) There is a canonical equivalence of categories

$$\mathcal{M}: B\mathcal{T}_p(\mathfrak{X}) \xrightarrow{\sim} \text{Vect}_{[0,1]}(\mathfrak{X}^{\text{syn}})$$

compatible with Cartier duality.

(2) There is a natural equivalence $T_p \simeq T_{\text{ét}} \circ \mathcal{M}$, where T_p is the functor associating to a p -divisible group H its Tate module $T_p(H) = \varprojlim H[p^n]_{\eta}$, viewed as a $\mathbb{Z}_p$ -local system on $\mathfrak{X}_{\eta}$.

cohomology

2.5. F -gauges and cohomology. In this final subsection we recall the relationship between F -gauges, étale realization, and higher pushforwards. Throughout K denotes a complete mixed characteristic $(0, p)$ discrete valuation field with perfect residue field k , and C a completed algebraic closure of K .

Setup 2.5.1. Let $\mathfrak{X}$ be a formal $\mathcal{O}_K$ -scheme with generic fiber X . We denote by $D_{\text{lis}}^b(X, \mathbb{Z}_p)$ the category of derived p -complete locally bounded $\mathbb{Z}_p$ -modules $\mathbb{L}$ on $X_{\text{proét}}$ whose cohomology sheaves $H^i(\mathbb{L})$ are locally constant on $X_{\text{proét}}$ with finitely generated stalks. If $\mathfrak{X}$ is a smooth formal $\mathcal{O}_K$ -scheme, let $D_{\text{crys}}^b(X, \mathbb{Z}_p)$ be the full subcategory of those $\mathbb{L}$ such that $H^i(\mathbb{L})[1/p]$ is crystalline for all i .

Remark 2.5.2 (Étale realization for perfect F -gauges). By [5, Constructions 6.3.1 and 6.3.2] there is an étale realization functor

$$T_{\text{ét}} : \text{Perf}(\mathfrak{X}^{\text{syn}}) \rightarrow D_{\text{lis}}^b(X, \mathbb{Z}_p).$$

By [75, Corollary 4.1], if $\mathfrak{X}$ is smooth, then this functor takes values in $D_{\text{crys}}^b(X, \mathbb{Z}_p)$. Moreover, the étale realization functor is compatible with that from Remark 2.3.2 via the natural embedding of $\text{Vect}^{\text{an}, \varphi}(\mathfrak{X}_{\Delta})$ into $\text{Perf}(\mathfrak{X}^{\text{syn}})$ via the functor $\Pi_{\mathfrak{X}}$ from [33, Theorem 3.32] (see also [75, §3.2]).

Remark 2.5.3 (Crystalline realization functor). Suppose that $\mathfrak{X}$ is a p -quasisyntomic p -adic formal scheme. Then there is a natural functor

$$T_{\text{crys}} : \text{Perf}(\mathfrak{X}^{\text{syn}}) \rightarrow \text{Perf}^{\varphi}((\mathfrak{X}_{p=0})_{\text{crys}}),$$

where the target is the category of perfect complexes in F -crystals on the absolute crystalline site of $\mathfrak{X}_{p=0}$. More precisely, if $\mathcal{V}$ is a perfect F -gauge, then the construction from 2.2.4 gives us a perfect complex of prismatic F -crystals on $\mathfrak{X}$ and by restriction a perfect complex of prismatic F -crystals on $\mathfrak{X}_{p=0}$, and one then applies the equivalence in [34, Theorem 6.4].

Remark 2.5.4 (Perfect complexes in F -isocrystals on smooth formal $\mathcal{O}_K$ -schemes). Suppose now that $\mathfrak{X}$ is a smooth formal $\mathcal{O}_K$ -scheme. In this case, one has a natural identification

$$\text{Perf}^{\varphi}((\mathfrak{X}_{p=0})_{\text{crys}})[1/p] \simeq \text{Perf}^{\varphi}((\mathfrak{X}_k/W(k))_{\text{crys}})[1/p].$$

Here the left-hand side is the category of perfect complexes in F -isocrystals on the absolute crystalline site of $\mathfrak{X}_{p=0}$, and the right-hand side the category of perfect complexes in F -isocrystals on the relative crystalline site $(\mathfrak{X}_k/W(k))_{\text{crys}}$. This identification is a version of Dwork's trick (see [34, Remark 2.15] and the references therein). In the sequel we will often implicitly make this identification.

Remark 2.5.5 (The filtered crystalline realization functor on smooth formal $\mathcal{O}_K$ -scheme). Suppose again that $\mathfrak{X}$ is a smooth formal $\mathcal{O}_K$ -scheme. Then, the crystalline realization functor admits a filtered upgrade

$$T_{\text{crys}}^+ : \text{Perf}(\mathfrak{X}^{\text{syn}}) \rightarrow \text{Perf}^{\varphi}(\mathfrak{X}_{\text{crys}})[1/p],$$

where the target is the category of perfect complexes in *filtered* F -isocrystals on $\mathfrak{X}$ as in [75, Definition 4.18], and where T_{crys}^+ is denoted by Beil in *op. cit.*

The following shows that all of these constructions are compatible with pushforwards. For the proof (due to combining work from [34], [33], and [75]) one can see [75, Corollary 4.7] and [75, Proposition 4.23], their proofs, and the references therein.

pushforward

Theorem 2.5.6. Suppose that $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ is a smooth proper map of smooth formal $\mathcal{O}_K$ -schemes with generic fiber map $f_{\eta} : X \rightarrow Y$. Then, pushforward defines a well-defined commutative square

$$\begin{array}{ccc} \text{Perf}(\mathfrak{X}^{\text{syn}}) & \xrightarrow{T_{\text{ét}}} & D_{\text{crys}}^b(X, \mathbb{Z}_p) \\ Rf_* \downarrow & & \downarrow R(f_{\eta})_* \\ \text{Perf}(\mathfrak{Y}^{\text{syn}}) & \xrightarrow{T_{\text{ét}}} & D_{\text{crys}}(Y, \mathbb{Z}_p). \end{array}$$

Moreover, for any object $\mathcal{E}$ of $\text{Perf}(\mathfrak{X}^{\text{syn}})$ and $i \geq 0$ there is a natural identification of filtered F -isocrystals

$$D_{\text{crys}}(R^i(f_{\eta})_* T_{\text{ét}}(\mathcal{E})[1/p]) \simeq R^i(f_k)_* T_{\text{crys}}^+(\mathcal{E})[1/p]$$

rd_general

Remark 2.5.7 (Pushforward over a general base). In fact, for any smooth proper map $\mathfrak{X} \rightarrow \mathfrak{Y}$ of formal $\mathcal{O}_K$ -schemes, we always have a well-defined pushforward map

$$Rf_* : \text{Perf}(\mathfrak{X}^{\text{syn}}) \rightarrow \text{Perf}(\mathfrak{Y}^{\text{syn}}).$$

This has the property that it carries perfect F -gauges over $\mathfrak{X}$ with Hodge–Tate weights in $[a, b]$ to those over $\mathfrak{Y}$ with Hodge–Tate weights in $[a - d, b]$, where d is the relative dimension of $\mathfrak{X}$ over $\mathfrak{Y}$; see [67, Proposition 8.2.6].

Corollary 2.5.8. *Suppose that $\mathfrak{X}$ is a smooth proper formal $\mathcal{O}_K$ -scheme and $\mathcal{V}$ is a prismatic F -gauge on $\mathfrak{X}$. Then, for all $i \geq 0$ the $\text{Gal}(\overline{K}/K)$ -representation $H_{\text{et}}^i(\mathfrak{X}_C, T_{\text{et}}(\mathcal{V}))[1/p]$ is crystalline and there is a natural identification of filtered F -isocrystals on K :*

$$D_{\text{crys}}(H_{\text{et}}^i(\mathfrak{X}_C, T_{\text{et}}(\mathcal{V}))[1/p]) \simeq H_{\text{crys}}^i((\mathfrak{X}_k/W(k)_{\text{crys}}, T_{\text{crys}}^+(\mathcal{E})).$$

Moreover, for all $1 \leq n \leq \infty$ we have a natural identification

$$T_{\text{et}}(Rf_*(\mathcal{V}/p^n)) \simeq R(f_\eta)_* T_{\text{et}}(\mathcal{V}/p^n).$$

Definition 2.5.9 (Bloch–Kato’s finite part of Galois cohomology). Suppose now that K is a finite extension of $\mathbb{Q}_p$. In [10, §3] one finds the definition, for a Galois $\mathbb{Q}_p$ -representation V of K , a subspace $H_f^1(K, V)$ of the Galois cohomology of V called its *finite part*. When V is crystalline one naturally has

$$H_f^1(K, V) \simeq \text{Ext}^1(\mathbb{Q}_p, V)$$

where these extensions are taken in the category $\text{Rep}_{\mathbb{Q}_p}^{\text{crys}}(\text{Gal}(\overline{K}/K))$ (e.g., see [69, Proposition 4.5.2]).

The following is a result of combining Theorem 2.5.6 and [75, Proposition 4.9].

Proposition 2.5.10. *Suppose that K is a finite extension of $\mathbb{Q}_p$, and let $\mathfrak{X}$ is a smooth proper formal $\mathcal{O}_K$ -scheme and $\mathcal{V}$ a prismatic F -gauge on $\mathfrak{X}$. Then, for all $i \geq 0$ we have a natural identification*

$$H_f^1(H_{\text{et}}^i(\mathfrak{X}_C, T_{\text{et}}(\mathcal{V}))[1/p]) \simeq H^1(\mathcal{O}_K^{\text{syn}}, R^i f_* \mathcal{V})[1/p].$$

3. $(\mathcal{G}, \mu)$ -APERTURES

The purpose of this section is to recall the results of [26], and to record some complements that will be useful in the sequel.

3.1. The main representability result. This subsection is a quick review of [26, §9].

ec:gmudisp

in_results

setup:g_mu

Setup 3.1.1. In the following, $\mathcal{G}$ will be a smooth connected affine group scheme over $\mathbb{Z}_p$, $\hat{\mathcal{O}}$ the ring of integers in a finite unramified extension of $\mathbb{Q}_p$, and $\mu : \mathbb{G}_{m, \hat{\mathcal{O}}} \rightarrow \mathcal{G}_{\hat{\mathcal{O}}}$ a **1-bounded** cocharacter, whose weights on $\text{Lie}(\mathcal{G})_{\hat{\mathcal{O}}}$ via the adjoint action are bounded above by 1. Let $\mathcal{P}_\mu^- \subset \mathcal{G}$ be the smooth subgroup scheme whose Lie algebra is identified with the sum of the weight- i spaces in $\text{Lie}(\mathcal{G})_{\hat{\mathcal{O}}}$ for $i \leq 0$. When $\mathcal{G}$ is reductive, μ is minuscule and $\mathcal{P}_\mu^-$ is a parabolic subgroup associated with μ .

torsor_Qmu

Remark 3.1.2. Associated with μ is the map of classifying stacks $B\mathbb{G}_{m, \hat{\mathcal{O}}} \rightarrow B\mathcal{G}_{\hat{\mathcal{O}}}$. This classifies a $\mathcal{G}$ -torsor $\mathcal{Q}_\mu$ over $B\mathbb{G}_{m, \hat{\mathcal{O}}}$.

Definition 3.1.3. For $R \in \text{CRing}_{\hat{\mathcal{O}}}^{p\text{-comp}}$, an n -**truncated $(\mathcal{G}, \mu)$ -aperture** over R is a $\mathcal{G}$ -torsor $\mathfrak{Q}$ over $R^{\text{syn}} \otimes \mathbb{Z}/p^n \mathbb{Z}$ ¹¹ satisfying the following condition: For every geometric point $R \rightarrow \kappa$ of $\text{Spf } R$, the restriction of $(x_{dR}^\vee)^* \mathfrak{Q}$ (see Remark 2.1.4) over $B\mathbb{G}_m \times \text{Spec } \kappa$ is isomorphic to that of $\mathcal{Q}_\mu$. These organize into an ∞ -groupoid $\text{BT}_n^{\mathcal{G}, \mu}(R)$, and the assignment $R \mapsto \text{BT}_n^{\mathcal{G}, \mu}(R)$ is a derived p -adic formal prestack. We will also set

$$\text{BT}_\infty^{\mathcal{G}, \mu} = \varprojlim_n \text{BT}_n^{\mathcal{G}, \mu}.$$

Objects in $\text{BT}_\infty^{\mathcal{G}, \mu}(R)$ will be called **$(\mathcal{G}, \mu)$ -apertures** over R . They classify $\mathcal{G}$ -torsors on R^{syn} that are bounded by μ in the sense explained above.

¹¹By this, we mean a map of p -adic formal stacks $R^{\text{syn}} \otimes \mathbb{Z}/p^n \mathbb{Z} \rightarrow B\mathcal{G}$.

Remark 3.1.4. Note that the isomorphism class of $\mathcal{Q}_\mu$ depends only on the conjugacy class of μ . This implies that the stacks $\mathrm{BT}_n^{\mathcal{G},\mu}$ also only depend on this conjugacy class and not on the particular choice of representative.

The next result is [26, Theorem 9.3.2].

Theorem 3.1.5. *The formal prestack $\mathrm{BT}_n^{\mathcal{G},\mu}$ is represented by a zero-dimensional connected quasi-compact smooth p -adic formal Artin stack over $\hat{\mathcal{O}}$ with affine diagonal. Moreover, for any pro-nilpotent divided power thickening $R' \rightarrow R$ in $\mathrm{CRing}_{\hat{\mathcal{O}}}^{p\text{-comp}}$, we have a Cartesian Grothendieck–Messing square¹²*

$$(3.1.5.1) \quad \begin{array}{ccc} \mathrm{BT}_n^{\mathcal{G},\mu}(R') & \longrightarrow & B\mathcal{P}_\mu^-(R'/\mathbb{L}p^n) \\ \downarrow & & \downarrow \\ \mathrm{BT}_n^{\mathcal{G},\mu}(R) & \longrightarrow & B\mathcal{P}_\mu^-(R/\mathbb{L}p^n) \times_{B\mathcal{G}(R/\mathbb{L}p^n)} B\mathcal{G}(R'/\mathbb{L}p^n). \end{array}$$

The transition maps $\mathrm{BT}_{n+1}^{\mathcal{G},\mu} \rightarrow \mathrm{BT}_n^{\mathcal{G},\mu}$ are smooth and surjective.

Example 3.1.6 (The case of GL_n and truncated p -divisible groups). A basic example is (GL_h, μ_d) where $d \leq h$ and μ_d is a minuscule cocharacter of GL_h defined over $\mathbb{Z}_p$ and splitting the direct summand $\mathbb{Z}^{h-d} \times \{0\} \subset \mathbb{Z}^h$. In this case, it is shown in [26, Theorem 11.2.7] that $\mathrm{BT}_n^{\mathrm{GL}_h, \mu_d}$ is canonically isomorphic to the stack of n -truncated Barsotti–Tate (or p -divisible) groups of height h and dimension d . This is a refinement of Theorem 2.4.2.

Remark 3.1.7 (Deformation rings). For any point $x \in \mathrm{BT}_\infty^{\mathcal{G},\mu}(\kappa)$ valued in a perfect field κ , the deformation functor for $\mathrm{BT}_\infty^{\mathcal{G},\mu}$ at x takes discrete values and is represented by a complete local formally smooth $W(\kappa)$ -algebra of relative dimension $\dim \mathcal{G} - \dim \mathcal{P}_\mu^-$. This is explained in [26, Lemma 10.2.5]. It can also be seen from the results of Ito [40], combined with *a priori* knowledge of the formal smoothness of $\mathrm{BT}_\infty^{\mathcal{G},\mu}$; see [39, Proposition 3.32].

Remark 3.1.8 (Classicality). There is a somewhat interesting phenomenon here. When R is a discrete p -complete ring, even though R^{syn} is in general only a derived stack that is covered flat locally by the spectra of animated (not necessarily discrete) rings, $\mathrm{BT}_n^{\mathcal{G},\mu}(R)$ is still just a 1-groupoid (or a 1-truncated ∞ -groupoid). When R/pR satisfies a mild finiteness condition, this is now explained by [67, Remark 9.9.11], which tells us that $\mathrm{BT}_n^{\mathcal{G},\mu}(R)$ can be computed using only the classical truncation of R^{syn} .

3.2. Frames and windows. In [26, §5 and §10], we find a general discussion of frames and windows, which allow us to give more concrete descriptions of the category of $(\mathcal{G}, \mu)$ -apertures in certain situations. Here, we will isolate a particular instance that will prove very useful to us.

Definition 3.2.1 (Breuil–Kisin frames). Let R be a p -complete ring. A **Breuil–Kisin frame** for R is a prism $\underline{A} = (A, I')$ such that A is p -complete and p -torsion free, $I' \subset A$ is an ideal and $R = A/I'$.

Example 3.2.2 (Frames for base $\mathcal{O}_K$ -algebras). Suppose that K is a complete discrete valuation field with perfect residue field κ , and set $K_0 = W(\kappa)[1/p] \subset K$. Fix a monic Eisenstein polynomial $E(u) \in W(\kappa)[u]$ such that $E(\pi) = 0$ for some uniformizer $\pi \in \mathcal{O}_K$. Suppose that R is a base $\mathcal{O}_K$ -algebra in the sense of [37, §1.1]: Then $R = \mathcal{O}_K \otimes_{W(\kappa)} R_0$ where R_0 is a flat $W(\kappa)$ -algebra that admits a Frobenius lift and hence a δ -structure. The prism $(\mathfrak{S}_R, (E(u)))$ where $\mathfrak{S}_R = R_0[[u]]$ is equipped with the δ - R_0 -algebra structure satisfying $\delta(u) = 0$ is now a Breuil–Kisin frame for R .

Example 3.2.3 (Frames for perfectoid rings). If R is a perfectoid ring, then $\underline{A}_R = (A_R, \mathrm{Fil}_{\mathcal{N}}^1 A_R)$ is a Breuil–Kisin frame for R . Moreover, one has an identification $A_R \simeq A_{\mathrm{inf}}(R)$ under which $\mathrm{Fil}_{\mathcal{N}}^1 A_R$ corresponds to $\varphi^{-1}(\ker \theta)$ where $\theta: A_{\mathrm{inf}}(R) \rightarrow R$ is Fontaine’s map.

¹²This diagram does depend on the choice of PD structure on $\ker(R' \rightarrow R)$, as well as the fact that it is (pro-)nilpotent. For example, one may apply it to the usual trivial PD structure on $\ker(\mathbb{Z}/4\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z})$ but not to the one inherited from the unique PD structure on $2\mathbb{Z}_2 \subseteq \mathbb{Z}_2$.

Example 3.2.4 (Frames via Frobenius liftings). Suppose that R is a flat $\hat{\mathcal{O}}$ -algebra equipped with a Frobenius lift $\varphi: R \rightarrow R$. We can then view $(R, (p))$ as a Breuil–Kisin frame $\underline{R}$ for R/pR .

Remark 3.2.5 (Liftings of étale covers). Suppose that $\underline{A}$ is a Breuil–Kisin frame for R . Then any p -completely étale map $R \rightarrow \tilde{R}$ lifts uniquely to a (p, I') -completely étale map $A \rightarrow \tilde{A}$ of δ -rings, and $(\tilde{A}, I' \tilde{A})$ is a Breuil–Kisin frame for $\tilde{R}$. See [26, Proposition 5.4.23].

Remark 3.2.6 (The Rees stack associated with a Breuil–Kisin frame). Suppose⁵ that $\underline{A}$ is a Breuil–Kisin frame for R . Set $I = \varphi^* I' \subset A$. We then have the (p, I') -complete formal Rees stack $\mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$ associated with the I' -adic filtration on A , and the Frobenius lift on A extends to a map of Rees stacks $\mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A) \rightarrow \mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$. We obtain two maps $\tau, \sigma: \mathrm{Spf} A \rightarrow \mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$, where τ is the pullback of $\mathbb{G}_m/\mathbb{G}_m \rightarrow \mathbb{A}^1/\mathbb{G}_m$ and σ is obtained from the filtered Frobenius lift and the isomorphism

$$\mathrm{Spf} A \xrightarrow{\sim} \mathcal{R}(\mathrm{Fil}_{I, \pm}^\bullet, A).$$

Note also that we have a canonical map $x_{\underline{A}}: \mathbb{A}^1/\mathbb{G}_m \times \mathrm{Spf} R \rightarrow \mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$ associated with the map $\mathrm{Fil}_{I'}^\bullet A \rightarrow \mathrm{Fil}_{\mathrm{triv}}^\bullet R$ of filtered rings.

Definition 3.2.7 ($\mathcal{G}$ -torsors over the Rees stack bounded by μ). A $\mathcal{G}$ -torsor $\mathcal{Q}_{\mathrm{Rees}}$ over $\mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$ is **bounded by μ** if, for any map $R \rightarrow \kappa$ to an algebraically closed field κ , the restriction of $x_{\underline{A}}^* \mathcal{Q}_{\mathrm{Rees}}$ along the composition

$$B\mathbb{G}_m \times \mathrm{Spec} \kappa \rightarrow B\mathbb{G}_m \times \mathrm{Spf} R \rightarrow \mathbb{A}^1/\mathbb{G}_m \times \mathrm{Spf} R$$

is isomorphic to $\mathcal{Q}_\mu$.

Remark 3.2.8 (Consequences of 1-boundedness). Since we have assumed that μ is 1-bounded, a $\mathcal{G}$ -torsor $\mathcal{Q}_{\mathrm{Rees}}$ over $\mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$ bounded by μ is determined up to isomorphism by the following data:

- The $\mathcal{G}$ -torsor $\mathcal{Q} \stackrel{\mathrm{def}}{=} \tau^* \mathcal{Q}_{\mathrm{Rees}}$ over $\mathrm{Spf} A$;
- The $\mathcal{G}$ -torsor $x_{\underline{A}}^* \mathcal{Q}_{\mathrm{Rees}}$ over $\mathbb{A}^1/\mathbb{G}_m \times \mathrm{Spf} R$, which, by the bounded-by- μ condition, is equivalent to giving a reduction of structure $Q^- \subset Q$ to a $\mathcal{P}_\mu^-$ -torsor for the $\mathcal{G}$ -torsor Q over $\mathrm{Spf} R$ obtained via restriction from $\mathcal{Q}$.

This can be deduced for instance from [26, Proposition 4.12.3].

Remark 3.2.9 (Description in terms of modifications). As a particular consequence of the description in Remark 3.2.8, we find that we can associate with the pair $(\mathcal{Q}, Q^- \subset Q)$ the $\mathcal{G}$ -torsor $\sigma^* \mathcal{Q}_{\mathrm{Rees}}$ over $\mathrm{Spf} A$. This can be obtained directly from the pair. For simplicity, we will restrict ourselves to the case where $I' = (E)$ is principal with a distinguished choice of generator E . Consider the sheaf of groups

$$H_\mu: \tilde{R} \mapsto \mathcal{G}(\tilde{A}) \times_{\mathcal{G}(\tilde{R})} \mathcal{P}_\mu^-(\tilde{R}) \subset \mathcal{G}(\tilde{A}) \stackrel{\mathrm{def}}{=} (L^+ \mathcal{G})(\tilde{R})$$

on the étale site of $\mathrm{Spf} R$, and note that conjugation by $\mu(E) \in \mathcal{G}(A[1/E])$ yields a map

$$\mathrm{int}(\mu(E)): H_\mu \rightarrow L^+ \mathcal{G}.$$

We can view $\mathcal{Q}$ as an $L^+ \mathcal{G}$ -torsor, and Q^- as yielding a reduction of structure group to an H_μ -torsor $Q^- \subset \mathcal{Q}$. Pushforward of Q^- along $\mathrm{int}(\mu(E))$ now yields a modification $\mathcal{Q}'$ of $\mathcal{Q}$ and a further pushforward along the endomorphism of $L^+ \mathcal{G}$ induced by the Frobenius lift φ on A now yields the $\mathcal{G}$ -torsor $\sigma^* \mathcal{Q}_{\mathrm{Rees}}$.

Definition 3.2.10 ($(\mathcal{G}, \mu)$ -windows over Breuil–Kisin frames). Suppose that $\underline{A}$ is a Breuil–Kisin frame for R . A $(\mathcal{G}, \mu)$ -**window** over $\underline{A}$ is a $\mathcal{G}$ -torsor $\mathcal{Q}$ over $\mathcal{R}(\mathrm{Fil}_{I'}^\bullet, A)$ bounded by μ and equipped with an isomorphism $\alpha: \sigma^* \mathcal{Q} \xrightarrow{\sim} \tau^* \mathcal{Q}$. Write $\mathrm{Wind}_{\underline{A}, \infty}^{\mathcal{G}, \mu}(R)$ for the groupoid of $(\mathcal{G}, \mu)$ -windows over $\underline{A}$.

Example 3.2.11 (The case of $\mathcal{G} = \mathrm{GL}_h$). Suppose that $(\mathcal{G}, \mu) = (\mathrm{GL}_h, \mu_d)$ from Example 3.1.6. Then, by [26, Proposition 5.6.8 and Corollary 5.7.4] and the argument from [67, Proposition 9.5.9], there is a canonical equivalence of groupoids

$$\bigsqcup_{d \leq h \in \mathbb{Z}_{\geq 0}} \mathrm{Wind}_{\underline{A}, \infty}^{\mathrm{GL}_h, \mu_d}(R) \xrightarrow{\sim} \mathrm{BK}_{\underline{A}, \infty}(R)$$

where the right-hand side is the groupoid of triples $(\mathcal{M}, \varphi_{\mathcal{M}}, V_{\mathcal{M}})$ where:

[5] **Alex:** May should also give context about it is confusing talked about from the 'class (e.g., as in Bh notes).

- $\mathcal{M}$ is a finite locally free A -module;
- $\varphi_{\mathcal{M}}: \varphi^* \mathcal{M} \rightarrow \mathcal{M}$ and $V_{\mathcal{M}}: I' \otimes_A \mathcal{M} \rightarrow \varphi^* \mathcal{M}$ such that the maps

$$V_{\mathcal{M}} \circ (1 \otimes \varphi_{\mathcal{M}}): I' \otimes_A \varphi^* \mathcal{M} \rightarrow \varphi^* \mathcal{M}; \quad \varphi_{\mathcal{M}} \circ V_{\mathcal{M}}: I' \otimes_A \mathcal{M} \rightarrow \mathcal{M}$$

are the maps induced by the inclusion $I' \subset A$;

It is easy to see that in fact the datum of $V_{\mathcal{M}}$ is redundant and can be replaced with the *condition* that $\varphi_{\mathcal{M}}$ is injective with cokernel a finite locally free module over R . In the case of Example 3.2.2, this is the category of *Breuil windows* over $\mathfrak{S}_R$ introduced in [92].

_alternate

Remark 3.2.12 (Alternate description via modifications). The groupoid $\text{Wind}_{\underline{A}, \infty}^{\mathcal{G}, \mu}(R)$ can also be described as follows using Remark 3.2.9 (after having chosen a generator $E \in I'$): An object here is a triple $(\mathcal{Q}, Q^-, \alpha)$ where $\mathcal{Q}$ is a $\mathcal{G}$ -torsor over $\text{Spf } A$, $Q^- \subset \mathcal{Q}$ is a reduction of structure group of $\mathcal{Q} \stackrel{\text{defn}}{=} \mathcal{Q}|_{\text{Spf } R}$ to a $\mathcal{P}_{\mu}^-$ -torsor, and $\alpha: \varphi^* \mathcal{Q}' \xrightarrow{\sim} \mathcal{Q}$ is an isomorphism of $\mathcal{G}$ -torsors, where $\mathcal{Q}'$ is the modification of $\mathcal{Q}$ along Q^- . More explicitly, the sheaf of groupoids

$$\tilde{R} \mapsto \text{Wind}_{\underline{A}, \infty}^{\mathcal{G}, \mu}(\tilde{R})$$

on the étale site of $\text{Spf } R$ is the quotient $[L^+ \mathcal{G}/H_{\mu}]$ where H_{μ} acts on $L^+ \mathcal{G}$ via the right action

$$g \cdot h = \tau(h)^{-1} g \varphi(\text{int}(\mu(E))(h)),$$

where $\tau: H_{\mu} \rightarrow L^+ \mathcal{G}$ is the tautological map.

representations

Remark 3.2.13 (Modifications in terms of representations). Suppose that we have $\Lambda \in \text{Rep}_{\mathbb{Z}_p}(\mathcal{G})$ and let $\mathcal{V}$ be its twist by $\mathcal{Q}$. Then the associated twist $\mathcal{V}'$ by $\mathcal{Q}'$ can be described as follows: Let $V = R \otimes_A \mathcal{V}$. The reduction of structure Q^- yields a filtration $\text{Fil}^{\bullet} V$. Working étale locally on R if necessary we can assume that $\text{Fil}^{\bullet} V$ is split by a cocharacter of $\mathcal{G}_R$ that is geometrically conjugate to μ : This gives us a splitting $V = \bigoplus_i V^i$ where $\text{Fil}^i V = \bigoplus_{j \geq i} V^j$. Lift this to a cocharacter of $\mathcal{G}_A$ and consider the associated splitting $\mathcal{V} = \bigoplus_i \mathcal{V}^i$ (cf. [48, Proposition (1.1.5)]). Then we have the equality

$$\mathcal{V}' = \bigoplus_i E^{-i} \mathcal{V}^i = \sum_i E^{-i} \text{Fil}^i \mathcal{V} \subset \mathcal{V}[E^{-1}],$$

where $\text{Fil}^i \mathcal{V} = \bigoplus_{j \geq i} \mathcal{V}^j$. In particular, the isomorphism α yielding a window gives an isomorphism

$$\varphi^* \left(\sum_i E^{-i} \text{Fil}^i \mathcal{V} \right) \xrightarrow{\sim} \mathcal{V}.$$

pn_tensors

Setup 3.2.14 (Tensor packages). It will be convenient to make the following choices. Let $\mathcal{G} \rightarrow \text{GL}(\Lambda)$ be a faithful representation. We choose finitely many tensors $\{s_{\alpha}\} \subset \Lambda^{\otimes}$ such that $\mathcal{G}$ is their pointwise stabilizer: This is always possible by [13, Theorem 1.1].

r_packages

Remark 3.2.15 (Explicit description using tensor packages). Using Remark 3.2.9, we see that giving $\mathfrak{Q}$ in $\text{Wind}_{\underline{A}, \infty}^{\mathcal{G}, \mu}(R)$ is equivalent to the groupoid of triples $(\mathcal{Q}, Q^-, \xi)$, where:

- $\mathcal{Q}$ is a $\mathcal{G}$ -torsor over A ;
- $Q^- \subset \mathcal{Q} \stackrel{\text{defn}}{=} \mathcal{Q}|_{\text{Spec } R}$ is a reduction of structure group to a $\mathcal{P}_{\mu}^-$ -torsor;
- ξ is an isomorphism $\varphi^* \mathcal{Q}' \xrightarrow{\sim} \mathcal{Q}$ of $\mathcal{G}$ -torsors over A , where $\mathcal{Q}'$ is the modification of $\mathcal{Q}$ along Q^- .

This can be made even more concrete using the tensor package from Setup 3.2.14.

- (1) Giving $\mathfrak{Q}$ is equivalent to specifying a finite locally free A -module $\mathcal{L}$ and tensors $\{s_{\alpha, \mathcal{L}}\} \subset \mathcal{L}^{\otimes}$ such that, for some p -completely étale and faithfully flat map $R \rightarrow \tilde{R}$ with corresponding lift $A \rightarrow \tilde{A}$, there is an isomorphism

$$\eta: \tilde{A} \otimes_{\mathbb{Z}_p} \Lambda \xrightarrow{\sim} \tilde{A} \otimes_A \mathcal{L}$$

carrying $\{1 \otimes s_{\alpha}\}$ to $\{1 \otimes s_{\alpha, \mathcal{L}}\}$.

- (2) The reduction of structure group $Q^- \subset \mathcal{Q}$ amounts to giving a filtration $\text{Fil}^{\bullet} L$ on $L \stackrel{\text{defn}}{=} R \otimes_A \mathcal{L}$ such that for some $R \rightarrow \tilde{R}$ as above the isomorphism η can be chosen so that the induced filtration on $\tilde{R} \otimes_{\mathbb{Z}_p} \Lambda$ is split by μ .

- (3) If we set $\mathcal{L}^m \subset \mathcal{L}$ to be the pre-image of $\text{Fil}^m L$, the modification $\mathcal{Q}'$ now corresponds to the A -module

$$\mathcal{L}' = \sum_m E^{-m} \mathcal{L}^m \subset \mathcal{L}[1/E]$$

along with the same collection of tensors $\{s_{\alpha, \mathcal{L}}\}$, now viewed inside $\mathcal{L}', \otimes$.

- (4) The isomorphism ξ corresponds to an isomorphism of A -modules

$$\varphi^* \mathcal{L}' = \sum_m \varphi(E)^{-m} \varphi^* \mathcal{L}^m \xrightarrow{\sim} \mathcal{L}$$

carrying $\{\varphi^* s_{\alpha, \mathcal{L}}\}$ to $\{s_{\alpha, \mathcal{L}}\}$.

matization

Proposition 3.2.16 (Mapping the Rees stack to the filtered prismatization). *There is a canonical map of formal stacks $\iota_{\underline{A}}^{\mathcal{N}}: \mathcal{R}(\text{Fil}_I^\bullet, A) \rightarrow R^{\mathcal{N}}$ such that*

$$\iota_{\underline{A}}^{\mathcal{N}} \circ \tau = j_{\text{dR}} \circ \iota_{(A, I)}; \quad \iota_{\underline{A}}^{\mathcal{N}} \circ \sigma = j_{\text{HT}} \circ \iota_{(A, I)}; \quad \iota_{\underline{A}}^{\mathcal{N}} \circ x_{\underline{A}} = x_{\text{dR}}^{\mathcal{N}}.$$

Proof. See [26, Example 6.10.5]. □

to_windows

Corollary 3.2.17 (Apertures to windows). *Suppose that R is a p -complete $\hat{\mathcal{O}}$ -algebra. Then there is a canonical functor*

$$\text{BT}_{\infty}^{\mathcal{G}, \mu}(R) \rightarrow \text{Wind}_{\underline{A}, \infty}^{\mathcal{G}, \mu}(R).$$

The next result is a special case of [26, Lemma 9.2.3].

oid_rings

Proposition 3.2.18 (Description for perfectoid rings). *Suppose that R is perfectoid. Then the functor*

$$\text{BT}_{\infty}^{\mathcal{G}, \mu}(R) \rightarrow \text{Wind}_{\underline{\Delta}_R, \infty}^{\mathcal{G}, \mu}(R)$$

is an equivalence.

perfectoid

Remark 3.2.19 (Quotient description for perfectoid rings). Let $\xi \in \underline{\Delta}_R$ be a generator for $\text{Fil}_{\mathcal{N}}^1 \underline{\Delta}_R$. Another way of phrasing the proposition, using Remark 3.2.12, is that the restriction of the stack $\text{BT}_{\infty}^{\mathcal{G}, \mu}$ to perfectoid R -algebras is the quotient $[L^+ \mathcal{G}/H_{\mu}]$ (computed in the p -completely étale topology) where H_{μ} acts on $L^+ \mathcal{G}$ via the right action

$$g \cdot h = \tau(h)^{-1} g \varphi(\text{int}(\mu(\xi))(h)).$$

Here, $\tau: H_{\mu} \rightarrow L^+ \mathcal{G}$ is the tautological map.

_truncated

Remark 3.2.20 (Quotient description for truncated apertures). In fact a similar description holds for the n -truncated versions as well, which can be deduced by combining Remarks 5.5.5 and 5.5.6 and Lemma 9.2.3 of [26]. One now looks at the group

$$H_{\mu}^{(n)}(R) = \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R) \times_{\mathcal{G}(R/p^n R)} \mathcal{P}_{\mu}^{-}(R/p^n R).$$

This admits two maps

$$\tau, \sigma: H_{\mu}^{(n)}(R) \rightarrow \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R),$$

where τ is simply the projection onto the first coordinate, and σ is another map, which we will only describe here for R p -torsion free. In this case, we have

_free_desc

$$(3.2.20.1) \quad H_{\mu}^{(n)}(R) = \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R) \times_{\mathcal{G}(R/p^n R)} \mathcal{P}_{\mu}^{-}(R/p^n R) = \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R) \cap \mu(\xi)^{-1} \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R) \mu(\xi) \subset \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R).$$

and σ is the map $h \mapsto \varphi(\mu(\xi)h\mu(\xi)^{-1})$.

The restriction of $\text{BT}_{\infty}^{\mathcal{G}, \mu}$ to perfectoid $\hat{\mathcal{O}}$ -algebras is now given by the étale sheafification of $R \mapsto \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R)/H_{\mu}^{(n)}(R)$, where the action is given by

$$H_{\mu}^{(n)}(R) \times \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R) \xrightarrow{(h, g) \mapsto \tau(h)^{-1} g \sigma(h)} \mathcal{G}(\underline{\Delta}_R/p^n \underline{\Delta}_R).$$

3.3. Isocrystals with G -structure. Here, we recall the relationship between apertures and F -isocrystals with G -structure.

Construction 3.3.1 (Apertures to isocrystals). Let R be a perfect $\mathbb{F}_p$ -algebra. Then by Remark 3.2.19 we have

$$\mathrm{BT}_{\infty}^{\mathcal{G},\mu}(R) = [L^+\mathcal{G}/H_{\mu}](R),$$

where H_{μ} is the étale sheaf on perfect $\mathbb{F}_p$ -algebras given by $H_{\mu} = L^+\mathcal{G} \times_{\mathcal{G}} \mathcal{P}_{\mu}^-$. The action is via

$$\begin{aligned} L^+\mathcal{G} \times H_{\mu} &\rightarrow L^+\mathcal{G} \\ (g, h) &\mapsto h^{-1}g\varphi(\mu(p)h\mu(p)^{-1}). \end{aligned}$$

Set $G = \mathcal{G}_{\mathbb{Q}_p}$ and let $LG: R \mapsto G(W(R)[1/p])$ be the Witt vector loop group associated with G . Then we also have the étale quotient

$$\mathrm{Isoc}_G \stackrel{\mathrm{defn}}{=} [LG/\mathrm{Ad}_{\varphi} LG],$$

associated with the φ -twisted adjoint action

$$\begin{aligned} \mathrm{Ad}_{\varphi}: LG \times LG &\rightarrow LG \\ (g, h) &\mapsto h^{-1}g\varphi(h). \end{aligned}$$

The map $L^+\mathcal{G} \xrightarrow{g \mapsto g\varphi(\mu(p))} LG$ now descends to a functorial map

$$\mathrm{BT}_{\infty}^{\mathcal{G},\mu}(R) \rightarrow \mathrm{Isoc}_G(R)$$

for perfect $\mathbb{F}_p$ -algebras R .

Remark 3.3.2. When $R = \kappa$ is an algebraically closed field, then the quotient description of $\mathrm{BT}_{\infty}^{\mathcal{G},\mu}(\kappa)$ appearing above shows that we have

$$\mathrm{BT}_{\infty}^{\mathcal{G},\mu}(\kappa) = [\mathcal{G}(W(\kappa))/H_{\mu}(\kappa)] \xrightarrow[\sim]{g \mapsto g\varphi(\mu(p))} [\mathcal{G}(W(\kappa))\varphi(\mu(p))\mathcal{G}(W(\kappa))/\mathrm{Ad}_{\varphi}\mathcal{G}(W(\kappa))].$$

The set of isomorphism classes on the right hand side is the set $C(\mathcal{G}, \{\varphi(\mu)\})$ from [93, §8.2], which is in fact independent of the choice of the algebraically closed field κ . The set of isomorphism classes in $\mathrm{Isoc}_G(\kappa)$ can be identified with Kottwitz's set $B(G)$, which is also independent of the choice of κ .¹ Within $B(G)$, we have the μ -admissible subset $B(G, \{\mu\})$, which depends only on the Galois orbit of the conjugacy class $\{\mu\}$. By [79, Theorem 4.2], the natural map

$$C(\mathcal{G}, \{\varphi(\mu)\}) \rightarrow B(G)$$

induced by inclusion $G(W(\kappa)) \subset G(W(\kappa)[1/p])$ lands inside $B(G, \{\mu\}) = B(G, \{\varphi(\mu)\})$

Lemma 3.3.3. For an algebraically closed field κ , the map $\mathrm{BT}_{\infty}^{\mathcal{G},\mu}(\kappa) \rightarrow B(G)$ induced from Construction 3.3.1 has image $B(G, \{\mu\}) = B(G, \{\varphi(\mu)\})$.

Proof. Given the previous remark, we only need to know that every $[b] \in B(G, \{\varphi(\mu)\})$ is in the image of $C(\mathcal{G}, \{\varphi(\mu)\})$. For this, we need a result of Wintenberger [95, Corollaire 3] (see also the more general non-emptiness result of Gashi [27, Theorem 5.2]): This tells us that there exists $g \in G(W(\kappa)[1/p])$ and $h_1, h_2 \in G(W(\kappa))$ such that $g^{-1}b\varphi(g) = h_1\varphi(\mu(p))h_2$. $\square$

3.4. The de Rham realization.

Construction 3.4.1 (The Hodge-filtered de Rham realization). Suppose that we have $\mathfrak{Q} \in \mathrm{BT}_{\infty}^{\mathcal{G},\mu}(R)$. Then pulling back along the filtered de Rham point $x_{\mathrm{dR}}^{\mathcal{N}}$ from Remark 2.1.4 gives us a $\mathcal{G}$ -bundle $\mathrm{Fil}_{\mathrm{Hdg}}^{\bullet} T_{\mathrm{dR}}(\mathfrak{Q})$ over $\mathbb{A}^1/\mathbb{G}_m \times \mathrm{Spf} R$.

Remark 3.4.2. It follows from the definition of $\mathrm{BT}_{\infty}^{\mathcal{G},\mu}(R)$ and [26, Remark 4.9.7] that $\mathrm{Fil}_{\mathrm{Hdg}}^{\bullet} T_{\mathrm{dR}}(\mathfrak{Q})$ is étale locally on $\mathrm{Spf} R$ isomorphic to the pullback to $\mathbb{A}^1/\mathbb{G}_m \times \mathrm{Spf} R$ of the $\mathcal{G}$ -torsor $\mathcal{Q}_{\mu}$ over $B\mathbb{G}_m \times \mathrm{Spf} \hat{\mathcal{O}}$ from Remark 3.1.2. Moreover, *loc. cit.* also shows that giving such a $\mathcal{G}$ -bundle over $\mathbb{A}^1/\mathbb{G}_m \times \mathrm{Spf} R$ is equivalent to giving a $\mathcal{P}_{\mu}^-$ -torsor over R . That is, we have constructed a canonical map $\mathrm{BT}_{\infty}^{\mathcal{G},\mu} \rightarrow B\mathcal{P}_{\mu}^-$. This is precisely the one showing up in the deformation theory explained in Theorem 3.1.5.

[1] Keerthi: this.

Remark 3.4.3 (A period map and versality). Here is a reformulation of the deformation theory in Theorem 3.1.5 that will be useful. Let Gr_μ be the Grassmannian scheme over $\hat{\mathcal{O}}$ associated with μ : It depends only on the conjugacy class of μ and can be presented as the fppf quotient $\mathcal{G}_{\hat{\mathcal{O}}}/\mathcal{P}_\mu^-$ for our choice of representative μ . We can view the classifying stack $B\mathcal{P}_\mu^-$ from this perspective as the fppf quotient $\mathrm{Gr}_\mu/\mathcal{G}_{\hat{\mathcal{O}}}$. Now, $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet T_{\mathrm{dR}}(\mathfrak{Q})$ can be viewed as arising from a canonical map $\mathrm{BT}_\infty^{\mathcal{G},\mu} \rightarrow \mathrm{Gr}_\mu/\mathcal{G}_{\hat{\mathcal{O}}}$. Then Grothendieck–Messing theory tells us that the resulting map on cotangent complexes of p -adic formal stacks

$$\mathbb{L}_{\mathrm{Gr}_\mu/\mathcal{G}_{\hat{\mathcal{O}}}}^\wedge|_{\mathrm{BT}_\infty^{\mathcal{G},\mu}} \rightarrow \mathbb{L}_{\mathrm{BT}_\infty^{\mathcal{G},\mu}}^\wedge$$

factors through an isomorphism

$$\mathbb{L}_{(\mathrm{Gr}_\mu/\mathcal{G}_{\hat{\mathcal{O}}})/B\mathcal{G}_{\hat{\mathcal{O}}}}^\wedge|_{\mathrm{BT}_\infty^{\mathcal{G},\mu}} \xrightarrow{\sim} \mathbb{L}_{\mathrm{BT}_\infty^{\mathcal{G},\mu}}^\wedge.$$

In particular, note that we have a canonical isomorphism⁶

3.5. Realization in F -zips with additional structure. We now quickly review the relationship between apertures and F -zips as in [78].

Notation 3.5.1. Let $\mathcal{P}_{\varphi(\mu)}^+ \subset \mathcal{G}$ be the smooth subgroup scheme whose Lie algebra is identified with the sum of the weight- i spaces in $\mathrm{Lie}(\mathcal{G})_{\hat{\mathcal{O}}}$ for $\varphi(\mu)$ with $i \geq 0$. Let $\mathcal{U}_\mu^-$ (resp. $\mathcal{U}_{\varphi(\mu)}^+$) be the smooth subgroup scheme of $\mathcal{P}_\mu^-$ (resp. $\mathcal{P}_{\varphi(\mu)}^+$) whose Lie algebra is the sum of the weight- i spaces for μ (resp. for $\varphi(\mu)$) with $i < 0$ (resp. $i > 0$). Set $\mathcal{M}_\mu = \mathcal{P}_\mu^-/\mathcal{U}_\mu^-$ and $\mathcal{M}_{\varphi(\mu)} = \mathcal{P}_{\varphi(\mu)}^+/\mathcal{U}_{\varphi(\mu)}^+$. We have a canonical isomorphism of group schemes $\varphi^*\mathcal{M}_\mu \xrightarrow{\sim} \mathcal{M}_{\varphi(\mu)}$.

Definition 3.5.2 ($\mathcal{G}$ -zips). Let R be a k -algebra where k is the residue field of $\hat{\mathcal{O}}$. An **F -zip with $\mathcal{G}$ -structure** or **$\mathcal{G}$ -zip** of type μ over R (see [78, Definition 1.4]) is a tuple

$$(\mathcal{Q}, \mathcal{Q}^-, \mathcal{Q}^+, \iota)$$

where $\mathcal{Q}$ is a $\mathcal{G}$ -torsor over R , $\mathcal{Q}^- \subset \mathcal{Q}$ is a reduction of structure to a $\mathcal{P}_\mu^-$ -torsor, $\mathcal{Q}^+ \subset \mathcal{Q}$ is a reduction of structure to a $\mathcal{P}_{\varphi(\mu)}^+$ -torsor and

$$\alpha: \varphi^*(\mathcal{Q}^-/\mathcal{U}_\mu^-) \xrightarrow{\sim} \mathcal{Q}^+/\mathcal{U}_{\varphi(\mu)}^+$$

is an isomorphism of $\mathcal{M}_{\varphi(\mu)}$ -torsors. We denote the groupoid of such objects by $\mathcal{G}\text{-zip}_\mu(R)$.

Remark 3.5.3. By [78, Theorem 1.5], $R \mapsto \mathcal{G}\text{-zip}_\mu(R)$ is a smooth 0-dimensional Artin stack over k .

Remark 3.5.4 (Stratification of $\mathcal{G}\text{-zip}_\mu$). Suppose that $\mathcal{G}$ is reductive. Then by [78, Theorem 3.20], the stack $\mathcal{G}\text{-zip}_\mu$ has an underlying finite T_0 -topological space corresponding to a certain subset ${}^JW \subset W$ of the Weyl group of $\mathcal{G}$ endowed with a refinement of the Bruhat ordering; see the explanation in [96, §5.3].¹³ This yields a stratification of $\mathcal{G}\text{-zip}_\mu$ —explained in §3.6 of *loc. cit.*—where each stratum is a smooth, locally closed substack.

The next result follows from the proof of [26, Theorem 9.3.2].

Proposition 3.5.5 (Apertures to $\mathcal{G}$ -zips). *There is a natural smooth surjective map*

$$\mathrm{BT}_1^{\mathcal{G},\mu} \otimes_{\hat{\mathcal{O}}} k \rightarrow \mathcal{G}\text{-zip}_\mu$$

of 0-dimensional Artin stacks over k that is a gerbe for a connected p -torsion finite flat commutative group scheme. Moreover, the resulting $\mathcal{P}_\mu^-$ -torsor over $\mathrm{BT}_1^{\mathcal{G},\mu} \otimes_{\hat{\mathcal{O}}} k$ agrees with the one described in Remark 3.4.2.

Remark 3.5.6 (Set-theoretic version). A particular consequence of the proposition is that, for any algebraically closed field κ over k , the map

$$\mathrm{BT}_1^{\mathcal{G},\mu}(\kappa) \rightarrow \mathcal{G}\text{-zip}_\mu(\kappa)$$

is an isomorphism of groupoids.

If $\mathcal{G}$ is reductive, then, for any algebraically closed field κ over k , looking at isomorphism classes of κ -points for the above map gives a map $\tilde{\zeta}: C(\mathcal{G}, \{\varphi(\mu)\}) \rightarrow {}^JW$, which is the one described explicitly in [96, Proposition 6.7].

¹³Note that what Wortmann denotes by μ corresponds to our $\varphi(\mu)$ here.

[6] **Alex:** Kee was an empty here—did you put something

nary_locus

3.6. The μ -ordinary locus. In this subsection, we will suppose that $\mathcal{G}$ is reductive. We'll now look at the μ -ordinary locus of $\mathrm{BT}_n^{\mathcal{G},\mu}$. Our treatment will be somewhat terse; more details will appear in [65].

rd_stratum

Definition 3.6.1 (The μ -ordinary stratum). Recall the map $\tilde{\zeta}$ from Remark 3.5.6. The element $\tilde{\zeta}(\varphi(\mu(p))) \in {}^JW$ corresponds to the unique open stratum of $\mathcal{G}\text{-zip}_\mu$; see for instance the proof of [96, Theorem 6.10]. Write $\mathcal{G}\text{-zip}_\mu^{\mathrm{ord}}$ for this open stratum. For $n \in \mathbb{N} \cup \{\infty\}$, we will write $\mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{ord}} \subset \mathrm{BT}_n^{\mathcal{G},\mu}$ for the unique open formal substack characterized by the property that its mod- p fiber is

$$\mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{ord}} \otimes k = (\mathrm{BT}_n^{\mathcal{G},\mu} \otimes k) \times_{\mathcal{G}\text{-zip}_\mu} \mathcal{G}\text{-zip}_\mu^{\mathrm{ord}}.$$

Setup 3.6.2. Choose a maximal torus $\mathcal{T} \subset \mathcal{G}$ contained in a Borel subgroup of $\mathcal{G}$. We can assume that μ has been chosen within its conjugacy class to factor through $\mathcal{T}_\mathcal{O}$.¹⁴ The sum of the Galois conjugates of μ now gives a cocharacter ν of $\mathcal{T}$ that is defined over $\mathbb{Z}_p$. Let $\mathcal{P}_\nu^+ \subset \mathcal{G}$ be the parabolic subgroup whose Lie algebra is the sum of the non-negative eigenspaces for ν , and let $\mathcal{U}_\nu^+$ be its unipotent radical.

nu_central

Lemma 3.6.3. *Let $\mathcal{M}_\nu \subset \mathcal{G}$ and $\mathcal{M}_\mu \subset \mathcal{G}_\mathcal{O}$ be the centralizers of ν and μ , respectively. Then we have $\mathcal{M}_{\nu,\mathcal{O}} \subset \mathcal{M}_\mu$.*

Proof. It is enough to know that μ is central in $\mathcal{M}_{\nu,\mathcal{O}}$. By [84, Lemma 2.7], this would follow if the average $\bar{\mu}$ of the Galois conjugates of μ is central in $\mathcal{M}_{\nu,\mathcal{O}}$. But this is clear, since ν is an integer multiple of $\bar{\mu}$. $\square$

TPnu_BTmu

Remark 3.6.4. We can view μ as a minuscule (in fact, central) cocharacter of $\mathcal{M}_\nu$, and as a 1-bounded cocharacter of $\mathcal{P}_\mu^-$. This gives us formal algebraic stacks $\mathrm{BT}_n^{\mathcal{M}_\nu,\mu}$ and $\mathrm{BT}_n^{\mathcal{P}_\nu^+,\mu}$ over $\hat{\mathcal{O}}$. Viewing $\mathcal{M}_\nu$ as the Levi quotient of $\mathcal{P}_\nu^-$ gives us maps

$$\mathrm{BT}_n^{\mathcal{M}_\nu,\mu} \rightarrow \mathrm{BT}_n^{\mathcal{P}_\nu^+,\mu} \rightarrow \mathrm{BT}_n^{\mathcal{M}_\nu,\mu}$$

whose composition is the identity.

_structure

Proposition 3.6.5. *Suppose that $n \in \mathbb{N}$. The formal stack $\mathrm{BT}_n^{\mathcal{M}_\nu,\mu}$ is étale over $\hat{\mathcal{O}}$ and is non-canonically isomorphic to the formal classifying stack for the finite group $\mathcal{M}_\nu(\mathbb{Z}/p^n\mathbb{Z})$. Moreover, there is a canonical (non-commutative) finite flat group scheme $\mathcal{F}_\nu(n)$ over $\mathrm{BT}_n^{\mathcal{M}_\nu,\mu}$ such that:*

- (1) $\mathcal{F}_\nu(n)$ admits a canonical filtration by finite flat subgroup schemes over $\mathrm{BT}_n^{\mathcal{M}_\nu,\mu}$, whose associated graded pieces $\mathcal{F}_{\nu,i}(n)$ are n -truncated Barsotti-Tate group schemes (and are in particular commutative);
- (2) For $n > 1$ and all i , there is a canonical isomorphism

$$\mathcal{F}_{\nu,i}(n-1) \simeq \mathcal{F}_{\nu,i}(n)[p^{n-1}]$$

of $(n-1)$ -truncated Barsotti-Tate group schemes;

- (3) The map $\mathrm{BT}_n^{\mathcal{P}_\nu^+,\mu} \rightarrow \mathrm{BT}_n^{\mathcal{M}_\nu,\mu}$ is canonically isomorphic to the classifying stack of $\mathcal{F}_\nu(n)$.

Proof. The first assertion follows from [26, Proposition 10.3.4].

For the rest, suppose that we have a finite étale $\hat{\mathcal{O}}$ -algebra R , and $\mathfrak{Q} \in \mathrm{BT}_n^{\mathcal{M}_\nu,\mu}(R)$ corresponding to a $\mathcal{G}$ -torsor over $R^{\mathrm{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}$. We can twist $\mathcal{U}_\nu^+$ by this torsor to get a unipotent (relative) group scheme $(\mathcal{U}_\mu^+)_{\mathfrak{Q}}$ over $R^{\mathrm{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}$. Unwinding definitions shows that the fiber of $\mathrm{BT}_n^{\mathcal{P}_\nu^+,\mu}$ over $\mathfrak{Q}$ is the functor on p -nilpotent R -algebras C given by

u_explicit

$$(3.6.5.1) \quad C \mapsto \mathrm{Map}_{/R^{\mathrm{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}}(C^{\mathrm{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}, B(\mathcal{U}_\nu^+)_{\mathfrak{Q}}).$$

Now, let $\mathcal{U}_{\mu,\bullet}^+$ be the ascending central series filtration on $\mathcal{U}_\mu^+$: This is stable under the action of $\mathcal{M}_\mu$, and so this twists to a filtration on $(\mathcal{U}_\mu^+)_{\mathfrak{Q}}$, whose associated graded are twists of the *commutative* unipotent group schemes $\mathcal{U}_{\mu,i}^+/\mathcal{U}_{\mu,i-1}^+$. Each such twist is therefore a vector group scheme $\mathbf{V}(\mathcal{M}_i)$ associated with a vector bundle $\mathcal{M}_i$ over

¹⁴By [54, Lemma 1.1.3], μ can be conjugated to a cocharacter factoring through $\mathcal{T}_{\mathcal{O}[1/p]}$ dominant with respect to the choice of Borel. This will necessarily be a cocharacter of $\mathcal{T}_\mathcal{O}$.

$R^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}$ —in other words, with a p^n -torsion F -gauge over R . The minuscule-ness of μ implies that $\mathcal{M}_i$ has Hodge-Tate weights in $\{0, 1\}$. By [26, Theorem 11.3.3] and [67, Proposition 4.3.2], the functors

$$C \mapsto \tau^{\leq 0} R\Gamma(C^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}, \mathcal{M}_i);$$

$$C \mapsto \tau^{\leq 0} R\Gamma(C^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}, \mathcal{M}_i[1]) \simeq \text{Map}_{/R^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}}(C^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}, B\mathbf{V}(\mathcal{M}_i))$$

on p -nilpotent R -algebras are represented by an n -truncated Barsotti-Tate group scheme $\mathcal{F}_{\nu, i}(n)_R$ over R and its classifying stack, respectively. This construction is compatible with change in n in the obvious way.

One now sees that the assignment

$$C \mapsto \text{Map}_{/R^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}}(C^{\text{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z}, (\mathcal{U}_{\nu}^+)_{\Omega})$$

is represented by a finite flat group scheme $\mathcal{F}_{\nu}(n)_R$ over R , and that the functor (3.6.5.1) is the classifying stack of the finite flat group scheme. By varying R and Ω , one now gets the desired finite flat group scheme over $\text{BT}_n^{\mathcal{M}_{\nu}, \mu}$. $\square$

local_ring

Corollary 3.6.6. *Suppose that we have $\mathfrak{P}_0 \in \text{BT}_{\infty}^{\mathcal{P}_{\nu}^+, \mu}(\kappa)$. Then the universal deformation space $\widehat{U}_{\mathfrak{P}_0}$ of $\text{BT}_{\infty}^{\mathcal{P}_{\nu}^+, \mu}$ at $\mathfrak{P}_0$ admits a canonical structure of a tower*

$$\widehat{U}_{\mathfrak{P}_0} = \widehat{U}_n \rightarrow \widehat{U}_{n-1} \rightarrow \cdots \rightarrow \widehat{U}_0 = \text{Spf } W(\kappa)$$

of formal schemes over $W(\kappa)$, where, for each $i > 0$, $\widehat{U}_i$ is a formal p -divisible group over $\widehat{U}_{i-1}$;

Proof. Given Proposition 3.6.5, this is implied by Lemma 3.6.7 below. $\square$

lete_local

Lemma 3.6.7. *Let R be a complete local ring with residue field κ , and let Art_R be the category of Artin local R -algebras with residue field κ . Let $\mathcal{H}$ be a p -divisible group over R , and let $0 \in (B\mathcal{H})(\kappa) = \varprojlim_n (B\mathcal{H}[p^n])(\kappa)$ be the trivial torsor for $\mathcal{H}$. Then the deformation functor*

$$\text{Def}_{B\mathcal{H}, 0} : C \mapsto \text{fib}_0((B\mathcal{H})(C) \rightarrow (B\mathcal{H})(\kappa))$$

on Art_R is pro-represented by the p -divisible formal group $\widehat{\mathcal{H}}$.

Proof. This is classical and can be deduced from [42]. $\square$

The connection of the above discussion with the ordinary locus is made by:

escription

Proposition 3.6.8. *For $n \in \mathbb{N}$, the natural map $\text{BT}_n^{\mathcal{P}_{\nu}^+, \mu} \rightarrow \text{BT}_n^{\mathcal{G}, \mu}$ is étale and its image is precisely the μ -ordinary locus $\text{BT}_n^{\mathcal{G}, \mu, \text{ord}}$.*

Proof. That the natural map is étale is shown in [26, Proposition 9.5.1]. For the description of the image, it is enough to actually check the case where $n = \infty$. Here, Proposition 3.6.5 shows that we have

$$\text{BT}_{\infty}^{\mathcal{P}_{\nu}^+, \mu}(\kappa) \xrightarrow{\sim} \text{BT}_{\infty}^{\mathcal{M}_{\nu}, \mu}(\kappa) \simeq [\mathcal{M}_{\nu}(W(\kappa))\varphi(\mu(p))/_{\text{Ad}_{\varphi}} \mathcal{M}_{\nu}(W(\kappa))].$$

Lang's lemma now tells us that the right hand side has a single isomorphism class represented by $\varphi(\mu(p))$, and so we conclude now using Lemma 3.6.12 below. $\square$

nary_locus

Remark 3.6.9. Combining Proposition 3.6.8 with Corollary 3.6.6 tells us that, for any $\Omega_0 \in \text{BT}_{\infty}^{\mathcal{G}, \mu, \text{ord}}(W(\kappa))$, the deformation space Def_{Ω_0} over $W(\kappa)$ admits a canonical section obtained as the successive composition of identity sections of a tower of relative formal p -divisible groups.

lift_local

Definition 3.6.10 (Canonical lift for μ -ordinary apertures). The lift of Ω_0 to $\text{BT}_{\infty}^{\mathcal{G}, \mu}(W(\kappa))$ corresponding to the canonical section of the deformation space Def_{Ω_0} is called the **canonical lift** of Ω_0 and is denoted Ω_0^{can} .

Notation 3.6.11. Let κ be an algebraically closed field over k . Following [96], given $g \in \mathcal{G}(W(\kappa))\varphi(\mu(p))\mathcal{G}(W(\kappa))$, we will write $\langle g \rangle$ for the associated class in $C(\mathcal{G}, \{\varphi(\mu)\})$, $[g]$ for the class in $B(\mathcal{G}, \{\mu\})$, and $\llbracket g \rrbracket \subset C(\mathcal{G}, \{\varphi(\mu)\})$ for the subset $\tilde{\zeta}^{-1}(\zeta(\langle g \rangle))$. Here, $\tilde{\zeta}$ is the map from Remark 3.5.6.

witz_lemma

Lemma 3.6.12. *With the above notation, we have equivalences*

$$[g] = [\varphi(\mu(p))] \Leftrightarrow \langle g \rangle = \langle \varphi(\mu(p)) \rangle \Leftrightarrow \llbracket g \rrbracket = \llbracket \varphi(\mu(p)) \rrbracket.$$

Proof. See [96, Proposition 7.2]. $\square$

isocrystals

3.7. Filtered F -crystal realization. Here, we review some material on the crystalline realizations of apertures (cf. [38], which recovers much of the material via Tannakian considerations.).

semiperfect

Remark 3.7.1 (Crystalline realization over semiperfectoid rings). For any semiperfect (animated commutative) $\mathbb{F}_p$ -algebra R , we have a canonical comparison isomorphism $\Delta_R \xrightarrow{\sim} A_{\text{crys}}(R)$ (see for instance [26, Remark 6.9.5]). Hence a $\mathcal{G}$ -bundle over $R^\Delta = \text{Spf } \Delta_R$ is the same as one over $A_{\text{crys}}(R)$. Suppose now that we have $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R)$: Then just as in [5, Remark 6.3.4], the $\mathcal{G}$ -bundle $T_{\text{crys}}(\mathfrak{Q})$ in crystals associated with $j_{\Delta}^* \mathfrak{Q}$ is equipped with an isomorphism $\varphi^* T_{\text{crys}}(\mathfrak{Q})[1/p] \xrightarrow{\sim} T_{\text{crys}}(\mathfrak{Q})[1/p]$ of $\mathcal{G}$ -bundles over $A_{\text{crys}}(R)$.

realization

Construction 3.7.2 (Crystalline realization). By quasisyntomic descent, Remark 3.7.1 tells us that for any $\mathbb{F}_p$ -algebra R , every $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R)$ yields a $\mathcal{G}$ -bundle in F -crystals over R , which we denote by $T_{\text{crys}}(\mathfrak{Q})$. See [38, §1.1.1] for an explanation in the context of vector bundle F -gauges.

Remark 3.7.3 (F -isocrystals over perfect rings). Suppose that R is a perfect $\mathbb{F}_p$ -algebra. Then, $\text{Isoc}_G(R)$ is the groupoid of pairs $(\mathcal{E}, \varphi_{\mathcal{E}})$, where $\mathcal{E}$ is a G -torsor over $W(R)[1/p]$ and $\varphi_{\mathcal{E}}: \varphi^* \mathcal{E} \xrightarrow{\sim} \mathcal{E}$ is an isomorphism of G -bundles. The functor $\text{BT}_{\infty}^{\mathcal{G},\mu}(R) \rightarrow \text{Isoc}_G(R)$ from Construction 3.3.1 can now be understood as follows. Given $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R)$, one considers the G -bundle in F -isocrystals underlying the $\mathcal{G}$ -bundle in F -crystals $T_{\text{crys}}(\mathfrak{Q})$: This corresponds exactly to a G -bundle over $W(R)[1/p]$ equipped with an isomorphism from its Frobenius twist, and hence an object of $\text{Isoc}_G(R)$.

nd_de_rham

Remark 3.7.4 (Crystals and the de Rham realization). Suppose that R is a p -complete $\hat{\mathcal{O}}$ -algebra. Then every crystal in $\mathcal{G}$ -bundles over R/pR yields a $\mathcal{G}$ -bundle over R ; here, we are using the fact that $R \rightarrow R/pR$ is a pro-divided power thickening. In particular, if we have $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R)$ reducing to $\mathfrak{Q}_0 \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R/pR)$, then we obtain a $\mathcal{G}$ -bundle over R associated with $T_{\text{crys}}(\mathfrak{Q}_0)$. This $\mathcal{G}$ -bundle is in fact canonically isomorphic to $T_{\text{dR}}(\mathfrak{Q})$; see [38, Theorem 1.19] and its proof.

nius_lifts

Remark 3.7.5 (Frobenius lifts and filtrations). Suppose that R is a flat $\hat{\mathcal{O}}$ -algebra equipped with a Frobenius lift $\varphi: R \rightarrow R$. As in Example 3.2.4, this gives us a Breuil–Kisin frame $\underline{R}$ for R , and we obtain a functor $\Phi_{\underline{R}}: \text{BT}_{\infty}^{\mathcal{G},\mu}(R/pR) \rightarrow \text{Wind}_{\underline{R},\infty}^{\mathcal{G},\mu}(R/pR)$. Suppose that we have $\mathfrak{Q}_0 \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R/pR)$. One finds that in the notation of Remark 3.2.12, $\Phi_{\underline{R}}(\mathfrak{Q}_0)$ corresponds to the tuple

$$(T_{\text{crys}}(\mathfrak{Q}_0), \text{Fil}_{\text{Hdg}}^{\bullet} T_{\text{dR}}(\mathfrak{Q}_0), \alpha)$$

with α an isomorphism

$$\varphi^* T_{\text{crys}}(\mathfrak{Q}_0)' \xrightarrow{\sim} T_{\text{crys}}(\mathfrak{Q}_0),$$

where $T_{\text{crys}}(\mathfrak{Q}_0)'$ is the modification along $\text{Fil}_{\text{Hdg}}^{\bullet} T_{\text{dR}}(\mathfrak{Q}_0)$ from Remark 3.2.13.

visibility

Remark 3.7.6 (limpid divisibility). In the situation of Remark 3.7.5, suppose that $\mathfrak{Q}_0$ lifts to $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R)$. Then we have a lift $\text{Fil}_{\text{Hdg}}^{\bullet} T_{\text{dR}}(\mathfrak{Q})$ of the filtered bundle $\text{Fil}_{\text{Hdg}}^{\bullet} T_{\text{dR}}(\mathfrak{Q}_0)$. For $\Lambda \in \text{Rep}_{\mathbb{Z}_p}(\mathcal{G})$, let $\text{Fil}_{\text{Hdg}}^{\bullet} V$ be the filtered finite locally free R -module obtained by twisting Λ by $\text{Fil}_{\text{Hdg}}^{\bullet} T_{\text{dR}}(\mathfrak{Q})$. Then Remark 3.2.13 tells us that we have a functorial-in- Λ isomorphism

$$\varphi^* \left(\sum_i p^{-i} \text{Fil}_{\text{Hdg}}^i V \right) \xrightarrow{\sim} V$$

of finite locally free R -modules.

F-crystal

Construction 3.7.7 (Filtered F -crystals). Let κ be a perfect field, and suppose that $\text{Spf } R$ is a base formal $W(\kappa)$ -scheme. Suppose that we have $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R)$ lifting $\mathfrak{Q}_0 \in \text{BT}_{\infty}^{\mathcal{G},\mu}(R/pR)$. Then $T_{\text{dR}}(\mathfrak{Q})$ is equipped with a topologically nilpotent integrable connection ∇ , which completely determines the $\mathcal{G}$ -bundle in crystals $T_{\text{crys}}(\mathfrak{Q}_0)$. Now, if R is equipped with a Frobenius lift $\varphi: R \rightarrow R$, then the F -crystal structure on $T_{\text{crys}}(\mathfrak{Q}_0)$ gives us an isomorphism

$$\varphi^* T_{\text{dR}}(\mathfrak{Q})[1/p] \xrightarrow{\sim} T_{\text{dR}}(\mathfrak{Q})[1/p]$$

of $\mathcal{G}$ -bundles over $R[1/p]$ that is parallel for ∇ .

For any representation Λ , this gives us an isomorphism $\varphi^*V[1/p] \xrightarrow{\sim} V[1/p]$, which restricts to the integral isomorphism explained in Remark 3.7.6. Moreover, the filtration $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet T_{\mathrm{dR}}(\Omega)$ satisfies Griffiths transversality (in the sense that the associated filtration on the twist of any representation of $\mathcal{G}$ by $T_{\mathrm{dR}}(\Omega)$ satisfies Griffiths transversality); see [38, Theorem 2.10]. In sum, in the language of [38, §2.1.2], for any base formal $W(\kappa)$ -scheme $\mathfrak{X}$, and $\Omega \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu}(\mathfrak{X})$, we can associate with Ω an exact and monoidal functor from $\mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G})$ to the category of *strongly divisible filtered F -crystals* over $\mathfrak{X}$.

realization

3.8. The étale realization and p -adic comparison. Here, we recall the construction of the étale realization of apertures and some p -adic comparison results for it that make precise its relationship with the objects defined in the previous subsection.

_apertures

Construction 3.8.1 (The étale realization for apertures). Let R be a derived p -complete animated commutative ring. To an object Ω of $B\mathcal{G}(R^{\mathrm{syn}} \otimes \mathbb{Z}/p^n\mathbb{Z})$, we will attach an object $T_{\mathrm{ét}}(\Omega)$ that specializes to the construction given in Corollary 2.3.12 when R is p -quasisyntomic and p -torsion free.

For every object $(A, I, R \rightarrow \bar{A})$ in R_{Δ} , Remark 2.1.1 tells us that $j_{\Delta}^*\Omega \in B\mathcal{G}(R^{\Delta} \otimes \mathbb{Z}/p^n\mathbb{Z})$ yields a $\mathcal{G}$ -torsor $\mathcal{P}_{\Omega}(A, I, R \rightarrow \bar{A})$ over $A/\mathbb{L}p^n$. Furthermore, just as in [5, Remark 6.3.4], the fact that we have the underlying syntomic torsor Ω implies that there is a natural isomorphism of $\mathcal{G}$ -torsors

$$\varphi_P: \varphi^*\mathcal{P}_{\Omega}(A, I, R \rightarrow \bar{A})[1/I] \xrightarrow{\sim} \mathcal{P}_{\Omega}(A, I, R \rightarrow \bar{A})[1/I]$$

over $(A/\mathbb{L}p^n)[1/I]$. Now, just as in Constructions 6.3.1 and 6.3.2 of *loc. cit.*, we obtain an object $T_{\mathrm{ét}}(\Omega)$ in $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R[1/p])$ characterized by the equality

$$\Gamma(\mathrm{Spec} T[1/p], T_{\mathrm{ét}}(\Omega)) = \mathcal{P}_{\Omega}(\Delta_T, I_T, R \rightarrow T)[1/I_T]^{\varphi_P = \mathrm{id}},$$

for p -torsion free perfectoid R -algebras T . Taking the limit over n gives us a realization functor

$$T_{\mathrm{ét}}: B\mathcal{G}(R^{\mathrm{syn}}) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(R[1/p]).$$

This construction globalizes to a general derived p -adic formal scheme $\mathfrak{X}$ in the obvious way.

socrystals

Remark 3.8.2 (Filtered F -isocrystals and crystalline comparison). Let κ be a perfect field over $\hat{\mathcal{O}}$ of characteristic p and let $\mathfrak{X}$ be a base formal $W(\kappa)$ -scheme. Suppose that we have $\Omega \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu}(\mathfrak{X})$. For any $\Lambda \in \mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G})$, we can now associate two filtered F -isocrystals¹⁵ over the special fiber $\mathfrak{X}_{\kappa}$:

- The filtered F -isocrystal associated with the filtered F -crystal $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet V$ from Construction 3.7.7;
- The filtered F -isocrystal associated with the crystalline $\mathbb{Z}_p$ -local system $(\Lambda)_{T_{\mathrm{ét}}(\Omega)}$.

By [38, Theorem 2.10], these two filtered F -isocrystals are canonically isomorphic.

comparison

Remark 3.8.3 (Filtered bundles and de Rham comparison). As usual, let $\mathcal{O}_K$ be a complete DVR over $\hat{\mathcal{O}}$ with perfect residue field, and let $\mathfrak{X}$ be a base formal $\mathcal{O}_K$ -scheme. Suppose that we are given $\Omega \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu}(\mathfrak{X})$ lifting $Q \in \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\mathrm{crys}}(\mathfrak{X}_{\eta})$. We can now associate two filtered G -bundles over $\mathfrak{X}_{\eta}$ equipped with integrable connections:

- The generic fiber of the Hodge-filtered de Rham realization $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet T_{\mathrm{dR}}(\Omega)$;
- Since Q is crystalline, and therefore in particular de Rham, we have the filtered G -bundle $\mathrm{Fil}^\bullet D_{\mathrm{dR}}(Q)$ associating with each $V \in \mathrm{Rep}_{\mathbb{Q}_p}(G)$, the filtered vector bundle $\mathrm{Fil}^\bullet D_{\mathrm{dR}}((V)_Q)$ from [58, Theorem 3.7(iv)].

comparison

Lemma 3.8.4. *The two filtered bundles above with integrable connections are canonically isomorphic.*

Proof. When $W(\kappa) = \mathcal{O}_K$, this is immediate from Remark 3.8.2. The proof of Imai–Kato–Younis cited there works also in this context, if one ignores F -structures. Indeed, everything after the first paragraph in *loc. cit.* works exactly the same, replacing $\mathbb{D}_{\mathrm{crys}}[1/p]$ and D_{crys} with $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet T_{\mathrm{dR}}(\Omega)[1/p]$ and $\mathrm{Fil}^\bullet D_{\mathrm{dR}}(Q)$, respectively. $\square$

¹⁵See [37, p. 28] for this notion

perfectoid

3.9. Étale realization over perfectoid rings. In this subsection, we will see that the étale realization on $\mathrm{BT}_n^{\mathcal{G}, \mu}$ is faithful for base formal $\mathcal{O}_K$ -schemes.

perfectoid

Remark 3.9.1 (Étale $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -torsors over perfectoid rings). Suppose that R is perfectoid and p -torsion free. By Remark 2.3.8, giving a $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -local system over $R[1/p]$ is equivalent to giving an exact symmetric monoidal functor

$$\mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \rightarrow \mathrm{Loc}_{\mathbb{Z}/p^n\mathbb{Z}}(R[1/p]).$$

Now, tilting combined with Katz's Riemann-Hilbert equivalence (see [8, Proposition 3.6]) tells us that the right-hand side is equivalent to the category of pairs $(\mathcal{V}, \eta)$, where $\mathcal{V}$ is a finite locally free module over $\Delta_R/p^n\Delta_R[1/\varepsilon]$ and $\eta: \varphi^*\mathcal{V} \xrightarrow{\sim} \mathcal{V}$ is an isomorphism of $\Delta_R/p^n\Delta_R[1/\varepsilon]$ -modules. Therefore, we conclude that giving a $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -local system over $R[1/p]$ is equivalent to giving a $\mathcal{G}$ -torsor $\mathcal{Q}$ over $(\Delta_R/p^n\Delta_R)[1/\varepsilon]$ along with an isomorphism $\varphi_{\mathcal{Q}}: \varphi^*\mathcal{Q} \xrightarrow{\sim} \mathcal{Q}$.

al_systems

Remark 3.9.2 (Quotient description of the image of the étale realization). In terms of the description of the target from Remark 3.9.1, the étale realization functor $\mathrm{BT}_n^{\mathcal{G}, \mu}(R) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R[1/p])$ factors through the fully faithful sub-groupoid spanned by pairs $(\mathcal{Q}, \varphi_{\mathcal{Q}})$ where $\mathcal{Q}$ can be trivialized over $(\Delta_{\tilde{R}}/p^n\Delta_{\tilde{R}})[1/\varepsilon]$ for some p -completely faithfully flat and étale map $R \rightarrow \tilde{R}$ of perfectoid rings.

Observe that this sub-groupoid is the evaluation at R of the étale sheafification of the functor on p -torsion free perfectoid rings given by

$$R \mapsto [\mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]) / \mathrm{Ad}_{\varphi}(\mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]))].$$

Here, the right-hand side is the quotient by the φ -semilinear adjoint action

$$\begin{aligned} \mathrm{Ad}_{\varphi}: \mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]) \times \mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]) &\rightarrow \mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]) \\ (g, h) &\mapsto h^{-1}g\varphi(h). \end{aligned}$$

From this perspective, the étale realization map from Construction 3.8.1 can be viewed as the étale sheafification of the map

$$[\mathcal{G}(\Delta_R/p^n\Delta_R)/H_{\mu}^{(n)}(R)] \rightarrow [\mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]) / \mathrm{Ad}_{\varphi}(\mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]))]$$

induced by

$$\mathcal{G}(\Delta_R/p^n\Delta_R) \xrightarrow{g \mapsto g\varphi(\mu(\xi))} \mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon])$$

and the map of groups

hmu_to_LG

$$(3.9.2.1) \quad H_{\mu}^{(n)}(R) = \mathcal{G}(\Delta_R/p^n\Delta_R) \times_{\mathcal{G}(R/p^nR)} \mathcal{P}_{\mu}^{-}(R/p^nR) \subset \mathcal{G}(\Delta/p^n\Delta_R) \rightarrow \mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]).$$

description

Remark 3.9.3. In the situation above, the group $H_{\mu}^{(n)}(R)$ can be described more simply: We have

$$H_{\mu}^{(n)}(R) = \mathcal{G}(\Delta_R/p^n\Delta_R) \cap \mu(\xi)\mathcal{G}(\Delta_R/p^n\Delta_R)\mu(\xi)^{-1} \leq \mathcal{G}(\Delta_R/p^n\Delta_R[1/\varepsilon]).$$

To see this, note that by [55, Remark 6.3.3] the map

$$\mathcal{P}_{\mu}^{-}(\Delta_R/p^n\Delta_R) \times (\Delta_R/p^n\Delta_R \otimes_{\mathcal{O}} \mathfrak{g}_1) \xrightarrow{(h, X) \mapsto h \exp(\xi X)} \mathcal{G}(\Delta_R/p^n\Delta_R)$$

is a bijection onto $H_{\mu}^{(n)}(R)$. Here $\mathfrak{g}_1 \subset \mathfrak{g}_{\mathcal{O}}$ is the weight-1 eigenspace for μ .

d_faithful

Proposition 3.9.4. Suppose that R is perfectoid and p -torsion free. Then the étale realization functor

$$\mathrm{BT}_n^{\mathcal{G}, \mu}(R) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R[1/p])$$

is faithful.

Proof. Given Remark 3.9.2 and the description of $H_{\mu}^{(n)}(R)$ in (3.2.20.1), this reduces to the easy assertion that the map (3.9.2.1) is injective. $\square$

h_faithful

Corollary 3.9.5. Suppose that $\mathfrak{X}$ is a base formal $\mathcal{O}_K$ -scheme. Then the étale realization functor

$$\mathrm{BT}_n^{\mathcal{G}, \mu}(\mathfrak{X}) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\mathfrak{X}_{\eta})$$

is faithful.

Proof. This is because $\mathfrak{X}$ admits a quasisyntomic cover $\{\mathrm{Spf}(R_i) \rightarrow \mathfrak{X}\}$ with R_i perfectoid and p -torsion free for all i ; see [37, Lemma 1.15]. $\square$

4. TATE'S FULL FAITHFULNESS FOR APERTURES AND ITS CONSEQUENCES

In this section, we prove a generalization Tate's full faithfulness theorem for p -divisible groups to the context of apertures. We also derive some corollaries to this that will be of importance for the main theorems of this paper.

4.1. Algebraization and Tate's full faithfulness. Let $\mathcal{O}$ be a mixed characteristic $(0, p)$ discrete valuation ring with completion $\hat{\mathcal{O}}$. We will now define an algebraic variant of $\mathrm{BT}_n^{\mathcal{G}, \mu}$ over $\mathcal{O}$ by gluing it to the stack of $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -local systems over E , and we will use it to prove the aperture analogue of Tate's full faithfulness theorem for p -divisible groups by reducing it ultimately to Corollary 2.3.12.

Construction 4.1.1 (Algebraic BT). For any animated commutative $\mathcal{O}$ -algebra R with derived p -adic completion $\hat{R}$, set

$$\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R) \stackrel{\mathrm{defn}}{=} \mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R}) \times_{T_{\mathrm{ét}}, \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\hat{R}[1/p])} \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R[1/p]),$$

where $T_{\mathrm{ét}}$ is the étale realization functor (which we henceforth suppress from the notation). Also, set

$$\mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(R) = \varprojlim_n \mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R).$$

Remark 4.1.2. For any R and $1 \leq n \leq \infty$, we obtain a canonical étale realization map

$$T_{\mathrm{ét}}: \mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R[1/p])$$

obtained by projecting to the second component.

This globalizes: For any Deligne–Mumford stack $\mathcal{X}$ over $\mathcal{O}$ with generic fiber X , we obtain a functor

$$T_{\mathrm{ét}}: \mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{X}) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(X).$$

Theorem 4.1.3 (Algebraic representability and affineness of diagonal). *The functor $\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}$ is represented by a smooth 0-dimensional Artin stack over $\mathcal{O}$ with affine diagonal.*

The rest of this subsection and the next will be devoted to the proof of this theorem, with the completion of the proof of the affineness of the diagonal postponed to the next subsection. For now, let us record an important consequence.

Definition 4.1.4 (η -normality). Suppose that $\mathcal{X}$ is a Deligne–Mumford stack over $\mathcal{O}$. We will say that $\mathcal{X}$ is η -normal¹⁶ if it admits an étale cover by affine schemes of the form $\mathrm{Spec} R$ with R both p -torsion free and integrally closed in $R[1/p]$. Note that, if $\mathcal{X}$ is normal then it is automatically η -normal.

Theorem 4.1.5 (Tate full faithfulness for apertures). *Let $\mathcal{X}$ be a Noetherian, η -normal Deligne–Mumford stack over $\mathcal{O}$ with generic fiber X . Then the functor*

$$T_{\mathrm{ét}}: \mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{X}) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(X)$$

is fully faithful.

Proof. By étale descent, we reduce to the case where $\mathcal{X} = \mathrm{Spec} R$ is affine. We essentially follow part of the proof by Tate of his theorem for p -divisible groups [89]. By Theorem 4.1.3, for any pair

$$(\Omega_1, \Omega_2) \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(R) \times \mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(R),$$

the scheme of isomorphisms from Ω_1 to Ω_2 is represented by an affine scheme $\mathcal{I} \rightarrow \mathrm{Spec} R$. The theorem comes down to knowing that every $R[1/p]$ -point of $\mathcal{I}$ extends to an R -valued point.

For this, by the η -normality of R , it is enough to know that an $R[1/p]$ -valued point of $\mathcal{I}$ extends over the localization at every height 1 prime of R that is minimal over pR .¹⁷ This reduces us to the case where R is a mixed

¹⁶See [1, Appendix A]

¹⁷The argument for this is identical to that of algebraic Hartogs lemma for normal rings; see [87, Tag 031T].

characteristic discrete valuation ring with maximal ideal $\mathfrak{m}$. Now, let κ be the residue field of R . For any purely inseparable finite extension κ'/κ , there exists a finite flat quasisyntomic map of discrete valuation rings $R \rightarrow R'$ such that $R'/\mathfrak{m}R' \simeq \kappa'^{18}$. Using this, one sees that there exists an ind-finite flat quasisyntomic cover $R \rightarrow R_\infty$ where R_∞ is a discrete valuation ring with perfect residue field. Via flat descent, it is now enough to show:

- (1) $\mathcal{I}(R_\infty) \rightarrow \mathcal{I}(R_\infty[1/p])$ is a bijection.
- (2) $\mathcal{I}(R_\infty^{\otimes_{R^m}}) \rightarrow \mathcal{I}(R_\infty^{\otimes_{R^m}}[1/p])$ is injective.

By the definition of $\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}$, to prove assertion (1) (resp. (2)), we can replace R_∞ (resp. $R_\infty^{\otimes_{R^m}}$) with its (derived) p -completion. Note that in assertion (2) the algebras involved are quasisyntomic over R_∞ . Therefore, both assertions follow from Theorem 2.3.6. $\square$

faithfulness

Corollary 4.1.6. *Suppose that $\mathcal{X}$ is a normal flat Noetherian Deligne-Mumford stack over $\mathcal{O}$ and $\mathcal{X}' \rightarrow \mathcal{X}$ is a smooth cover. Then, for $\mathcal{Q} \in \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(\mathcal{X}_\eta)$, there exists a $\mathfrak{Q} \in \mathrm{BT}_\infty^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{X})$ with $T_{\mathrm{ét}}(\mathfrak{Q})$ isomorphic to $\mathcal{Q}$ if and only if there is a $\mathfrak{Q}' \in \mathrm{BT}_\infty^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{X}')$ with $T_{\mathrm{ét}}(\mathfrak{Q}')$ isomorphic to $\mathcal{Q}|_{\mathcal{X}'[1/p]}$.*

Proof. The only if direction is obvious. For the if direction, since $\mathcal{X}' \rightarrow \mathcal{X}$ is smooth each k -fold self-fiber product $(\mathcal{X}')^{\times_{\mathcal{X}} k}$ is still a normal flat Noetherian DM stack over $\mathcal{O}$. Therefore, Theorem 4.1.5 tells us that the tautological descent datum for $\mathcal{Q}|_{\mathcal{X}'}$ yields a descent datum for $\mathfrak{Q}'$, which is effective. $\square$

Remark 4.1.7. Combining the theorem with Theorem 2.4.2, we obtain a proof of Tate's seminal full faithfulness theorem for p -divisible groups [89, Theorem 4]. Unwinding everything, one sees that the proof obtained here is not very different from that of Tate's, except that his final reduction is to the case where there exists a homomorphism of p -divisible groups over a complete valuation ring inducing a given isomorphism of Tate modules. Since we do not have any direct way of defining a map between apertures that is not an isomorphism, we end up appealing to the faithfulness assertion of Theorem 2.3.6 instead.

Let us now begin preparations for the proof of Theorem 4.1.3.

presentation

Lemma 4.1.8 (Completed finite presentation). *Suppose that $\{R_i\}_{i \in I}$ is an inductive system of derived p -complete animated commutative rings with colimit R , and let $\hat{R}$ be the derived p -completion of R . Then the natural map*

$$\varinjlim_i \mathrm{BT}_n^{\mathcal{G}, \mu}(R_i) \rightarrow \mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R})$$

is an equivalence.

Proof. If the inductive system is one of $\mathbb{Z}/p^m\mathbb{Z}$ -algebras, then the conclusion holds because $\mathrm{BT}_n^{\mathcal{G}, \mu}$ is a finitely presented formal Artin stack.

For the general case, set $\bar{R}_i \stackrel{\mathrm{defn}}{=} R_i/\mathbb{L}p^2$ and $\bar{R} = R/\mathbb{L}p^2$,¹⁹ and note that by Grothendieck–Messing theory we have a Cartesian square

$$\begin{array}{ccc} \mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R}) & \longrightarrow & B\mathcal{P}_\mu^-(R/\mathbb{L}p^n) \\ \downarrow & & \downarrow \\ \mathrm{BT}_n^{\mathcal{G}, \mu}(\bar{R}) & \longrightarrow & B\mathcal{P}_\mu^-(\bar{R}/\mathbb{L}p^n) \times_{B\mathcal{G}(\bar{R}/\mathbb{L}p^n)} B\mathcal{G}(R/\mathbb{L}p^n), \end{array}$$

and similarly with $\hat{R}$ replaced by R_i . We can now conclude using the previous paragraph, and the finite presentation of the stacks $B\mathcal{P}_\mu^-$ and $B\mathcal{G}$.²⁰ $\square$

presentation

Lemma 4.1.9 (Algebraic finite presentation). *The prestack $\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}$ over $\mathcal{O}$ is locally of finite presentation.*

¹⁸If $\kappa' = \kappa(a^{1/p})$ for some $a \in \kappa \setminus \kappa^p$, then we can take $R' = R[X]/(X^p - \pi X - \tilde{a})$, for some uniformizer $\pi \in R$ and some lift $\tilde{a} \in R$ of a . The general case follows by induction on $[\kappa' : \kappa]$.

¹⁹The choice of p^2 here is to ensure that $R \rightarrow \bar{R}$ is an inverse limit of *nilpotent* divided power extensions even when $p = 2$: For p odd, one could also choose to work with $R/\mathbb{L}p$ instead.

²⁰We are using the fact that filtered colimits of ∞ -groupoids commute with finite limits; see [63, Proposition 5.3.3.3].

Proof. We need to know that the natural map

$$\begin{array}{c} \varinjlim_i \left[\mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R}_i) \times_{\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\hat{R}_i[1/p])} \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R_i[1/p]) \right] \\ \downarrow \\ \mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R}) \times_{\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\hat{R}[1/p])} \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R[1/p]) \end{array}$$

is an isomorphism for any inductive system $\{R_i\}$ of animated commutative $\mathcal{O}$ -algebras with colimit R .

Now, since $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}$ is a finite Deligne–Mumford stack the natural map

resentation (4.1.9.1)
$$\varinjlim_i \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(R_i[1/p]) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\varinjlim_i R_i[1/p]),$$

is an isomorphism. Therefore, it is enough to verify that the following commutative square is Cartesian

$$\begin{array}{ccc} \varinjlim_i \mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R}_i) & \longrightarrow & \mathrm{BT}_n^{\mathcal{G}, \mu}(\hat{R}) \\ \downarrow & & \downarrow \\ \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\varinjlim_i \hat{R}_i[1/p]) & \longrightarrow & \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}(\hat{R}[1/p]). \end{array}$$

The ring $\varinjlim_i \hat{R}_i$ is p -Henselian with p -adic completion $\hat{R}$. Therefore, by [12, Corollary 2.1.20], the bottom arrow is an isomorphism. The top arrow is an isomorphism by Lemma 4.1.8. $\square$

The next result is immediate from the definitions, the (formal) étaleness of the Deligne–Mumford stack $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}$ over $\mathcal{O}[1/p]$, and Theorem 3.1.5.

ion_theory **Lemma 4.1.10** (Algebraic deformation theory). *For every nilpotent divided power extension $R' \twoheadrightarrow R$ of animated commutative $\mathcal{O}$ -algebras, there is a canonical Cartesian diagram*

$$\begin{array}{ccc} \mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R') & \longrightarrow & B\mathcal{P}_\mu^-(R'/\mathbb{L}p^n) \\ \downarrow & & \downarrow \\ \mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R) & \longrightarrow & B\mathcal{P}_\mu^-(R/\mathbb{L}p^n) \times_{B\mathcal{G}(R/\mathbb{L}p^n)} B\mathcal{G}(R'/\mathbb{L}p^n). \end{array}$$

Proof of Theorem 4.1.3. The proof will be via verifying the conditions for Artin–Lurie representability [62, Theorem 7.1.6]. It is important for this that we allow animated inputs, though with this caveat the reader will note that the argument itself (except for the parts establishing finite presentation and affineness of the diagonal) is quite formal. In what follows, we will use without comment the fact that $\mathrm{BT}_n^{\mathcal{G}, \mu} \otimes \mathbb{Z}/p^m\mathbb{Z}$ (for $m \geq 1$) and $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}$ are both locally finitely presented algebraic stacks over $\mathcal{O}$, and so satisfy the conditions of *loc. cit.*

Condition (1: local finite presentation) is verified by Lemma 4.1.9. Conditions (2: being an étale sheaf), (3: integrability), (5: infinitesimal cohesiveness), and (6: nilcompleteness) all involve behavior with respect to limits and are easily verified.

For condition (4: existence of a cotangent complex), note that, for any $\mathcal{O}$ -algebra R and a map $x: \mathrm{Spf} \hat{R} \rightarrow \mathrm{BT}_n^{\mathcal{G}, \mu}$, the cotangent complex $\mathbb{L}_{\mathrm{BT}_n^{\mathcal{G}, \mu}/\mathcal{O}_{E_v}, x}$ is a $\mathbb{Z}/p^n\mathbb{Z}$ -module object in the ∞ -category $\mathrm{Perf}(\hat{R})$. In particular, it algebraizes canonically to a perfect complex over R . If x lifts to a point $(x, x') \in \mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R)$, then Lemma 4.1.10 shows that the assignment $(x, x') \mapsto \mathbb{L}_{\mathrm{BT}_n^{\mathcal{G}, \mu}/\mathcal{O}_{E_v}, x}$ is the desired cotangent complex for $\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}$ over $\mathcal{O}$. Note that this is still a perfect complex over $\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}$ with Tor amplitude in $[0, 1]$.

Condition (7: 1-truncatedness) is verified by seeing that $\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}}(R)$ is 1-truncated for any discrete $\mathcal{O}$ -algebra R : Indeed, this is true for all the spaces involved in the fiber product defining it.

Thus we have verified all the conditions of Lurie's theorem, and so can conclude using this, the description of the cotangent complex, and the quasi-compactness of both $\mathrm{BT}_n^{\mathcal{G},\mu}$ and $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})}$ that $\mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}}$ is a smooth 0-dimensional Artin stack over $\mathcal{O}$.

We are now tasked with showing that the diagonal Δ , which we now know is a locally of finite type algebraic space, is actually affine. For this, note that $\Delta[1/p]$ is affine (in fact, finite étale) and the p -adic completion $\hat{\Delta}$ is also affine by Theorem 3.1.5. Moreover, it follows from Proposition 3.9.4 that the diagonal is a *separated* algebraic space: Indeed, the cited result shows that the map $\Delta(R) \rightarrow \Delta(F)$ is injective for any perfectoid valuation ring R of mixed characteristic with fraction field F . Since every mixed characteristic discrete valuation ring is dominated by a perfectoid valuation ring, we see that the injectivity also holds for such discrete valuation rings. For discrete valuation rings of equal characteristic, this injectivity is immediate from the already noted facts about $\Delta[1/p]$ and $\hat{\Delta}$.

Choose a smooth cover

$$\mathrm{Spec} R \rightarrow \mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}} \times \mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}}$$

by a smooth affine scheme over $\mathcal{O}$. Let B be the finite étale $R[1/p]$ -algebra representing the pullback of $\Delta[1/p]$, and let S be the topologically of finite type $\hat{R}$ -algebra representing the pullback of $\hat{\Delta}$. Then the criterion of Achinger–Yousis from [2, Proposition 3.6] tells us that $\Delta|_{\mathrm{Spec} R}$ is affine if and only if the natural map $B \rightarrow S[1/p]$ has dense image. This follows from Proposition 4.2.4 below. $\square$

4.2. Analytic properties of the diagonal. The purpose of this subsection is to prove Proposition 4.2.4, and so to complete the proof of the affineness of the diagonal of $\mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}}$.

Setup 4.2.1. Suppose that we have a smooth covering $\mathrm{Spec} R \rightarrow \mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}} \times \mathrm{BT}_n^{\mathcal{G},\mu,\mathrm{alg}}$ as in the proof of Theorem 4.1.3. This classifies a pair of n -truncated apertures $(\mathfrak{Q}_1, \mathfrak{Q}_2)$ over the p -completion $\hat{R}$ and a pair of $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -local systems (Q_1, Q_2) over $R[1/p]$ whose restrictions over $\hat{R}[1/p]$ are isomorphic to $(T_{\mathrm{ét}}(\mathfrak{Q}_1), T_{\mathrm{ét}}(\mathfrak{Q}_2))$. Write $\Delta_R \rightarrow \mathrm{Spec} R$ for the restriction of the diagonal, and let $\hat{\Delta}_R \rightarrow \mathrm{Spf} \hat{R}$ be the associated formal algebraic space. As we observed in the proof of Theorem 4.1.3, Δ_R is a finitely presented *separated* algebraic space over R .

Remark 4.2.2. By definition, $\hat{\Delta}_R$ parameterizes isomorphisms between the n -truncated apertures $\mathfrak{Q}_1$ and $\mathfrak{Q}_2$. The diagonal of $\mathrm{BT}_n^{\mathcal{G},\mu}$ is an affine, finitely presented map, and so $\hat{\Delta}_R$ is represented by a topologically of finite type $\hat{R}$ -algebra S . Similarly, $\Delta_R[1/p]$ is the finite étale scheme over $R[1/p]$ parameterizing isomorphisms between the $\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})$ -local systems Q_1 and Q_2 , and so is represented by a finite étale $R[1/p]$ -algebra B .

Remark 4.2.3. We can associate with Δ_R the generic fiber $X \stackrel{\mathrm{defn}}{=} (\hat{\Delta}_R)_\eta$ of the formal scheme $\hat{\Delta}_R$: This is a finitely presented affinoid adic space over $Y \stackrel{\mathrm{defn}}{=} \mathrm{Spa}(\hat{R}[1/p], \hat{R})$. We can also consider the affinoid adic space X' over Y obtained by taking the analytification (relative to R in the sense of [2, Definition 2.15]) of the finite étale $\hat{R}[1/p]$ -scheme $\hat{R}[1/p] \otimes_R \Delta_R$.

Concretely, we have $X = \mathrm{Spa}(S[1/p], S)$ and $X' = \mathrm{Spa}(\hat{B}, \hat{B}^+)$, where $\hat{B} = B \otimes_{R[1/p]} \hat{R}[1/p]$, and $\hat{B}^+ \subset \hat{B}$ is the integral closure of $\hat{R}$. We then obtain a diagram of adic spaces

$$\begin{array}{ccc} X & \xrightarrow{\quad} & X' \\ & \searrow & \downarrow \\ & & Y \end{array}$$

where the vertical arrow is finite [36, (1.4.2)], and where the top horizontal map is an open immersion [36, Proposition 1.9.6]: For the latter, see also [2, Proposition 2.16].

Note that the map $X \rightarrow X'$ from Remark 4.2.3 yields maps of $R[1/p]$ -algebras $B \rightarrow \hat{B} \rightarrow S[1/p]$. Our goal is to prove the following:

Proposition 4.2.4. *The map $B \rightarrow S[1/p]$ has dense image.*

Remark 4.2.5 (The diagonal as an analytic diamond). Suppose that T is a perfectoid R -algebra and fix a generator ξ of $\mathrm{Fil}_{\mathcal{N}}^1 \Delta_T$. Unwinding Remark 3.2.20, and after étale localization on $\mathrm{Spf} T$ if necessary, we can assume that the restriction of $(\mathfrak{Q}_1, \mathfrak{Q}_2)$ over T arises from a pair (x_1, x_2) belonging to the set $\mathcal{G}(\Delta_T/p^n \Delta_T) \times \mathcal{G}(\Delta_T/p^n \Delta_T)$. Then, for any p -torsion free perfectoid T -algebra T' that is integrally closed in $T'[1/p]$, we have canonical identifications⁷

$$\begin{aligned} X(T'[1/p], T') &= \{h \in H_{\mu}^{(n)}(T') : x_2 = h^{-1} x_1 \varphi(\mu(\xi) h \mu(\xi)^{-1})\} \\ &= \{h \in H_{\mu}^{(n)}(T') : h = x_1 \varphi(\mu(\xi) h \mu(\xi)^{-1}) x_2^{-1}\}, \end{aligned}$$

and

$$\begin{aligned} X'(T'[1/p], T') &= \{h \in \mathcal{G}(\Delta_{T'}/p^n \Delta_{T'}[1/\xi]) : x_2 = h^{-1} x_1 \varphi(\mu(\xi) h \mu(\xi)^{-1})\} \\ &= \{h \in \mathcal{G}(\Delta_{T'}/p^n \Delta_{T'}[1/\xi]) : h = x_1 \varphi(\mu(\xi) h \mu(\xi)^{-1}) x_2^{-1}\} \end{aligned}$$

In other words, we have concretely described the diamonds $X_T^{\diamond}, X_T'^{\diamond}$ underlying the adic spaces $\mathrm{Spa}(T[1/p], T) \times_Y X$ and $\mathrm{Spa}(T[1/p], T) \times_Y X'$ as in [83, §10.1].

Remark 4.2.6 (Linearization). Keep the setup from the previous remark. Choose a faithful representation $\mathcal{G} \hookrightarrow \mathrm{GL}(\Lambda)$, and set $M = \mathrm{End}(\Lambda) \otimes_{\mathbb{Z}_p} \Delta_T/p^n \Delta_T$: This is equipped with a canonical φ -semilinear bijection $1 \otimes \varphi$, which we will denote simply by φ . Write $\Phi : M[1/\xi] \xrightarrow{\sim} M[1/\xi]$ for the operator $m \mapsto x_1 \varphi(\mu(\xi)) \varphi(m) \varphi(\mu(\xi))^{-1} x_2^{-1}$, and define functors $Z^{\diamond}$ and $Z'^{\diamond}$ on affinoid perfectoid spaces over $(T[1/p], T)$ by

$$\begin{aligned} Z^{\diamond}(T'[1/p], T') &= \{m \in \Delta_{T'} \otimes_{\Delta_T} M : m = \Phi(m) \in (\Delta_{T'} \otimes_{\Delta_T} M)[1/\xi]\} \\ Z'^{\diamond}(T'[1/p], T') &= \{m \in (\Delta_{T'} \otimes_{\Delta_T} M)[1/\xi] : m = \Phi(m) \in (\Delta_{T'} \otimes_{\Delta_T} M)[1/\xi]\} \end{aligned}$$

We then have an isomorphism $X_T^{\diamond} \xrightarrow{\sim} Z^{\diamond} \times_{Z'^{\diamond}} X_T'^{\diamond}$ of diamonds over $\mathrm{Spd}(T[1/p], T) = \mathrm{Spa}(T^{\flat}[1/\varpi], T^{\flat})$, where $\varpi \in T^{\flat}$ is the pseudouniformizer given by the image of ξ under the natural map $\Delta_T \rightarrow T^{\flat}$.

Lemma 4.2.7. *The diamonds $Z^{\diamond}$ and $Z'^{\diamond}$ are represented by affinoid perfectoid spaces $Z = \mathrm{Spa}(D, D^+)$ and $Z' = \mathrm{Spa}(D', D'^+)$ over $\mathrm{Spa}(T[1/p], T)$. Moreover, the map $Z \rightarrow Z'$ is a rational open embedding, and the map $D \rightarrow D'$ has dense image.*

Proof. Via the usual dévissage, we reduce to the case where $n = 1$. Choose $k \geq 0$ with $\Phi(M) \subset \varpi^{-k} M$, and set $\tilde{\Phi} = \varpi^k \Phi$. In the following we identify $\Delta_T/p \Delta_T[1/\xi]$ with $T^{\flat}[1/\varpi]$.

By Artin–Schreier theory, the functor $C \mapsto (C \otimes_{T^{\flat}[1/\varpi]} M[1/\varpi])^{\Phi=1}$ is represented by a finite étale group scheme over $T^{\flat}[1/\varpi]$ locally isomorphic to $\mathbb{F}_p^{\mathrm{rank}_{T^{\flat}} M}$. Therefore, there exists $r \geq 0$ such that, for all affinoid perfectoids $\mathrm{Spa}(T'[1/p], T')$ over T , we have

$$\begin{aligned} (4.2.7.1) \quad Z^{\diamond}(T'[1/p], T') &= \left\{ m \in \varpi^{-r} (T'^{\flat} \otimes_{T^{\flat}} M) : \Phi(m) = m \right\} \\ &\xrightarrow{\sim} \left\{ n \in T'^{\flat} \otimes_{T^{\flat}} M : \Phi(n) = \varpi^{pr} n \right\} \\ &= \left\{ n \in T'^{\flat} \otimes_{T^{\flat}} M : \tilde{\Phi}(n) = \varpi^{k+pr} n \right\}, \end{aligned}$$

where the middle identification is given by $m \mapsto \varpi^r m$. Moreover, via this identification, we have

$$(4.2.7.2) \quad Z'^{\diamond}(T'[1/p], T') \simeq Z'^{\diamond}(T'[1/p], T') \cap \varpi^r (T'^{\flat} \otimes_{T^{\flat}} M) \subset T'^{\flat} \otimes_{T^{\flat}} M.$$

Now, (4.2.7.1) tells us that $Z'^{\diamond}$ is represented by an affinoid perfectoid over $\mathrm{Spa}(T^{\flat}[1/\varpi], T^{\flat})$: It is a closed subsheaf of the perfectoid closed unit disk

$$\mathbf{D}_{\mathrm{perf}}(M) : (T'^{\flat}[1/\varpi], T'^{\flat}) \mapsto T'^{\flat} \otimes_{T^{\flat}} M,$$

associated with M . Furthermore, (4.2.7.2) tells us that $Z^{\diamond}$ is a rational open in $Z'^{\diamond}$, obtained as the pre-image of the rational open immersion $\mathbf{D}_{\mathrm{perf}}(M) \xrightarrow{\varpi^r} \mathbf{D}_{\mathrm{perf}}(M)$.

By the tilting equivalence [83, Theorem 6.2.7], we conclude that $Z = \mathrm{Spa}(D, D^+)$ and $Z' = \mathrm{Spa}(D', D'^+)$ are represented by affinoid perfectoids over $\mathrm{Spa}(T[1/p], T)$. It remains to see that the properties of being a rational open

[7] **Alex:** Should cite Remark 3.2.20 somewhere here

embedding and of $D^b \rightarrow D'^b$ having dense image are preserved by untilting: The first follows from [44, Theorem 3.6.14], and the second can be deduced from the explicit description of untilts of certain rational open embeddings in Lemma 3.6.13 of *loc. cit.* $\square$

Proof of Proposition 4.2.4. Note that $\hat{R}$ is p -completely smooth over $\mathcal{O}$ and so admits a p -completely flat map $\hat{R} \rightarrow \hat{R}_\infty$ where $\hat{R}_\infty$ is perfectoid and is the p -completed filtered colimit of finite flat $\hat{R}$ -algebras. Lemma 4.2.7 and Remark 4.2.6 together tell us that the map

$$(4.2.7.3) \quad B \hat{\otimes} \hat{R}_\infty \rightarrow (S \hat{\otimes} \hat{R}_\infty)[1/p]$$

has dense image. This implies that $B \rightarrow S[1/p]$ has dense image. Indeed, it suffices to show that, for all $n \geq 1$, if Q_n denotes the cokernel of the map of R -modules $B \rightarrow S[1/p]/p^n S$ then $Q_n = 0$. Let us note that Q_n is p^∞ -torsion, and as $R \rightarrow \hat{R}_\infty$ is p -completely faithfully flat, we see that Q_n is zero if and only if $Q_n \otimes_R \hat{R}_\infty$ is zero. But, this follows from the density of the image in (4.2.7.3). $\square$

4.3. An extension property. The following proposition will be used to prove Theorem A.

Proposition 4.3.1. *Suppose that we have two η -normal separated algebraic spaces $\mathcal{X}_1$ and $\mathcal{X}_2$ over $\mathcal{O}$, and that we are given $\mathfrak{Q}_i \in \mathrm{BT}_\infty^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{X}_i)$ lifting $Q_i \in \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(X_i)$ where $X_i \stackrel{\mathrm{defn}}{=} \mathcal{X}_{i, \eta}$. Suppose that:*

- (1) *The stack $\mathcal{X}_2$ is of finite type over $\mathcal{O}$ and the stack $\mathcal{X}_1$ is excellent.*
- (2) *The formal classifying map $\hat{\mathcal{X}}_2 \rightarrow \mathrm{BT}_\infty^{\mathcal{G}, \mu}$ for $\mathfrak{Q}_2$ is formally unramified.*
- (3) *There is a map $f: X_1 \rightarrow X_2$ and an isomorphism $\zeta: Q_2|_{X_1} \xrightarrow{\sim} Q_1$.*
- (4) *Given any mixed characteristic $(0, p)$ complete discrete valuation field F over $\mathcal{O}$ with perfect residue field, f maps $\mathcal{X}_1(\mathcal{O}_F) \subset X_1(F)$ to $\mathcal{X}_2(\mathcal{O}_F) \subset X_2(F)$.*

Then there are unique extensions of f to a map $\mathcal{X}_1 \rightarrow \mathcal{X}_2$ and of ζ to an isomorphism $\mathfrak{Q}_1|_{\mathcal{X}_1} \xrightarrow{\sim} \mathfrak{Q}_2$.

Proof. The proof we are about to see is essentially a generalization of an argument of Pappas from [72, Theorem 7.1.7] (see also [39, Theorem 3.13]).

Let $\mathcal{Y} \rightarrow \mathcal{X}_1 \times_{\mathrm{Spec} \mathcal{O}} \mathcal{X}_2$ be constructed as follows: Take the scheme-theoretic image $\mathcal{Y}'$ of the graph

$$X_1 \xrightarrow{\mathrm{id} \times f} X_1 \times_{\mathrm{Spec} E} X_2 \hookrightarrow \mathcal{X}_1 \times_{\mathrm{Spec} \mathcal{O}} \mathcal{X}_2.$$

Since $\mathcal{X}_2$ is of finite type over $\mathcal{O}$, we see $\mathcal{Y}'$ is of finite type over $\mathcal{X}_1$. Now take $\mathcal{Y}$ to be the η -normalization of $\mathcal{Y}'$ ²¹: This is of finite type over $\mathcal{X}_1$ because of the excellence; see hypothesis [87, Tag 035S]. It suffices to show $\mathcal{Y} \xrightarrow{\sim} \mathcal{X}_1$, as this implies that $\mathcal{Y}$ is the graph of a unique extension $\mathcal{X}_1 \rightarrow \mathcal{X}_2$ with the desired properties.

The restrictions of $\mathfrak{Q}_1$ and $\mathfrak{Q}_2$ yield two objects in $\mathrm{BT}_\infty^{\mathcal{G}, \mu}(\mathcal{Y})$ equipped with an isomorphism of the underlying local systems in $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(\mathcal{Y}_\eta)$. Therefore, by Theorem 4.1.5, we see that there is an isomorphism $\mathfrak{Q}_1|_{\mathcal{Y}} \xrightarrow{\sim} \mathfrak{Q}_2|_{\mathcal{Y}}$. In other words, the classifying maps

$$\mathcal{Y} \rightarrow \mathcal{X}_1 \rightarrow \mathrm{BT}_\infty^{\mathcal{G}, \mu, \mathrm{alg}}; \quad \mathcal{Y} \rightarrow \mathcal{X}_2 \rightarrow \mathrm{BT}_\infty^{\mathcal{G}, \mu, \mathrm{alg}},$$

are isomorphic.

Suppose that we have a closed geometric point $y \in \mathcal{Y}(\kappa)$ of the special fiber mapping to $x_i \in \mathcal{X}_i(\kappa)$. We claim that the map of complete local Noetherian rings $\hat{\mathcal{O}}_{\mathcal{X}_1, x_1} \rightarrow \hat{\mathcal{O}}_{\mathcal{Y}, y}$ is finite.

Write $z \in \mathrm{BT}_\infty^{\mathcal{G}, \mu}(W(\kappa))$ for the point corresponding to $\mathfrak{Q}_{1, x_1} \simeq \mathfrak{Q}_{2, x_2}$. Let $\hat{\mathcal{O}}_y$ (resp. $\hat{\mathcal{O}}_{x_1}$, $\hat{\mathcal{O}}_{x_2}$, and $\hat{\mathcal{O}}_z$) be the universal deformation rings over $W(\kappa)$ for $\mathcal{Y}$ (resp. $\mathcal{X}_1$, $\mathcal{X}_2$, and $\mathrm{BT}_\infty^{\mathcal{G}, \mu, \mathrm{alg}}$) at their respective κ -points. We now have factorizations

$$\hat{\mathcal{O}}_z \rightarrow \hat{\mathcal{O}}_{x_1} \rightarrow \hat{\mathcal{O}}_y; \quad \hat{\mathcal{O}}_z \rightarrow \hat{\mathcal{O}}_{x_2} \rightarrow \hat{\mathcal{O}}_y,$$

of the map $\hat{\mathcal{O}}_z \rightarrow \hat{\mathcal{O}}_y$. By our formal unramifiedness hypothesis, the first map in the second factoring is surjective. By construction, the combined map

$$\hat{\mathcal{O}}_{x_1} \hat{\otimes}_{W(\kappa)} \hat{\mathcal{O}}_{x_2} \rightarrow \hat{\mathcal{O}}_y,$$

²¹More specifically, take $\mathcal{Y}$ to be relative spectrum over $\mathcal{Y}'$ of the quasi-coherent $\mathcal{O}_{\mathcal{Y}'}$ -algebra given by integral closure of $\mathcal{O}_{\mathcal{Y}'}$ in $j_* \mathcal{O}_{\mathcal{Y}'_\eta}$ where $j: \mathcal{Y}'_\eta \rightarrow \mathcal{Y}'$ is the open inclusion.

whose image is now the same as that of the first factor $\widehat{\mathcal{O}}_{x_1}$, is finite (since $\mathcal{Y} \rightarrow \mathcal{X}_1 \times_{\text{spf } \mathcal{O}} \mathcal{X}_2$ is finite). We conclude that the map $\widehat{\mathcal{O}}_{x_1} \rightarrow \widehat{\mathcal{O}}_y$ is already finite.

What we have seen above shows that the map $\mathcal{Y} \rightarrow \mathcal{X}_1$ is quasi-finite; see for instance [80, Ch. IV, Propositions 2 and 3]. Since it is also an isomorphism after inverting p , it follows from Zariski's Main Theorem and the η -normality of $\mathcal{X}_1$ that it is in fact an open immersion: one may reduce by étale descent to [30, Corollary 12.88].

We now claim that every algebraically closed point $x_1 \in \mathcal{X}_1(\kappa)$ of the special fiber is inside the open $\mathcal{Y}$, which implies that $\mathcal{Y} \rightarrow \mathcal{X}_1$ is an isomorphism as desired. Indeed, by the flatness of $\mathcal{X}_1$, we can find a mixed characteristic complete discrete valuation ring $\mathcal{O}_F$ with residue field κ such that x_1 lifts to $\mathcal{X}_1(\mathcal{O}_F)$; see [87, Tag 0CM2]. But hypothesis (4) tells us that this lift lies in $\mathcal{Y}(\mathcal{O}_F)$. $\square$

5. CRITERIA FOR THE EXISTENCE OF APERTURES

The purpose of this section is to develop criteria for proving the existence of apertures when $\mathcal{G}$ is reductive. These are mainly of a 'pointwise' nature, and will be employed in the next section to prove the canonicity of integral models. We will maintain the notation from Setup 3.1.1. Most of our results here will carry the additional assumption that $\mathcal{G}$ is in fact reductive.

5.1. Apertures over p -completely smooth $\mathcal{O}_K$ -algebras. Let K be a complete mixed characteristic $(0, p)$ discrete valuation with perfect residue field κ and absolute ramification index e , and fix a uniformizer $\pi \in K$. Our goal here is to characterize the image of the fully faithful étale realization functor for base formal $\mathcal{O}_K$ -schemes. We will assume that $\mathcal{G}$ is *reductive*.

Definition 5.1.1 (The type of a crystalline local system). Suppose that G is reductive and that we have $\mathbf{Q} \in \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}}(K)$. Then, for every representation V of $\mathbf{Q}$ we obtain a filtered K -vector space $\text{Fil}^\bullet D_{\text{dR}}((V)_{\mathbf{Q}})$. By [47, Lemma (1.4.5)], there exists a cocharacter of G_K that splits these filtrations simultaneously for all V . In particular, this gives us a well-defined geometric conjugacy class of cocharacters of G . This conjugacy class is the **type** of $\mathbf{Q}$. If λ is a representative of the conjugacy class defined over K , then we will abuse terminology and say that $\mathbf{Q}$ has type λ . More generally, if $\mathfrak{X}$ is a base formal $\mathcal{O}_K$ -scheme, a crystalline $\mathcal{G}(\mathbb{Z}_p)$ -local systems $\mathbf{Q}$ over $\mathfrak{X}$ is of **type** λ if, for all $V \in \text{Rep}_{\mathbb{Q}_p}(G)$, the filtration $\text{Fil}_{\text{Hdg}}^\bullet D_{\text{dR}}((V)_{\mathbf{Q}})$ is, étale locally on $\mathfrak{X}_\eta$, split by the cocharacter λ .

Notation 5.1.2. For a cocharacter λ of G_K , let $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}, \lambda}(K)$ be the groupoid of crystalline $\mathcal{G}(\mathbb{Z}_p)$ -local systems of type λ on K . More generally, for any base formal $\mathcal{O}_K$ -scheme $\mathfrak{X}$, we define $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}, \mu}(\mathfrak{X}_\eta)$ to be the groupoid of crystalline $\mathcal{G}(\mathbb{Z}_p)$ -local systems of type λ on $\mathfrak{X}_\eta$.

The two following results characterize the image of the fully faithful étale realization functor on $\text{BT}_\infty^{\mathcal{G}, \mu}(R)$ in some cases.

Proposition 5.1.3. *The fully faithful functor*

$$T_{\text{ét}}: \text{BT}_\infty^{\mathcal{G}, \mu}(\mathcal{O}_K) \rightarrow \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(K)$$

has essential image $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}, \mu}(K)$.

Theorem 5.1.4. *Suppose that $\mathfrak{X}$ is a base formal $\mathcal{O}_K$ -scheme and that $2e < p - 1$. Then the fully faithful functor*

$$T_{\text{ét}}: \text{BT}_\infty^{\mathcal{G}, \mu}(\mathfrak{X}) \rightarrow \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(\mathfrak{X}_\eta)$$

has essential image $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}, \mu}(\mathfrak{X}_\eta)$.

Remark 5.1.5 (Crystallinity from pointwise crystallinity). The work of Guo-Yang [35], as explained in Remark 2.3.5, tells us that, when $\mathfrak{X}$ is p -completely smooth over $\mathcal{O}_K$, then the following conditions are equivalent for a $\mathcal{G}(\mathbb{Z}_p)$ -local system $\mathbf{Q}$ over $\mathfrak{X}_\eta$:

- (1) $\mathbf{Q}$ belongs to $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}, \mu}(\mathfrak{X}_\eta)$;
- (2) For every classical point $x: \text{Spa}(K') \rightarrow \mathfrak{X}_\eta$, one has that $\mathbf{Q}_x$ is in $\text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\text{crys}, \mu}(K')$.

Thus, in this case, the second pointwise condition can be used to describe the essential image of $T_{\text{ét}}$.

ialization

Remark 5.1.6 (Crystalline specialization functors). In the situation of either Theorem 5.1.4 or Proposition 5.1.3 (the latter with $\mathfrak{X} = \mathrm{Spf} \mathcal{O}_K$), we obtain functors

$$\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\mathrm{crys}, \mu}(\mathfrak{X}_\eta) \xleftarrow{\sim} \mathrm{BT}_\infty^{\mathcal{G}, \mu}(\mathfrak{X}) \rightarrow \mathrm{BT}_\infty^{\mathcal{G}, \mu}(\mathfrak{X}_\kappa),$$

where $\mathfrak{X}_\kappa$ is the special fiber of $\mathfrak{X}$. Using this, we can functorially associate with every $Q \in \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\mathrm{crys}, \mu}(\mathfrak{X}_\eta)$ a $(\mathcal{G}, \mu)$ -aperture $\mathbb{D}_{\mathrm{crys}}(Q)$ over $\mathfrak{X}_\kappa$. In particular, for every $\Lambda \in \mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G})$, we obtain the F -gauge

$$\mathbb{D}_{\mathrm{crys}}((\Lambda)_Q) \stackrel{\mathrm{defn}}{=} (\Lambda)_{\mathbb{D}_{\mathrm{crys}}(Q)} \in \mathrm{Vect}(\mathfrak{X}_\kappa^{\mathrm{syn}}).$$

If we take the underlying F -crystal over $\mathfrak{X}_\kappa$, then we are essentially getting a special case of the similarly denoted functor from [38, §2].⁸²

To prove our main results, we will first need a bit of preparation. Theorem 5.1.4 quickly reduces to the affine case $\mathfrak{X} = \mathrm{Spf}(R)$, and so we will focus on this below.

e_of_bt_mu

Lemma 5.1.7. *The étale realization functor $T_{\mathrm{ét}}: \mathrm{BT}_\infty^{\mathcal{G}, \mu}(\mathcal{O}_K) \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\mathrm{crys}}(K)$ lands in $\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\mathrm{crys}, \mu}(K)$*

Proof. Given $\Omega \in \mathrm{BT}_\infty^{\mathcal{G}, \mu}(\mathcal{O}_K)$ lifting $Q \in \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}^{\mathrm{crys}}(K)$, Lemma 3.8.4 shows that $\mathrm{Fil}^\bullet D_{\mathrm{dR}}(Q)$ is canonically isomorphic to $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet T_{\mathrm{dR}}(\Omega)$. This, combined with the definition of $\mathrm{BT}_\infty^{\mathcal{G}, \mu}$, implies the claim. $\square$

gebraicity

Remark 5.1.8 (Reduction to a question of algebraicity). Theorem 2.3.6 tells us that the tensor functor $\mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \rightarrow \mathrm{Loc}_{\mathbb{Z}_p}(R[1/p])$ associated with Q factors through an exact symmetric monoidal functor

$$\mathrm{Rep}_{\mathbb{Z}_p}(\mathcal{G}) \rightarrow \mathrm{Vect}^{\mathrm{an}, \varphi}(R_\Delta, \mathcal{O}_\Delta).$$

By [39, Propositions 1.28 and 1.39], the theorem now amounts to showing that this functor factors through the fully faithful subcategory $\mathrm{Vect}^\varphi(R_\Delta, \mathcal{O}_\Delta)$. Indeed, the cited results would then imply that Q arises from a $\mathcal{G}$ -torsor Ω over R^{syn} , and our pointwise assumption on the type, combined with Lemma 5.1.7, implies that Ω must be in $\mathrm{BT}_\infty^{\mathcal{G}, \mu}(R)$.

To establish the desired factorization, it is now sufficient to show that the analytic prismatic F -crystal associated with $(\Lambda)_Q$ is in fact a prismatic F -crystal for *any* faithful representation of $\mathcal{G}$.

_criterion

Remark 5.1.9 (Reduction to a question of finite freeness). Working étale locally on $\mathrm{Spf} R$, we can assume that R is a base $\mathcal{O}_K$ -algebra admitting a Breuil–Kisin frame $\mathfrak{S}_R$ as in Example 3.2.2. The analytic prismatic F -crystal associated with $(\Lambda)_Q$ gives a vector bundle $\mathcal{V}^{\mathrm{an}}$ over $\mathrm{Spec} \mathfrak{S}_R \setminus V(p, E(u))$. Knowing that we in fact have a prismatic F -crystal amounts to knowing that the reflexive $\mathfrak{S}_R$ -module

$$\mathcal{L} = H^0\left(\mathrm{Spec} \mathfrak{S}_R \setminus V(p, E(u)), \mathcal{V}^{\mathrm{an}}\right)$$

is in fact finite locally free (cf. [37, Proposition 1.26]).

Proof of Proposition 5.1.3. In this case, $\mathfrak{S}_{\mathcal{O}_K}$ is a regular local Noetherian ring of dimension 2, and so every reflexive finitely generated module over it is finite free. $\square$

e_of_torus

Lemma 5.1.10. *Suppose that $\mathcal{G} = \mathcal{T}$ is a torus over $\mathbb{Z}_p$. Then, Theorem 5.1.4 holds.*

Proof. We can reduce to the situation in Remark 5.1.9, and so need to check that the reflexive $\mathfrak{S}_R$ -module $\mathcal{L}$ associated with a faithful representation Λ is finite locally free. Since this can be checked after a faithfully flat base change, we can assume without loss of generality that $\mathcal{T}$ is split over $\mathfrak{S}_R$. In particular, $\mathcal{L}^{\mathrm{an}}$ is now a direct sum of line bundles, and this implies that $\mathcal{L}$ is finite locally free; see [31, Exposé XI, Théorème 3.18]. $\square$

lete_local

Lemma 5.1.11 (Reduction to the complete local case). *Suppose that e is arbitrary and that the following equivalent conditions hold for every maximal ideal $\mathfrak{m} \subset R$:*

- (1) *The restriction of Q over $\mathrm{Spec} \hat{R}_{\mathfrak{m}}[1/p]$ lifts to $\mathrm{BT}_\infty^{\mathcal{G}, \mu}(\hat{R}_{\mathfrak{m}})$.*
- (2) *The analytic prismatic F -crystal over $\hat{R}_{\mathfrak{m}}$ associated with $(\Lambda)_Q$ is in fact a prismatic F -crystal.*

Then Q lifts to $\mathrm{BT}_\infty^{\mathcal{G}, \mu}(R)$.

[8] **Alex:** Cou
maybe be a g
to say that, in
eral sense, $\mathbb{D}_{\mathrm{crys}}$
specialization
for the formal
 $\mathrm{BT}_\infty^{\mathcal{G}, \mu}$?
[2] **Keerthi:**
more than the
We're saying
tube around t
reduction locu
described exp
least for smoo

Proof. As in Remark 5.1.9, we can assume that R is a base $\mathcal{O}_K$ -algebra of the form $R_0 \otimes_{W(\kappa)} \mathcal{O}_K$, and are therefore reduced to checking that the $\mathfrak{S}_R$ -module $\mathcal{L}$ is finite locally free. For this, it is enough to check that, for every maximal ideal $\mathfrak{m} \subset R$, lifting to a maximal ideal $\mathfrak{n} \subset \mathfrak{S}_R$, the completion of $\mathcal{L}$ at $\mathfrak{n}$ is finite locally free. However, the completion $(\mathfrak{S}_R)_{\mathfrak{n}}^{\wedge}$ underlies a Breuil–Kisin frame for $\hat{R}_{\mathfrak{m}} = \hat{R}_{0, \mathfrak{m}_0} \otimes_{W(\kappa)} \mathcal{O}_K$, where $\mathfrak{m}_0 = \mathfrak{m} \cap R_0$. Therefore, our equivalent hypotheses tell us that $\hat{\mathcal{L}}_{\mathfrak{n}}$ is obtained from the evaluation of a prismatic F -crystal over $\hat{R}_{\mathfrak{m}}$, and is thus finite free, as desired. $\square$

Notation 5.1.12. Suppose now that R is a base $\mathcal{O}_K$ -algebra and that T is the completion of R at a maximal ideal. Let $\underline{\mathfrak{S}}_T$ be the Breuil–Kisin frame for T obtained as a completion of $\mathfrak{S}_R$ as in the proof above, and set $\mathcal{L}_T = T \otimes_R \mathcal{L}$.

Lemma 5.1.13. *Suppose that μ acts on Λ with weights in the interval $[a, b]$ with $|b - a| = r$. Then, if $er < p - 1$, $\mathcal{L}_T$ is finite free over T . In particular, if $\mathcal{G}$ is adjoint, then Theorem 5.1.4 holds.*

Proof. The first assertion follows from the argument in [59, Proposition 4.3]. The second now follows from the fact that we can choose Λ to be the adjoint representation, which has μ -weights $\{-1, 0, 1\}$. $\square$

Remark 5.1.14. Suppose that κ is algebraically closed. Then one sees that a $\mathcal{G}$ -torsor over $U \stackrel{\text{defn}}{=} \text{Spec } \mathfrak{S}_T \setminus V(p, E(u))$ extends to a $\mathcal{G}$ -torsor over $\mathfrak{S}_T$ if and only if it is actually trivial. This is because every $\mathcal{G}$ -torsor over $\mathfrak{S}_T$ is trivial, since $\mathfrak{S}_T$ is now a strictly Henselian local ring. Moreover, the restriction map for $\mathcal{G}$ -torsors from $\text{Spec } \mathfrak{S}_R$ to U is fully faithful (e.g., see [87, Tag 0EBJ]).

Proof of Theorem 5.1.4. Without loss of generality, we can assume that κ is algebraically closed and that R is a base $\mathcal{O}_K$ -algebra with Breuil–Kisin frame $\underline{\mathfrak{S}}_R$. We only have to show that for every completion T of R , the $\mathcal{G}$ -torsor $\mathcal{Q}_T^{\text{an}}$ over $U \stackrel{\text{defn}}{=} \text{Spec } \mathfrak{S}_T \setminus V(p, E(u))$ obtained from the analytic prismatic F -crystal realization of $\mathbf{Q}$ extends to a $\mathcal{G}$ -torsor over $\mathfrak{S}_T$ or, equivalently, that it is trivial.

Consider the map $\mathcal{G} \rightarrow \bar{\mathcal{G}} \stackrel{\text{defn}}{=} \mathcal{G}^{\text{ad}} \times \mathcal{G}^{\text{ab}}$. This is a finite flat cover of reductive group schemes, and Lemmas 5.1.10 and 5.1.13 together tell us that the theorem is valid with $\mathcal{G}$ replaced with $\bar{\mathcal{G}}$ and $\mathbf{Q}$ replaced with the associated $\bar{\mathcal{G}}(\mathbb{Z}_p)$ -torsor over $R[1/p]$. Therefore, $\mathcal{Q}_T^{\text{an}}$ is trivial after change of structure group to $\bar{\mathcal{G}}$, and so admits a reduction of structure group to a $\mathcal{Z}$ -torsor over U , where $\mathcal{Z} = \ker(\mathcal{G} \rightarrow \bar{\mathcal{G}})$. However, by purity of branch locus for torsors under finite flat group schemes [68, Theorem 3.1], all $\mathcal{Z}$ -torsors over U extend uniquely to $\mathcal{Z}$ -torsors over $\mathfrak{S}_T$. This implies that $\mathcal{Q}_T^{\text{an}}$ also extends to a $\mathcal{G}$ -torsor over $\mathfrak{S}_T$, as desired. $\square$

Remark 5.1.15 (The Hodge type case). If $(\mathcal{G}, \mu)$ is of *Hodge type*, i.e., there exists a map $(\mathcal{G}, \mu) \rightarrow (\text{GL}_h, \mu_d)$ (see Example 3.1.6) for some d such that the underlying map $\mathcal{G} \rightarrow \text{GL}_h$ is a closed immersion, then we can improve the bound to $p < e - 1$: Indeed, we can replace the use of the adjoint representation with the faithful representation given by such a map, and this will have μ -weights in $\{0, 1\}$. When $\mathcal{G} = \text{GL}_h$ this amounts to the assertion that any crystalline local system over $R[1/p]$ with Hodge–Tate weights in $\{0, 1\}$ arises from a p -divisible group. This is a result of Liu–Moon [59, Theorem 1.2].

5.2. Versality. When we obtain an aperture using Theorem 5.1.4, versality is not automatic. In applications, it can be understood via a Kodaira–Spencer map, which we now discuss.

Remark 5.2.1 (The formal de Rham realization and a versality condition). Suppose that we have a map $f: \mathcal{X} \rightarrow \text{BT}_{\infty}^{\mathcal{G}, \mu, \text{alg}}$ of stacks over $\mathcal{O}$. After completion, we obtain a filtered de Rham realization

$$\hat{\mathcal{X}} \rightarrow \text{BT}_{\infty}^{\mathcal{G}, \mu} \xrightarrow{T_{\text{dR}}} \text{Gr}_{\mu} / \mathcal{G}_{\mathcal{O}_{E, (v)}}.$$

By the discussion in Remark 3.4.3, this gives a map of p -complete complexes

$$\mathbb{L}_{(\text{Gr}_{\mu} / \mathcal{G}_{\mathcal{O}_{E, (v)}}) / B\mathcal{G}_{\mathcal{O}}}^{\wedge} |_{\hat{\mathcal{X}}} \rightarrow \mathbb{L}_{\mathcal{X} / \mathcal{O}_{E, (v)}}^{\wedge},$$

which is an isomorphism if and only if the map $\hat{\mathcal{X}} \rightarrow \text{BT}_{\infty}^{\mathcal{G}, \mu}$ is formally étale.

a--Spencer

Remark 5.2.2 (The Kodaira–Spencer map). Suppose that $\mathcal{X}$ is a smooth scheme over $\mathcal{O}_{E,(v)}$. Dualizing the map from the previous remark, we obtain a map

_space_map

$$(5.2.2.1) \quad \mathbb{T}_{\mathcal{X}/\mathcal{O}_{E,(v)}}^{\wedge} \rightarrow \mathbb{T}_{(\mathrm{Gr}_{\mu}/\mathcal{G}_{\mathcal{O}_{E,(v)}})/B\mathcal{G}_{\mathcal{O}}}|_{\widehat{\mathcal{X}}}$$

of vector bundles over $\widehat{\mathcal{X}}$. The left-hand side here is the p -completed tangent bundle of $\mathcal{X}$ while the right-hand side can be described explicitly: The aperture $\mathfrak{Q}$ classified by f admits a Hodge-filtered de Rham realization $\mathrm{Fil}_{\mathrm{Hdg}}^{\bullet} T_{\mathrm{dR}}(\mathfrak{Q})$ equipped with a topologically nilpotent integrable connection. Let $\mathfrak{g}$ be the Lie algebra of $\mathcal{G}$, equipped with the adjoint representation. Twisting this by $\mathrm{Fil}_{\mathrm{Hdg}}^{\bullet} T_{\mathrm{dR}}(\mathfrak{Q})$, we obtain a filtered vector bundle $\mathrm{Fil}^{\bullet} \mathfrak{g}_{\mathrm{twist}}$ over $\widehat{\mathcal{X}}$. There is now a canonical isomorphism

$$\mathbb{T}_{(\mathrm{Gr}_{\mu}/\mathcal{G}_{\mathcal{O}_{E,(v)}})/B\mathcal{G}_{\mathcal{O}}}|_{\widehat{\mathcal{X}}} \simeq \mathfrak{g}_{\mathrm{twist}} / \mathrm{Fil}^0 \mathfrak{g}_{\mathrm{twist}}.$$

Now, (5.2.2.1) now gives a map $\mathbb{T}_{\mathcal{X}/\mathcal{O}_{E,(v)}}^{\wedge} \rightarrow \mathfrak{g}_{\mathrm{twist}} / \mathrm{Fil}^0 \mathfrak{g}_{\mathrm{twist}}$. This is just the Kodaira–Spencer map associated with the integrable connection on $\mathrm{Fil}_{\mathrm{Hdg}}^{\bullet} T_{\mathrm{dR}}(\mathfrak{Q})$, and so versality comes down to knowing that this is an isomorphism.

ture_group

5.3. Apertures via reduction of structure group.

Setup 5.3.1. Let κ be an algebraically closed characteristic p field over $\mathcal{O}$, and let $R_1 \rightarrow R_2$ be a map of complete local normal rings that are flat and formally of finite type over $W(\kappa)$. We will assume that both R_1 and R_2 have residue field κ . Suppose that we have a closed immersion $(\mathcal{G}_2, \mu_2) \rightarrow (\mathcal{G}_1, \mu_1)$ of two pairs as in Setup 3.1.1, with μ_1 and μ_2 both defined over $\mathcal{O}$, and with both $\mathcal{G}_1$ and $\mathcal{G}_2$ *reductive*. Suppose also that we have $\mathbf{Q}_i \in \mathrm{Loc}_{\mathcal{G}_i(\mathbb{Z}_p)}(R_i[1/p])$ such that the restriction of $\mathbf{Q}_1$ over $\mathrm{Spec} R_2[1/p]$ is isomorphic to the change of structure group of $\mathbf{Q}_2$ along the map $\mathcal{G}_2(\mathbb{Z}_p) \rightarrow \mathcal{G}_1(\mathbb{Z}_p)$. We will assume that $\mathbf{Q}_1$ lifts to $\mathfrak{Q}_1 \in \mathrm{BT}_{\infty}^{\mathcal{G}_1, \mu_1}(R_1)$. Write $\mathfrak{Q}_{1,0} = \mathfrak{Q}_1|_{\kappa^{\mathrm{syn}}}$ for the resulting aperture over κ .

via_lambda

Remark 5.3.2. Let us put ourselves in the situation of Setup 3.2.14: We will choose a faithful representation Λ of $\mathcal{G}_1$, and take $\{s_{\alpha}\} \subset \Lambda^{\otimes}$ to be a finite collection of tensors whose pointwise stabilizer is $\mathcal{G}_2$. As in Remark 3.2.15, the F -gauge $(\Lambda)_{\mathfrak{Q}_1}|_{\kappa^{\mathrm{syn}}}$ corresponds to a p -adically filtered $W(\kappa)$ -module $\mathrm{Fil}^{\bullet} L_0$ equipped with an isomorphism

$$\sum_{i \in \mathbb{Z}} p^{-i} \varphi^* \mathrm{Fil}^i L_0 \xrightarrow{\sim} L_0.$$

For every $x: R_2 \rightarrow \mathcal{O}_K$, the existence of the reduction of structure $\mathbf{Q}_2$ for $\mathbf{Q}_1|_{\mathrm{Spec} R_2[1/p]}$ tells us that we have global sections

$$\{s_{\alpha,x}\} \subset H^0(\mathrm{Spec} K, (\Lambda)_{\mathfrak{Q}}^{\otimes}).$$

Via the specialization functor $\mathbb{D}_{\mathrm{crys}}$, these map to φ -invariant tensors

$$\{\mathbb{D}_{\mathrm{crys}}(s_{\alpha,x})\} \in \mathbb{D}_{\mathrm{crys}}((\Lambda)_{\mathfrak{Q}_2|_x})^{\otimes} \simeq L_0^{\otimes}.$$

The isomorphism $\mathbb{D}_{\mathrm{crys}}((\Lambda)_{\mathfrak{Q}_2|_x}) \simeq L_0$ used here is the canonical one induced by the specialization functor.

_limpidity

Remark 5.3.3 (Families of F -isocrystals). Suppose that R_1 is formally smooth over $W(\kappa)$, equipped with a Frobenius lift $\varphi: R_1 \rightarrow R_1$. Then the rigid analytic space $\widehat{U}_{1,\eta}$ associated with $\widehat{U}_1 = \mathrm{Spf} R_1$ is an open ball. Moreover, the F -gauge $\mathfrak{L}$ lifting $(\Lambda)_{\mathfrak{Q}_1}$ yields a filtered F -isocrystal over R_1/pR_1 , which realizes to a filtered vector bundle $\mathrm{Fil}^{\bullet} \mathcal{L}_{\mathbb{Q}}$ over $\widehat{U}_{1,\eta}$, equipped with an integrable connection, and specializing to $\mathrm{Fil}^{\bullet} L_x[1/p]$ at every classical point $x \in \widehat{U}_{1,\eta}(K)$. Set

$$\mathcal{L}_{\mathbb{Q}}^{\nabla} \stackrel{\mathrm{defn}}{=} H^0(\widehat{U}_{1,\eta}, \mathcal{L}_{\mathbb{Q}})^{\nabla=0}.$$

This is a finite dimensional vector space over $W(\kappa)[1/p]$, and inherits the structure of an F -isocrystal from that on $\mathcal{L}_{\mathbb{Q}}$. Moreover, we have a canonical isomorphism of F -isocrystals

$$\delta: \mathcal{O}_{\widehat{U}_{1,\eta}} \otimes_{W(\kappa)[1/p]} \mathcal{L}_{\mathbb{Q}}^{\nabla} \xrightarrow{\sim} \mathcal{L}_{\mathbb{Q}}$$

over $\widehat{U}_{1,\eta}$. This is a consequence of Dwork’s trick; see [43, Proposition 3.1].

_limpidity

Remark 5.3.4 (Application of the de Rham functor of Liu–Zhu). By Remark 3.8.2, $\mathrm{Fil}^\bullet \mathcal{L}_\mathbb{Q}$ is also obtained from the filtered F -isocrystal associated with the $\mathbb{Z}_p$ -local system $(\Lambda)_{\mathbb{Q}_1}$. If $\widehat{U}_{2,\eta}$ is a *smooth* rigid analytic space over $W(\kappa)[1/p]$, then the restriction of $\mathrm{Fil}^\bullet \mathcal{L}_\mathbb{Q}$ over $\widehat{U}_{2,\eta}$ is once again a filtered vector bundle with integrable connection, and is *associated* with $(\Lambda)_{\mathbb{Q}_2}$ via the D_{dR} functor of Liu–Zhu [58, Theorem 3.9]. In particular, this functor carries the global sections

$$\{s_{\alpha, \mathbb{Q}_2}\} \subset H^0(\mathrm{Spec} R_2[1/p], (\Lambda)_{\mathbb{Q}_2}^{\otimes})$$

to a collection

$$\{s_{\alpha, \mathcal{L}_\mathbb{Q}}\} \subset H^0(\widehat{U}_{2,\eta}, \mathrm{Fil}^0 \mathcal{L}_\mathbb{Q}^{\otimes})^{\nabla=0}.$$

_reduction

Remark 5.3.5 (Type of reduction at every point). Suppose that $\mathbb{Q}_1$ lifts to $\mathfrak{Q}_1 \in \mathrm{BT}_\infty^{\mathcal{G}_1, \mu_1}(R_1)$, and that we have a finite extension $K/W(\kappa)[1/p]$, and a map $x: R_2 \rightarrow \mathcal{O}_K$ of complete local rings. The $\mathcal{G}_1(\mathbb{Z}_p)$ -local system $\mathbb{Q}_1|_x$ has a reduction of structure to a $\mathcal{G}_2(\mathbb{Z}_p)$ -local system $\mathbb{Q}_2|_x$, which is in $\mathrm{Loc}_{\mathcal{G}_2(\mathbb{Z}_p)}^{\mathrm{crys}, \mu'_2}(K)$ for some cocharacter μ'_2 mapping to the $\mathcal{G}_1$ -conjugacy class of μ_1 . We will refer to the conjugacy class of μ'_2 as the **type** of x .

_specializations

Remark 5.3.6 (Compatibility between specializations of points). Suppose that we have two points $x: R_2 \rightarrow \mathcal{O}_K$ and $y: R_2 \rightarrow \mathcal{O}_L$ as in Remark 5.3.5 for finite extensions K and L of $W(\kappa)$. Then, via the specialization functor from Remark 5.1.6, we obtain $\mathcal{G}_2$ -bundles $\mathbb{D}_{\mathrm{crys}}(\mathbb{Q}_2|_x)$ and $\mathbb{D}_{\mathrm{crys}}(\mathbb{Q}_2|_y)$ over κ , which are both canonically reductions of structure group for the $\mathcal{G}_1$ -torsor $\mathfrak{Q}_2$. In particular, it makes sense to ask for them to be *equal*; in this case, it follows that x and y have the same type. Explicitly, this equality amounts to asking that, for every index α , we have

$$\mathbb{D}_{\mathrm{crys}}(s_{\alpha, x}) = \mathbb{D}_{\mathrm{crys}}(s_{\alpha, y}) \in L_0^{\otimes}$$

_structure_group

Proposition 5.3.7. *In the situation of Remark 5.3.3, suppose further that the rigid analytic space $\widehat{U}_{2,\eta}$ is also smooth over $W(\kappa)[1/p]$, and that some x as above has type μ_2 . Then:*

- (1) *There exists a lift of $\mathbb{Q}_2$ to $\mathfrak{Q}_2 \in \mathrm{BT}_\infty^{\mathcal{G}_2, \mu_2}(R_2)$.*
- (2) *Suppose that the following conditions hold:*
 - (a) *The classifying map $\mathrm{Spf} R_1 \rightarrow \mathrm{BT}_\infty^{\mathcal{G}_1, \mu_1}$ for $\mathfrak{Q}_1$ is formally unramified;*
 - (b) *If $I = \ker(R_1 \rightarrow R_2)$, then R_2 is finite over R_1/I and we have an equality of fraction fields $K(R_1/I) = K(R_2)$;*
 - (c) $\dim R_2 = 1 + \dim \mathcal{G}_2 - \dim \mathcal{P}_{\mu_2}^-$.*Then $R_1 \rightarrow R_2$ is in fact surjective and the classifying map $\mathrm{Spf} R_2 \rightarrow \mathrm{BT}_\infty^{\mathcal{G}_2, \mu_2}$ for $\mathfrak{Q}_2$ is the universal deformation ring over $W(\kappa)$ for $\mathrm{BT}_\infty^{\mathcal{G}_2, \mu_2}$ at $\mathfrak{Q}_{2,0}$.*

Proof. To begin, note that there is a canonical isomorphism of F -isocrystals $L_0[1/p] \xrightarrow{\sim} \mathcal{L}_\mathbb{Q}^\nabla$. Suppose that we have $x: R_2 \rightarrow \mathcal{O}_K$ as in our hypothesis. The tensors $\{s_{\alpha, \mathcal{L}_x}\}$ map to the φ -invariant tensors

$$\{\mathbb{D}_{\mathrm{crys}}(s_{\alpha, x})\} \in L_0^{\otimes},$$

and these in turn yield a collection of parallel φ -invariant tensors

$$\{\tilde{s}_{\alpha, \mathcal{L}_\mathbb{Q}}\} \stackrel{\mathrm{defn}}{=} \delta(\{1 \otimes \mathbb{D}_{\mathrm{crys}}(s_{\alpha, x})\}) \subset H^0(\widehat{U}_{1,\eta}, \mathcal{L}_\mathbb{Q}^{\otimes})^{\nabla=0}.$$

The restriction of this collection over $\widehat{U}_{2,\eta}$ agrees with $\{s_{\alpha, \mathcal{L}_x}\}$ at x and so, by horizontality, must agree everywhere.²² Thus, the tensors $\{\mathbb{D}_{\mathrm{crys}}(s_{\alpha, x})\}$ are independent of x , and so we see using Remark 5.3.6 that there is a canonical reduction of structure $\mathfrak{Q}_{2,0} \subset \mathfrak{Q}_{1,0}$ to a $(\mathcal{G}_2, \mu_2)$ -aperture over κ such that, for all finite extensions $K'/W(\kappa)[1/p]$ and all maps $x': R_2 \rightarrow \mathcal{O}_{K'}$, we have

$$\mathbb{D}_{\mathrm{crys}}(\mathbb{Q}_2|_{x'}) = \mathfrak{Q}_{2,0} \subset \mathfrak{Q}_{1,0}.$$

Let $\tilde{R}_i$ be the universal deformation ring for $\mathfrak{Q}_{i,0}$ for $i = 1, 2$ (see Remark 3.1.7). The map $\tilde{R}_1 \rightarrow \tilde{R}_2$ is a surjective map of formally smooth complete local Noetherian $W(\kappa)$ -algebras: This amounts to the observation (which follows from the Grothendieck–Messing theory explained in Theorem 3.1.5) that $\mathrm{BT}_\infty^{\mathcal{G}_2, \mu_2} \rightarrow \mathrm{BT}_\infty^{\mathcal{G}_1, \mu_1}$ is a formally unramified map of formally smooth formal stacks.

²²We are using the connectedness of $\widehat{U}_{2,\eta}$ here; see [18, Lemma 7.3.5].

We have a map $\tilde{R}_1 \rightarrow R_1$ classifying $\mathfrak{Q}_1$ as a deformation of $\mathfrak{Q}_{1,0}$. Now, by what we have seen above, for every $x': R_2 \rightarrow \mathcal{O}_{K'}$, the composition $\tilde{R}_1 \rightarrow R_1 \rightarrow R_2 \rightarrow \mathcal{O}_{K'}$ factors through $\tilde{R}_2$. Since $\tilde{R}_1 \rightarrow \tilde{R}_2$ is surjective, this implies that in fact the composition $\tilde{R}_1 \rightarrow R_1 \rightarrow R_2$ factors through $\tilde{R}_2$: In other words, we have a deformation $\mathfrak{Q}_2$ over R_2 of $\mathfrak{Q}_{2,0}$ specializing to the various $T_{\text{ét}}^{-1}(\mathfrak{Q}_2|_{x'})$. To finish, we need to know that $T_{\text{ét}}(\mathfrak{Q}_2) = \mathfrak{Q}_2$ as reductions of structure to $\mathcal{G}_2(\mathbb{Z}_p)$ -torsors within the $\mathcal{G}_1(\mathbb{Z}_p)$ -local system $\mathfrak{Q}_1|_{\text{Spec } R_2[1/p]}$, but this equality can be checked over the single given point x , where it holds by construction.

Let us now verify (2): Under the hypotheses here, the map $\tilde{R}_1 \rightarrow R_1$ is surjective, and the map $\tilde{R}_2 \rightarrow R_2$ is finite and its image has the same field of fractions as R_2 . Since we have the equality

$$\dim \tilde{R}_2 = 1 + \dim \mathcal{G}_2 - \dim \mathcal{P}_{\mu_2}^- = \dim R_2$$

by hypothesis, and since $\tilde{R}_2$ is normal, we now find that $\tilde{R}_2 \xrightarrow{\sim} R_2$. This of course shows that $R_1 \rightarrow R_2$ is surjective, and identifies R_2 with the deformation ring of $\text{BT}_{\infty}^{\mathcal{G}_2, \mu_2}$ at $\mathfrak{Q}_{2,0}$. $\square$

ture_group

Setup 5.3.8. $\mathcal{X}$ will be a normal algebraic space flat of finite type over $\mathcal{O}$ with generic fiber X , equipped with $\mathfrak{Q} \in \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(X)$. We will assume $\mathcal{X}$ is equidimensional of relative dimension $\dim \mathcal{G} - \dim \mathcal{P}_{\mu}^-$ over $\mathcal{O}$. Also, we will require that there exist a closed immersion $(\mathcal{G}, \mu) \hookrightarrow (\mathcal{G}^{\sharp}, \mu^{\sharp})$ of pairs as in Setup 3.1.1 with $\mathcal{G}, \mathcal{G}^{\sharp}$ both reductive, and that there exists a commuting diagram

$$\begin{array}{ccc} X & \longrightarrow & \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)} = \text{BT}_{\infty}^{\mathcal{G}, \mu, \text{alg}} \otimes \mathbb{Q}_p \\ \downarrow & & \downarrow \\ \mathcal{X} & & \text{BT}_{\infty}^{\mathcal{G}, \mu} \\ \downarrow & & \downarrow \\ \mathcal{X}^{\sharp} & \longrightarrow & \text{BT}_{\infty}^{\mathcal{G}^{\sharp}, \mu^{\sharp}, \text{alg}} \end{array}$$

where the top horizontal arrow classifies $\mathfrak{Q}$.

ion_normal

Proposition 5.3.9. *In the situation of Setup 5.3.8, suppose that the following condition holds: For all closed points x_0 of $\mathcal{X}$, the tuple*

$$(R_1 = \hat{\mathcal{O}}_{\mathcal{X}^{\sharp}, x_0}, R_2 = \hat{\mathcal{O}}_{\mathcal{X}, x_0}, \mathfrak{Q}_1 = \mathfrak{Q}^{\sharp}|_{R_1^{\text{syn}}}, \mathfrak{Q}_2 = \mathfrak{Q}_2|_{\text{Spec } R_2[1/p]})$$

satisfies the hypotheses of Proposition 5.3.7. Then $\mathfrak{Q}$ lifts to $\mathfrak{Q} \in \text{BT}_{\infty}^{\mathcal{G}, \mu, \text{alg}}(\mathcal{X})$, and the classifying map $\hat{\mathcal{X}} \rightarrow \text{BT}_{\infty}^{\mathcal{G}, \mu}$ is formally étale.

Proof. We can assume that $\mathcal{X}$ is affine. Proposition 5.3.7 tells us that the complete local ring $\hat{R}_x$ of $\mathcal{X}$ at any geometric closed point x is formally smooth over $\mathcal{O}$ and that in fact $\mathfrak{Q}$ lifts to a reduction of structure $\mathfrak{Q}_x \in \text{BT}_{\infty}^{\mathcal{G}, \mu}(\hat{R}_x)$ such that the classifying map $\text{Spf } \hat{R}_x \rightarrow \text{BT}_{\infty}^{\mathcal{G}, \mu}$ is formally étale. In particular, $\hat{\mathcal{X}}$ is p -completely smooth over $\hat{\mathcal{O}}$, and the conditions of Lemma 5.1.11 have been verified. This shows the existence of $\mathfrak{Q}$.

That the resulting classifying map on $\hat{\mathcal{X}}$ is formally étale now follows from Proposition 5.3.7 once again, and Lemma 5.3.10 below. $\square$

_etale ness

Lemma 5.3.10. *Suppose that we have a p -complete ring R and:*

- A smooth p -adic formal algebraic space $\mathfrak{X}$ over $\text{Spf } R$;
- A formally cohesive²³ and integrable p -adic formal prestack $\mathfrak{Y}$ over $\text{Spf } R$ admitting a p -completed cotangent complex $\mathbb{L}_{\mathfrak{Y}/R}^{\wedge}$ that is in fact a vector bundle of finite rank over $\mathfrak{Y}$; and
- A map $\varpi: \mathfrak{X} \rightarrow \mathfrak{Y}$;

with the following property: For every closed point $x: \text{Spec } \kappa \rightarrow \mathfrak{X}$ with κ algebraically closed, the map of complete local rings $\hat{\mathcal{O}}_{\mathfrak{Y}, \varpi(x)} \rightarrow \hat{\mathcal{O}}_{\mathfrak{X}, x}$ is an isomorphism. Then ϖ is formally étale: that is, the formal cotangent complex $\mathbb{L}_{\mathfrak{X}/\mathfrak{Y}}^{\wedge}$ is nullhomotopic.

²³See [62, Definition 6.2.1].

Proof. Note that our hypotheses imply that $\mathfrak{Y}$ admits universal deformation rings, so the given property is sensible; see [62, Corollary 6.2.14], which is just a classical criterion of Schlessinger. Moreover, by our smoothness hypothesis, the cotangent complex $\mathbb{L}_{\mathfrak{X}/R}^\wedge$ is a vector bundle of finite rank over $\mathfrak{X}$.

We can of course assume that $\mathfrak{X} = \mathrm{Spf} S$ for some p -complete R -algebra S with $\mathbb{L}_{(S/pS)/(R/pR)}$ finite locally free over S/pS . Via the fundamental sequence

$$\varpi^* \mathbb{L}_{\mathfrak{Y}/R}^\wedge \rightarrow \mathbb{L}_{\mathfrak{X}/R}^\wedge \rightarrow \mathbb{L}_{\mathfrak{X}/\mathfrak{Y}}^\wedge$$

of p -completed cotangent complexes and our hypotheses, we find that $\mathbb{L}_{\mathfrak{X}/\mathfrak{Y}}^\wedge$ is a perfect complex over S with Tor amplitude in $[-1, 0]$. Therefore, to prove the lemma, it is enough to know that, for any map $x: S \rightarrow \kappa$ with κ algebraically closed and with kernel a maximal ideal, we have $x^* \mathbb{L}_{\mathfrak{X}/\mathfrak{Y}}^\wedge \simeq 0$ as perfect complexes over κ .

Let $\mathcal{C}_\kappa$ be the ∞ -category of animated Artin $W(\kappa)$ -algebras with residue field κ ²⁴. Then, by the definition of universal deformation rings in the derived context, for any $(A, \mathfrak{m})$ in $\mathcal{C}_\kappa$ equipped with a local map from $\widehat{\mathcal{O}}_{\mathfrak{Y}, \varpi(x)}$, we have⁹³

$$\mathcal{M}_x(A) \stackrel{\mathrm{defn}}{=} \mathrm{fib}_x(\mathrm{Map}_{/\mathfrak{Y}}(\mathrm{Spec} A, \mathfrak{X}) \rightarrow \mathrm{Map}_{/\mathfrak{Y}}(\mathrm{Spec} \kappa, \mathfrak{X})) \simeq \mathrm{Map}_{\widehat{\mathcal{O}}_{\mathfrak{Y}, \varpi(x)}/}(\widehat{\mathcal{O}}_{\mathfrak{X}, x}, A)$$

where the mapping space on the right is taken in the ∞ -category of animated complete local $W(\kappa)$ -algebras with residue field κ . Our hypothesis now says that the right-hand side is contractible.

On the other hand, by the definition of the cotangent complex, if we take $A = \kappa \oplus M$ to be an animated square-zero extension of κ by a bounded above complex M of κ -vector spaces, we have

$$\mathcal{M}_x(A) \simeq \tau^{\leq 0} \mathrm{RHom}_\kappa(x^* \mathbb{L}_{\mathfrak{X}/\mathfrak{Y}}^\wedge, M).$$

Thus, in total, we see that $x^* \mathbb{L}_{\mathfrak{X}/\mathfrak{Y}}^\wedge$ is nullhomotopic, as desired. □

5.4. Behavior under central isogenies.

Setup 5.4.1. Suppose that we have a central extnsion $\mathcal{G} \rightarrow \overline{\mathcal{G}}$ of reductive group schemes over $\mathbb{Z}_p$. By this we mean that there is a faithfully flat homomorphism $\mathcal{G} \rightarrow \overline{\mathcal{G}}$ of reductive $\mathbb{Z}_p$ -group schemes whose kernel $\mathcal{Z}$ is central in $\mathcal{G}$. For instance, $\overline{\mathcal{G}} = \mathcal{G}^{\mathrm{ad}}$ could be its adjoint quotient. The cocharacter μ induces a cocharacter $\overline{\mu}$ of $\overline{\mathcal{G}}$, and we have the associated zero dimensional algebraic stacks $\mathrm{BT}_n^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{alg}}$ over $\mathcal{O}$ with p -completion $\mathrm{BT}_n^{\overline{\mathcal{G}}, \overline{\mu}}$ over $\hat{\mathcal{O}}$ and with generic fiber $\mathrm{Loc}_{\overline{\mathcal{G}}(\mathbb{Z}/p^n\mathbb{Z})}$ viewed as a stack over $\mathcal{O}[1/p]$.

Proposition 5.4.2. *The natural map*

$$\mathrm{BT}_n^{\mathcal{G}, \mu, \mathrm{alg}} \rightarrow \mathrm{BT}_n^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{alg}}$$

is an étale surjection of stacks over $\mathcal{O}$. In fact, the map factors as a gerbe banded by $\mathcal{Z}(\mathbb{Z}/p^n\mathbb{Z})$ followed by a torsor for $H_{\mathrm{fppf}}^1(\mathrm{Spec} \mathbb{Z}/p^n\mathbb{Z}, \mathcal{Z})$.

Proof. The deformation theory from Lemma 4.1.10 tells us that this map is étale.

In the generic fiber, the map factors as

$$\mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})} \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})/\mathcal{Z}(\mathbb{Z}/p^n\mathbb{Z})} \rightarrow \mathrm{Loc}_{\overline{\mathcal{G}}(\mathbb{Z}/p^n\mathbb{Z})}$$

where the first map is a gerbe banded by $\mathcal{Z}(\mathbb{Z}/p^n\mathbb{Z})$ and the second map parameterizes reductions of structure group for $\overline{\mathcal{G}}(\mathbb{Z}/p^n\mathbb{Z})$ -local systems to $(\mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})/\mathcal{Z}(\mathbb{Z}/p^n\mathbb{Z}))$ -local systems. The exact sequence

$$1 \rightarrow \mathcal{G}(\mathbb{Z}/p^n\mathbb{Z})/\mathcal{Z}(\mathbb{Z}/p^n\mathbb{Z}) \rightarrow \overline{\mathcal{G}}(\mathbb{Z}/p^n\mathbb{Z}) \rightarrow H_{\mathrm{fppf}}^1(\mathrm{Spec} \mathbb{Z}/p^n\mathbb{Z}, \mathcal{Z}) \rightarrow 1$$

obtained from the vanishing of $H_{\mathrm{fppf}}^1(\mathrm{Spec} \mathbb{Z}/p^n\mathbb{Z}, \mathcal{G})$ —a consequence of Lang’s lemma—tells us that the second map is a torsor for $H_{\mathrm{fppf}}^1(\mathrm{Spec} \mathbb{Z}/p^n\mathbb{Z}, \mathcal{Z})$.

In particular, this description tells us that the map is surjective in the generic fiber. The surjectivity in the special fiber follows from the quotient description for algebraically closed points from Remark 3.2.20. □

⁴

²⁴See [62, §6.2].

[9] **Alex:** Per
we should rem
slashes/backsl
the subscript
that’s inconsi
the notation v
elsewhere, e.g
next centered
[3] **Keerthi:**
isn’t really an
sistency becau
next centered
involves a diff
of mapping sp
not one in a c
category.

[4] **Keerthi:**

hfulness_2

Proposition 5.4.3 (Descending along central covers). *Suppose that $\overline{\mathcal{X}}$ is a normal flat Noetherian algebraic space over $\mathcal{O}$ with $\overline{\mathbf{Q}} \in \mathrm{Loc}_{\overline{\mathcal{G}}(\mathbb{Z}_p)}(\overline{\mathcal{X}}_\eta)$. Suppose that there exist:*

- (1) *An étale cover $\mathcal{X} \rightarrow \overline{\mathcal{X}}$ with a lift $\mathbf{Q} \in \mathrm{Loc}_{\mathcal{G}}(\mathcal{X}_\eta)$ of the restriction of $\overline{\mathbf{Q}}$;*
- (2) *A lift $\mathbf{\Omega} \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{X})$ of $\mathbf{Q}$.*

Then $\overline{\mathbf{Q}}$ lifts to $\overline{\mathbf{\Omega}} \in \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{alg}}(\overline{\mathcal{X}})$, and the p -completed classifying map $\widehat{\overline{\mathcal{X}}} \rightarrow \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}}$ for $\overline{\mathbf{\Omega}}$ is formally étale if and only if the corresponding one $\widehat{\mathcal{X}} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}, \mu}$ is so.

Proof. The first part of the conclusion is immediate from Corollary 4.1.6, while the second is a consequence of Proposition 5.4.2. $\square$

tral_cover

Setup 5.4.4. Suppose that we have the following data:

- (1) A flat normal algebraic space $\mathcal{X}$ over $\mathcal{O}$ with generic fiber X ;
- (2) An aperture $\overline{\mathbf{\Omega}} \in \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{alg}}(\mathcal{X})$ lifting $\overline{\mathbf{Q}} \in \mathrm{Loc}_{\overline{\mathcal{G}}(\mathbb{Z}_p)}(X)$;
- (3) A finite Galois cover $Y \rightarrow X$ with Galois group Δ such that $\overline{\mathbf{Q}}$ lifts to $\mathbf{Q} \in \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}(Y)$.

Let $\mathcal{Y} \rightarrow \mathcal{X}$ be the normalization of $\mathcal{X}$ in Y .

Notation 5.4.5. Let $\mathcal{X}^{\mathrm{ord}} \subset \mathcal{X}$ be the open subspace with the same generic fiber but whose p -adic completion is $\widehat{\mathcal{X}} \times_{\mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}}} \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{ord}}$.

cal_lift_X

Remark 5.4.6. Suppose that $\mathcal{X}^{\mathrm{ord}} \rightarrow \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{alg}}$ is formally étale after p -adic completion. Then, for every perfect field κ over k and $x \in \mathcal{X}(\kappa)$, we have a *canonical lift* $x^{\mathrm{can}} \in \mathcal{X}(W(\kappa))$ corresponding to the canonical lift $\overline{\mathbf{\Omega}}_x^{\mathrm{can}}$ of the $\overline{\mu}$ -ordinary aperture $\overline{\mathbf{\Omega}}_x$ (Definition 3.6.10).

_condition

Remark 5.4.7. In the situation of the previous remark, suppose $K/W(\kappa)[1/p]$ is a finite extension with residue field κ' and that $\tilde{y} \in \mathcal{Y}(\mathcal{O}_K)$ is a lift of x^{can} , mapping to $y \in \mathcal{Y}(\kappa)$. If the $\mathcal{G}(\mathbb{Z}_p)$ -local system $\mathbf{Q}|_{\tilde{y}}$ admits a lift $\mathbf{\Omega}_{\tilde{y}} \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu}(\mathcal{O}_K)$, we obtain a lift $\mathbf{\Omega}_y \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu}(\kappa')$ of $\mathbf{\Omega}_x \in \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}}$, and this is in the μ -ordinary locus of $\mathrm{BT}_{\infty}^{\mathcal{G}, \mu}$. It is therefore sensible to ask if $\mathbf{\Omega}_{\tilde{y}}$ is the canonical lift of $\mathbf{\Omega}_y$. If in addition we in fact have $\tilde{y} \in \mathcal{Y}(W(\kappa'))$, we will say that $\tilde{y}$ is the *canonical lift* of its mod- p reduction.

tral_covers

Proposition 5.4.8 (Ascending along central covers). *Fix an algebraic closure κ of k , and suppose that the following is true:*

- (1) *$\mathcal{X}$ is smooth over $\mathcal{O}$;*
- (2) *The classifying map $\mathcal{X} \rightarrow \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}, \mathrm{alg}}$ is formally étale after p -adic completion;*
- (3) *Given $x \in \mathcal{X}(\kappa)$ with canonical lift $x^{\mathrm{can}} \in \mathcal{X}(W(\kappa))$, every lift of x^{can} to $\mathcal{Y}$ is the canonical lift of its mod- p reduction.*

Then:

- (1) *$\mathcal{Y} \rightarrow \mathcal{X}$ is once again a Galois cover;*
- (2) *$\mathbf{Q}$ lifts to $\mathbf{\Omega} \in \mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{Y})$.*

Proof. Hypotheses (2) tells us that $\mathcal{X}^{\mathrm{ord}} \subset \mathcal{X}$ is open and fiberwise dense.

To prove assertion (1), by purity of the branch locus [32, Corollaire 3.3], it is enough to know that $\mathcal{Y}^{\mathrm{ord}} \stackrel{\mathrm{defn}}{=} \mathcal{Y} \times_{\mathcal{X}} \mathcal{X}^{\mathrm{ord}} \rightarrow \mathcal{X}^{\mathrm{ord}}$ is a Galois cover.

Moreover, once assertion (1) is known, we see in particular that $\mathcal{Y}$ is also smooth over $\mathcal{O}$. Therefore, to prove assertion (2), we reduce, using Lemma 5.4.9, Proposition 5.4.2 and purity of branch locus once again, to showing that $\mathbf{Q}|_{\mathcal{Y}^{\mathrm{ord}}}$ lifts to $\mathrm{BT}_{\infty}^{\mathcal{G}, \mu, \mathrm{alg}}(\mathcal{Y}^{\mathrm{ord}})$.

Therefore, we see that, without loss of generality, we can assume that $\mathcal{X}^{\mathrm{ord}} = \mathcal{X}$ and $\mathcal{Y}^{\mathrm{ord}} = \mathcal{Y}$.

Now, suppose that we have $y \in \mathcal{Y}(\kappa)$ mapping to $x \in \mathcal{X}(\kappa)$, and let $\widehat{Y}_y$ be the completion of $\mathcal{Y}$ at y . $\square$

_extending

Lemma 5.4.9. *Suppose that $\mathcal{X}$ is a smooth algebraic space over $\mathcal{O}$ with generic fiber X and that $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ is a gerbe banded by a finite abelian group Δ . Suppose that there is a section $s : X \rightarrow \tilde{\mathcal{X}}$ and an open dense subset $U \subset \mathcal{X}_k$ of*

the special fiber, with complement $Z \subset \mathcal{X}_k$, such that s extends to a section over the open $\mathcal{X} \setminus Z$. Then s extends to a section $\mathcal{X} \rightarrow \tilde{\mathcal{X}}$.

Proof. To begin, if we have two extensions of s over $\mathcal{X}$, then the space of isomorphisms between them is a Δ -torsor over $\mathcal{X}$, admitting a canonical trivialization over X , and so is itself canonically trivial (see for instance [87, Tag 0BQG]). In particular, to prove the existence of such an extension, we can work étale locally and assume that the gerbe is trivial. The section s now corresponds to a Δ -torsor $Y \rightarrow X$, and the extension property amounts to the concrete assertion that the normalization of $\mathcal{X}$ in Y is a finite Galois cover $\mathcal{Y} \rightarrow \mathcal{X}$. This follows from our hypotheses and purity of branch locus [32, Corollaire 3.3]. $\square$

6. INTEGRAL CANONICAL MODELS: DEFINITIONS AND FIRST PROPERTIES

We will be following the conventions for Shimura varieties as laid out in [20].

Setup 6.0.1. Given a Shimura datum (G, X) , for $x \in X$, the conjugacy class $\{\mu_x\}$ of the associated Shimura cocharacter μ_x ²⁵ is defined over the reflex field $E \subset \mathbb{C}$ and is independent of the choice of x . We will denote this E -rational class by $\{\mu\}$. For a compact open subgroup $K \subset G(\mathbb{A}_f)$, let Sh_K be the canonical model over E for the complex Shimura variety with complex points given by the orbifold

$$\text{Sh}_K(\mathbb{C}) = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K.$$

Note that Sh_K is a Deligne–Mumford stack over E that is a quasi-projective scheme when K is *neat* in the sense of [77, §0.6], and that in general Sh_K is the stack quotient of $\text{Sh}_{K'}$ by the finite group K/K' , for any choice of neat compact open subgroup $K' \subset K$.

Remark 6.0.2 (Comparison with the work of Lovering). Since¹⁰ the characterization given here is at finite level, it works without any additional argument to give integral canonical models of pre-abelian type over $\mathcal{O}_E[1/N]$ where N is the set of primes where K is unramified; see Theorem 8.5.6. It also immediately gives a functorial way of obtaining filtered F -crystals lifting automorphic vector bundles (see Remark ??). In particular, it recovers Lovering’s results from [61, 60] and extends them to the pre-abelian context.

Remark 6.0.3 (Comparison with Shen). In¹¹ [85], Shen defines a notion of de Rham F -gauge, and constructs such an object on $\mathcal{S}_K \otimes \overline{\mathbb{F}}_p$ where $\mathcal{S}_K$ is the integral canonical model (a la Kisin) of a Shimura variety of Hodge type. This essentially boils down to the morphism $\mathcal{S}_K \otimes \overline{\mathbb{F}}_p \rightarrow \text{BT}_1^{\mathcal{G}, -\mu_v}$ obtained from the universal aperture over $\mathcal{S}_K$.

6.1. Étale realizations on Shimura varieties.

Definition 6.1.1 ([51, Definition 1.5.4, Lemma 1.5.5]). A torus T over $\mathbb{Q}$ is **cuspidal** if it is isogenous to a product of a $\mathbb{Q}$ -split torus and a $\mathbb{Q}$ -torus that is anisotropic over $\mathbb{R}$.

For a general $\mathbb{Q}$ -torus T , the **anti-cuspidal part** $T_{ac} \subset T$ is the minimal $\mathbb{Q}$ -subtorus such that T/T_{ac} is cuspidal.

Remark 6.1.2. The anti-cuspidal part is also characterized as the smallest subtorus of the maximal anisotropic subtorus $T_a \subset T$ that contains the maximal $\mathbb{R}$ -split subtorus of $T_{a, \mathbb{R}}$: in other words, it is the $\mathbb{Q}$ -Zariski closure in T of the maximal $\mathbb{R}$ -split subtorus of $T_{a, \mathbb{R}}$.

Definition 6.1.3 ([51, §1.5.6]). We will define $G^c = G/Z(G)_{ac}$, where $Z(G) \subset G$ is the center, and where $Z(G)_{ac}$ is the anti-cuspidal part of the connected component $Z(G)^\circ \subset Z(G)$: we will refer to it as the **cuspidal quotient** of G .

Remark 6.1.4. As noted in the footnote on p. 34 of *loc. cit.*, this definition differs from the one used for instance by Lovering in [61], since we are not assuming Milne’s axiom requiring that $Z(G)^\circ$ split over a CM field. If we made this assumption, then we could identify $Z(G)_{ac}$ with the maximal anisotropic subtorus of $Z(G)$ that is $\mathbb{R}$ -split.

Lemma 6.1.5. Set $Z(G)_K \stackrel{\text{def}}{=} Z(G)(\mathbb{Q}) \cap K$. Then:

²⁵For any representation W of $G_{\mathbb{R}}$, x yields a Hodge structure W_x on W and μ_x acts on $W_x^{p,q}$ by $z \mapsto z^{-p}$.

:canonical

m:lovering

rorem:Shen

ion_global

i-cuspidal

lem:GQK

[10] Alex: K
we need to m
moved it from

[11] Alex: K
we need to m
moved it from

- (1) The map of complex analytic spaces $X \times G(\mathbb{A}_f) \rightarrow \mathrm{Sh}_K(\mathbb{C})$ is Galois with Galois group given by $(G(\mathbb{Q}) \times K)/Z(G)_K$ where $Z(G)_K$ is embedded diagonally.
- (2) For any normal subgroup $K' \subset K$ $\mathrm{Sh}_{K'} \rightarrow \mathrm{Sh}_K$ is a finite Galois cover with Galois group $K/K'Z(G)_K$.
- (3) For K neat, we have $Z(G)_K \subset Z(G)_{ac}(\mathbb{Q})$; in particular, the map $G(\mathbb{Q}) \rightarrow G^c(\mathbb{Q})$ factors through $Z(G)_K \backslash G(\mathbb{Q})$, and the map $K \rightarrow G^c(\mathbb{A}_f)$ factors through $K/Z(G)_K^-$, where $Z(G)_K^- \subset K$ is the closure of $Z(G)_K$ in K .

Proof. Only the last assertion needs proof. This follows from [51, Lemma 1.5.7]. $\square$

realization

Construction 6.1.6 (Étale realization). By assertion (2) of Lemma 6.1.5, we have a cofiltered system of Galois covers

$$\{\mathrm{Sh}_{K'} \rightarrow \mathrm{Sh}_K\}_{K' \triangleleft K}$$

where each cover has Galois group $K/K'Z(G)_K$. The inverse limit of this system is a pro-Galois cover of Sh_K with Galois group $K/Z(G)_K^-$, which by assertion (3) of the same lemma admits a map to $G^c(\mathbb{A}_f)$. Let $K^c \subset G^c(\mathbb{A}_f)$ be the compact open subgroup determined as follows: Let S be the finite set of primes such that $K \cap G(\mathbb{Q}_\ell)$ is not hyperspecial. The group K is now of the form $K^S K_S$, where $K^S = \prod_{\ell \notin S} K_\ell$ is a product of hyperspecial compact open subgroups $K_\ell \subset G(\mathbb{Q}_\ell)$, and $K_S \subset \prod_{\ell \in S} G(\mathbb{Q}_\ell)$ is also compact open. We now take $K^c = K^{c,S} K_S^c$ where $K^{c,S} = \prod_{\ell \notin S} K_\ell^c$ with $K_\ell^c \subset G^c(\mathbb{Q}_\ell)$ is the unique hyperspecial subgroup containing the image of K_ℓ , and K_S^c is the image of K_S in $\prod_{\ell \in S} G^c(\mathbb{Q}_\ell)$. In particular, we see that $K/Z(G)_K^-$ maps into K^c , and so we obtain a canonical pro-étale K^c -torsor over Sh_K , which we denote by $\mathbf{Et}_K$. For every prime ℓ , if K_ℓ^c is the projection of K^c into $G^c(\mathbb{Q}_\ell)$, then we will write $\mathbf{Et}_{K,\ell}$ for the K_ℓ^c -local system obtained from $\mathbf{Et}_K$.

ctoriality

Lemma 6.1.7. If $f: (G_1, X_1) \rightarrow (G_2, X_2)$ is a map of Shimura data, then $f(Z(G_1)_{ac}) \subseteq Z(G_2)_{ac}$, and thus f induces a map $f^c: G_1^c \rightarrow G_2^c$.

Proof. See [39, Lemma 3.8]. $\square$

alizations

Remark 6.1.8 (Functoriality of étale realizations). In the situation of the lemma above, suppose that we have compact open subgroups $K_i \subset G_i(\mathbb{A}_f)$ and $K_i^c \subset G_i^c(\mathbb{A}_f)$ for $i = 1, 2$ with $f(K_1) \subset K_2$, $f^c(K_1^c) \subset f^c(K_2^c)$, and with K_i mapping into K_i^c under $G_i(\mathbb{A}_f) \rightarrow G_i^c(\mathbb{A}_f)$. Then by the theory of canonical models of Shimura varieties, we obtain a map

$$\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{K_2} \otimes_{E_2} E_1$$

of stacks over E_1 . Here, E_i is the reflex field for (G_i, X_i) . Then the construction in Construction 6.1.6, combined with Lemma 6.1.7, shows that there is a canonical K_1^c -equivariant map

$$\mathbf{Et}_{K_1} \rightarrow \mathbf{Et}_{K_2} |_{\mathrm{Sh}_{K_1}}.$$

_varieties

6.2. The de Rham realization on Shimura varieties. We recall here some properties of the de Rham realization on Shimura varieties. We will maintain the notation and hypotheses stemming from Definition 6.3.1.

Construction 6.2.1. By definition of the reflex field, the geometric conjugacy class $[\mu]^{125}$ in G of the Shimura cocharacters

$$\mu_x: \mathbb{G}_{m,\mathbb{C}} \xrightarrow{z \mapsto (z,1)} \mathbb{G}_{m,\mathbb{C}} \times \mathbb{G}_{m,\mathbb{C}} \xleftarrow[\simeq]{(az, a\bar{z}) \mapsto a \otimes z} \mathbb{S}_{\mathbb{C}} \xrightarrow{h_x} G_{\mathbb{C}}$$

is defined over $E \subset \mathbb{C}$. This implies the following: There exists a canonical projective homogeneous variety $\mathrm{Gr}_{[-\mu]}$ over E parameterizing filtrations on representations¹³⁶ of G^{26} that are étale locally split by a cocharacter of G in the geometric conjugacy class $[-\mu]$.

Remark 6.2.2 (The de Rham realization). As explained in [21, §5.2], there is a canonical functor from $\mathrm{Rep}_{\mathbb{Q}}(G^c)$ to the category of variations of $\mathbb{Q}$ -Hodge structures over the complex analytic space $\mathrm{Sh}_{K,\mathbb{C}}^{\mathrm{an}}$: Given $V \in \mathrm{Rep}_{\mathbb{Q}}(G^c)$, the fiber of the associated variation of $\mathbb{Q}$ -Hodge structure at any lift $(x, g) \in X \times G(\mathbb{A}_f)$ is canonically isomorphic to the Hodge structure on V arising from the Deligne cocharacter $h_x: \mathbb{S} \rightarrow G_{\mathbb{R}}$ at x .²⁷ Moreover, the underlying

²⁶More precisely, filtrations on the canonical fiber functor $\mathrm{Rep}_E(G) \rightarrow \mathrm{Vect}_E$.

²⁷Here, we follow Deligne's convention from [20] that $h_x(z)$ acts on the (p, q) -part of the Hodge decomposition via $z^{-p}\bar{z}^{-q}$.

[12] Alex: M
is just a new c
tion, but I th
were denoting
Or was that o
the rational co
class?
[5] Keerthi:
to decide whic
preferable. W
you think?
[13] Alex: In
note that follo
we might cite
[6] Keerthi:
citing for this
should actual
Saavedra-Riva
sis, but I'd ra
let it be.

filtered vector bundle with integrable connection for each object in the image of the functor is algebraic. That is, with every $V \in \text{Rep}_{\mathbb{Q}}(G^c)$, we can associate a filtered vector bundle $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,\mathbb{C}}(V)$ over $\text{Sh}_{K,\mathbb{C}}$, equipped with an integrable connection satisfying Griffiths transversality. This functorial association yields in particular a map $\text{Sh}_{K,\mathbb{C}} \rightarrow \text{Gr}_{-\mu,\mathbb{C}}/G_{\mathbb{C}}$ classifying a filtered G -torsor $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,\mathbb{C}}$ over $\text{Sh}_{K,\mathbb{C}}$, and this filtered G -torsor is equipped with an integrable connection satisfying Griffiths transversality; compare with Remark 3.4.3. By [70, Theorems 4.3 and 5.1], this descends canonically to a filtered G -torsor $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,\mathbb{Q}}$ over Sh_K , and the integrable connection also descends. Therefore, for any $V \in \text{Rep}_{\mathbb{Q}}(G^c)$, we obtain a canonical filtered vector bundle $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_K(V)$ over Sh_K with an integrable connection satisfying Griffiths transversality.

Remark 6.2.3 (The Kodaira–Spencer map). Let $\mathbb{T}_{\text{Sh}_K/E}$ be the tangent bundle for Sh_K . By Griffiths transversality, for any $V \in \text{Rep}_{\mathbb{Q}}(G^c)$, we obtain a map of vector bundles

$$\mathbb{T}_{\text{Sh}_K/E} \rightarrow \oplus_i \underline{\text{Hom}} \left(\text{gr}_{\text{Hdg}}^i \mathbf{dR}_K(V), \text{gr}_{\text{Hdg}}^{i-1} \mathbf{dR}_K(V) \right).$$

This arises from a single map $\mathbb{T}_{\text{Sh}_K/E} \rightarrow \mathbf{dR}_K(\mathfrak{g}_{\mathbb{Q}})/\text{Fil}^0 \mathbf{dR}_K(\mathfrak{g}_{\mathbb{Q}})$, where $\mathfrak{g}$ is the Lie algebra of $\mathcal{G}_{(p)}$ equipped with the adjoint action, and we refer to this latter map as the **Kodaira–Spencer map** for Sh_K . By the construction of the Shimura variety, this map is an isomorphism. See [20, Proposition 1.1.14] for a discussion of this over $\mathbb{C}$.

Remark 6.2.4 (de Rham-ness of the p -adic étale realization). Fix a place $v|p$ of E , and let $\text{Sh}_{K,E_v}^{\text{an}}$ be the rigid analytification of Sh_K over E_v . Any $V \in \text{Rep}_{\mathbb{Q}}(G^c)$, viewed as a $\mathbb{Q}_p$ -representation of G , can be twisted by $\mathbf{Et}_{K,p}$ to obtain a $\mathbb{Q}_p$ -local system $\mathbf{Et}_{K,p}(V)$ over Sh_K . In [58], Liu and Zhu showed that the restriction over $\text{Sh}_{K,E_v}^{\text{an}}$ of $\mathbf{Et}_{K,p}(V)$ is *de Rham*. More precisely, they showed that there is a tensor-functorially associated filtered vector bundle $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,v}^{\text{LZ},\text{an}}(V)$ over $\text{Sh}_{K,E_v}^{\text{an}}$ equipped with an integrable connection satisfying Griffiths transversality, such that $\mathbf{Et}_{K,p}(V)$ is *associated* with it in the sense of [82, Definition 8.3]. In particular, for every classical point x of $\text{Sh}_{K,E_v}^{\text{an}}$, $\mathbf{Et}_{K,p,x}(V)$ corresponds to a de Rham Galois representation, and the fiber of $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,v}^{\text{LZ},\text{an}}(V)$ at x is canonically isomorphic to the image of $\mathbf{Et}_{K,p,x}(V)$ under Fontaine’s D_{dR} functor.

Remark 6.2.5 (Work of Diao–Lan–Liu–Zhu). In [21], we find the following improvements to the results of [58]:

- First, using toroidal compactifications, it is shown that $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,v}^{\text{LZ},\text{an}}(V)$ admits a canonical algebraization $\text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,v}^{\text{LZ}}(V)$ over Sh_{K,E_v} compatible with tensor structures; see the discussion on pp. 536–537 of *loc. cit.*
- Next, the authors show in Theorem 5.3.1 of *loc. cit.* that the tensor functors

$$\begin{aligned} V &\mapsto \text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_{K,v}^{\text{LZ}}(V) \\ V &\mapsto \text{Fil}_{\text{Hdg}}^{\bullet} \mathbf{dR}_K(V)|_{\text{Sh}_{K,E_v}} \end{aligned}$$

are canonically isomorphic. Here, we view both as functors from $\text{Rep}_{\mathbb{Q}}(G^c)$ to filtered vector bundles over Sh_{K,E_v} with integrable connection.

6.3. Unramified tuples. From here on, we will fix a prime p .

Definition 6.3.1 (Unramified tuples). A p -**unramified Shimura tuple** or simply **unramified tuple** (since p will remain fixed), is a tuple $(G, \mathcal{G}, X, K)$ where (G, X) is a Shimura datum, $\mathcal{G}$ is a reductive model for G over $\mathbb{Z}_p$, and $K \subset G(\mathbb{A}_f)$ is a compact open subgroup of the form $K^p K_p$ where $K_p = \mathcal{G}(\mathbb{Z}_p)$. Such a tuple is **neat** if K is neat.

Setup 6.3.2. Fix an unramified tuple $(G, \mathcal{G}, X, K)$. This yields a reductive model $\mathcal{G}_{(p)}$ for G over $\mathbb{Z}_{(p)}$. We will write $\mathcal{G}_{(p)}^c$ for the induced reductive model for G^c and $\mathcal{G}^c$ for its base-change over $\mathbb{Z}_p$. Concretely, $\mathcal{G}_{(p)}^c$ is obtained by taking the quotient of $\mathcal{G}_{(p)}$ by the Zariski closure of $Z(G)_{ac}$.

Remark 6.3.3. If $(G, \mathcal{G}, X, K)$ is an unramified tuple, the compact open $K^c \subset G^c(\mathbb{A}_f)$ as in Construction 6.1.6 will always be of the form $K^c = K_p^c K^{c,p}$ where $K_p^c \stackrel{\text{defn}}{=} \mathcal{G}^c(\mathbb{Z}_p) \subset G^c(\mathbb{Q}_p)$.

ied_tuples

Definition 6.3.4 (Maps between unramified tuples). A map $(G_1, \mathcal{G}_1, X_1, K_1) \rightarrow (G_2, \mathcal{G}_2, X_2, K_2)$ between unramified tuples is the data of a map of $\mathbb{Q}$ -groups $f: G_1 \rightarrow G_2$ and an element $g \in G_2(\mathbb{A}_f^p)$ with the following properties:

- $f_{\mathbb{Q}_p}$ extends to a map of $\mathbb{Z}_p$ -group schemes $\mathcal{G}_1 \rightarrow \mathcal{G}_2$;
- f induces a map of Shimura data $(G_1, X_1) \rightarrow (G_2, X_2)$;
- We have $gf(K_1^p)g^{-1} \subset K_2^p$.

rom_tuples

Remark 6.3.5. Let $E_2 \subset E_1 \subset \mathbb{C}$ be the reflex fields of (G_2, X_2) and (G_1, X_1) , respectively. There is a unique map of Shimura varieties

$$\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{K_2} \otimes_{E_2} E_1$$

whose evaluation on $\mathbb{C}$ -points is given by

$$G_1(\mathbb{Q}) \backslash X_1 \times G_1(\mathbb{A}_f) / K_1 \xrightarrow{[(x, h)] \mapsto [(f(x), f(h)g^{-1})]} G_2(\mathbb{Q}) \backslash X_2 \times G_2(\mathbb{A}_f) / K_2.$$

It now follows from Remark 6.1.8 that we have a canonical K_1^c -equivariant map

$$\mathbf{Et}_{K_1} \rightarrow \mathbf{Et}_{K_2|_{\mathrm{Sh}_{K_1}}}.$$

This requires the additional observation that the map of Shimura varieties factors as

$$\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{g^{-1}K_2g} \otimes_{E_2} E_1 \xrightarrow[\simeq]{\eta_g} \mathrm{Sh}_{K_2} \otimes_{E_2} E_1$$

where the second map is the isomorphism of Shimura varieties whose evaluation on $\mathbb{C}$ -points is

$$G_2(\mathbb{Q}) \backslash X_1 \times G_1(\mathbb{A}_f) / g^{-1}K_2g \xrightarrow{[(x, h)] \mapsto [(x, hg^{-1})]} G_2(\mathbb{Q}) \backslash X_2 \times G_2(\mathbb{A}_f) / K_2.$$

Moreover, we have a canonical isomorphism of $K_2^c \simeq g^{-1}K_2^cg$ -local systems

$$\eta_g^* \mathbf{Et}_{K_2} \xrightarrow[\simeq]{} \mathbf{Et}_{g^{-1}K_2g}$$

over $\mathrm{Sh}_{g^{-1}K_2g}$.

to-p_hecke

Remark 6.3.6 (Prime-to- p Hecke correspondences and connected components). There is an action of $G(\mathbb{A}_f^p)$ on the underlying cofiltered system of E -schemes $\{\mathrm{Sh}_{K', pK_p}\}$: Given $g \in G(\mathbb{A}_f^p)$, we obtain a canonical isomorphism $\mathrm{Sh}_{K', pK_p} \xrightarrow[\simeq]{} \mathrm{Sh}_{gK', pg^{-1}K_p}$ induced over $\mathbb{C}$ by the endomorphism of $X \times G(\mathbb{A}_f)$ given by $(x, h) \mapsto (x, hg^{-1})$. This gives an action of $G(\mathbb{A}_f^p)$ on Sh_{K_p} . By [20, p. 2.1.3] and [47, Lemma (2.2.6)], the induced action on the connected components of $\mathrm{Sh}_{K_p, \overline{\mathbb{Q}}}$ factors through a simply transitive action of the quotient $G(\mathbb{A}_f^p) / (\bigcap_{K', p} \mathcal{G}_{(p)}(\mathbb{Z}_{(p)})_+ K'^{', p})$, which is in fact an abelian group.

l_aperture

6.4. Consequences of having a universal aperture.

Setup 6.4.1. Fix an unramified tuple $(G, \mathcal{G}, X, K)$, and a place $v|p$ of the reflex field E . Set $\hat{\mathcal{O}} = \mathcal{O}_{E_v}$, and let k be the residue field of $\hat{\mathcal{O}}$. If we view $\{\mu\}$ as an E_v -rational conjugacy class, then we in fact have a representative $\mu_v: \mathbb{G}_{m, \mathcal{O}} \rightarrow \mathcal{G}_{\hat{\mathcal{O}}}$ for $\{\mu\}$: The existence of a representative over E_v follows from the quasi-splitness of G_{E_v} [54, Lemma 1.1.3], while that over $\hat{\mathcal{O}}$ now follows from [47, Proposition 1.1.4]. We will view μ_v as a cocharacter of $\mathcal{G}_{\hat{\mathcal{O}}}^c$. Recall from §4.1 the pro-algebraic stack $\mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}$ over $\mathcal{O}_{E, (v)}$ with generic fiber $\mathrm{Loc}_{G^c(\mathbb{Z}_p)}$ and v -adic completion $\mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v}$.

rmal_model

Definition 6.4.2. Suppose that we have a map of schemes $U \rightarrow \mathrm{Sh}_K$, and suppose that $\mathcal{U}$ is a flat integral model for U over $\mathcal{O}_{E, (v)}$: By this, we mean that $\mathcal{U}$ is a flat separated algebraic space over $\mathcal{O}_{E, (v)}$ equipped with an identification $\mathcal{U} \otimes_{\mathcal{O}_{E, (v)}} E \xrightarrow[\simeq]{} U$. We will say that $\mathcal{U}$ is **apertile** if there is an extension $\mathcal{U} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}$ of the classifying map $U \rightarrow \mathrm{Loc}_{G^c(\mathbb{Z}_p)}$ for $\mathbf{Et}_{K, p}|_U$.

m:apertile

Remark 6.4.3. By Theorem 4.1.5, such an extension is unique up to unique isomorphism if $\mathcal{U}$ is η -normal. In particular, in this case, being apertile is a *property* of the integral model $\mathcal{U}$.

[14] **Alex:** No bit deal either but I feel like the phrase ‘pro-crystalline’ could be clarificatory and points in the

wise_lifts

Remark 6.4.4 (Type of $\mathbf{Et}_{K,p}$). The existence of the aperture implies in particular that the restriction of $\mathbf{Et}_{K,p}$ over $\hat{\mathcal{U}}_\eta$ is a crystalline $\mathcal{G}(\mathbb{Z}_p)$ -local system of type $-\mu_v$ at every classical point.¹⁴ Conversely, if we knew *a priori* that $\mathbf{Et}_{K,p}$ is crystalline at every classical point of $\hat{\mathcal{U}}_\eta$, then, by Remarks 6.2.4 and 6.2.5, we see that its type at any such point has to be $-\mu_v$.

al_de_rham

Construction 6.4.5 (The integral de Rham realization). Consider the composition

e_rham_map

$$(6.4.5.1) \quad \hat{\mathcal{U}} \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v} \rightarrow \mathrm{Gr}_{-\mu_v} / \mathcal{G}_{\mathcal{O}_{E_v}}$$

where the second map is as in Remark 3.4.3. This map classifies a filtered $\mathcal{G}$ -bundle over $\hat{\mathcal{U}}$, whose restriction over $\hat{\mathcal{U}}_\eta$ is canonically isomorphic to the restriction of $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet \mathbf{dR}_{K,\mathbb{Q}}$: One checks this using Remarks 6.2.5 and 3.8.3. This tells us that there exists a map

$$\mathcal{U} \rightarrow \mathrm{Gr}_{-\mu_v} / \mathcal{G}_{\mathcal{O}_{E_v}}$$

whose formal completion is (6.4.5.1) and whose generic fiber classifies $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet \mathbf{dR}_{K,\mathbb{Q}}$.¹⁵ Denote the associated filtered $\mathcal{G}$ -bundle over $\mathcal{S}_K$ by $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet \mathbf{dR}_K$.

[15] **Alex:** M state that this Beauville–Las ing?

braization

Remark 6.4.6 (The integral Kodaira–Spencer map). Suppose that $\mathcal{U}$ is *smooth* over $\mathcal{O}_{E,(v)}$. Then the filtered $\mathcal{G}$ -bundle from the previous remark gives us a Kodaira–Spencer map

pencer_map

$$(6.4.6.1) \quad \mathbb{T}_{\mathcal{U}/\mathcal{O}_{E,(v)}} \rightarrow \mathbf{dR}_K(\mathfrak{g}) / \mathrm{Fil}^0 \mathbf{dR}_K(\mathfrak{g})$$

whose restriction over the generic fiber is described in Remark 6.2.3 and whose v -adic completion is described in Remark 5.2.2.

_condition

Lemma 6.4.7 (A versality condition). *The following are equivalent:*

- (1) *The map $\hat{\mathcal{U}} \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}$ is formally étale*
- (2) *$\mathcal{U}$ is smooth over $\mathcal{O}_{E,(v)}$ and the map (6.4.6.1) is an isomorphism.*

Proof. Immediate from Remark 5.2.2. □

cal_models

6.5. Integral canonical models. *For the rest of this section and the next, we will assume that the level K^p is neat. We will return to the general situation in §8.4 below. We can now give the definition of integral canonical models and formulate their mapping properties.*

defn:ICM

Definition 6.5.1 (Integral canonical models). An apertile integral model $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_{E,(v)}$ is a **crystalline integral canonical model**—or **crystalline ICM** for short—if it satisfies the equivalent conditions of Lemma 6.4.7, and if, for any mixed characteristic $(0, p)$ complete discrete valuation field F over $\mathcal{O}_{E_v}$ with perfect residue field, the following are equivalent for $x \in \mathrm{Sh}_K(F)$:

- (1) $x \in \mathcal{S}_K(\mathcal{O}_F)$;
- (2) $\mathbf{Et}_{K,p,x}$ is crystalline;
- (3) $\mathbf{Et}_{K,p,x}$ is potentially crystalline.

It is an **étale integral canonical model**—or **étale ICM** for short—if, instead, we replace the last condition with: For any mixed characteristic $(0, p)$ discrete valuation field F over $\mathcal{O}_{E_v}$, the following are equivalent for $x \in \mathrm{Sh}_K(F)$:

- (1) $x \in \mathcal{S}_K(\mathcal{O}_F)$;
- (2) $\mathbf{Et}_{K,\ell,x}$ is unramified for all $\ell \neq p$;
- (3) $\mathbf{Et}_{K,\ell,x}$ is potentially unramified for all $\ell \neq p$.

If the context is agnostic to the precise type of integral canonical model involved, we simply call $\mathcal{S}_K$ an **ICM** for Sh_K . In either case, we call the morphism $\varpi: \mathcal{S}_K \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}$ the *syntomic realization*.

n_ell-adic

Remark 6.5.2 (Extension of ℓ -adic local systems). The pointwise condition in the definition of an étale ICM implies that, for all $\ell \neq p$, $\mathbf{Et}_{K,\ell}$ extends to a local system over $\mathcal{S}_K$: One applies the unramifiedness to the local ring at any codimension 1 point and then uses purity.

Let us restate Theorem A in the following form, which is now immediate from Proposition 4.3.1.

g_property

Theorem 6.5.3 (Mapping property of ICMs). *Let $\mathcal{Y}$ be an excellent, separated algebraic space flat over $\mathcal{O}_{E,(v)}$ with generic fiber Y . Suppose that $\hat{\mathcal{Y}}$ is η -normal in the sense of [1, Appendix A].*

- (1) *If $\mathcal{S}_K$ is a crystalline ICM for Sh_K over $\mathcal{O}_{E,(v)}$, then giving a map $\tilde{f}: \mathcal{Y} \rightarrow \mathcal{S}_K$ is equivalent to giving maps $\eta: \mathcal{Y} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}$ and $f: Y \rightarrow \mathrm{Sh}_K$ such that the generic fiber $Y \rightarrow \mathrm{Loc}_{\mathcal{G}^c(\mathbb{Z}_p)}$ of η lifts f along the map $\mathrm{Sh}_K \rightarrow \mathrm{Loc}_{\mathcal{G}^c(\mathbb{Z}_p)}$ classifying $\mathbf{Et}_{K,p}$.*
- (2) *If $\mathcal{S}_K$ is an étale ICM, then giving a map $\tilde{f}: \mathcal{Y} \rightarrow \mathcal{S}_K$ is equivalent to giving maps $\eta: \mathcal{Y} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}$ and $f: Y \rightarrow \mathrm{Sh}_K$ such that:*
 - (a) *The generic fiber $Y \rightarrow \mathrm{Loc}_{\mathcal{G}^c(\mathbb{Z}_p)}$ of η lifts f along the map $\mathrm{Sh}_K \rightarrow \mathrm{Loc}_{\mathcal{G}^c(\mathbb{Z}_p)}$ classifying $\mathbf{Et}_{K,p}$;*
 - (b) *For all $\ell \neq p$, $\mathbf{Et}_{K,\ell|Y}$ extends to a K_{ℓ}^c -local system over $\mathcal{Y}$.*

In both cases, we have canonical isomorphisms of classifying maps $\varpi \circ \tilde{f} \simeq \eta$.

uniqueness

Corollary 6.5.4 (Uniqueness). *An ICM $\mathcal{S}_K$ for Sh_K is unique up to unique isomorphism.*

functoriality

Corollary 6.5.5 (Functoriality). *Suppose that we have a map of tuples as in Definition 6.3.4, and for $i = 1, 2$ ICMs $\mathcal{S}_{K_i}$ for Sh_{K_i} over $\mathcal{O}_{E_i,(v_i)}$. Then the map $\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{K_2} \otimes_{E_2} E_1$ extends to a map $\mathcal{S}_{K_1} \rightarrow \mathcal{S}_{K_2} \otimes_{\mathcal{O}_{E_2,(v_2)}} \mathcal{O}_{E_1,(v_1)}$.*

Proof. The only thing to observe here is that, under the map of Shimura varieties, for every prime ℓ , the restriction of $\mathbf{Et}_{K_2,\ell}$ to Sh_{K_1} is canonically obtained from $\mathbf{Et}_{K_1,\ell}$ via pushforward along the map $K_{1,\ell}^c \rightarrow K_{2,\ell}^c$ induced by

$$G_1^c(\mathbb{Q}_{\ell}) \xrightarrow{f} G_2^c(\mathbb{Q}_{\ell}) \xrightarrow{h \mapsto g \ell h g^{-1}} G_2^c(\mathbb{Q}_{\ell}).$$

This was observed in Remark 6.3.5. □

-to-p_part

Corollary 6.5.6 (Prime-to- p Hecke correspondences). *Suppose that $K'^{\cdot p} \subset K^p$ is a compact open subgroup and set $K' = K_p K'^{\cdot p}$.*

- (1) *If Sh_K admits an ICM $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$, then the normalization of $\mathcal{S}_K$ in $\mathrm{Sh}_{K'}$ is an ICM $\mathcal{S}_{K'}$ for $\mathrm{Sh}_{K'}$;*
- (2) *If (1) holds, and $g \in G(\mathbb{A}_f^p)$ is such that $gK'^{\cdot p}g^{-1} \subset K^p$, then the map $i_g: \mathrm{Sh}_{K'} \rightarrow \mathrm{Sh}_K$ determined on complex points by*

$$G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K' \xrightarrow{[(x,h)] \mapsto [(x,hg^{-1})]} G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K,$$

extends to a finite étale map of ICMs $\mathcal{S}_{K'} \rightarrow \mathcal{S}_K$.

Proof. The first assertion follows from the fact that $\mathbf{Et}_{K,\ell}$ extends to a local system over $\mathcal{S}_K$, for all $\ell \neq p$, and the second is immediate from Corollary 6.5.5. □

comparison

Remark 6.5.7 (Comparison with the definition of Milne–Moonen). In references such as [70], [71], and [47], one finds a definition of integral canonical models that involves the inverse limit

$$\mathrm{Sh}_{K_p} = \varprojlim_K \mathrm{Sh}_K,$$

where the limit is over compact opens of the form $K_p K^p$. An integral model $\mathcal{S}_{K_p}$ for Sh_{K_p} has the *extension property* if, for any regular and formally smooth $\mathcal{O}_{E,(v)}$ -scheme S , any map $S_{\eta} \rightarrow \mathrm{Sh}_{K_p}$ of E -schemes extends to a map $S \rightarrow \mathcal{S}_{K_p}$. If $\mathcal{S}_{K_p}$ is itself regular and formally smooth, this property characterizes it as an integral canonical model in the sense of Milne.

Suppose that $(G, \mathcal{G}, X)$ admits étale ICMs over $\mathcal{O}_{E,(v)}$. Then we can take the inverse limit over the resulting inverse system $\{\mathcal{S}_K\}$ of finite étale maps to get a model $\mathcal{S}_{K_p}$ for Sh_{K_p} . We claim that $\mathcal{S}_{K_p}$ satisfies Milne's extension property when $p > 3$.

First, we can assume that $S = \mathrm{Spec} R$ is affine. Now, for any mixed characteristic complete discrete valuation field F over E_v with perfect residue field, and an $x: \mathrm{Spec}(F) \rightarrow \mathrm{Sh}_{K_p}$, the projection of x_K to Sh_K lifts to an $\mathcal{O}_F$ -point of $\mathcal{S}_K$: Indeed, the existence of the lift x shows that, for all $\ell \neq p$, the local system $\mathbf{Et}_{K,\ell,x_K}$ is unramified (in fact, trivial). Therefore, the desired assertion follows from the definition of an étale ICM.

Next, by Popescu desingularization [87, Tag 07GC], we can present R as a filtered colimit of smooth $\mathcal{O}_{E,(v)}$ -algebras implying $\mathrm{Spec} S$ is an inverse limit of smooth affine $\mathcal{O}_{E,(v)}$ -schemes S_i . Given a map $S_{\eta} \rightarrow \mathrm{Sh}_{K_p}$, the

projection onto Sh_K for any finite level K factors through $S_{i,\eta}$ for some i . The previous paragraph shows that the restriction of $\mathbf{Et}_{K,p}$ over $S_{i,\eta}$ is pointwise crystalline, and so Proposition 7.9.2 now tells us that we have an extension $S_i \rightarrow \mathcal{S}_K$, and this implies our claim.

Remark 6.5.8 (Comparison with the definition of Bakker–Shankar–Tsimmerman). Combining Remark 6.5.7 with the discussion in [4, §1.3] tells us that when $p > 3$, étale ICMs also satisfy the canonicity property of Bakker–Shankar–Tsimmerman.

7. INTEGRAL CANONICAL MODELS: CONSTRUCTIONS AND FURTHER PROPERTIES

We will maintain the notation from above. In this section, we will prove our main theorems on the existence of ICMs, as well as prove a few deeper properties of such models.

7.1. CM points and canonical models for tori.

Construction 7.1.1 (CM Shimura varieties). Let T be a torus over $\mathbb{Q}$ equipped with a map $h_T: \mathbb{S} \rightarrow T_{\mathbb{R}}$ and let $\mu_T \in X_*(T)$ be the Shimura cocharacter associated with h . Let $E_T \subset \overline{\mathbb{Q}}$ be the field of definition of μ_T . Then, for any neat compact open subgroup $K_T \subset T(\mathbb{A}_f)$, the tuple $(T, \{h\}, K_T)$ yields a zero-dimensional Shimura variety Sh_{K_T} over E with

$$\mathrm{Sh}_{K_T}(\mathbb{C}) \simeq T(\mathbb{Q}) \backslash T(\mathbb{A}_f) / K.$$

Fix a place $w|p$ of E_T , and let $\mathcal{S}_{K_T}$ be the normalization of $\mathrm{Spec} \mathcal{O}_{E_T, (w)}$ in Sh_{K_T} . We will assume that K_T is of the form $K_{T,p} K_T^p$, where $K_{T,p} \subset T(\mathbb{Q}_p)$ and $K_T^p \subset T(\mathbb{A}_f^p)$ are compact open subgroups.

Remark 7.1.2 (Galois structure of CM Shimura variety). As explained in [20] (see also the discussion in [51, §1.5.3]), the theory of canonical models is controlled by that for CM Shimura varieties. Consider the reflex norm

$$r_{\mu_T}: \mathrm{Res}_{E_T/\mathbb{Q}} \mathbb{G}_m \xrightarrow{\mathrm{Res}_{E_T/\mathbb{Q}} \mu_T} \mathrm{Res}_{E_T/\mathbb{Q}} T \xrightarrow{\mathrm{Nm}_{E_T/\mathbb{Q}}} T.$$

This induces a map

$$(7.1.2.1) \quad E_T^\times \backslash \mathbb{A}_{E_T}^\times \xrightarrow{r_{\mu}(\mathbb{A}_f)} T(\mathbb{Q}) \backslash T(\mathbb{A}) \rightarrow T(\mathbb{Q}) \backslash T(\mathbb{A}) / K_T$$

that factors via the global reciprocity map (with the geometric normalization) through a homomorphism

$$\tau_K(\mu_T): \mathrm{Gal}(E_T^{\mathrm{ab}}/E_T) \rightarrow T(\mathbb{Q}) \backslash T(\mathbb{A}) / K_T$$

giving the target the structure of a $\mathrm{Gal}(E_T^{\mathrm{ab}}/E_T)$ -equivariant finite set. The Shimura variety Sh_{K_T} is now canonically isomorphic to the étale scheme over E_T associated with this datum.

Remark 7.1.3 (Étale realizations for tori). Suppose that we have a finite extension $F/E_{T,w}$ and a point $x \in \mathrm{Sh}_{K_T}(F)$. As discussed in [51, §4.3.13, 5.7], for each prime ℓ , we can describe the Galois representation¹⁶⁷ associated with the local system $\mathbf{Et}_{K_T, \ell, x}$ as follows: The reflex norm r_{μ_T} yields a map

$$(7.1.3.1) \quad E_T^\times \backslash \mathbb{A}_{E_T}^\times \xrightarrow{r_{\mu}(\mathbb{A}_f)} T(\mathbb{Q}) \backslash T(\mathbb{A}) \rightarrow \pi_0(T^c(\mathbb{Q}) \backslash T^c(\mathbb{A})) \rightarrow T^c(\mathbb{Q}) \backslash T^c(\mathbb{A}_f)$$

that factors via the global reciprocity map through a homomorphism

$$\tau(\mu_T): \mathrm{Gal}(E_T^{\mathrm{ab}}/E_T) \rightarrow T^c(\mathbb{Q}) \backslash T^c(\mathbb{A}_f)$$

The composition

$$\mathrm{Gal}(\overline{\mathbb{Q}_p}/F) \rightarrow \mathrm{Gal}(E_T^{\mathrm{ab}}/E_T) \rightarrow T^c(\mathbb{Q}) \backslash T^c(\mathbb{A}_f) \rightarrow T^c(\mathbb{Q}_\ell)$$

factors through $K_{T, \ell}^c$, and the local system associated with it is canonically isomorphic to $\mathbf{Et}_{K_T, \ell, x}$.

Remark 7.1.4 (Crystallinity at p). The description in Remark 7.1.3, combined with [51, Proposition 4.3.14], shows that the $T(\mathbb{Q}_p)$ -local system underlying $\mathbf{Et}_{K_T, p, x}$ is crystalline of type $-\mu_T$.

[16] Alex: As haven't really this terminology maybe it would to clarify you $T^c(\mathbb{Q}_\ell)$ -valued rep

[7] Keerthi: clarified by this sion.

unramified

Remark 7.1.5 (Unramifiedness at $\ell \neq p$). For $\ell \neq p$, the description from Remark 7.1.3 shows that the local system $\mathbf{Et}_{K_T, \ell, x}$ is unramified. Indeed, by local reciprocity, it is enough to know that the restriction of the map (7.1.3.1) along

$$\mathcal{O}_F^\times \rightarrow \mathcal{O}_{E_T, w}^\times \rightarrow E_T^\times \backslash \mathbb{A}_{E_T}^\times$$

is trivial. However, this restriction has image in the subgroup $T^c(\mathbb{Q}) \backslash T^c(\mathbb{Q})K_{T, p}^c$, and the projection of this subgroup onto $T^c(\mathbb{A}_f^p)$ has trivial intersection with the image $K_T^{p, c}$ of K_T^p . This is because, by the discreteness of $T^c(\mathbb{Q})$ in $T^c(\mathbb{A}_f)$ and the neatness of K_T —and therefore that of its image $K_T^c \subset T^c(\mathbb{A}_f)$ —the intersection $T^c(\mathbb{Q}) \cap K_T^c$ is trivial.

Integral canonical models always exist for CM Shimura varieties:

s_for_tori

Proposition 7.1.6 (ICMs for tori). *Suppose that $T_{\mathbb{Q}_p}$ extends to a torus $\mathcal{T}$ over $\mathbb{Z}_p$ ¹⁷⁸ such that $K_{T, p} = \mathcal{T}(\mathbb{Z}_p)$. Then $\mathcal{S}_{K_T}$ is both a crystalline ICM and an étale ICM for Sh_{K_T} .*

Proof. In this case, one finds using Remark 7.1.2 that $\mathcal{S}_{K_T}$ is a finite étale scheme over $\mathcal{O}_{E_T, (w)}$. Moreover, the stacks $\mathrm{BT}_n^{\mathcal{T}, -\mu_T}$ are étale over $\mathcal{O}_{E_T, w}$ and are in fact non-canonically isomorphic to the $\mathcal{O}_{E_T, w}$ -stack $\mathrm{Loc}_{\mathcal{T}(\mathbb{Z}/p^n\mathbb{Z})}$ of $\mathcal{T}(\mathbb{Z}/p^n\mathbb{Z})$ -local systems; see [26, Proposition 10.4.1]. Therefore, any map $\widehat{\mathcal{S}}_{K_T, w} \rightarrow \mathrm{BT}_\infty^{\mathcal{T}, -\mu_T}$ will be formally étale. Proposition 5.1.3 now reduces us to checking that, for every finite extension $L/E_{T, w}$ and every $x \in \mathcal{S}_K(\mathcal{O}_L)$, the following statements hold:

- (1) $\mathbf{Et}_{K_T, p, x}$ belongs to $\mathrm{Loc}_{\mathcal{T}(\mathbb{Z}_p)}^{\mathrm{crys}, -\mu_T}(L)$;
- (2) For $\ell \neq p$, $\mathbf{Et}_{K_T, \ell, x}$ is unramified.

These follow from Remarks 7.1.4 and 7.1.5. □

cm_points

Construction 7.1.7. Suppose that we have an unramified tuple $(G, \mathcal{G}, X, K)$ and an ICM $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_{E, (v)}$ for a place $v|p$. As a special case of Construction 7.1.1, suppose that we have a torus T over $\mathbb{Q}$ and a closed immersion $i : T \rightarrow G$ over $\mathbb{Q}$ such that, for some $x \in X$, the map $h_x : \mathbb{S} \rightarrow G_{\mathbb{R}}$ factors through $i_{\mathbb{R}}$ via a map $h_T : \mathbb{S} \rightarrow T_{\mathbb{R}}$. Given $g \in G(\mathbb{A}_f^p)$, set $K_{T, g} = i^{-1}(gKg^{-1}) \subset T(\mathbb{A}_f)$. Then, for any place $w|p$ of E_T above v , we obtain an integral model $\mathcal{S}_{K_{T, g}, (w)}$ for $\mathrm{Sh}_{K_{T, g}}$ over $\mathcal{O}_{E_T, (w)}$ as the normalization of $\mathrm{Spec} \mathcal{O}_{E, (v)}$ in $\mathrm{Sh}_{K_{T, g}}$.

cm_points

Lemma 7.1.8 (Extension of CM points). *The map $\mathrm{Sh}_{K_{T, g}} \rightarrow \mathrm{Sh}_K \otimes_E E_T$ of canonical models given on $\mathbb{C}$ -points by*

$$T(\mathbb{Q}) \backslash T(\mathbb{A}_f) / K_{T, g} \xrightarrow{[t] \mapsto [(x_T, i(t)g)]} G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K$$

extends canonically to a map of integral models $\mathcal{S}_{K_{T, g}, (w)} \rightarrow \mathcal{S}_K \otimes_{\mathcal{O}_{E, (v)}} \mathcal{O}_{E_T, (w)}$.

Proof. To alleviate notation, set $K_T = K_{T, g}$. Given Theorem 6.5.3, it is enough to know the following:

- (1) The restriction of $\mathbf{Et}_{K, p}$ over Sh_{K_T} lifts to a map

$$\mathcal{S}_{K_T, (w)} \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}.$$

- (2) For all $\ell \neq p$, the restriction of $\mathbf{Et}_{K, \ell}$ over Sh_{K_T} extends to a local system over $\mathcal{S}_{K_T}$.

Since $\mathcal{S}_{K_T, (w)}$ is a product of spectra of localizations of rings of integers of number fields, we reduce to knowing (by Lemma 5.1.11 in the p -adic case) that, for every complete local ring $\hat{\mathcal{O}}$ of $\mathcal{S}_{K_T, (w)}$, the restriction of $\mathbf{Et}_{K, p}$ (resp. $\mathbf{Et}_{K, \ell}$) over $\hat{\mathcal{O}}[1/p]$ lifts to $\mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}(\hat{\mathcal{O}})$ (resp. extends to a local system over $\hat{\mathcal{O}}$).

By Proposition 5.1.3, the p -adic condition is equivalent to checking that the restriction of $\mathbf{Et}_{K, p}$ over $\hat{\mathcal{O}}[1/p]$ is crystalline of type $-\mu_v$, which is immediate from Remark 7.1.4. The unramifiedness of the ℓ -adic realizations is similarly immediate from Remark 7.1.5.¹⁸ □

Definition 7.1.9 (CM points). For any $\mathcal{O}_{E, (v)}$ -algebra R ⁹, a point $x \in \mathcal{S}_K(R)$ is a **CM point** if there exist a finite extension F/E , a place $w'|p$ of F above v , a tuple (T, i, h_T, g) as in Construction 7.1.7, a place $w|p$ of E_T , a

[17] Alex: M
it's patently c
to everyone re
but should we
“(i.e., T splits
unramified ex

[8] Keerthi:
overkill to me

[18] Alex: I t
again it would
ful to cite the
which specifie
relationship b
pullbacks of e
realizations un
morphism of S
varieties.

[9] Keerthi:
if this definiti
unnecessarily

faithfully flat map $R \rightarrow R'$, and a commuting diagram of schemes

$$\begin{array}{ccccc} \mathrm{Spec} R' & \longrightarrow & \mathrm{Spec} \mathcal{O}_{F,(w')} & \longrightarrow & \mathcal{S}_{K_{T,g},(w)} \\ & \searrow & \downarrow & & \downarrow \\ & & \mathrm{Spec} R & \xrightarrow{x} & \mathcal{S}_K \end{array}$$

where the right vertical arrow is the obtained from Lemma 7.1.8.¹⁹

7.2. Spaces of isogenies. The purpose of this section is to recall how Hecke correspondences can be used to define spaces of isogenies between points on an ICM $\mathcal{S}_K$.

Proposition 7.2.1 (Descent along prime-to- p étale covers). *Suppose that we have a map of unramified tuples $(G, \mathcal{G}, X, K_1) \rightarrow (G, \mathcal{G}, X, K_2)$ that is the identity on the underlying Shimura data. If Sh_{K_1} admits an ICM $\mathcal{S}_{K_1}$ over $\mathcal{O}_{E,(v)}$, then Sh_{K_2} admits an ICM $\mathcal{S}_{K_2}$, and the map $\mathcal{S}_{K_1} \rightarrow \mathcal{S}_{K_2}$ of $\mathcal{O}_{E,(v)}$ -algebraic spaces is finite étale.*

Proof. Let $\Delta = K_2/K_1 Z(G)_{K_2}$ be the Galois group of the cover $\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{K_2}$. By Theorem 6.5.3, the action of Δ on Sh_{K_1} extends to an action on $\mathcal{S}_{K_1}$. If we know that this extended action is free, then we obtain a finite étale cover

$$\mathcal{S}_{K_1} \rightarrow \mathcal{S}_{K_2} \stackrel{\mathrm{defn}}{=} \mathcal{S}_{K_1}/\Delta$$

giving us an integral model $\mathcal{S}_{K_2}$ for Sh_{K_2} .

Assuming this freeness, we claim that $\mathcal{S}_{K_2}$ is an ICM for Sh_{K_2} . The only possibly non-trivial fact to be verified is that it is apertile, which follows from Corollary 4.1.6. □

7.3. Reduction of structure group. We can now prove Theorem D, whose statement we recall:

Theorem 7.3.1 (Reduction of structure group). *Suppose that $(G, \mathcal{G}, X, K) \rightarrow (G^\sharp, \mathcal{G}^\sharp, X^\sharp, K^\sharp)$ is a map of neat unramified tuples such that $\mathcal{G} \rightarrow \mathcal{G}^\sharp$ is a closed immersion. Let $E^\sharp$ be the reflex field for $(G^\sharp, X^\sharp)$, and let $v|p$ be a place for E lying above a place $v^\sharp$ for $E^\sharp$. Suppose that $\mathrm{Sh}_{K^\sharp}$ admits an ICM $\mathcal{S}_{K^\sharp}$ over $\mathcal{O}_{E^\sharp,(v^\sharp)}$. Then the normalization of $\mathcal{O}_{E,(v)} \otimes_{\mathcal{O}_{E^\sharp,(v^\sharp)}} \mathcal{S}_{K^\sharp}$ in Sh_K gives an ICM for Sh_K over $\mathcal{O}_{E,(v)}$.*

Proof. Suppose first that $\mathrm{Sh}_K \rightarrow \mathrm{Sh}_{K^\sharp}$ is a closed immersion. We then have the following diagram

$$\begin{array}{ccc} \mathrm{Sh}_K & \longrightarrow & \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)} = \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}[1/p] \\ \downarrow & & \downarrow \\ \mathcal{S}_K & \overset{? \exists}{\dashrightarrow} & \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v, \mathrm{alg}} \\ \downarrow & & \downarrow \\ \mathcal{S}_{K^\sharp} & \longrightarrow & \mathrm{BT}_\infty^{\mathcal{G}^\sharp, -\mu_{v^\sharp}, \mathrm{alg}} \end{array}$$

The bottom left vertical arrow is the normalization of a closed immersion by construction, and the bottom horizontal arrow is formally étale. Therefore, to fill in the middle dotted arrow and also know that it is formally étale, we can appeal to Proposition 5.3.9, whose hypotheses are verified by Remark 6.4.4.

To finish, we need to know that $\mathcal{S}_K$ satisfies the pointwise conditions for ICMs: This just amounts to the observation that by construction the equality

$$\mathcal{S}_K(\mathcal{O}_F) = \mathrm{Sh}_K(F) \cap \mathcal{S}_{K^\sharp}(\mathcal{O}_F) \subset \mathrm{Sh}_{K^\sharp}(F)$$

holds for any discrete valuation field F over E .

[19] Alex: To might make it to just say the faithfully flat factorizes through of the maps. . not spell this explicitly.

Now, on to the general case: By [47, Lemma (2.1.2)], we can find compact open subgroups $K_1^p \subset K^p$ and $K_1^{\sharp,p} \subset K^{\sharp,p}$ such that, with $K_1 = K_1^p K_p$ and $K_1^{\sharp} = K_1^{\sharp,p} K_p^{\sharp}$, we have $K_1 = K_1^{\sharp} \cap G(\mathbb{A}_f^p)$ and the map $\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{K_1^{\sharp}}$ is a closed immersion. By the first paragraph, and Proposition 7.2.1, it is enough to know that $\mathrm{Sh}_{K_1^{\sharp}}$ admits an ICM whenever $\mathrm{Sh}_{K^{\sharp}}$ does so, and this is (1) of Corollary 6.5.6. $\square$

7.4. The global μ -ordinary locus and canonical lifts. Here, we will find some results about the μ -ordinary loci in ICMs. We will fix an unramified tuple $(G, \mathcal{G}, X, K)$ with an ICM $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$.

Definition 7.4.1 (The global μ -ordinary locus). Let $\mathcal{S}_K$ be an ICM for Sh_K over $\mathcal{O}_{E,(v)}$ with special fiber $\mathcal{S}_{K,k(v)}$. The μ -ordinary locus $\mathcal{S}_{K,k(v)}^{\mathrm{ord}}$ is the pre-image of the open and dense locus $\mathrm{BT}_1^{\mathcal{G}^c, -\mu_v, \mathrm{ord}} \subset \mathrm{BT}_1^{\mathcal{G}^c, -\mu_v}$ from Definition 3.6.1.

Theorem 7.4.2 (Properties of the μ -ordinary locus). *The following statements hold:*

- (1) *The μ -ordinary locus is open and fiberwise dense in $\mathcal{S}_K$.*
- (2) *For any algebraically closed field κ in characteristic p and $x \in \mathcal{S}_{K,k(v)}^{\mathrm{ord}}(\kappa)$, the deformation space Def_x for $\mathcal{S}_K$ at x admits a canonical structure of a tower*

$$\mathrm{Def}_x = \widehat{U}_{x,n} \rightarrow \cdots \rightarrow \widehat{U}_{x,0} = \mathrm{Spf} W(\kappa)$$

where each map in the tower is represented by a formal p -divisible group;

- (3) *In particular, x admits a **canonical lift** $x^{\mathrm{can}} \in \mathcal{S}_K(W(\kappa))$ corresponding to the successive composition of identity sections of the tower.*
- (4) *If κ is algebraic over $k(v)$ and $\mathcal{S}_K$ is a crystalline ICM, then the canonical lift is a CM point of $\mathcal{S}_K$.*

Proof. The first assertion is clear from the definition and the smoothness of the classifying map $\hat{\mathcal{S}}_K \rightarrow \mathrm{BT}_1^{\mathcal{G}^c, -\mu_v}$. The next two assertions are immediate from the definition of an ICM and the results of §3.6. The rest of this subsection will be devoted to the proof of the last assertion. $\square$

7.5. Ascent for ICMs.

Setup 7.5.1. Suppose that $(G, \mathcal{G}, X, K) \rightarrow (\overline{G}, \overline{\mathcal{G}}, \overline{X}, \overline{K})$ is a map of unramified tuples such that $\mathcal{G} \rightarrow \overline{\mathcal{G}}$ is a surjective map of reductive group schemes with central kernel. Let $\overline{E}$ be the reflex field for $(\overline{G}, \overline{X})$, and let $v|p$ be a place for E lying above a place $\overline{v}$ for $\overline{E}$.

Remark 7.5.2 (CM points and central covers). Suppose²⁰¹⁰ that we have (T, i, h_T) as in Construction 7.1.7 with $\overline{T} \subset \overline{G}$ the image of T , and $\overline{g} \in \overline{G}(\mathbb{A}_f^p)$. Then we can consider the fiber product

$$\mathrm{Sh}_K \times_{\mathrm{Sh}_{\overline{K}}} \mathrm{Sh}_{\overline{K}_{\overline{T}, \overline{g}}} \rightarrow \mathrm{Sh}_{\overline{K}_{\overline{T}, \overline{g}}}.$$

By looking at $\mathbb{C}$ -points, one finds that this is a disjoint union of CM Shimura varieties of the form $\mathrm{Sh}_{K_{T', g'}}$, where $i' : (T', h_{T'}) \hookrightarrow (G, X)$ is a $\overline{G}(\mathbb{Q})$ -conjugate of i .

The next result proves the most non-trivial part of Theorem ??, whose notation we will be using here.

Proposition 7.5.3. *Suppose that $\mathcal{G} \rightarrow \overline{\mathcal{G}}$ has finite kernel and that we have a finite p -unramified extension E'/E and an étale map $V \rightarrow \mathrm{Sh}_{\overline{K} \otimes_{\overline{E}}} E'$. If V admits a crystalline ICM $\mathcal{V}$ over $\mathcal{O}_{\overline{E}, (\overline{v})}$ (relative to $\mathrm{Sh}_{\overline{K}}$), then $V \times_{\mathrm{Sh}_{\overline{K}}} \mathrm{Sh}_K$ also admits a crystalline ICM over $\mathcal{O}_{E,(v)}$ (relative to Sh_K).*

Proof. The map $\tilde{V} \stackrel{\mathrm{def}}{=} V \times_{\mathrm{Sh}_{\overline{K}}} \mathrm{Sh}_K \rightarrow V$ is finite étale. Let $\tilde{\mathcal{V}}$ be the normalization of $\mathcal{V}$ in $\tilde{V}$. We claim that $\tilde{\mathcal{V}}$ is the desired ICM for $\tilde{V}$.

We begin by checking that $\tilde{\mathcal{V}} \rightarrow \mathcal{V}$ is finite étale and that the classifying map for $\mathbf{Et}_{K,p}|_{\tilde{\mathcal{V}}}$ admits an extension $\tilde{\mathcal{V}} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v, \mathrm{alg}}$ lifting the composition

$$\tilde{\mathcal{V}} \rightarrow \mathcal{V} \rightarrow \mathrm{BT}_{\infty}^{\overline{\mathcal{G}}, \overline{\mu}_v^{-1}, \mathrm{alg}}.$$

To do this, we apply the criteria from Proposition ??. We will see that the μ -ordinary locus is the dense open subspace $U \subset \mathcal{V} \otimes k(v)$ with the required properties.¹¹

[20] **Alex:** Do use this remark

[10] **Keerthi** is essential for proposition.

[11] **Keerthi** about canonic

At this point, we know that $\tilde{\mathcal{V}} \rightarrow \mathcal{V}$ is finite étale and that $\tilde{\mathcal{V}}$ is an apertile integral model for V satisfying the conditions of Lemma 6.4.7. It only remains to check that, for F as in the definition of integral canonical models, $\tilde{\mathcal{V}}(\mathcal{O}_F) \subset \tilde{V}(F)$ contains all potentially crystalline points, but this is immediate from the corresponding assertion for $\mathcal{V}$.²¹¹² $\square$

7.6. Abstract twisting.

Definition 7.6.1 (Modified Weil restriction). Suppose that F is a totally real field in which p is unramified. The **modified Weil restriction** $\text{Res}'_{F/\mathbb{Q}}(G, \mathcal{G}, X)$ is the unramified tuple with underlying reductive group

$$\text{Res}'_{F/\mathbb{Q}} G \stackrel{\text{defn}}{=} \ker(\text{Res}_{F/\mathbb{Q}} G \rightarrow \text{Res}_{F/\mathbb{Q}} G^{\text{ab}}/G^{\text{ab}})$$

with reductive model

$$\text{Res}'_{(\mathcal{O}_F \otimes \mathbb{Z}_p)/\mathbb{Z}_p} \mathcal{G} = \ker(\text{Res}_{(\mathcal{O}_F \otimes \mathbb{Z}_p)/\mathbb{Z}_p} \mathcal{G} \rightarrow \text{Res}_{(\mathcal{O}_F \otimes \mathbb{Z}_p)/\mathbb{Z}_p} \mathcal{G}^{\text{ab}}/\mathcal{G}^{\text{ab}})$$

and Hermitian symmetric domain $\text{Res}'_{F/\mathbb{Q}} X$ given by the orbit under

$$(\text{Res}'_{F/\mathbb{Q}} G)(\mathbb{R}) \subset (\text{Res}_{F/\mathbb{Q}} G)(\mathbb{R}) = \prod_{\tau:F \rightarrow \mathbb{R}} G(\mathbb{R})$$

of the diagonal embedding $X \hookrightarrow \prod_{\tau:F \rightarrow \mathbb{R}} X$. Note that the reflex field of $\text{Res}'_{F/\mathbb{Q}}(G, \mathcal{G}, X)$ is still E .

Assumption 7.6.2. For all totally real fields F in which p is unramified, every unramified tuple with underlying unramified Shimura datum $\text{Res}'_{F/\mathbb{Q}}(G, \mathcal{G}, X)$ admits an ICM over $\mathcal{O}_{E,(v)}$.

7.7. Descent for ICMs. We will put ourselves in the situation of Setup 7.5.1. Our goal is to prove part (2) of Theorem F in the following formulation:

Theorem 7.7.1 (Descent for ICMs). *Suppose that Assumption 7.6.2 holds. Then $\text{Sh}_{\overline{K}}$ admits an ICM $\mathcal{S}_{\overline{K}}$ over $\mathcal{O}_{\overline{E},(\overline{v})}$ and the map $\mathcal{S}_K \rightarrow \mathcal{S}_{\overline{K}} \otimes_{\mathcal{O}_{\overline{E},(\overline{v})}} \mathcal{O}_{E,(v)}$ is finite étale.*

For the proof of the theorem, it will be useful to have a generalization of the notion of an ICM to étale schemes over Sh_K . This is purely for technical reasons, and is not necessarily the ‘correct’ such definition for for all such schemes, so we would suggest that the reader not take it too seriously.

Definition 7.7.2 ((Generalized) Integral canonical models). Suppose that we have a finite extension E'/E in which p is still unramified, a place $v'|v$ of E , and an étale map $U \rightarrow E' \otimes_E \text{Sh}_K$. An apertile²⁸ integral model $\mathcal{U}$ for U over $\mathcal{O}_{E',(v')}$ is a **crystalline integral canonical model**—or **crystalline ICM** for short—if it satisfies the equivalent conditions of Lemma 6.4.7, and if, for any mixed characteristic $(0, p)$ complete discrete valuation field F over $\mathcal{O}_{E',v'}$, with perfect residue field, the following are equivalent for $x \in U(F)$:

- (1) $x \in \mathcal{U}(\mathcal{O}_F)$;
- (2) $\mathbf{Et}_{K,p,x}$ is crystalline;
- (3) $\mathbf{Et}_{K,p,x}$ is potentially crystalline.

It is an **étale integral canonical model**—or **étale ICM** for short—if, instead, we replace the last condition with: For any mixed characteristic $(0, p)$ discrete valuation field F over $\mathcal{O}_{E',v'}$, the following are equivalent for $x \in U(F)$:

- (1) $x \in \mathcal{U}(\mathcal{O}_F)$;
- (2) $\mathbf{Et}_{K,\ell,x}$ is unramified for all $\ell \neq p$;
- (3) $\mathbf{Et}_{K,\ell,x}$ is potentially unramified for all $\ell \neq p$.

If the context is agnostic to the precise type of integral canonical model involved, we will simply call $\mathcal{U}$ an **ICM** for U . If we want to emphasize the dependence on the map to Sh_K , we will say that it is an **ICM relative to Sh_K** .

Remark 7.7.3 (Properties of generalized ICMs). The proofs from §6.5 work *mutatis mutandis* to show:

- (1) An ICM for U is unique up to unique isomorphism.

²⁸By this, we mean that $\mathcal{U}$ is apertile when viewed as an integral model for the E -scheme U over $\mathcal{O}_{E,(v)}$.

[21] Alex: M would be clear say the point cause f : wt $\mathcal{V}$ is finite one h $f^{-1}(\hat{\mathcal{V}}_\eta) = \hat{\mathcal{V}}$ of the pot. cr locus is pot. c [12] Keerthi what else coul It quickly gets boring even fo reader to spel obvious thing

- (2) Suppose that we have a map of unramified tuples $(G_1, \mathcal{G}_1, X_1, K_1) \rightarrow (G_2, \mathcal{G}_2, X_2, K_2)$, p -unramified extensions E'_2/E_2 and E'_1/E_1 with $E'_2 \subset E'_1$, and that we have a commuting diagram of E'_2 -schemes

$$\begin{array}{ccc} U_1 & \longrightarrow & U_2 \\ \downarrow & & \downarrow \\ E'_1 \otimes_{E_1} \mathrm{Sh}_{K_1} & \longrightarrow & E'_2 \otimes_{E_2} \mathrm{Sh}_{K_2} \end{array}$$

where the vertical arrows are étale, and where the bottom arrow is obtained as a base-change of the map from Remark 6.3.5. Suppose that $v'_1|p$ is a place of E'_1 lying above places v'_2 of E'_2 , v_2 of E_2 , and v_1 of E_1 . Suppose also that $\mathcal{U}_1$ (resp. $\mathcal{U}_2$) is an ICM for U_1 over $\mathcal{O}_{E'_1, (v'_1)}$ relative to Sh_{K_1} (resp. for U_2 over $\mathcal{O}_{E'_2, (v'_2)}$ relative to Sh_{K_2}). Then the map $U_1 \rightarrow U_2$ extends to a map $\mathcal{U}_1 \rightarrow \mathcal{U}_2$ such that we have a commuting diagram of $\mathcal{O}_{E_2, (v_2)}$ -stacks

$$\begin{array}{ccc} \mathcal{U}_1 & \longrightarrow & \mathcal{U}_2 \\ \downarrow & & \downarrow \\ \mathrm{BT}_\infty^{\mathcal{G}_1, -\mu_{v_1}, \mathrm{alg}} & \longrightarrow & \mathrm{BT}_\infty^{\mathcal{G}_2, -\mu_{v_2}, \mathrm{alg}} \end{array}$$

where the vertical arrows are formally étale.

s_gen_ICMs

Lemma 7.7.4 (Descent for coefficients). *Suppose that E''/E' is a further p -unramified extension, and that $v''|v'$ is a place of E'' such that $E'' \otimes_{E'} U$ admits an ICM $\mathcal{U}''$ over $\mathcal{O}_{E'', (v'')}$. Then U admits an ICM already over $\mathcal{O}_{E', (v')}$.*

Proof. We can assume without loss of generality that E''/E' is finite Galois.

It suffices to show that $\widehat{\mathcal{U}}''$, along with the classifying map $\widehat{\mathcal{U}}'' \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}$, descends to a formal algebraic space $\widehat{\mathcal{U}}$ over $\mathcal{O}_{E', (v')}$. Indeed, this would show—by the Beauville-Laszlo gluing results of [2] for instance—the existence of an apertile integral model $\mathcal{U}$ over $\mathcal{O}_{E', (v')}$ satisfying the equivalent conditions of Lemma 6.4.7, and the pointwise conditions would follow immediately from those for $\mathcal{U}''$.

To show that $\widehat{\mathcal{U}}''$ descends, it suffices to establish that the action of $\mathrm{Gal}(E''_{v''}/E'_{v'}) \leq \mathrm{Gal}(E''/E')$ on $E'' \otimes_{E'} U$ extends to an action on $\mathcal{U}''$, which follows from (2) of Remark 7.7.3. $\square$

Setup 7.7.5. In the situation of (2) of Remark 7.7.3, suppose that $\mathcal{G}_1 \rightarrow \mathcal{G}_2$ is a central cover, and that the map $U_1 \rightarrow U_2 \otimes_{E'_2} E'_1$ is a finite Galois cover with Galois group Δ . Suppose that we have an ICM $\mathcal{U}_1$ for U_1 over $\mathcal{O}_{E'_1, (v'_1)}$ relative to Sh_{K_1} . Then $\mathcal{U}_1$ is also an ICM over $\mathcal{O}_{E'_1, (v'_1)}$ for U_1 relative to Sh_{K_2} . Therefore, by the functoriality explained in (2) of Remark 7.7.3, the action of Δ on U_1 extends to one on $\mathcal{U}_1$ as an algebraic space over $\mathcal{O}_{E'_1, (v'_1)}$.

scent_icms

Lemma 7.7.6 (Galois descent for ICMs). *In the above situation, suppose that the action of Δ on $\mathcal{U}_1$ is free. Then U_2 also admits an ICM over $\mathcal{O}_{E_2, (v_2)}$.*

Proof. By Lemma 7.7.4, we can assume that $E'_2 = E'_1$. By our hypothesis, the Deligne–Mumford stack $[\mathcal{U}_1/\Delta]$ is an algebraic space $\mathcal{U}_2$ over $\mathcal{O}_{E'_1, (v'_1)}$ with generic fiber U_2 . Moreover, by Proposition 5.4.3, we obtain an extension $\mathcal{U}_2 \rightarrow \mathrm{BT}_\infty^{\mathcal{G}_2, -\mu_{v_2}, \mathrm{alg}}$ whose p -adic completion is formally étale.

We now claim that $\mathcal{U}_2$ is an ICM for U_2 . Let us consider the situation of a crystalline ICM; the argument for an étale ICM is similar. We have to verify that if $F/E'_{1, v'_1}$ is a complete discrete valuation field with perfect residue field, and $x \in U_2(F)$ is such that the $\mathcal{G}_2^c(\mathbb{Z}_p)$ -local system $\mathbf{Et}_{K_2, p, x}$ is potentially crystalline, then $x \in \mathcal{U}_2(\mathcal{O}_F)$. For this, after replacing F with a finite extension if necessary we can assume that x lifts to $y \in U_1(F)$. We claim that the $\mathcal{G}_1^c(\mathbb{Z}_p)$ -local system $\mathbf{Et}_{K_1, p, y}$ is also potentially crystalline: This would imply that we have $y \in \mathcal{U}_1(\mathcal{O}_F)$ and hence that its image x lies in $\mathcal{U}_2(\mathcal{O}_F)$.

To prove the claim, consider the map $\mathcal{G}_1 \rightarrow \mathcal{G}_2 \times \mathcal{G}^{\mathrm{ab}}$: This has finite kernel, and so to know that a $\mathcal{G}_1^c(\mathbb{Z}_p)$ -local system over F is potentially crystalline, it is enough to know that the associated local systems for $\mathcal{G}_2^c(\mathbb{Z}_p)$ and $\mathcal{G}^{\mathrm{ab}, c}(\mathbb{Z}_p)$ -local systems are potentially crystalline. For $\mathbf{Et}_{K_1, p, y}$, we are guaranteed the potential crystallinity

of the first by hypothesis (and the functoriality explained in Remark 6.1.8), and that of the second follows from Remark 7.1.4. $\square$

gral_model

Construction 7.7.7. Suppose that we are given a finite p -unramified extension $E^\heartsuit/E$, a place $v^\heartsuit|v$, and an open and closed subscheme

$$\mathrm{Sh}_K^\heartsuit \subset \mathrm{Sh}_K \otimes_E E^\heartsuit.$$

Given a normal integral model $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_{E,(v)}$, we obtain a normal integral model $\mathcal{S}_K^\heartsuit$ for $\mathrm{Sh}_K^\heartsuit$ over $\mathcal{O}_{E^\heartsuit,(v^\heartsuit)}$ by taking the Zariski closure of $\mathrm{Sh}_K^\heartsuit$ in $\mathcal{S}_K \otimes_{\mathcal{O}_{E,(v)}} \mathcal{O}_{E^\heartsuit,(v^\heartsuit)}$. It is easy to see that $\mathcal{S}_K^\heartsuit$ is an ICM for $\mathrm{Sh}_K^\heartsuit$ if $\mathcal{S}_K$ is one for Sh_K .

canonical

Lemma 7.7.8 (Canonical models via Hecke translates). *If $\mathrm{Sh}_K^\heartsuit$ admits an ICM $\mathcal{S}_K^\heartsuit$ over $\mathcal{O}_{E^\heartsuit,v^\heartsuit}$, then Sh_K admits an ICM $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$ such that $\mathcal{S}_K^\heartsuit$ is obtained from $\mathcal{S}_K$ by Construction 7.7.7.*

Proof. We will argue as in [61, Lemma 2.5.16]. Let us first note that, for all $K' \subset K$ of the form $K_p K'^p$, $\mathrm{Sh}_{K'}^\heartsuit \stackrel{\mathrm{defn}}{=} \mathrm{Sh}_K^\heartsuit \times_{\mathrm{Sh}_K} \mathrm{Sh}_{K'}$ admits an ICM $\mathcal{S}_{K'}^\heartsuit$: For étale ICMs, this is immediate from the fact that the local systems $\mathbf{Et}_{K,\ell}|_{\mathrm{Sh}_K^\heartsuit}$ extend over $\mathcal{S}_K^\heartsuit$. For crystalline ICMs, this is a consequence of Proposition 7.5.3 applied to the maps $(G, \mathcal{G}, X, K') \rightarrow (G, \mathcal{G}, X, K)$ and $V = \mathrm{Sh}_K^\heartsuit \subset \mathrm{Sh}_K \otimes_E E^\heartsuit$.

For any $g \in G(\mathbb{A}_f^p)$, we have an isomorphism of Shimura varieties $\iota_{K',g}: \mathrm{Sh}_{K'} \xrightarrow{\sim} \mathrm{Sh}_{gK'g^{-1}}$ given on $\mathbb{C}$ -points by

$$[(x, h)] \mapsto [(x, hg^{-1})].$$

This isomorphism carries the restriction of $\mathbf{Et}_{K,p}$ to the source to that over the target.

Then we see that $\iota_{K',g}(\mathrm{Sh}_{K'}^\heartsuit) \subset \mathrm{Sh}_{gK'g^{-1}} \otimes_E E^\heartsuit$ also admits an ICM over $\mathcal{O}_{E^\heartsuit,v^\heartsuit}$, which is in fact isomorphic to $\mathcal{S}_{K'}^\heartsuit$. If $K'_g = K' \cap gK'g^{-1}$, this in turn implies that

$$\mathrm{Sh}_{K'_g}^\heartsuit \stackrel{\mathrm{defn}}{=} \iota_{K',g}(\mathrm{Sh}_{K'}^\heartsuit) \times_{\mathrm{Sh}_{gK'g^{-1}}} \mathrm{Sh}_{K'_g}$$

admits an ICM model $\mathcal{S}_{K'_g}^\heartsuit$; see the argument from the first paragraph of the proof.

By Lemma 7.7.6, we find that the image of $\mathrm{Sh}_{K'_g}^\heartsuit$ in $\mathrm{Sh}_K \otimes_E E^\heartsuit$ admits an ICM over $\mathcal{O}_{E^\heartsuit,v^\heartsuit}$.

Now, using Remark 6.3.6, we find that there is a finite set of pairs $\Xi = \{(K', g)\}$ such that the map

$$\bigsqcup_{(K',g) \in \Xi} \mathrm{Sh}_{K'_g}^\heartsuit \rightarrow \mathrm{Sh}_K \otimes_E E^\heartsuit$$

is surjective.

Therefore, $\mathrm{Sh}_K \otimes_E E^\heartsuit$ is covered by open and closed subschemes, each of which admits an ICM over $\mathcal{O}_{E^\heartsuit,v^\heartsuit}$. Using (1) of Remark 7.7.3 and Zariski descent, we see that $\mathrm{Sh}_K \otimes_E E^\heartsuit$ admits an ICM over $\mathcal{O}_{E^\heartsuit,v^\heartsuit}$. We now conclude using Lemma 7.7.4. $\square$

Proof of Theorem 7.7.1. The map $\mathrm{Sh}_K \rightarrow E \otimes_{\overline{E}} \mathrm{Sh}_{\overline{K}}$ is a finite Galois cover over its image $\mathrm{Sh}_{\overline{K}}^\heartsuit \subset E \otimes_{\overline{E}} \mathrm{Sh}_{\overline{K}}$. Therefore, Lemma 7.7.6 tells us²² that $\mathrm{Sh}_{\overline{K}}^\heartsuit$ admits an ICM over $\mathcal{O}_{E,(v)}$, and Lemma 7.7.8 tells us that Sh_K admits an ICM over $\mathcal{O}_{\overline{E},(\overline{v})}$. $\square$

abelian_type

7.8. Shimura varieties of pre-abelian type.

Definition 7.8.1. A Shimura datum (G, X) is of **Hodge type** if G admits a faithful symplectic representation H on which some (hence any) element of X induces a Hodge structure with weights $(-1, 0), (0, -1)$. It is of **abelian type** if there exists a Shimura datum (G_1, X_1) of Hodge type and a map $G_1^{\mathrm{der}} \rightarrow G_2^{\mathrm{der}}$ of derived groups inducing an isomorphism on the corresponding adjoint Shimura data. It is of **pre-abelian type** if the adjoint Shimura datum $(G^{\mathrm{ad}}, X^{\mathrm{ad}})$ is of abelian type.

pre-abelian

Remark 7.8.2 (Classification of pre-abelian Shimura data). By the classification of Shimura data by Deligne [20, §1.3], a Shimura datum (G, X) is of pre-abelian type if and only if each simple factor of the adjoint Shimura datum $(G^{\mathrm{ad}}, X^{\mathrm{ad}})$ ²³ is classical of type $A, B, C, D^{\mathbb{R}}, D^{\mathbb{H}}$.

[22] Alex: Re this is where modify things

[23] Alex: Fo nience of use, it's worth just "G^{ad} is classical type...".

Note that a simple adjoint Shimura datum has underlying group of the form $\text{Res}_{F/\mathbb{Q}} H$ where F is a totally real field and H is an absolutely simple adjoint group over F . The meaning of type $D^{\mathbb{R}}$ is that, for each embedding $\tau : F \rightarrow \mathbb{R}$, $H_{\tau} \stackrel{\text{defn}}{=} H \otimes_{F,\tau} \mathbb{R}$ is the adjoint quotient of an orthogonal group associated with a quadratic form of signature either $(2n, 2)$, $(2n, n)$ or $(2n + 2, 0)$. The meaning of type $D^{\mathbb{H}}$ is instead that H_{τ} is the adjoint quotient of $\text{O}^*(2n)$, the group of isometries of a skew-Hermitian space over the ring of quaternions $\mathbb{H}$. The Shimura datum is of *abelian* type if the derived subgroup of every simple factor of $G_{\mathbb{R}}$ of type $D^{\mathbb{H}}$ is a quotient of $\text{SO}^*(2n)$.

pre-abelian

Example 7.8.3 (A non-abelian but pre-abelian Shimura datum). The basic example of a pre-abelian-type (but not abelian-type) Shimura datum is obtained as follows. Take a totally real field F_0 and a quaternion algebra B over F_0 such that for some distinguished archimedean place v_0 of F_0 we have $B_{v_0} \simeq \mathbb{H}$, and $B_v \simeq \text{Mat}_2(\mathbb{R})$ for archimedean places $v \neq v_0$. We then take a left B -module V of rank $n \geq 4$ with a Hermitian pairing which is definitive at $v \neq v_0$ and signature $(n, 0)$ at v_0 . Set H^{ad} to be $\text{PU}(V, h)$ and $G^{\text{ad}} = \text{Res}_{F_0/\mathbb{Q}} H^{\text{ad}}$. Define $h : \mathbb{S} \rightarrow G_{\mathbb{R}}^{\text{ad}}$ to have trivial projection to each archimedean place $v \neq v_0$, be such that if $W = V \otimes_{F_0, v_0} \mathbb{R}$ then the projection to v_0 defines a Hodge structure of weight $\{(-1, 0), (0, -1)\}$ where $W_{\mathbb{C}}^{(-1, 0)}$ is the maximal isotropic subspace. This defines a Shimura datum $(G^{\text{ad}}, X^{\text{ad}})$. Let H be the central cover of H^{ad} determined by the property that $H_{\mathbb{C}}$ is isomorphic to the GSpin cover of $H_{\mathbb{C}}^{\text{ad}} \simeq \text{PSO}(2n)$. Then X^{ad} admits a lift X to $G = \text{Res}_{F_0/\mathbb{Q}} H$, and (G, X) is of pre-abelian but not of abelian type.

hm:abelian

Theorem 7.8.4 (Canonical models of pre-abelian type). *If $(G, \mathcal{G}, X, K)$ is of pre-abelian type, then it admits a crystalline ICM over $\mathcal{O}_{E, (v)}$, and this is also an étale ICM.*

Proof. Suppose first that $(G, \mathcal{G}, X)$ is of Hodge type, meaning that (G, X) admits a closed immersion into a Siegel Shimura datum $(G^{\sharp}, X^{\sharp})$. Using Zarhin's trick, we can assume that the latter can be extended to an unramified tuple $(G^{\sharp}, \mathcal{G}^{\sharp}, X^{\sharp})$: Concretely, this means that we can choose a symplectic representation V of G such that $V_{\mathbb{Q}_p}$ admits a self-dual lattice $V_{\mathbb{Z}_p}$ stabilized by $\mathcal{G}$, and such that $-\mu_v$ acts on V with weights $0, 1$.

Now, for a suitable choice of level $K^{\sharp, p} \subset G^{\sharp}(\mathbb{A}_f^p)$, $\text{Sh}_{K^{\sharp}}$ admits an integral model $\mathcal{S}_{K^{\sharp}}$ over $\mathbb{Z}_{(p)}$ given by the moduli of abelian varieties equipped with a prime-to- p polarization and some prime-to- p level structure. By Theorem 7.3.1, it is enough to know that $\mathcal{S}_{K^{\sharp}}$ is both a crystalline ICM and an étale ICM. Therefore, we can assume that $(G, \mathcal{G}, X)$ is itself of Siegel type.

Here, by Theorem 2.4.2, we have an F -gauge over $\mathcal{S}_K$ associated with the p -divisible group of the universal abelian scheme $\mathcal{A} \rightarrow \mathcal{S}_K$. We can actually view this as a (GL_{2g}, μ_g) -aperture over $\mathcal{S}_K$, where g is the relative dimension of $\mathcal{A}$. Now, Proposition 5.3.9 applies with

$$\mathcal{X} = \mathcal{X}^{\sharp} = \mathcal{S}_K; (\mathcal{G}^{\sharp}, \mu^{\sharp}) = (\text{GL}_{2g}, \mu_g).$$

The pointwise condition is immediate from Remarks 1.3.4 and 6.4.4; that the map $\widehat{\mathcal{S}}_{K, v} \rightarrow \text{BT}_{\infty}^{\text{GL}_{2g}, \mu_g}$ is formally unramified follows from classical Serre–Tate theory and Theorem 2.4.2 once again.

Now, the modified Weil restriction from a totally real field F of a Siegel Shimura datum associated with a symplectic space V is once again of Hodge type: It embeds into the Siegel Shimura datum associated with $[F : \mathbb{Q}]$ -copies of V . Therefore, we see that Assumption 7.6.2 is valid whenever $(G, \mathcal{G}, X)$ is of Hodge type. Theorem 7.7.1 now tells us that $(G, \mathcal{G}, X)$ admits étale and crystalline ICMs whenever (G, X) is of *adjoint* abelian type, and we conclude the proof of the theorem using Theorem E. $\square$

Remark 7.8.5. The alert reader will note that polarized p -divisible groups do not make an appearance in the above proof (and they do not in Kisin's original argument in [47] either). One could shorten the proof slightly by noting that the moduli stack of prime-to- p degree polarized abelian schemes of dimension g carries a canonical principally polarized p -divisible group of height $2g$, which can then be reinterpreted as a (GSp_{2g}, μ_g) -aperture; see [26, §11.6]. However, this reinterpretation has not been carried out fully satisfactorily (see footnote 43 of *loc. cit.*), and so we have preferred to avoid using it in our argument above.

properties

7.9. Mapping properties for smooth inputs. In this subsection, we will give more concrete mapping properties for ICMs in terms of pointwise criteria.

conditions

Definition 7.9.1 (Pointwise conditions). Suppose that we have an algebraic space $\mathcal{X}$ over $\mathcal{O}_{E,(v)}$ equipped with a map $f : \mathcal{X}_\eta \rightarrow \mathrm{Sh}_K$. Let F/E_v be a finite extension and let $x \in \mathcal{X}(\mathcal{O}_F)$. Then, we say that f is **crystalline at x** if $\mathbf{Et}_{K,p}|_x$ is crystalline. We say that f is **pointwise crystalline** if it is crystalline at all such x . It is **pointwise unramified** if, for all such x , we instead require that $\mathbf{Et}_{K,\ell}|_x$ is unramified for all $\ell \neq p$.

canonical

Proposition 7.9.2 (A pointwise mapping property for smooth inputs). *Suppose that one of the following conditions holds:*

- (a) $p > 3$;
- (b) $p > 2$ and (G, X) is of pre-abelian type.

Suppose that $\mathcal{X}$ is a smooth $\mathcal{O}_{E,(v)}$ -scheme with generic fiber X , equipped with a map $f : X \rightarrow \mathrm{Sh}_K$.

- (1) If $\mathcal{S}_K$ is a crystalline ICM for Sh_K over $\mathcal{O}_{E,(v)}$, f extends to a map $\mathcal{X} \rightarrow \mathcal{S}_K$ if and only if f is pointwise crystalline.
- (2) If $\mathcal{S}_K$ is an étale ICM, then f extends to a map $\mathcal{X} \rightarrow \mathcal{S}_K$ if and only if f is both pointwise crystalline and pointwise unramified.

Proof. Suppose first that $p > 2$. Given Theorem 6.5.3, it is enough to know that $\mathbf{Et}_{K,p}|_X$ lifts to a $(\mathcal{G}^c, -\mu_v)$ -aperture over $\mathcal{X}$. But this is immediate from Theorem 5.1.4 and Remark 6.4.4.

Next, suppose that (G, X) is of Hodge type and that $p > 2$. Then the same argument applies after taking Remark 5.1.15 into account. The pre-abelian case can be deduced from this via standard arguments that we leave to the reader. $\square$

r_neronian

Corollary 7.9.3 (Néronian property for proper models). *The following statements are equivalent for an integral model $\mathcal{S}_K$ for Sh_K :*

- (1) The integral model $\mathcal{S}_K$ is proper over $\mathcal{O}_{E,(v)}$ and is a crystalline ICM for Sh_K .
- (2) The integral model $\mathcal{S}_K$ is proper over $\mathcal{O}_{E,(v)}$ and is an étale ICM for Sh_K .

If these equivalent conditions hold, along with one of the conditions from Proposition 7.9.2, then $\mathcal{S}_K$ is in fact a Nèron model for Sh_K : In the notation of that proposition, every map $X \rightarrow \mathrm{Sh}_K$ extends to a map $\mathcal{X} \rightarrow \mathcal{S}_K$.

Proof. Suppose that Sh_K admits a proper integral model $\mathcal{S}_K$. Then, for any finite extension F/E_v , we have $\mathcal{S}_K(\mathcal{O}_F) = \mathrm{Sh}_K(F)$. Therefore, if $\mathcal{S}_K$ is apertile, then for all $x \in \mathrm{Sh}_K(F)$, $\mathbf{Et}_{K,p,x}$ is crystalline. If $\mathcal{S}_K$ is now an étale ICM, then it follows immediately that it is also a crystalline ICM. Conversely, if $\mathcal{S}_K$ is a crystalline ICM, then Corollary ?? tells us that, for all $x \in \mathrm{Sh}_K(F)$ and $\ell \neq p$, $\mathbf{Et}_{K,\ell,x}$ is also unramified. Therefore $\mathcal{S}_K$ is also an étale ICM.

That $\mathcal{S}_K$ is a Nèron model for Sh_K is now immediate from Proposition 7.9.2. $\square$

Remark 7.9.4 (Proper models of pre-abelian type). A necessary condition for properness of $\mathcal{S}_K$ is that G^{ad} be anisotropic, since this is known to be equivalent to the generic fiber Sh_K being proper. With this condition in hand, it follows from [66, Theorem 1] that, when (G, X) is of pre-abelian type, and $p > 2$, the ICMs from Theorem 7.8.4 are proper and so are Nèron models of their generic fibers.

or-nothing

Remark 7.9.5 (Sufficient to check crystallinity at one point). Suppose one knew that $\mathbf{Et}_{K,p}$ is a *pointwise semistable* local system on Sh_K : In the abelian type case, this follows from the structure of the boundary of the integral model of a toroidal compactification; see [97, Theorem 4.39] and earlier results in [66]. We expect that this is true in general as well.

Under such a pointwise semistability condition, recent results of Guo–Yang [35, Theorem 1.1] and Diao–Yao [22, Theorem 1.1] allow us to considerably weaken the hypotheses of Proposition 7.9.2 as they pertain to crystalline ICMs. Namely, in this case, statement (1) from this proposition can be upgraded to only require f to be crystalline at x for *some* $x \in \mathcal{X}(\mathcal{O}_F)$ in every connected component of $\mathcal{X}$.

n-diao-yao

Remark 7.9.6. Suppose that $p > 2$ and $(G, \mathcal{G}, X, K)$ is an unramified Shimura tuple of *pre-abelian type*. We do not know yet if $\mathbf{Et}_{K,p}$ is semistable over Sh_K . However, since it suffices to know that the composition $X \rightarrow \mathrm{Sh}_K \rightarrow \mathrm{Sh}_{K^{\mathrm{ad}}}$ extends to a map $\mathcal{X} \rightarrow \mathcal{S}_{K^{\mathrm{ad}}}$, the strengthened mapping property from Remark 7.9.5 still holds for $\mathcal{S}_K$ itself.

sub:bst

7.10. Canonicity of the models of Bakker–Shankar–Tsimmerman. Here, we will find a proof of Theorem G. We can assume that the rank of G is at least 2, since otherwise the Shimura datum is of pre-abelian type and so is already addressed by Theorem 7.8.4. By Remark 1.2.8, we can also assume that G is *adjoint*.

Proof of Theorem G. We will take a prime $p > 3$ large enough such that the following holds:

- (1) (Log smooth compactification) First, we can assume that p is sufficiently large so that Sh_K admits a smooth model $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$ with a log smooth compactification;
- (2) (Versal Kodaira–Spencer map) Furthermore, by increasing the lower bound for p , we can also assume that the filtered G -bundle with integrable connection $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet \mathbf{dR}_K$ extends to a filtered $\mathcal{G}_{(p)}$ -bundle with integrable connection on $\mathcal{S}_K$, which we denote by the same symbol. We can also assume that this satisfies Griffiths’ transversality and that the associated Kodaira–Spencer map

$$\mathbb{T}_{\mathcal{S}_K/\mathcal{O}_{E,(v)}} \rightarrow \mathbf{dR}_K(\mathfrak{g})/\mathrm{Fil}_{\mathrm{Hdg}}^0 \mathbf{dR}_K(\mathfrak{g})$$

is an isomorphism.

- (3) (Existence of ℓ -adic local systems) By work of Klevdal-Patrikis [52, Theorem 1.3] and Patrikis [74, Corollary 3.4], we can assume that, for all $\ell \neq p$, $\mathbf{Et}_{K,\ell}$ extends to an ℓ -adic local system on $\mathcal{S}_K$, which again we denote by the same symbol.
- (4) (Pointwise crystallinity) By [76, Theorem 7.1], we can also assume that, for some choice of faithful representation Λ of $\mathcal{G}$, the filtered de Rham realization $\mathrm{Fil}_{\mathrm{Hdg}}^\bullet \mathbf{dR}_K(\Lambda)$ lifts to a Fontaine-Laffaille module (up to suitable twist) over $\mathcal{S}_K$. This implies in particular that $\mathbf{Et}_{K,p}$ is crystalline at every classical point of $\widehat{\mathcal{S}}_{K,\eta}$.

At this point, we are almost there. Theorem 5.1.4, combined with Remark 6.4.4 and Lemma 6.4.7, tells us that we have an extension $\mathcal{S}_K \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, \sim \mu_v, \mathrm{alg}}$ whose v -adic completion is formally étale.

Let us now check that $\mathcal{S}_K$ satisfies condition (1) for being an étale ICM. That is, we must know that, if $x \in \mathrm{Sh}_K(F)$ is such that $\mathbf{Et}_{K,\ell,x}$ is potentially unramified for all $\ell \neq p$, then $x \in \mathcal{S}_K(\mathcal{O}_F)$. This follows from the argument in [76, Lemma 8.4].

Finally, suppose that G^{ad} is anisotropic. In this case, Sh_K is a proper variety over E , and, since $\mathcal{S}_K$ admits a log smooth compactification, it follows that $\mathcal{S}_K$ is also proper over $\mathcal{O}_{E,(v)}$. We now conclude using Corollary 7.9.3. $\square$

Remark 7.10.1. Since we only require the above conditions for *adjoint* Shimura data, and also do not require any ampleness conditions, but only the versality of the Kodaira–Spencer map, the lower bound on p we arrive at here is logically smaller than that in [4]. Of course, one expects that the bound is 2!

8. APPLICATIONS AND FURTHER RESULTS

In this section, we collect some immediate applications and extensions of our theory of ICMs. In addition to discussing cohomological applications and integral models at (some) parahoric levels, we prove Theorems H, J, and I from the introduction. We freely use notation and terminology from §6–7.

8.1. Surjectivity. The following result of Vasiu [90, Main Theorem A], will be used in the proof of Theorem H, which is the aim of this subsection.

Theorem 8.1.1 (Vasiu). *Let the notation be as in Setup 3.1.1. Then there exists $m(\mathcal{G}, \mu) \geq 1$ such that, for any algebraically closed field κ over $\mathcal{O}$, and any $n \geq m(\mathcal{G}, \mu)$, the map of groupoids*

$$\mathrm{BT}_\infty^{\mathcal{G}, \mu}(\kappa) \rightarrow \mathrm{BT}_n^{\mathcal{G}, \mu}(\kappa)$$

induces a bijection on isomorphism classes.

Proof. This is just a reformulation of the cited result of Vasiu, using the quotient descriptions of $\mathrm{BT}_n^{\mathcal{G}, \mu}(\kappa)$ and $\mathrm{BT}_\infty^{\mathcal{G}, \mu}(\kappa)$ given to us by Remarks 3.2.20 and 3.2.19. $\square$

er_results

rjectivity

oundedness

Remark 8.1.2. An equicharacteristic version of Theorem 8.1.1 is given by [94, Theorem 6.31]. The argument in *loc. cit.* can be combined with work of Zhu on the Witt vector affine Grassmannian [99] to give an alternate proof of Theorem 8.1.1.

Definition 8.1.3. We will say that two unramified Shimura data $(G, \mathcal{G}, X)$ and $(G', \mathcal{G}', X')$ are in the same **central equivalence class** if the associated adjoint unramified Shimura data are isomorphic.

g_property

Definition 8.1.4. We will say that an unramified Shimura tuple $(G, \mathcal{G}, X, K)$ admitting a crystalline ICM $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$ satisfies the **de Jong extension property (at v)** if the following condition holds: Suppose that C is a smooth curve over an algebraically closed field κ over $k(v)$ with smooth compactification $\overline{C}$. Then a map $C \rightarrow \mathcal{S}_K$ extends to $\overline{C} \rightarrow \mathcal{S}_K$ if and only if the composition with $\mathcal{S}_K \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v}$ extends to a map $h : C \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}^c, -\mu_v}$.

automatic

Remark 8.1.5. The extension property is automatic if Sh_K admits a proper ICM $\mathcal{S}_K$ over $\mathcal{O}_{E,(v)}$.

ong_siegel

Remark 8.1.6. The extension property holds for the Siegel Shimura datum by results of de Jong [19, Corollaries 1.2 and 2.5], justifying the terminology.

properness

Proposition 8.1.7. *Suppose that there exists $(G', \mathcal{G}', X')$ in the central equivalence class of $(G, \mathcal{G}, X)$ and a closed immersion $(G', \mathcal{G}', X') \hookrightarrow (G^{\sharp}, \mathcal{G}^{\sharp}, X^{\sharp})$ into another unramified Shimura datum such that, for all places $v^{\sharp} | p$ of its reflex field $E^{\sharp}$, the unramified Shimura tuple $(G^{\sharp}, \mathcal{G}^{\sharp}, X^{\sharp}, K^{\sharp})$ has the de Jong property at $v^{\sharp}$ for some $K^{\sharp}$. Then $(G, \mathcal{G}, X, K)$ has the de Jong property at all places $v | p$ of E .*

Proof. First, let us see that $(G, \mathcal{G}, X)$ admits crystalline ICMs over $\mathcal{O}_{E,(v)}$. To begin, in the notation of the main assumption above, Theorem 7.3.1 implies that $(G', \mathcal{G}', X')$ admits crystalline ICMs. Next, Theorem ?? successively implies that the adjoint triple $(G^{\mathrm{ad}}, \mathcal{G}^{\mathrm{ad}}, X^{\mathrm{ad}})$ and hence $(G, \mathcal{G}, X)$ itself admits crystalline ICMs over $\mathcal{O}_{E,(v)}$.

Now, let C be a smooth curve over an algebraically closed field κ over $k(v)$ and a map $C \rightarrow \mathcal{S}_K$ such that the resulting $(\mathcal{G}^c, -\mu_v)$ -aperture over C extends to one over its smooth compactification $\overline{C}$.

Suppose first that we have a map of tuples $(G, \mathcal{G}, X, K) \rightarrow (G^{\sharp}, \mathcal{G}^{\sharp}, X^{\sharp}, K^{\sharp})$ with $\mathcal{G} \rightarrow \mathcal{G}^{\sharp}$ a closed immersion, and with $(G^{\sharp}, \mathcal{G}^{\sharp}, X^{\sharp}, K^{\sharp})$ satisfying the de Jong property. In this case, the map $\mathcal{S}_K \rightarrow \mathcal{S}_{K^{\sharp}}$ is finite by Theorem 7.3.1. By assumptions, we have an extension $\overline{C} \rightarrow \mathcal{S}_{K^{\sharp}}$ of the composition $C \rightarrow \mathcal{S}_K \rightarrow \mathcal{S}_{K^{\sharp}}$ and this must necessarily lift to the desired extension to $\mathcal{S}_K$.

Now, by our hypotheses, the assumption in the previous paragraph holds, up to replacing our unramified triple with another triple $(G', \mathcal{G}', X')$ in its central isogeny class. Suppose now that $(G^{\mathrm{ad}}, \mathcal{G}^{\mathrm{ad}}, X^{\mathrm{ad}}, K^{\mathrm{ad}})$ is an unramified tuple associated with the adjoint group with both K and K' mapping into K^{ad} , and consider the composition

$$f^{\mathrm{ad}} : C \rightarrow \mathcal{S}_K \rightarrow \mathcal{S}_{K^{\mathrm{ad}}}.$$

Since the second map here is finite étale by Theorem ??, it is enough to show f^{ad} extends to a map $\overline{C} \rightarrow \mathcal{S}_{K^{\mathrm{ad}}}$. Now, we can find a level $K' \subset G'(\mathbb{A}_f)$ and a map of unramified tuples

$$(G', \mathcal{G}', X', K') \rightarrow (G^{\mathrm{ad}}, \mathcal{G}^{\mathrm{ad}}, X^{\mathrm{ad}}, K^{\mathrm{ad}})$$

such that the image of C is contained in the image of the associated finite étale map $\mathcal{S}_{K'} \rightarrow \mathcal{S}_{K^{\mathrm{ad}}}$ (e.g., by Remark 6.3.6). By pulling back, we can then find a smooth proper curve $\overline{C}'$ over κ and a finite map $\pi : \overline{C}' \rightarrow \overline{C}$ such that f^{ad} lifts to a map $f' : \pi^{-1}(C) \rightarrow \mathcal{S}_{K'}$, which can be completed to a map $\overline{C}' \rightarrow \mathcal{S}_{K'}$. This then evidently implies that f^{ad} admits an extension to all of $\overline{C}$. $\square$

Theorem H is now implied by combining Proposition 8.1.7, Remark 8.1.5, Remark 8.1.6, and the following result:

rjectivity

Theorem 8.1.8. *Suppose that $(G, \mathcal{G}, X, K)$ admits a crystalline ICM over $\mathcal{O}_{E,(v)}$ satisfying the de Jong extension property. Then, for all $1 \leq n \leq \infty$, the map of formal Artin stacks*

$$\widehat{\mathcal{S}}_K \rightarrow \mathrm{BT}_n^{\mathcal{G}^c, -\mu_v}$$

is surjective, and smooth for finite n .

Proof. Note that, for every finite n , the classifying map $\widehat{\mathcal{S}}_K \rightarrow \mathrm{BT}_n^{\mathcal{G}^c, -\mu_v}$ is a formally smooth map of smooth formal Artin stacks over $\mathcal{O}$, and is therefore smooth.

By smoothness, the map $|\widehat{\mathcal{S}}_K| \rightarrow |\mathrm{BT}_n^{\mathcal{G}^c, -\mu_v}|$ of the underlying topological spaces is open, and its surjectivity amounts to saying that its image intersects every connected component and that its image is closed under specialization. The first is easy, since $\mathrm{BT}_n^{\mathcal{G}^c, -\mu_v}$ is connected (see Theorem 3.1.5).

For the surjectivity at finite level, observe the following consequence of Theorem 8.1.1: the map $|\mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}| \rightarrow |\mathrm{BT}_n^{\mathcal{G}^c, -\mu_v}|$ is a homeomorphism for n sufficiently large. Thus, to complete the proof, it is sufficient to know that the map $|\widehat{\mathcal{S}}_K| \rightarrow |\mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}|$ has image closed under specialization, and this is immediate from the de Jong extension property. Surjectivity at infinite level then follows from Theorem 8.1.1. $\square$

-emptiness

8.2. Non-emptiness of strata. Throughout the following, we fix $\mathcal{S}_K$ be an ICM for an unramified Shimura tuple $(G, \mathcal{G}, X, K)$ over $\mathcal{O}_{E,(v)}$.

Construction 8.2.1 (Newton strata). By Construction 3.3.1, for any perfect $k(v)$ -algebra R , we obtain canonical maps

newton-map

$$(8.2.1.1) \quad \mathcal{S}_K(R) \rightarrow \mathrm{BT}_\infty^{\mathcal{G}, \mu}(R) \rightarrow \mathrm{Isoc}_G(R).$$

For any class $[b] \in B(G)$, this gives a topological subspace $\mathcal{S}_K[b] \subset \mathcal{S}_K \otimes k(v)$ defined by the property that, for any perfect field κ , the κ -points of $\mathcal{S}_K$ contained in this subset are the ones whose image under the composition of (8.2.1.1) with the natural map $\mathrm{Isoc}_G(R) \rightarrow B(G)$ is $[b]$. We call $\mathcal{S}_K[b]$ the **Newton stratum** associated with $[b]$. Newton strata do form a stratification by [79, Theorem 3.6].

The next result proves Theorem J.

-emptiness

Theorem 8.2.2. *The Newton stratum $\mathcal{S}_K[b]$ is non-empty if and only if $[b] \in B(G_{\mathbb{Q}_p}^c, -\mu_v)$.*

Proof. By Lemma 7.1.8, every special point $x: \mathrm{Spec} F \rightarrow \mathrm{Sh}_K$ extends to a map $\mathrm{Spec} \mathcal{O}_{F,(w)} \rightarrow \mathcal{S}_K$ for some place $w|v$ of F . The argument in [49, Proposition 1.3.10] then gives the desired conclusion. $\square$

Remark 8.2.3. Note that if $(G, \mathcal{G}, X, K)$ satisfies the de Jong extension property, then, by Theorem H, it suffices to verify that there exists $\mathcal{Q} \in \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}(\mathbb{F}_p)$ with $[b_{\mathcal{Q}}] = [b]$. This follows from Lemma 3.3.3.

:EO_strata

Definition 8.2.4. The target of the map from Construction ?? admits a stratification by smooth locally closed substacks as in Remark 3.5.4. An **Ekedahl–Oort stratum** of $\mathcal{S}_K \otimes k(v)$ is the pre-image of any such stratum under the above map.

-emptiness

Proposition 8.2.5. *Suppose that $\mathcal{S}_K$ satisfies the de Jong extension property. Then every Ekedahl–Oort stratum of $\mathcal{S}_K \otimes k(v)$ is non-empty.*

Proof. Given the surjectivity of the syntomic realization (Theorem H), this comes down to the surjectivity assertion in Proposition 3.5.5. $\square$

cohomology

8.3. Applications to cohomology. In this subsection, we explicitly state the obvious cohomological results about integral canonical models which follow from the results of §2.5. See *loc. cit.* for the terminology and notation used below.

Notation 8.3.1. Let $\mathcal{S}_K$ be an ICM for $(G, \mathcal{G}, X, K)$. For a $\mathbb{Z}_p$ -representation Λ of $\mathcal{G}^c$ we write $\mathbf{Et}_K(\Lambda)$ for the $\mathbb{Z}_p$ -local system on Sh_K given by $(\Lambda)_{\mathbf{Et}_{K,p}}$. We uniquely extend this to an association $V \mapsto \mathbf{Et}_K(V)$ for $V \in \mathrm{Rep}_{\mathbb{Q}_p}(G^c)$ so that $\mathbf{Et}_K(\Lambda)[1/p]$ when $V = \Lambda[1/p]$.

Similarly, denote by $\mathbf{Syn}_K(\Lambda)$ the F -gauge on $\widehat{\mathcal{S}}_K$ obtained by pulling back the universal F -gauge along the syntomic realization $\varpi: \widehat{\mathcal{S}}_K \rightarrow \mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}$ and evaluating it at Λ . Finally, we shorten the notation $T_{\mathrm{crys}}^+(\mathbf{Syn}_K(\Lambda))$ to $\mathbf{Crys}_K^+(\Lambda)$. We again uniquely extend this to associations $V \mapsto \mathbf{Syn}_K(V)$ and $V \mapsto \mathbf{Crys}_K^+(V)$ in the obvious way.

cal-system

Proposition 8.3.2. *Suppose that $\mathcal{S}_K$ is a proper ICM over $\mathcal{O}_{E,(v)}$, and let C_v denote a completed algebraic closure of E_v . Then, for all $i \geq 0$ and $\mathbb{Q}_p$ -representations V of G^c , the $\text{Gal}(\overline{E}_v/E_v)$ -representation $H_{\text{et}}^i(\text{Sh}_K \otimes \overline{E}_v, \mathbf{Et}_K(V))$ is crystalline and there is a natural identification of filtered F -isocrystals:*

$$D_{\text{crys}}\left(H_{\text{et}}^i(\text{Sh}_K \otimes C_v, \mathbf{Et}_K(V))\right) \simeq H_{\text{crys}}^i((\mathcal{S}_{k(v)}/W(k(v)))_{\text{crys}}, \mathbf{Crys}_K^+(V)).$$

Moreover, for all $1 \leq n \leq \infty$ we have a natural identification

$$T_{\text{et}}(Rf_*(\mathbf{Et}_K(\Lambda)/p^n)) \simeq R(f_\eta)_*T_{\text{et}}(\mathbf{Syn}_K(\Lambda)/p^n).$$

Proposition 8.3.3. *Fix notation as in Proposition 8.3.2. For all $i \geq 0$ we have a natural identification*

$$H_f^1\left(H_{\text{et}}^i(\text{Sh}_K \otimes C_v, \mathbf{Et}_K(V))\right) \simeq H^1(\mathcal{O}_{E_v}^{\text{syn}}, R^i f_* \mathbf{Syn}_K(V)).$$

ura_stacks

8.4. Integral canonical models for Shimura stacks. In this subsection, we will drop the assumption that K be neat, but still assume that $(G, \mathcal{G}, X, K)$ is an unramified tuple. In this generality, Sh_K is a Deligne–Mumford stack and not necessarily a scheme over E . However, we still have the map $\text{Sh}_K \rightarrow \text{Loc}_{\mathcal{G}(\mathbb{Z}_p)}$ classifying the local system $\mathbf{Et}_{K,p}$.

tacky_icms

Definition 8.4.1 (Stacky integral canonical models). A **crystalline** (resp. **étale**) **ICM** for Sh_K over $\mathcal{O}_{E,(v)}$ is a separated Deligne–Mumford stack $\mathcal{S}_K$ of finite type over $\mathcal{O}_{E,(v)}$ with generic fiber Sh_K such that, for some neat compact open subgroup $K' \subset K$ with $K'_p = K_p$, normalization of $\mathcal{S}_K$ in $\text{Sh}_{K'}$ is a crystalline (resp. étale) ICM for $\text{Sh}_{K'}$. As before, if the context is agnostic to the type, we will simply call $\mathcal{S}_K$ an **ICM** for Sh_K .

We can now collect the properties of stacky ICMs.

erre-tate

Remark 8.4.2 (Serre–Tate property for stacky ICMs). If $\mathcal{S}_K$ is an ICM for Sh_K , by Corollary 4.1.6, we see that there is an extension $\mathcal{S}_K \rightarrow \text{BT}_{\infty}^{\mathcal{G}^c, -\mu_v, \text{alg}}$ of the classifying map for $\mathbf{Et}_{K,p}$, whose p -adic completion is formally étale.

non-stacky

Remark 8.4.3 (Stacky vs. non-stacky ICMs). Suppose that K is neat and Sh_K admits an ICM $\mathcal{S}_K$ in the stacky sense. Then by Proposition 7.2.1, $\mathcal{S}_K$ is in fact an algebraic space. In particular, the two notions of ICMs are not in conflict with each other.

rty_stack

Remark 8.4.4 (Mapping property for stacky ICMs). Using étale descent, the mapping property from Theorem 6.5.3 holds verbatim for stacky ICMs, and one can even take $\mathcal{Y}$ to be an excellent separated DM stack over $\mathcal{O}_{E,(v)}$, with the following caveat: One has to impose the condition that, for each $\ell \neq p$, the restriction of $\mathbf{Et}_{K,\ell}$ over $\mathcal{Y}_\eta$ extends to a K_ℓ^c -local system over $\mathcal{Y}$ represented by pro-algebraic spaces over $\mathcal{O}_{E,(v)}$.

As a consequence, we get the following generalization of Corollaries 6.5.4 and 6.5.5:

nct_stack

Proposition 8.4.5 (Uniqueness and functoriality for stacky ICMs). *Stacky ICMs are unique up to unique isomorphism and are functorial for maps of unramified tuples.*

nce_stack

Proposition 8.4.6 (Existence of stacky ICMs). *Suppose that for some neat open subgroup $K' \subset K$ with $K' = K'^p K_p$, $\text{Sh}_{K'}$ admits an ICM $\mathcal{S}_{K'}$ over $\mathcal{O}_{E,(v)}$ in the sense of Definition 6.5.1. Then Sh_K admits an ICM over $\mathcal{O}_{E,(v)}$. Therefore:*

- (1) *If p is large enough, then Sh_K admits an étale ICM over $\mathcal{O}_{E,(v)}$.*
- (2) *If (G, X) is of pre-abelian type or if p is large enough and G^{ad} is anisotropic, then Sh_K admits a crystalline ICM over $\mathcal{O}_{E,(v)}$.*

Proof. The map $\text{Sh}_{K'} \rightarrow \text{Sh}_K$ is a finite Galois cover of DM stacks over E . Let Δ be its Galois group: the action of Δ on $\text{Sh}_{K'}$ extends to an action on $\mathcal{S}_{K'}$ by Theorem 6.5.3 and the fact that $\mathbf{Et}_{K',p}$ is Δ -equivariant. Therefore, we can set $\mathcal{S}_K = [\mathcal{S}_{K'}/\Delta]$: This is a DM stack over $\mathcal{O}_{E,(v)}$, and is clearly a stacky ICM for Sh_K . $\square$

f_integers

8.5. Integral canonical models over global rings of integers.

Setup 8.5.1. In this subsection, we will drop our fixed prime p . Instead, we take an open subgroup $K \subset G(\mathbb{A}_f)$ and define Ξ_K to be the set of primes ℓ such that $K \cap G(\mathbb{Q}_\ell)$ is *not* hyperspecial. For $S \supseteq \Xi_K$ a set of primes, there exists a reductive group scheme $\mathcal{G}^S$ over $\mathbb{Z}[S^{-1}]$ with $\mathcal{G}^S(\mathbb{Z}_p) = K_p$ for $p \notin S$.

unramified

Remark 8.5.2. If G admits a reductive model $\mathcal{G}_{\mathbb{Z}}$ over $\mathbb{Z}$, and we take $K = \mathcal{G}_{\mathbb{Z}}(\widehat{\mathbb{Z}})$, then Ξ_K is empty. For instance if $G = \mathrm{GSp}_{2g, \mathbb{Q}}$, then we can take $\mathcal{G}_{\mathbb{Z}} = \mathrm{GSp}_{2g, \mathbb{Z}}$. In this case, the associated Shimura stack Sh_K is the moduli stack $\mathcal{A}_{g, \mathbb{Q}}$ over $\mathbb{Q}$ of principally polarized abelian varieties of dimension g .

Definition 8.5.3 (Global integral canonical models). Let S be a set of primes containing Ξ_K . An integral model $\mathcal{S}_K$ for Sh_K over $\mathcal{O}_E[S^{-1}]$ is a **crystalline** (resp. **an étale**) **integral canonical model** for Sh_K if, for every $p \notin S$ and every place $v|p$ of E , $\mathcal{S}_K \otimes_{\mathcal{O}_E} \mathcal{O}_{E, (v)}$ is a crystalline (resp. an étale) integral canonical model for Sh_K over $\mathcal{O}_{E, (v)}$. Either type of model will be referred to as an **integral canonical model** or **ICM**.

Let us list some properties of global ICMs that follow easily from the local versions.

global_ICMs

Proposition 8.5.4 (Mapping property for global ICMs). *Let $\mathcal{Y}$ be an excellent separated normal Deligne–Mumford stack flat over $\mathcal{O}_E[S^{-1}]$ with generic fiber Y over E . Suppose that we are given a map $f: Y \rightarrow \mathrm{Sh}_K$ with the following properties:*

- (1) *For every prime $p \notin S$ and every $v|p$, the map $Y \rightarrow \mathrm{Loc}_{\mathcal{G}(\mathbb{Z}_p)}$ classifying $\mathbf{Et}_{K, p}$ extends to a map $\mathcal{Y} \otimes_{\mathcal{O}_E} \mathcal{O}_{E, (v)} \rightarrow \mathrm{BT}_{\infty}^{\mathcal{G}_{\mathbb{Z}_p}^S, -\mu_v, \mathrm{alg}}$.*
- (2) *For all primes ℓ , $\mathbf{Et}_{K, \ell}|_Y$ extends to a K_ℓ^c -local system on $\mathcal{Y}[\ell^{-1}]$ represented by pro-algebraic spaces over $\mathcal{O}_{E, (v)}$.*

Then f extends to a map $\mathcal{Y} \rightarrow \mathcal{S}_K$.

ctoriality

Proposition 8.5.5 (Uniqueness and functoriality for global ICMs). *An ICM $\mathcal{S}_K$ over $\mathcal{O}_E[S^{-1}]$ is determined uniquely up to unique isomorphism. In fact, given a map of Shimura data $f: (G_1, X_1) \rightarrow (G_2, X_2)$ with $f(K_1) \subset K_2$, and ICMs $\mathcal{S}_{K_i}$ over $\mathcal{O}_{E_i}[S^{-1}]$, the canonical map $\mathrm{Sh}_{K_1} \rightarrow \mathrm{Sh}_{K_2} \otimes_{E_2} E_1$ extends to a map*

$$\mathcal{S}_{K_1} \rightarrow \mathcal{S}_{K_2} \otimes_{\mathcal{O}_{E_2}} \mathcal{O}_{E_1}.$$

_existence

Theorem 8.5.6 (Existence of global ICMs). (1) *If (G, X) is of pre-abelian type, then Sh_K admits an integral model $\mathcal{S}_K$ over $\mathcal{O}_E[\Xi_K^{-1}]$ that is both a crystalline and étale ICM.*
 (2) *In general, Sh_K admits an étale ICM over $\mathcal{O}_E[S^{-1}]$ for some large enough set of primes S containing Ξ_K . This model is also proper and a crystalline ICM if G^{ad} is anisotropic.*

lobal_icms

Remark 8.5.7 (Global ICMs of Hodge type). Suppose that (G, X) is of Hodge type and that we have an embedding $(G, X) \hookrightarrow (\mathrm{GSp}(V), S^\pm(V))$ into a Siegel Shimura datum associated with a symplectic space V over $\mathbb{Q}$. Then the proof of Theorem C shows that the global ICM $\mathcal{S}_K$ for Sh_K is obtained by taking the normalization of a moduli stack of polarized abelian varieties over $\mathcal{O}_E[S^{-1}]$ in Sh_K . In particular, $\mathcal{A}_g$, the stack of principally polarized abelian varieties, is a global ICM for its generic fiber.

Remark 8.5.8 (Global ICMs of abelian type). For Shimura varieties of abelian type, a version of statement (1) in the theorem above is due to Lovering [61]. His proof amounts to a careful finite-level analysis of a combination of Deligne’s quotient construction from [20] with Kisin’s twisting construction from [47], whereas the result here is an immediate consequence of our local characterization of ICMs at finite level.

ric_levels

8.6. Integral canonical models at parahoric level. Here we explain that any unramified Shimura datum $(G, \mathcal{G}, X)$ which admits an integral canonical in the sense of Definition 6.5.1 automatically admits a Pappas–Rapoport integral canonical model as studied in [73].

We then apply results of Takaya from [88] to obtain Pappas–Rapoport integral canonical models for subhyperspecial (e.g., Iwahori) level. Finally, we give an analogue of the mapping property from Theorem 6.5.3 in this parahoric situation.

Notation 8.6.1. In the following we denote by $\mathbf{Perf}_A$, for a p -adically complete ring A , the category of characteristic p perfectoid spaces S equipped with a map $S \rightarrow \mathrm{Spd}(A)$. For any other undefined piece of notation or terminology, we direct the reader to the comprehensive discussion in [73, §2-3].

Definition 8.6.2 (Tame local Shimura datum). A **tame local Shimura datum** is a triple $(\mathcal{G}, b, \{\mu\})$ where $\mathcal{G}$ is a parahoric group $\mathbb{Z}_p$ -scheme with generic fiber G , $\{\mu\}$ is a conjugacy class of minuscule cocharacters of $G_{\overline{\mathbb{Q}_p}}$, and b is an element of $G(\overline{\mathbb{Q}_p})$ inducing an element of $B(G, -\mu)$.

Definition 8.6.3 (Local Shimura variety). Given a tame local Shimura datum $(\mathcal{G}, b, \{\mu\})$ with reflex field F , we obtain a presheaf

$$\mathcal{M}_{\mathcal{G}, b, \{\mu\}}^{\mathrm{int}} : \mathbf{Perf}_{\mathcal{O}_{\tilde{F}}} \rightarrow \mathbf{Set},$$

assigning to $S \rightarrow \mathrm{Spd}(\mathcal{O}_{\tilde{F}})$ the set of isomorphism classes of tuples $(S^\sharp, \mathcal{P}, \phi_{\mathcal{P}}, i_r)$, where:

- $S^\sharp$ is the untile of S over $\mathcal{O}_{\tilde{E}}$ associated with $S \rightarrow \mathrm{Spd}(\mathcal{O}_{\tilde{F}})$,
- $(\mathcal{P}, \phi_{\mathcal{P}})$ is a $\mathcal{G}$ -shtuka on S with one leg along $S^\sharp$ bounded by $\{\mu\}$ (see [73, Definition 2.4.3]).

By [83, §25.1], the presheaf $\mathcal{M}_{\mathcal{G}, b, \{\mu\}}^{\mathrm{int}}$ is a small v -sheaf (in the sense of [81, Definition 12.1]), which is called the **integral local Shimura variety** associated to the tame local Shimura datum $(\mathcal{G}, b, \{\mu\})$.

Definition 8.6.4 (Tame Shimura datum/tuple). A **tame Shimura datum** is a triple $(G, \mathcal{G}, X)$ where (G, X) is a Shimura datum and $\mathcal{G}$ is a parahoric model of $G_{\mathbb{Q}_p}$. A **tame Shimura tuple** is a quadruple $(G, \mathcal{G}, X, K)$ where $(G, \mathcal{G}, X)$ is a tame Shimura datum and $K \subseteq G(\mathbb{A}_f)$ is a compact open subgroup with $K = K_p K^p$ and where $K_p = \mathcal{G}(\mathbb{Z}_p)$. We say $(G, \mathcal{G}, X, K)$ is **neat** if K is.

Remark 8.6.5. Fix a tame Shimura tuple $(G, \mathcal{G}, X, K)$ with reflex field E and a place v of E of lying over p . In the following we shall abusively write Sh_K for $\mathrm{Sh}_K(G, X)_{E_v}$. We will write $\{\mu_v\}$ for the conjugacy class of $G_{\overline{E_v}}$ determined by the Hodge cocharacter of (G, X) .

As in Construction 6.1.6 for one may build proétale $\mathcal{G}^c(\mathbb{Z}_p)$ -torsor $\mathbf{Et}_{K,p}$ on Sh_K . Here $\mathcal{G}^c$ is the parahoric model of G^c induced by the parahoric model $\mathcal{G}$ of G (e.g., see [17, §4.1]). As in the reductive case, we treat $\{\mu_v\}$ also as a conjugacy class of cocharacters of $G_{\overline{E_v}}^c$.

Definition 8.6.6 (Generic shtuka associated to a tame Shimura datum). We denote by $\mathcal{P}_{K,E_v}$ the $\mathcal{G}^c$ -shtuka bounded by $\{-\mu_v\}$ on $\mathrm{Sh}_K^\diamond \rightarrow \mathrm{Spd}(E)$ defined by $U_{\mathrm{sht}}(\mathbf{Et}_{K,p})$ with notation as in [37, §3.2.2].

Construction 8.6.7 (Formal neighborhoods of local Shimura varieties associated to $\overline{k}(v)$ -points of an integral model of Sh_K). Let $\mathcal{S}_K$ be flat normal $\mathcal{O}_{E_v}$ -model of Sh_K equipped with an extension $\mathcal{P}_K$ of $\mathcal{P}_{K,E_v}$ to $\mathcal{S}_K^\diamond$ (see [73, Definition 2.1.9]).²⁹ For any point x of $\mathcal{S}_K(\overline{k}(v))$, the pullback $x^* \mathcal{P}_K$ defines a $\mathcal{G}^c$ -shtuka over $\mathrm{Spd}(\overline{k}(v))$ which by [73, Example 2.4.9] naturally determines an object $(\mathcal{P}_x, \phi_x)$ of $\mathrm{BT}_\infty^{\mathcal{G}^c, -\mu_v}(\overline{k}(v))$ and by Construction 3.3.1 an element b_x in $G(\overline{\mathbb{Q}_p})$ inducing an element of $B(G^c, \mu_v^{-1})$.

The triple $(\mathcal{G}^c, b_x, \{\mu_v\})$ defines a parahoric local Shimura datum, and we can consider the integral local Shimura variety $\mathcal{M}_{\mathcal{G}^c, b_x, \{\mu_v\}}^{\mathrm{int}}$. Moreover, the pair $(\mathcal{P}_x, \mathrm{id})$ determines a point x_0 of the affine Deligne–Lusztig set $X_{\mathcal{G}^c}(b_x, \mu_v^{-1})(\overline{k}(v))$ (e.g., as in [73, Definition 3.3.1]), and so a point of $\mathcal{M}_{\mathcal{G}^c, b_x, \{\mu_v\}}^{\mathrm{int}}(\overline{k}(v))$ via the identification given in [28, Proposition 2.61.(1)]. One may then consider the formal neighborhood $(\mathcal{M}_{\mathcal{G}^c, b_x, \{\mu_v\}}^{\mathrm{int}})_{/x_0}^\wedge$ as in [29, Definition 4.18].

Definition 8.6.8 (Pappas–Rapoport integral models). Fix a tame Shimura datum $(G, \mathcal{G}, X)$. Then, a system $\{\mathcal{S}_K\}$ of normal flat $\mathcal{O}_{E_v}$ -models of Sh_K , as one ranges over all neat tame Shimura tuples $(G, \mathcal{G}, X, K)$, is a **Pappas–Rapoport integral canonical model** for $(G, \mathcal{G}, X)$ if it satisfies conditions (i), (ii), (iii) and (iv) of [17, Definition 4.3].

We say a Pappas–Rapoport integral canonical model is **crystalline** if, for any K , and for any complete discrete valuation field F with mixed characteristic $(0, p)$ with perfect residue field, the following are equivalent for $x \in \mathrm{Sh}_K(F)$:

²⁹Note that $\mathcal{P}_K$ is necessarily bounded by $\{\mu\}$ by [15, Lemma 2.1].

- (1) $x \in \mathcal{S}_K(\mathcal{O}_F)$;
- (2) $\mathbf{Et}_{K,p,x}$ is crystalline;
- (3) $\mathbf{Et}_{K,p,x}$ is potentially crystalline.

re-PR ICMS

Proposition 8.6.9. *Suppose that $(G, \mathcal{G}, X)$ is an unramified Shimura datum admitting an étale ICM $\mathcal{S}_K$ for each (equiv. any) neat unramified Shimura tuple $(G, \mathcal{G}, X, K)$. Then:*

- (1) *The collection $\{\mathcal{S}_K\}$ is a Pappas–Rapoport integral canonical model for $(G, \mathcal{G}, X)$.*
- (2) *If $(G, \mathcal{G}, X)$ is a crystalline ICM, then the Pappas–Rapoport integral model $\{\mathcal{S}_K\}$ is crystalline.*

Proof. For (1), we check the conditions from [17, Definition 4.3]. For Condition (i), see Remark 6.5.7. Condition (ii) follows from Theorem ???. For condition (iii), observe that by definition of being an étale ICM, each $\mathcal{S}_K$ carries a prismatic F -crystal in the sense of [37, Definition 3.18] with de Rham local system $\mathbf{Et}_{K,p}$. Thus, applying the shtuka realization functor from [37, Construction 3.19] to this prismatic F -crystal yields a shtuka $\mathcal{P}_K$ on $\mathcal{S}_K^{\diamond/}$. That $\mathcal{P}_K$ models $\mathcal{P}_{K,E}$ is clear as $(\mathcal{P}_K)_E$ is (by definition) $U_{\text{sht}}(\mathbf{Et}_{K,p}) = \mathcal{P}_{K,E}$. Finally, to check (iv), we may use the formal étaleness of the syntomic realization ϖ and [39, Proposition 3.32] (see also [26, §10.2]) to reduce ourselves to constructing

$$\Theta_x: \left(\mathcal{M}_{\mathcal{G}^c, b_x, \{\mu_v\}}^{\text{int}} \right)_{/x_0}^{\wedge} \xrightarrow{\sim} \text{Spf}(R_{\mathcal{G}, \{\mu_v\}}),$$

where the target is as in loc. cit., and with the property that pullback of $\mathcal{P}_K$ on the target is the universal $\mathcal{G}^c$ -shtuka on the source. But, this is the content of [40, Theorem 5.3.5].

Moreover, (2) is now immediate from the definition of a crystalline ICM; see also Corollary ??. $\square$

Definition 8.6.10 (Subhyperspecial parahoric). Let $F/\mathbb{Q}_p$ be a finite extension and let H be a reductive group over F . A subgroup of $H(F)$ is called **subhyperspecial parahoric** if it is a parahoric subgroup in the sense of [41, Definition 4.1.4] contained in $\mathcal{H}(\mathcal{O}_F)$ for a reductive $\mathcal{O}_F$ -model $\mathcal{H}$ of H .

Remark 8.6.11 (Explicit description of subhyperspecial parahorics). For a reductive model $\mathcal{H}$ of H over $\mathcal{O}_F$, the subhyperspecial parahorics contained in $\mathcal{H}(\mathcal{O}_F)$ are precisely those of the form $\text{red}^{-1}(P(k))$ where k is the residue field of F , $P \subseteq \mathcal{H}_k$ is a parabolic subgroup, and

$$\text{red}: \mathcal{H}(\mathcal{O}_F) \rightarrow \mathcal{H}(k)$$

is the reduction map. Equivalently, the subhyperspecial parahoric subgroups of $H(K)$ are $\mathcal{H}'(\mathcal{O}_K)$ for a parahoric group $\mathcal{O}_K$ -scheme $\mathcal{H}'$ (see [17, Definition 2.6]) obtained as the dilatation (see [41, §A.5]) of a reductive $\mathcal{O}_F$ -model $\mathcal{H}$ along a parabolic subgroup $P \subseteq \mathcal{H}_k$. We call these $\mathcal{O}_K$ -models $\mathcal{H}'$ of H **parabolic dilatations of $\mathcal{H}$** . We will just call $\mathcal{H}'$ a **parabolic dilatation** if $\mathcal{H}$ is implicit.

Example 8.6.12 (Iwahoris are subhyperspecial). Considering $\text{red}^{-1}(B(k))$ for $B \subseteq \mathcal{H}_k$ a Borel subgroup we obtain the (or more precisely a) Iwahori subgroup of $H(F)$ as in [41, Definition 4.1.3].

minorizing

Proposition 8.6.13. *Let $(G, \mathcal{G}, X)$ be an unramified Shimura datum satisfying (SV5) and admitting étale (resp. étale and crystalline) ICMs. Then, for any parabolic dilatation $\mathcal{G}'$ of $\mathcal{G}$, the tame Shimura datum $(G, \mathcal{G}', X)$ admits a Pappas–Rapoport integral canonical model (resp. crystalline such model).*

Proof. This is immediate from Proposition 8.6.9 and [88, Theorem 6.20]. $\square$

end-large-p

Corollary 8.6.14. *Suppose that $(G, \mathcal{G}', X)$ is a tame parahoric Shimura datum with $\mathcal{G}'$ a parabolic dilatation. Suppose further that either:*

- (1) *(G, X) is of pre-abelian type;*
- (2) *p is sufficiently large and G is anisotropic mod center.*

Then, $(G, \mathcal{G}', X)$ admits a crystalline Pappas–Rapoport integral canonical model.

Remark 8.6.15 (Agreement with Kisin–Pappas–Zhou). When $(G, \mathcal{G}', X)$ is of abelian type and $p > 3$ the models constructed by Corollary 8.6.14 agree with the Kisin–Pappas–Zhou models constructed in [46] and [50]. Indeed, this follows from the unicity of Pappas–Rapoport integral canonical models (see [73, Theorem 4.3.1]) and the results of [17], building on previous work from [73] and [16].

Definition 8.6.16 (Moduli stack of $(\mathcal{G}, \mu)$ -shtukas). Let notation be as in Definition 8.6.3. We denote by $\mathrm{Sht}_{\mathcal{G}, \{\mu\}}$ the v -stack on $\mathbf{Perf}_{\mathcal{O}_F}$ associating to $S \rightarrow \mathrm{Spd}(\mathcal{O}_F)$ the groupoid of $\mathcal{G}$ -shtukas over S with leg along $S^\sharp$ bounded by $\{\mu\}$.

The proof of the following result is along the same lines as that of Theorem 6.5.3. One has to replace the use of Theorem 4.1.5 with [73, Theorem 2.7.7], as well as use [83, Proposition 18.4.1].

Theorem 8.6.17. *Let $(G, \mathcal{G}, X, K)$ be a neat tame Shimura tuple admitting a crystalline Pappas–Rapoport integral canonical model $\{\mathcal{S}_K\}$. Suppose that $\mathcal{X}$ is a flat, η -normal algebraic space of finite type over $\mathcal{O}_{E, (v)}$. Then a map $\mathcal{X}_\eta \rightarrow \mathrm{Sh}_K$ extends to a map $\mathcal{X} \rightarrow \mathcal{S}_K$ if and only if the composition $\mathcal{X}_\eta^\diamond \rightarrow \mathrm{Sh}_K^\diamond \rightarrow \mathrm{Sht}_{\mathcal{G}, \{-\mu_v\}}$ lifts to a map $\mathcal{X}^\diamond \rightarrow \mathrm{Sht}_{\mathcal{G}, \{-\mu_v\}}$.*

REFERENCES

- [1] Piotr Achinger, Marcin Lara, and Alex Youcis. “Specialization for the pro-étale fundamental group”. In: *Compos. Math.* 158.8 (Aug. 2022), pp. 1713–1745. URL: <http://dx.doi.org/10.1112/S0010437X2200759X> (cited on pp. 26, 45).
- [2] Piotr Achinger and Alex Youcis. “Beauville-Laszlo gluing of algebraic spaces”. In: (2024). eprint: 2410.20500. URL: <https://arxiv.org/abs/2410.20500> (cited on pp. 29, 51).
- [3] Johannes Anschütz and Arthur-César Le Bras. “Prismatic Dieudonné theory”. In: *Forum Math. Pi* 11 (2023), Paper No. e2, 92 (cited on p. 12).
- [4] Benjamin Bakker, Ananth N Shankar, and Jacob Tsimmerman. “Integral canonical models of exceptional Shimura varieties”. In: (2025). eprint: 2405.12392. URL: <https://arxiv.org/abs/2405.12392> (cited on pp. 1, 5, 7, 46, 55).
- [5] Bhargav Bhatt. *Prismatic F -gauges*. URL: <https://www.math.ias.edu/~bhatt/teaching/mat549f22/lectures.pdf> (cited on pp. 8, 10, 13, 23–24).
- [6] Bhargav Bhatt and Jacob Lurie. “The prismatic formal schemes”. In: (2022). eprint: 2201.06124. URL: <https://arxiv.org/abs/2201.06124> (cited on p. 8).
- [7] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. “Topological Hochschild homology and integral p -adic Hodge theory”. In: *Publ. Math. Inst. Hautes Études Sci.* 129 (2019), pp. 199–310. ISSN: 0073-8301, 1618-1913. DOI: 10.1007/s10240-019-00106-9. URL: <https://doi.org/10.1007/s10240-019-00106-9> (cited on p. 9).
- [8] Bhargav Bhatt and Peter Scholze. “Prismatic F -crystals and crystalline Galois representations”. In: *Camb. J. Math.* 11.2 (2023), pp. 507–562 (cited on pp. 9–10, 25).
- [9] Bhargav Bhatt and Peter Scholze. “Prisms and prismatic cohomology”. In: *Ann. of Math. (2)* 196.3 (2022), pp. 1135–1275. ISSN: 0003-486X, 1939-8980. DOI: 10.4007/annals.2022.196.3.5. URL: <https://doi-org.myaccess.library.utoronto.ca/10.4007/annals.2022.196.3.5> (cited on p. 9).
- [10] Spencer Bloch and Kazuya Kato. “ L -functions and Tamagawa numbers of motives”. In: *The Grothendieck Festschrift, Vol. I*. Vol. 86. Progr. Math. Birkhäuser Boston, Boston, MA, 1990, pp. 333–400. ISBN: 0-8176-3427-4 (cited on p. 14).
- [11] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*. Vol. 21. Ergebnisse der Mathematik und ihrer Grenzgebiete. Berlin: Springer-Verlag, 1990 (cited on p. 3).
- [12] Alexis Bouthier and Kestutis Česnavičius. “Torsors on loop groups and the Hitchin fibration”. In: *Ann. Sci. Éc. Norm. Supér. (4)* 55.3 (2022), pp. 791–864 (cited on p. 28).
- [13] Michael Broshi. “ G -torsors over a Dedekind scheme”. In: *J. Pure Appl. Algebra* 217.1 (2013), pp. 11–19. ISSN: 0022-4049. DOI: 10.1016/j.jpaa.2012.01.011. URL: <https://doi.org/10.1016/j.jpaa.2012.01.011> (cited on p. 17).
- [14] Robert Coleman and Adrian Iovita. “The Frobenius and monodromy operators for curves and abelian varieties”. In: *Duke Math. J.* 97.1 (1999), pp. 171–215 (cited on p. 5).
- [15] Patrick Daniels. “Canonical integral models for Shimura varieties of toral type”. In: (2023). URL: <https://arxiv.org/abs/2207.09513> (cited on p. 60).

- [16] Patrick Daniels, Pol van Hoften, Dongryul Kim, and Mingjia Zhang. “On a conjecture of Pappas and Rapoport”. In: (2025). arXiv: [2403.19771](https://arxiv.org/abs/2403.19771) [math.NT]. URL: <https://arxiv.org/abs/2403.19771> (cited on pp. 4, 61).
- [17] Patrick Daniels and Alexander Youcis. “Canonical integral models for Shimura varieties of abelian type”. In: *Forum Math. Sigma* 13 (2025), Paper No. e69, 47. ISSN: 2050-5094. DOI: [10.1017/fms.2025.27](https://doi.org/10.1017/fms.2025.27). URL: <https://doi.org/10.1017/fms.2025.27> (cited on pp. 60–61).
- [18] Aise Johan de Jong. “Crystalline Dieudonné module theory via formal and rigid geometry”. In: *Inst. Hautes Études Sci. Publ. Math.* 82.1 (1995), pp. 5–96 (cited on p. 36).
- [19] Aise Johan de Jong. “Homomorphisms of Barsotti-Tate groups and crystals in positive characteristic”. In: *Invent. Math.* 134.2 (1998), pp. 301–333. URL: <http://dx.doi.org/10.1007/s002220050266> (cited on p. 56).
- [20] Pierre Deligne. “Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques”. In: *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*. Proc. Sympos. Pure Math., XXXIII. Amer. Math. Soc., Providence, R.I., 1979, pp. 247–289 (cited on pp. 40–43, 46, 52, 59).
- [21] Hansheng Diao, Kai-Wen Lan, Ruochuan Liu, and Xinwen Zhu. “Logarithmic Riemann–Hilbert correspondences for rigid varieties”. In: *J. Amer. Math. Soc.* 36.2 (2022), pp. 483–562. URL: <http://dx.doi.org/10.1090/jams/1002> (cited on pp. 41–42).
- [22] Hansheng Diao and Zijian Yao. “Monodromy and rigidity of crystalline local systems”. In: (2025). arXiv: [2509.19813](https://arxiv.org/abs/2509.19813) [math.AG]. URL: <https://arxiv.org/abs/2509.19813> (cited on p. 54).
- [23] Vladimir Drinfeld. “Prismatization”. In: (2022). eprint: [2005.04746](https://arxiv.org/abs/2005.04746). URL: <https://arxiv.org/abs/2005.04746> (cited on p. 8).
- [24] Vladimir Drinfeld. “Toward Shimurian analogs of Barsotti-Tate groups”. In: (2024). URL: <https://arxiv.org/abs/2309.02346> (cited on pp. 1–2).
- [25] Heng Du, Tong Liu, Yong Suk Moon, and Koji Shimizu. “Completed prismatic F -crystals and crystalline $\mathbb{Z}_p$ -local systems”. In: *Compositio Mathematica* 160.5 (2024), pp. 1101–1166. URL: <http://dx.doi.org/10.1112/S0010437X24007097> (cited on pp. 10–11).
- [26] Zachary Gardner and Keerthi Madapusi. “An algebraicity conjecture of Drinfeld and the moduli of p -divisible groups”. In: *arXiv preprint arXiv:2412.10226* (Dec. 2024) (cited on pp. 1–2, 7–9, 12, 14–16, 18–23, 47, 53, 61).
- [27] Qëndrim R. Gashi. “On a conjecture of Kottwitz and Rapoport”. In: *Ann. Sci. Éc. Norm. Supér. (4)* 43.6 (2010), pp. 1017–1038 (cited on p. 19).
- [28] Ian Gleason. *On the geometric connected components of moduli spaces of p -adic shtukas and local Shimura varieties*. 2025. arXiv: [2107.03579](https://arxiv.org/abs/2107.03579) [math.AG]. URL: <https://arxiv.org/abs/2107.03579> (cited on p. 60).
- [29] Ian Gleason. “Specialization maps for Scholze’s category of diamonds”. In: *Math. Ann.* 391.2 (2025), pp. 1611–1679. ISSN: 0025-5831, 1432-1807. DOI: [10.1007/s00208-024-02952-3](https://doi.org/10.1007/s00208-024-02952-3). URL: <https://doi.org/10.1007/s00208-024-02952-3> (cited on p. 60).
- [30] Ulrich Görtz and Torsten Wedhorn. *Algebraic geometry I. Schemes—with examples and exercises*. Springer Studium Mathematik—Master. Second edition [of 2675155]. Springer Spektrum, Wiesbaden, [2020] ©2020, pp. vii+625. ISBN: 978-3-658-30732-5; 978-3-658-30733-2. DOI: [10.1007/978-3-658-30733-2](https://doi.org/10.1007/978-3-658-30733-2). URL: <https://doi.org/10.1007/978-3-658-30733-2> (cited on p. 32).
- [31] Alexander Grothendieck. *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. Ed. by Yves Laszlo. Vol. 4. Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Société Mathématique de France, Paris, 2005, pp. x+208 (cited on p. 33).
- [32] Alexander Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*. Vol. 3. Documents Mathématiques (Paris). Paris: Société Mathématique de France, 2003 (cited on pp. 39–40).
- [33] Haoyang Guo and Shizhang Li. “Frobenius height of prismatic cohomology with coefficients”. In: (2025). arXiv: [2309.06663](https://arxiv.org/abs/2309.06663) [math.AG]. URL: <https://arxiv.org/abs/2309.06663> (cited on pp. 10, 13).

- [34] Haoyang Guo and Emanuel Reinecke. “A prismatic approach to crystalline local systems”. In: *Invent. Math.* 236.1 (2024), pp. 17–164. URL: <https://doi.org/10.1007/s00222-024-01238-4> (cited on pp. 10–11, 13).
- [35] Haoyang Guo and Ziquan Yang. “Pointwise criteria of p -adic local systems”. In: (2024). arXiv: [2409.19742](https://arxiv.org/abs/2409.19742) [math.AG]. URL: <https://arxiv.org/abs/2409.19742> (cited on pp. 11, 32, 54).
- [36] Roland Huber. *Étale cohomology of rigid analytic varieties and adic spaces*. Vol. E30. Aspects of Mathematics. Friedr. Vieweg & Sohn, Braunschweig, 1996, pp. x+450 (cited on p. 29).
- [37] Naoki Imai, Hiroki Kato, and Alex Youcis. “A Tannakian framework for prismatic F -crystals”. In: (2024). arXiv: [2406.08259](https://arxiv.org/abs/2406.08259) [math.NT]. URL: <https://arxiv.org/abs/2406.08259> (cited on pp. 6, 10–12, 15, 24, 26, 33, 60–61).
- [38] Naoki Imai, Hiroki Kato, and Alex Youcis. *An integral analogue of Fontaine’s crystalline functor*. 2025. arXiv: [2504.16282](https://arxiv.org/abs/2504.16282) [math.NT]. URL: <https://arxiv.org/abs/2504.16282> (cited on pp. 6, 23–24, 33).
- [39] Naoki Imai, Hiroki Kato, and Alex Youcis. “The prismatic realization functor for Shimura varieties of abelian type”. In: (2023). URL: <https://arxiv.org/abs/2310.08472> (cited on pp. 1–3, 5–6, 9–10, 15, 31, 33, 41, 61).
- [40] Kazuhiro Ito. “Deformation theory for prismatic G -displays”. In: (2023). eprint: [2306.05361](https://arxiv.org/abs/2306.05361). URL: <https://arxiv.org/abs/2306.05361> (cited on pp. 15, 61).
- [41] Tasho Kaletha and Gopal Prasad. *Bruhat-Tits theory—a new approach*. Vol. 44. New Mathematical Monographs. Cambridge University Press, Cambridge, 2023, pp. xxx+718. ISBN: 978-1-108-83196-3 (cited on p. 61).
- [42] Nicholas Katz. “Serre-Tate local moduli”. In: *Surfaces Algébriques (Orsay, 1976-78)*. Vol. 868. Lecture Notes in Mathematics. Berlin: Springer, 1981, pp. 138–202 (cited on p. 22).
- [43] Nicholas Katz. “Travaux de Dwork”. French. In: (1973) (cited on p. 35).
- [44] Kiran S. Kedlaya and Ruochuan Liu. “Relative p -adic Hodge theory: foundations”. In: *Astérisque* 371 (2015), p. 239. ISSN: 0303-1179,2492-5926 (cited on p. 31).
- [45] Wansu Kim and Keerthi Madapusi. “2-adic integral canonical models”. In: *Forum of Mathematics, Sigma* 4 (Jan. 2016) (cited on p. 3).
- [46] M Kisin and G Pappas. “Integral models of Shimura varieties with parahoric level structure”. In: *Publications mathématiques de l’IHÉS* 128.1 (Nov. 2018), pp. 121–218 (cited on p. 61).
- [47] Mark Kisin. “Integral models for Shimura varieties of abelian type”. In: *J. Amer. Math. Soc.* 23.4 (2010), pp. 967–1012 (cited on pp. 3, 5, 32, 43, 45, 49, 53, 59).
- [48] Mark Kisin. “Mod p points on Shimura varieties of abelian type”. In: *J. Amer. Math. Soc.* 30.3 (2017), pp. 819–914 (cited on pp. 4, 6–7, 17).
- [49] Mark Kisin, Keerthi Madapusi Pera, and Sug Woo Shin. “Honda–Tate theory for Shimura varieties”. In: *Duke Math. J.* 171.7 (2022), pp. 1559–1614 (cited on pp. 6, 57).
- [50] Mark Kisin, Georgios Pappas, and Rong Zhou. *Integral models of Shimura varieties with parahoric level structure, II*. 2024. arXiv: [2409.03689](https://arxiv.org/abs/2409.03689) [math.NT]. URL: <https://arxiv.org/abs/2409.03689> (cited on p. 61).
- [51] Mark Kisin, Sug Woo Shin, and Yihang Zhu. “The stable trace formula for Shimura varieties of abelian type”. In: (2021). URL: <https://arxiv.org/abs/2110.05381> (cited on pp. 40–41, 46).
- [52] Christian Klevdal and Stefan Patrikis. “Compatibility of canonical ℓ -adic local systems on adjoint Shimura varieties”. In: *Invent. Math.* 242.1 (2025), pp. 305–335. URL: <https://doi.org/10.1007/s00222-025-01357-6> (cited on pp. 7, 55).
- [53] Robert E Kottwitz. “Isocrystals with additional structure”. In: *Compos. Math.* 56.2 (1985), pp. 201–220. URL: <https://eudml.org/doc/89735> (cited on p. 7).
- [54] Robert E Kottwitz. “Shimura varieties and twisted orbital integrals”. In: *Math. Ann.* 269.3 (1984), pp. 287–300. URL: <http://link.springer.com/article/10.1007/BF01450697> (cited on pp. 21, 43).
- [55] Eike Lau. “Higher frames and G -displays”. In: *Algebra Number Theory* 15.9 (2021), pp. 2315–2355 (cited on pp. 8, 25).
- [56] Dong Uk Lee. “Nonemptiness of Newton strata of Shimura varieties of Hodge type”. In: *Algebra Number Theory* 12.2 (2018), pp. 259–283. URL: <http://dx.doi.org/10.2140/ant.2018.12.259> (cited on p. 6).

- | | | |
|------------|------|--|
| e_madapusi | [57] | Si Ying Lee and Keerthi Madapusi. “ p -isogenies with G -structure and their applications”. In: (). in preparation (cited on pp. 3, 7). |
| Liu2017-yz | [58] | Ruochuan Liu and Xinwen Zhu. “Rigidity and a Riemann–Hilbert correspondence for p -adic local systems”. In: <i>Invent. Math.</i> 207.1 (Jan. 2017), pp. 291–343. URL: http://dx.doi.org/10.1007/s00222-016-0671-7 (cited on pp. 24, 36, 42). |
| LiuMoon | [59] | Tong Liu and Yong Suk Moon. “Relative crystalline representations and p -divisible groups in the small ramification case”. In: <i>Algebra Number Theory</i> 14.10 (2020), pp. 2773–2789 (cited on p. 34). |
| ing2017-fy | [60] | Tom Lovering. “Filtered F -crystals on Shimura varieties of abelian type”. In: (Feb. 2017). arXiv: 1702.06611 [math.NT]. URL: http://arxiv.org/abs/1702.06611 (cited on p. 40). |
| ing2017-me | [61] | Tom Lovering. “Integral canonical models for automorphic vector bundles of abelian type”. In: <i>Algebra Number Theory</i> 11.8 (Oct. 2017), pp. 1837–1890 (cited on pp. 40, 52, 59). |
| rie_thesis | [62] | Jacob Lurie. <i>Derived algebraic geometry</i> . Thesis (Ph.D.)–Massachusetts Institute of Technology. ProQuest LLC, Ann Arbor, MI, 2004, (no paging) (cited on pp. 28, 37–38). |
| rie2009-oc | [63] | Jacob Lurie. <i>Higher Topos Theory</i> . en. Princeton University Press, July 2009. URL: https://www.degruyter.com/document/doi/10.1515/9781400830558/html (cited on p. 27). |
| rie2018-kh | [64] | Jacob Lurie. “Spectral algebraic geometry”. In: <i>preprint</i> (2018). URL: https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf (cited on p. 12). |
| erre-tate | [65] | Keerthi Madapusi. “Ordinary loci: a stack theoretic approach”. In: (2025). in preparation (cited on pp. 7, 21). |
| p:toroidal | [66] | Keerthi Madapusi. “Toroidal compactifications of integral models of Shimura varieties of Hodge type”. In: <i>Ann. Sci. Ec. Norm. Super.</i> 52.1 (2019), pp. 393–514 (cited on p. 54). |
| usi2025-ay | [67] | Keerthi Madapusi and Shubhodip Mondal. “Perfect F -gauges and finite flat group schemes”. In: <i>arXiv [math.NT]</i> (2025). URL: http://arxiv.org/abs/2509.01573 (cited on pp. 8, 14–16, 22). |
| ama2016-gc | [68] | Andrea Marrama. “A purity theorem for torsors”. MA thesis. Universiteit Duisburg-Essen, 2016. URL: https://www.math.u-bordeaux.fr/~ybilu/algant/documents/theses/Marrama.pdf (cited on p. 34). |
| ertonKisin | [69] | Emerton Matthew and Mark Kisin. “Extensions of crystalline representations”. URL: https://www.math.uchicago.edu/~emerton/pdffiles/ext.pdf (cited on p. 14). |
| :canonical | [70] | J S Milne. “Canonical models of (mixed) Shimura varieties and automorphic vector bundles”. In: <i>Automorphic forms, Shimura varieties, and L-functions, Vol. I (Ann Arbor, MI, 1988)</i> . Vol. 10. Perspect. Math. Boston, MA: Academic Press, 1990, pp. 283–414 (cited on pp. 42, 45). |
| nen:models | [71] | Ben Moonen. “Models of Shimura varieties in mixed characteristics”. In: <i>Galois representations in arithmetic algebraic geometry (Durham, 1996)</i> . Vol. 254. London Math. Soc. Lecture Note Ser. Cambridge: Cambridge Univ. Press, 1998, pp. 267–350 (cited on pp. 5, 45). |
| s_integral | [72] | Georgios Pappas. “On integral models of Shimura varieties”. In: <i>Math. Ann.</i> 385.3-4 (2023), pp. 2037–2097 (cited on pp. 1, 3, 5, 31). |
| s2023padic | [73] | Georgios Pappas and Michael Rapoport. “ p -adic shukas and the theory of global and local Shimura varieties”. In: (2023). URL: https://arxiv.org/abs/2106.08270 (cited on pp. 3, 5, 59–62). |
| patibility | [74] | Stefan Patrikis. “Compatibility of canonical ℓ -adic local systems on Shimura varieties, II”. In: (2025). eprint: 2504.00305. URL: https://arxiv.org/abs/2504.00305 (cited on p. 55). |
| Pentland | [75] | Dylan Pentland. <i>Syntomification and crystalline local systems</i> . 2025. arXiv: 2510.16961 [math.NT]. URL: https://arxiv.org/abs/2510.16961 (cited on pp. 13–14). |
| calheights | [76] | Jonathan Pila, Ananth Shankar, Jacob Tsimerman, Hélène Esnault, and Michael Groechenig. “Canonical Heights on Shimura Varieties and the André-Oort Conjecture”. In: (2024). eprint: 2109.08788. URL: https://arxiv.org/abs/2109.08788 (cited on p. 55). |
| ink:thesis | [77] | Richard Pink. <i>Arithmetical compactification of mixed Shimura varieties (Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, 1989)</i> . Bonner Mathematische Schriften. Bonn: Universität Bonn Mathematisches Institut, 1990 (cited on p. 40). |
| ink2015-ye | [78] | Richard Pink, Torsten Wedhorn, and Paul Ziegler. “ F -zips with additional structure”. In: <i>Pacific J. Math.</i> 274.1 (Mar. 2015), pp. 183–236. URL: http://msp.org/pjm/2015/274-1/p09.xhtml (cited on pp. 6, 20). |

- [79] Michael Rapoport and M Richartz. “On the classification and specialization of F -isocrystals with additional structure”. In: *Compos. Math.* 103.2 (1996), pp. 153–181 (cited on pp. 7, 19, 57).
- [80] Michel Raynaud. *Anneaux locaux henséliens*. 1970th ed. Lecture notes in mathematics. Berlin, Germany: Springer, 1970 (cited on p. 32).
- [81] Peter Scholze. “Etale cohomology of diamonds”. In: *arXiv [math.AG]* (2017). URL: <http://arxiv.org/abs/1709.07343> (cited on p. 60).
- [82] Peter Scholze. “ p -adic Hodge theory for rigid-analytic varieties”. In: *Forum Math. Pi* 1 (2013), p. 77 (cited on p. 42).
- [83] Peter Scholze and Jared Weinstein. *Berkeley Lectures on p -adic Geometry*. Princeton University Press, 2020. URL: <http://dx.doi.org/10.23943/princeton/9780691202082.003.0010> (cited on pp. 30, 60, 62).
- [84] Ananth N Shankar and Rong Zhou. “Serre-Tate theory for Shimura varieties of Hodge type”. In: *Math. Z.* 297.3 (Apr. 2021), pp. 1249–1271 (cited on p. 21).
- [85] Xu Shen. “De Rham F -gauges and Shimura varieties”. In: (2024). URL: <https://arxiv.org/abs/2403.01899> (cited on p. 40).
- [86] Xu Shen and Chao Zhang. “Stratifications in good reductions of Shimura varieties of abelian type”. In: *arXiv [math.AG]* (2017). URL: <http://arxiv.org/abs/1707.00439> (cited on p. 7).
- [87] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2022 (cited on pp. 26, 31–32, 34, 40, 45).
- [88] Yuta Takaya. *Relative representability and parahoric level structures*. 2025. arXiv: 2402.07135 [math.NT]. URL: <https://arxiv.org/abs/2402.07135> (cited on pp. 7, 59, 61).
- [89] J. T. Tate. “ p -divisible groups”. In: *Proc. Conf. Local Fields (Driebergen, 1966)*. Springer, Berlin-New York, 1967, pp. 158–183 (cited on pp. 26–27).
- [90] A Vasiu. “Crystalline Boundedness Principle”. In: *Annales Scientifiques De L Ecole Normale Supérieure* 39 (May 2002), pp. 245–300 (cited on p. 55).
- [91] Adrian Vasiu. “Integral canonical models of Shimura varieties of preabelian type”. In: *Asian J. Math.* 3.2 (1999), pp. 401–518 (cited on p. 5).
- [92] Adrian Vasiu and Thomas Zink. “Purity results for p -divisible groups and abelian schemes over regular bases of mixed characteristic”. In: *Doc. Math.* 15 (2010), pp. 571–599 (cited on p. 17).
- [93] Eva Viehmann and Torsten Wedhorn. “Ekedahl-Oort and Newton strata for Shimura varieties of PEL type”. In: *Math. Ann.* 356.4 (2013), pp. 1493–1550. ISSN: 0025-5831,1432-1807. DOI: 10.1007/s00208-012-0892-z. URL: <https://doi-org.myaccess.library.utoronto.ca/10.1007/s00208-012-0892-z> (cited on pp. 6–7, 19).
- [94] Eva Viehmann, Torsten Wedhorn, and Appendix by Christopher Lang. “Moduli of truncated shtukas and displays”. In: (2025). arXiv: 2506.01740 [math.AG]. URL: <https://arxiv.org/abs/2506.01740> (cited on p. 56).
- [95] J-P Wintenberger. “Existence de F -cristaux avec structures supplémentaires”. In: *Adv. Math.* 190.1 (2005), pp. 196–224 (cited on p. 19).
- [96] Daniel Wortmann. “The μ -ordinary locus for Shimura varieties of Hodge type”. In: (2013). URL: <http://arxiv.org/abs/1310.6444> (cited on pp. 7, 20–22).
- [97] Peihang Wu. “Arithmetic compactifications of integral models of Shimura varieties of abelian type”. In: (2025). eprint: 2505.09135. URL: <https://arxiv.org/abs/2505.09135> (cited on p. 54).
- [98] Chao Zhang. “Ekedahl-Oort strata for good reductions of Shimura varieties of Hodge type”. In: *Canadian Journal of Mathematics* 70.2 (2018), pp. 451–480 (cited on p. 7).
- [99] Xinwen Zhu. “Affine Grassmannians and the geometric Satake in mixed characteristic”. en. In: *Ann. Math.* 185.2 (Mar. 2017), pp. 403–492. URL: <https://doi.org/10.4007/annals.2017.185.2.2> (cited on p. 56).

KEERTHI MADAPUSI, DEPARTMENT OF MATHEMATICS, MALONEY HALL, BOSTON COLLEGE, CHESTNUT HILL, MA 02467, USA
Email address: `madapusi@bc.edu`

ALEX YUCIS, DEPARTMENT OF MATHEMATICS, BAHEN CENTRE, UNIVERSITY OF TORONTO, TORONTO, ON, M5S 2E4, CANADA
Email address: `alex.yucis@gmail.com`