

Ecosystem carbon balance in the Hawaiian Islands under different scenarios of future climate and land use change

Paul C. Selmants^{1,6}, Benjamin M. Sleeter², Jinxun Liu¹, Tamara S. Wilson¹, Parker C. Trauernicht³, Abby G. Frazier⁴, Gregory P. Asner⁵

Affiliations:

¹U.S. Geological Survey, Moffett Field, CA, USA

²U.S. Geological Survey, Seattle, WA, USA

³University of Hawai'i at Mānoa, Honolulu, HI, USA

⁴The East-West Center, Honolulu, HI, USA

⁵Arizona State University, Tempe, AZ, USA

⁶Author to whom correspondence should be addressed

Email: pselmants@usgs.gov

Running title: Hawaii carbon balance

Keywords: land use, climate change, carbon balance, Hawaii, scenarios, disturbance, ecosystem model

Date: July 29, 2020

Abstract

The State of Hawai'i passed legislation in 2018 setting a goal to be carbon neutral by 2045. Meeting this goal will partly depend on carbon sequestration by terrestrial ecosystems, yet the future direction and magnitude of the land carbon sink in the Hawaiian Islands is highly uncertain. We used simulation modeling to assess how projected future changes in climate and land use will influence ecosystem carbon balance in the Hawaiian Islands under four unique scenarios over a 90-year timespan. Net ecosystem carbon balance declined under all four scenarios. Moving from a high to a low radiative forcing scenario reduced net ecosystem carbon loss by ~21%, and net carbon losses were reduced by a total of ~55% under the combined scenario of low radiative forcing and low rates of land-use change. A sensitivity test of the CO₂ fertilization effect on plant productivity revealed it to be a major source of uncertainty in projections of ecosystem carbon balance. Reconciling this uncertainty in how net photosynthesis will respond to rising atmospheric CO₂ will be essential to better constraint of models used to evaluate the effectiveness of ecosystem-based climate mitigation strategies.

Introduction

Methods

We used the Land Use and Carbon Scenario Simulator (LUCAS), an integrated landscape change and carbon gain-loss model, to project changes in ecosystem carbon balance for the seven main Hawaiian Islands under four scenarios of climate and land-use change. The landscape change portion of LUCAS is a state-and-transition model that applies a Monte Carlo approach to track the state type and age of each simulation cell in response to a pre-determined set of transitions (Daniel *et al* 2016). The carbon gain-loss portion tracks carbon stocks within each simulation cell over time as continuous state variables, along with a pre-

defined set of continuous flows specifying stock level rates of change over time (Daniel *et al* 2018, Sleeter *et al* 2019). We parameterized the Hawai'i LUCAS model to estimate annual changes in carbon stocks and fluxes in response to land use, land use change, wildland fire, and long-term climate variability. Simulations were run for 90 years at an annual timestep, using initial conditions corresponding to the year 2010. All simulations were repeated for 30 Monte Carlo realizations.

Study area

The spatial extent of this study was the terrestrial portion of the seven main Hawaiian Islands (Figure 1), a total land area of 16,554 km². We subdivided this landscape into a grid of 264,870 simulation cells, each of which was 250 x 250 m in size. Each simulation cell was assigned to one of 210 possible state types based on the unique combination of three moisture zones (dry, mesic, and wet; Figure S1), seven islands, and ten discrete land cover classes (Figure 1).

States and transitions

Transition pathways were modified from Daniel *et al* (2016) and Selmants *et al* (2017). Transitions between state types were pre-defined to represent urbanization, agricultural contraction, agricultural expansion, harvesting of tree plantations, and wildfire. Agriculture, forest, grassland, tree plantation, and shrubland state types each had multiple transition pathways, while the barren state type could only transition to urban. Although several state types had urbanization transition pathways, there was no transition pathway from urban to any other state type. Water and wetland state types remained static throughout the simulation period.

We developed a high and a low land-use scenario, with transition targets based on historical trends of land use change and population projections. Annual transition targets for agricul-




Figure 1: Land cover classification of the seven main Hawaiian Islands, adapted from Jacobi et al (2017). Agriculture in this map combines herbaceous and woody crops, but these two crop types are treated as separate land cover classes in the simulation model. Water and Wetland land cover classes are not shown.

tural contraction and agricultural expansion were based on a time series of land use change from 1992-2011 (NOAA 2020).

Carbon stocks and flows

Initial conditions

Scenario simulations

Results

Discussion

Conclusion

Acknowledgements

This study was funded by the Biological Sequestration Program of the U.S. Geological Survey. Thanks to Nicholas Koch of Forest Solutions, Inc. for information on eucalyptus harvesting and to Leonardo Frid and Colin Daniel of ApexRMS for assistance with the SyncroSim software. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Data Availability

Tabular model output data are available from the USGS ScienceBase data repository at: <https://doi.org/10.5066/P9AWLFKZ>. Model input data and R code used to format input data, summarize output data, and compile this manuscript are available from the HI_Model GitHub repository at: https://github.com/selmants/HI_Model.

ORCID

Paul C. Selman <https://orcid.org/0000-0001-6211-3957>

Benjamin M. Sleeter <https://orcid.org/0000-0003-2371-9571>

Jinxun Liu <https://orcid.org/0000-0003-0561-8988>

Tamara S. Wilson <https://orcid.org/0000-0001-7399-7532>

Abby G. Frazier <https://orcid.org/0000-0003-4076-4577>

Gregory P. Asner <https://orcid.org/0000-0001-7893-6421>

References

Daniel C J, Frid L, Sleeter B M and Fortin M-J 2016 State-and-transition simulation models: A framework for forecasting landscape change *Methods in Ecology and Evolution* **7** 1413–23

Daniel C J, Sleeter B M, Frid L and Fortin M-J 2018 Integrating continuous stocks and flows into state-and-transition simulation models of landscape change *Methods in Ecology and Evolution* **9** 1133–43

Jacobi J, Price J, Gon III S and Berkowitz P 2017 Hawaii Land Cover and Habitat Status: U.S. Geological Survey data release Online: <https://doi.org/10.5066/F7DB80B9>

NOAA 2020 *Coastal Change Analysis Program (C-CAP) Regional Land Cover: Hawaii* (NOAA Office of Coastal Management) Online: <https://coast.noaa.gov/digitalcoast/data/>

Selman P C, Giardina C P, Jacobi J D and Zhu Z 2017 *Baseline and projected future carbon storage and carbon fluxes in ecosystems of Hawai‘i* (U.S. Geological Survey) Online: <https://doi.org/10.3133/pp1834>

Sleeter B M, Marvin D C, Cameron D R, Selman P C, Westerling A L, Kreidler J, Daniel C J, Liu J and Wilson T S 2019 Effects of 21st-century climate, land use, and disturbances

