

Ecosystem carbon balance in the Hawaiian Islands under different scenarios of future climate and land use change

Paul C. Selmants^{1,6}, Benjamin M. Sleeter², Jinxun Liu¹, Tamara S. Wilson¹, Parker C. Trauernicht³, Abby G. Frazier⁴, Gregory P. Asner⁵

Affiliations:

¹U.S. Geological Survey, Moffett Field, CA, USA

²U.S. Geological Survey, Seattle, WA, USA

³University of Hawai'i at Mānoa, Honolulu, HI, USA

⁴The East-West Center, Honolulu, HI, USA

⁵Arizona State University, Tempe, AZ, USA

⁶Author to whom correspondence should be addressed

Email: pselmants@usgs.gov

Running title: Hawaii carbon balance

Keywords: land use, climate change, carbon balance, Hawaii, scenarios, disturbance, ecosystem model

Date: September 14, 2020

Abstract

The State of Hawai‘i passed legislation in 2018 setting a goal to be carbon neutral by 2045. Meeting this goal will partly depend on carbon sequestration by terrestrial ecosystems, yet the future direction and magnitude of the land carbon sink in the Hawaiian Islands is highly uncertain. We used simulation modeling to assess how projected future changes in climate and land use will influence ecosystem carbon balance in the Hawaiian Islands under four unique scenarios over a 90-year timespan. Net ecosystem carbon balance declined under all four scenarios. Moving from a high to a low radiative forcing scenario reduced net ecosystem carbon loss by ~21%, and net carbon losses were reduced by a total of ~55% under the combined scenario of low radiative forcing and low rates of land-use change. A sensitivity test of the CO₂ fertilization effect on plant productivity revealed it to be a major source of uncertainty in projections of ecosystem carbon balance. Reconciling this uncertainty in how net photosynthesis will respond to rising atmospheric CO₂ will be essential to better constraint of models used to evaluate the effectiveness of ecosystem-based climate mitigation strategies.

Introduction

The main Hawaiian Islands are a complex mosaic of natural and human-dominated landscapes,

Methods

We used the Land Use and Carbon Scenario Simulator (LUCAS), an integrated landscape change and carbon gain-loss model, to project changes in ecosystem carbon balance for the seven main Hawaiian Islands under all combinations of two land-use scenarios (low and high) and two radiative forcing scenarios (RCP 4.5 and RCP 8.5). We also developed a separate

set of scenarios to test model sensitivity to different levels of a CO₂ fertilization effect (CFE). The landscape change portion of LUCAS is a state-and-transition model that applies a Monte Carlo approach to track the state type and age of each simulation cell in response to a pre-determined set of transitions (Daniel *et al* 2016). The carbon gain-loss portion tracks carbon stocks within each simulation cell over time as continuous state variables, along with a pre-defined set of continuous flows specifying stock level rates of change over time (Daniel *et al* 2018, Sleeter *et al* 2019). We parameterized the Hawai'i LUCAS model to estimate annual changes in carbon stocks and fluxes in response to land use, land use change, wildland fire, and long-term climate variability for the time period 2010-2100.

Study area

The spatial extent of this study was the terrestrial portion of the seven main Hawaiian Islands (figure 1), a total land area of 16,554 km². We subdivided this landscape into a grid of 264,870 simulation cells, each of which was 250 x 250 m in size. Each simulation cell was assigned to one of 210 possible state types based on the unique combination of three moisture zones (dry, mesic, and wet; figure S1), seven islands, and ten discrete land cover classes (figure 1).

States and transitions

We developed two land-use scenarios (low and high) with transition pathways modified from Daniel *et al* (2016). Transitions between state types were pre-defined to represent urbanization, agricultural contraction, agricultural expansion, harvesting of tree plantations, and wildfire. Agriculture, forest, grassland, tree plantation, and shrubland state types each had multiple transition pathways, while the barren state type could only transition to developed (i.e., urbanization). There was no transition pathway out of an urbanized (developed) state. Water and wetland state types remained static throughout the simulation period.




Figure 1: Land cover classification of the seven main Hawaiian Islands, adapted from Jacobi et al (2017). Agriculture in this map combines herbaceous and woody crops, but these two crop types are treated as separate land cover classes in the simulation model. Water and Wetland land cover classes are not shown.

Transition targets were based on historical trends of land use change in the Hawaiian Islands from 1992-2011 (NOAA 2020) and on population projections for the State of Hawai‘i (Kim and Bai 2018). For the high land-use scenario, transition rates for each timestep and Monte Carlo realization were sampled from uniform distributions bounded by the median and maximum historical rates of agricultural contraction, agricultural expansion, and urbanization for each island. For the low land-use scenario, rates of agricultural contraction and expansion were sampled from uniform distributions bounded by zero and the minimum historical rates for each island. Urbanization rates in the low land-use scenario were based on island-level population estimates and projections at five year intervals from 2010-2045 (Kim and Bai 2018). We converted population projections into urbanization transition targets following Sleeter *et al* (2017) by calculating population density for each island and then projecting future developed area based on the five-year incremental change in island population. The spatial extent of agricultural contraction, agricultural expansion, and urbanization was constrained in both land-use scenarios based on existing zoning maps (Daniel *et al* 2016). Transition targets for tree plantation harvest were set at ~75% of recent historical rates in the high land-use scenario and ~40% of recent historical rates in the low land-use scenario (Daniel *et al* 2016). In both land-use scenarios, approximately 60% of tree plantation harvests were replacement harvests resulting in conversion to agriculture. The remaining 40% were rotation harvests replanted to *Eucalyptus* spp.

The wildfire transition sub-model was modified from Daniel *et al* (2016) by incorporating a new 21-year historical wildfire spatial database of the Hawaiian Islands (figure S2). We used this new spatial database to calculate historical wildfire size distribution and ignition probabilities for each unique combination of moisture zone (figure S1), island, and state type (figure 1) for the years 1999-2019. Starting in 2020, the number and size of fires was randomly drawn from one of these historical year-sets for each timestep and Monte Carlo realization, using burn severity probabilities from Selmants *et al* (2017). Wildfire in the low land-use scenario was sampled from the subset of historical fire years at or below the median

area burned statewide from 1999-2019. The high land-use scenario sampled from historical fire years above the median area burned over the same 21-year period (Fig. S2a).

Carbon stocks and flows

The fate of carbon stocks was tracked for each simulation cell based on a suite of carbon flows (i.e., carbon fluxes) specifying the rates of change in these carbon stocks over time (Daniel *et al* 2018, Sleeter *et al* 2019). We defined carbon stocks as continuous state variables for each simulation cell, including live biomass, standing dead wood, down dead wood, litter, and soil organic carbon. We also included and tracked carbon in atmospheric, aquatic, and harvest product pools to enforce carbon mass balance (Daniel *et al* 2018). To transfer carbon between stocks, we defined baseline carbon flows as continuous variables resulting from growth, mortality, deadfall, woody decay, litter decomposition, and leaching (which includes runoff). We also defined carbon flows resulting from land use, land use change, and wildfire (Selmants *et al* 2017, Daniel *et al* 2018).

Initial carbon stocks and baseline carbon flows were estimated based on the moisture zone (figure S1), state type, and age of each simulation cell using a lookup table derived from the Integrated Biosphere Simulator (IBIS; Foley *et al* 1996, Liu *et al* 2020), a process-based dynamic global vegetation model. We initiated IBIS with minimal vegetation and simulated forward for 110 years using 30-year climate normals for the Hawaiian Islands (Giambelluca *et al* 2013, 2014). We calibrated IBIS carbon stocks with statewide gridded datasets of soil organic carbon (Soil Survey Staff 2016) and forest aboveground live biomass (Asner *et al* 2016). We also calibrated gross photosynthesis in IBIS using a Hawai‘i-specific gridded dataset derived from MODIS satellite imagery (Kimball *et al* 2017).

Carbon flow rates for each state type and moisture zone were estimated as the ratio of the IBIS-derived flux to the size of the originating carbon stock at each age (Sleeter *et al* 2018). A spatially explicit stationary growth multiplier was applied to each simulation cell

to reflect local variations in net primary productivity (NPP) driven by microclimate. This spatial growth multiplier was the NPP anomaly for each cell relative to mean values for each combination of state type and moisture zone (Sleeter *et al* 2019) calculated using empirical relationships between total annual NPP and mean annual rainfall or temperature (Schuur 2003, Del Grosso *et al* 2008). Climate change impacts on carbon flows were represented by temporal growth and decay multipliers applied to each simulation cell based on statistically downscaled CMIP5 climate projections for the Hawaiian Islands under each of the two radiative forcing scenarios (RCP 4.5 and RCP 8.5; Timm *et al* 2015, Timm 2017). The impact of future changes in rainfall and temperature on NPP were represented by annual growth multipliers calculated using empirical NPP models (Schuur 2003, Del Grosso *et al* 2008) and climate model projections of temperature and rainfall for each radiative forcing scenario. The effect of future warming on turnover rates of dead organic matter were represented by temporal decay multipliers calculated using Q10 functions and climate model temperature projections for each radiative forcing scenario. We applied a Q10 of 2.0 for wood and soil organic matter decay flows (Kurz *et al* 2009, Sleeter *et al* 2019) and a Q10 of 2.17 for litter decay flows (Bothwell *et al* 2014). Transition-triggered carbon flows resulting from disturbances associated with land use change, timber harvesting, and wildfire were based on values from Don *et al* (2011), Selmants *et al* (2017), and Daniel *et al* (2018).

CO₂ fertilization effect

Increasing atmospheric CO₂ concentrations stimulate leaf-level photosynthesis, potentially increasing NPP as well. However, the magnitude and persistence of this effect is highly uncertain, particularly across a range of climatic conditions and over long time spans. Following Sleeter *et al* (2019), we developed a separate set of scenarios designed to test the sensitivity of LUCAS model ecosystem carbon balance projections to the influence of differing rates of a CO₂ fertilization effect (CFE). We incorporated a NPP CFE multiplier representing the percent increase in NPP for every 100 ppm increase in atmospheric CO₂ concentration

under the high land use and high radiative forcing (RCP 8.5) scenario. We tested five CFE levels ranging from 5% to 15%, which is within the range of CFEs observed in free air CO₂ enrichment (FACE) experiments. For all levels, we assumed CFEs reached saturation at an atmospheric CO₂ concentration of 600 ppm, with no further increase in NPP despite a continued increase in CO₂ concentration to 930 ppm by 2100. This 600ppm threshold generally coincides with the upper limit from FACE experiments and is reached by the year 2060 under RCP 8.5.

Scenario simulations and analysis

Each of the four unique scenarios were run for 90 years at an annual timestep and repeated for 30 Monte Carlo realizations, using initial conditions corresponding to the year 2010. All simulations were performed within the SyncroSim (version 2.2.4) software framework with ST-Sim (version 3.2.13) and SF (version 3.2.10) add-on modules (Daniel *et al* 2016, 2018). Model inputs and outputs were prepared with the R statistical computing platform (R Core Team 2019) using the tidyverse (Wickham *et al* 2019), raster (Hijmans 2020), and rsyncrosim (Daniel *et al* 2020) packages. Carbon stocks and fluxes for the seven main Hawaiian Islands were calculated for each scenario by summing within each of the 30 Monte Carlo realizations on an annual basis and then calculating annual means as well as the annual upper and lower limits of the 30 Monte Carlo realizations. Carbon balance for the seven main Hawaiian Islands was calculated on annual basis for each scenario and Monte Carlo realization as net biome productivity (NBP), which was equal to annual carbon input in the form of net primary productivity (NPP) minus the annual sum of all carbon losses from terrestrial ecosystems, including heterotrophic respiration (R_h) from litter and soil, carbon fluxes to the atmosphere triggered by land use and land-use change, wildfire emissions, and aquatic carbon losses through leaching and overland flow. Positive NBP values indicated ecosystems of the seven main Hawaiian Islands were acting as a net sink for atmospheric CO₂, while negative NBP values indicated that these ecosystems were acting as a net carbon source to

the atmosphere (Chapin *et al* 2006).

Results

Terrestrial ecosystems of the seven main Hawaiian Islands stored an estimated 316 Tg of carbon at the beginning of the simulation period in 2010, with 58% in soil organic matter, 22% in living biomass, and 20% in surface dead organic matter (litter and dead wood; figure 2a). Ecosystems accumulated carbon in all scenarios but at different rates, with trajectories shaped primarily by climate change and to a lesser extent by land-use change. The highest and most consistent projected accumulation of ecosystem carbon occurred under the combination of low radiative forcing and low land use change, yielding a ~15% increase in ecosystem carbon to an average of 363 Tg by 2100 (figure 2a). In contrast, high radiative forcing and high land use change resulted in the lowest ecosystem carbon gain, reaching a peak of ~332 Tg in 2063 but declining to 327 Tg in 2100, resulting in a net increase of only 3% by the end of the simulation period (figure 2a). Ecosystem carbon accumulation was driven exclusively by increasing soil organic carbon across all four scenarios, all other stocks declined over time (figure 2b).

Net primary production (NPP) for the seven main Hawaiian Islands declined across all four scenarios, driven primarily by climate change and to a lesser extent by land use change (Fig. 3). The combination of high radiative forcing (RCP 8.5) and high land-use change led to the steepest decline in NPP over time, driven by intense long-term drying on the leeward sides of islands under RCP 8.5 (figure S4) and sustained losses of forest and shrubland land area in the high land-use scenario (figure S5). In contrast, climate change led to increased heterotrophic respiration (R_h) over time, such that more intense warming under RCP 8.5 (figure S4) resulted in R_h being ~3% higher by 2100 than under RCP 4.5 (figure 3). Land-use change substantially reduced R_h in the high land-use scenario (figure 3) because of long-term reductions in forest and shrubland land area (figure S5), similar to trends in NPP. Transition-




Figure 2: Projected changes in total ecosystem carbon storage (a) and individual carbon stocks (b) for the seven main Hawaiian Islands. Solid lines indicate the mean of 30 Monte Carlo realizations for each scenario, with shaded areas indicating the minimum and maximum range of Monte Carlo realizations.

triggered carbon fluxes to the atmosphere from land use, land-use change, and wildfire were largely independent of changes in climate, stabilizing by mid-century at an average of $\sim 0.4 \text{ Tg y}^{-1}$ in the high land-use scenario and an average of $\sim 0.2 \text{ Tg y}^{-1}$ in the low land-use scenario (figure 3). Uncertainty around transition-triggered carbon fluxes were higher in the high land-use scenario, driven by greater variability in wildland fire probabilities.

On average, terrestrial ecosystems of the seven main Hawaiian Islands collectively acted as a net carbon sink throughout the simulation period under RCP 4.5, but carbon sink strength by 2100 was $\sim 60\%$ greater under the low land-use scenario than under the high land-use scenario (figure 4). In contrast, ecosystems of the Hawaiian Islands became a net carbon source to the atmosphere toward the latter half of the simulation period under RCP 8.5, although the transition from sink to source occurred 15 years earlier on average under the high land-use scenario than under the low land-use scenario (figure 4). The magnitude of the carbon source was also $\sim 40\%$ greater under the high land-use scenario by the end of the simulation period.




Figure 3: Projected changes in net primary production (NPP), heterotrophic respiration (Rh) and carbon fluxes induced by land use and land-use change for the seven main Hawaiian Islands. Solid lines indicate the mean of 30 Monte Carlo realizations for each scenario, with shaded areas indicating the minimum and maximum range of Monte Carlo realizations.




Figure 4: Projected changes in net biome productivity (NBP) for the seven main Hawaiian Islands. Values above zero indicate terrestrial ecosystems are acting as a net carbon sink for atmospheric carbon and values below zero indicate ecosystems are acting as a net carbon source to the atmosphere. Solid lines indicate the mean of 30 Monte Carlo realizations for each scenario, with shaded areas indicating the minimum and maximum range of Monte Carlo realizations.

Discussion

Conclusion

Acknowledgements

This study was funded by the Biological Sequestration Program of the U.S. Geological Survey. Thanks to Nicholas Koch of Forest Solutions, Inc. for information on eucalyptus harvesting and to Leonardo Frid and Colin Daniel of ApexRMS for assistance with SyncroSim software. Thanks also to Christian Giardina and Zhiliang Zhu for making this project possible. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Data Availability

Tabular model output data and metadata are available in machine readable format from a USGS ScienceBase data repository at <https://doi.org/10.5066/P9AWLFKZ>. Model input data and R code used to format input data, summarize output data, and compile this manuscript are available from a GitHub repository at https://github.com/selmants/HI_Model.

ORCID

Paul C. Selman <https://orcid.org/0000-0001-6211-3957>
Benjamin M. Sleeter <https://orcid.org/0000-0003-2371-9571>
Jinxun Liu <https://orcid.org/0000-0003-0561-8988>
Tamara S. Wilson <https://orcid.org/0000-0001-7399-7532>

Abby G. Frazier <https://orcid.org/0000-0003-4076-4577>

Gregory P. Asner <https://orcid.org/0000-0001-7893-6421>

References

Asner G P, Sousan S, Knapp D E, Selmants P C, Martin R E, Hughes R F and Giardina C P 2016 Rapid forest carbon assessments of oceanic islands: A case study of the Hawaiian archipelago *Carbon Balance and Management* **11** Online: <http://www.cbmjournals.com/content/11/1/1>

Bothwell L D, Selmants P C, Giardina C P and Litton C M 2014 Leaf litter decomposition rates increase with rising mean annual temperature in Hawaiian tropical montane wet forests *PeerJ* **2** e685 Online: <https://peerj.com/articles/685>

Chapin F S, Woodwell G M, Randerson J T, Rastetter E B, Lovett G M, Baldocchi D D, Clark D A, Harmon M E, Schimel D S, Valentini R, Wirth C, Aber J D, Cole J J, Goulden M L, Harden J W, Heimann M, Howarth R W, Matson P A, McGuire A D, Melillo J M, Mooney H A, Neff J C, Houghton R A, Pace M L, Ryan M G, Running S W, Sala O E, Schlesinger W H and Schulze E-D 2006 Reconciling carbon-cycle concepts, terminology, and methods *Ecosystems* **9** 1041–50

Daniel C, Hughes J, Embrey A, Frid L and Lucet V 2020 *Rsyncrosim: The r interface to syncrosim* Online: <https://github.com/rsyncrosim/rsyncrosim>

Daniel C J, Frid L, Sleeter B M and Fortin M-J 2016 State-and-transition simulation models: A framework for forecasting landscape change *Methods in Ecology and Evolution* **7** 1413–23

Daniel C J, Sleeter B M, Frid L and Fortin M-J 2018 Integrating continuous stocks and flows into state-and-transition simulation models of landscape change *Methods in Ecology and Evolution* **9** 1133–43

251 Del Grosso S, Parton W, Stohlgren T, Zheng D, Bachelet D, Prince S, Hibbard K and Olson R
 252 2008 Global potential net primary production predicted from vegetation class, precipitation,
 253 and temperature *Ecology* **89** 2117–26

254 Don A, Schumacher J and Freibauer A 2011 Impact of tropical land-use change on soil
 255 organic carbon stocks – a meta-analysis *Global Change Biology* **17** 1658–70

256 Foley J A, Prentice I C, Ramankutty N, Levis S, Pollard D, Sitch S and Haxeltine A 1996
 257 An integrated biosphere model of land surface processes, terrestrial carbon balance, and
 258 vegetation dynamics *Global Biogeochemical Cycles* **10** 603–28

259 Giambelluca T W, Chen Q, Frazier A G, Price J P, Chen Y-L, Chu P-S, Eischeid J K and
 260 Delparte D M 2013 Online Rainfall Atlas of Hawai‘i *Bull. Amer. Meteor. Soc.* **94** 313–6

261 Giambelluca T W, Shuai X, Barnes M L, Alliss R J, Longman R J, Miura T, Chen Q,
 262 Frazier A G, Mudd R G, Cuo L and Businger A D 2014 *Evapotranspiration of Hawai‘i*
 263 Online: <http://evapotranspiration.geography.hawaii.edu/downloads.html>

264 Hijmans R J 2020 *Raster: Geographic data analysis and modeling* Online: [https://CRAN.R-](https://CRAN.R-project.org/package=raster)
 265 [project.org/package=raster](https://CRAN.R-project.org/package=raster)

266 Jacobi J, Price J, Gon III S and Berkowitz P 2017 Hawaii Land Cover and Habitat Status:
 267 U.S. Geological Survey data release Online: <https://doi.org/10.5066/F7DB80B9>

268 Kim Y-S and Bai J 2018 *Population and economic projections for the State of Hawaii to*
 269 *2045* (Hawaii Department of Business, Economic Development & Tourism) Online: [https:](https://dbedt.hawaii.gov/economic/economic-forecast/2045-long-range-forecast/)
 270 [//dbedt.hawaii.gov/economic/economic-forecast/2045-long-range-forecast/](https://dbedt.hawaii.gov/economic/economic-forecast/2045-long-range-forecast/)

271 Kimball H L, Selman P C, Moreno A, Running S W and Giardina C P 2017 Evaluating
 272 the role of land cover and climate uncertainties in computing gross primary production
 273 in Hawaiian Island ecosystems *PLOS ONE* **12** e0184466 Online: [http://journals.plos.org/](http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0184466)
 274 [plosone/article?id=10.1371/journal.pone.0184466](http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0184466)

275 Kurz W A, Dymond C C, White T M, Stinson G, Shaw C H, Rampley G J, Smyth C,
 276 Simpson B N, Neilson E T, Trofymow J A, Metsaranta J and Apps M J 2009 CBM-CFS3:
 277 A model of carbon-dynamics in forestry and land-use change implementing IPCC standards
 278 *Ecological Modelling* **220** 480–504 Online: [http://www.sciencedirect.com/science/article/
 279 pii/S0304380008005012](http://www.sciencedirect.com/science/article/pii/S0304380008005012)

280 Liu J, Sleeter B M, Zhu Z, Loveland T R, Sohl T, Howard S M, Key C H, Hawbaker T,
 281 Liu S, Reed B, Cochrane M A, Heath L S, Jiang H, Price D T, Chen J M, Zhou D, Bliss N
 282 B, Wilson T, Sherba J, Zhu Q, Luo Y and Poulter B 2020 Critical land change information
 283 enhances the understanding of carbon balance in the United States *Global Change Biology*
 284 **26** 3920–9 Online: <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15079>

285 NOAA 2020 *Coastal Change Analysis Program (C-CAP) Regional Land Cover: Hawaii*
 286 (NOAA Office of Coastal Management) Online: <https://coast.noaa.gov/digitalcoast/data/>

287 R Core Team 2019 *R: A language and environment for statistical computing* (Vienna, Austria:
 288 R Foundation for Statistical Computing) Online: <https://www.R-project.org/>

289 Schuur E A 2003 Productivity and global climate revisited: The sensitivity of tropical forest
 290 growth to precipitation *Ecology* **84** 1165–70

291 Selmants P C, Giardina C P, Jacobi J D and Zhu Z 2017 *Baseline and projected future*
 292 *carbon storage and carbon fluxes in ecosystems of Hawai'i* (U.S. Geological Survey) Online:
 293 <https://doi.org/10.3133/pp1834>

294 Sleeter B M, Liu J, Daniel C, Rayfield B, Sherba J, Hawbaker T J, Zhiliang Zhu, Selmants
 295 P C and Loveland T R 2018 Effects of contemporary land-use and land-cover change on the
 296 carbon balance of terrestrial ecosystems in the United States *Environ. Res. Lett.* **13** 045006
 297 Online: <http://stacks.iop.org/1748-9326/13/i=4/a=045006>

298 Sleeter B M, Marvin D C, Cameron D R, Selmants P C, Westerling A L, Kreitler J, Daniel
 299 C J, Liu J and Wilson T S 2019 Effects of 21st-century climate, land use, and disturbances

on ecosystem carbon balance in California *Global Change Biology* **25** 3334–53

Sleeter B M, Wilson T S, Sharygin E and Sherba J T 2017 Future scenarios of land change based on empirical data and demographic trends *Earth's Future* **5** 1068–83 Online: <http://doi.wiley.com/10.1002/2017EF000560>

Soil Survey Staff 2016 *Soil Survey Geographic (SSURGO) Database, Natural Resources Conservation Service, United States Department of Agriculture* (Natural Resources Conservation Service, USDA) Online: <https://sdmdataaccess.sc.egov.usda.gov>.

Timm O E 2017 Future warming rates over the Hawaiian Islands based on elevation-dependent scaling factors *International Journal of Climatology* **37** 1093–104 Online: <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.5065>

Timm O E, Giambelluca T W and Diaz H F 2015 Statistical downscaling of rainfall changes in Hawai'i based on the CMIP5 global model projections *Journal of Geophysical Research: Atmospheres* **120** 92–112

Wickham H, Averick M, Bryan J, Chang W, McGowan L D, François R, Grolemond G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen T L, Miller E, Bache S M, Müller K, Ooms J, Robinson D, Seidel D P, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K and Yutani H 2019 Welcome to the tidyverse *Journal of Open Source Software* **4** 1686