

1 Supplemental material: Ecosystem carbon balance in the  
2 Hawaiian Islands under different scenarios of future  
3 climate and land use change

4  
5 Paul C. Selmants<sup>1,6</sup>, Benjamin M. Sleeter<sup>2</sup>, Jinxun Liu<sup>1</sup>, Tamara S. Wilson<sup>1</sup>,  
6 Parker C. Trauernicht<sup>3</sup>, Abby G. Frazier<sup>4</sup>, Gregory P. Asner<sup>5</sup>

7 **Affiliations:**

8 <sup>1</sup>U.S. Geological Survey, Moffett Field, CA, USA

9 <sup>2</sup>U.S. Geological Survey, Seattle, WA, USA

10 <sup>3</sup>University of Hawai‘i at Mānoa, Honolulu, HI, USA

11 <sup>4</sup>The East-West Center, Honolulu, HI, USA

12 <sup>5</sup>Arizona State University, Tempe, AZ, USA

13 <sup>6</sup>Author to whom correspondence should be addressed

14 **Email:** [pselmants@usgs.gov](mailto:pselmants@usgs.gov)

15 **Running title:** Hawaii carbon balance

16 **Keywords:** land use, climate change, carbon balance, Hawaii, scenarios, disturbance, ecosystem  
17 model

18 **Date:** February 16, 2021

## Moisture Zones

State Classes for Agriculture, Forest, Grassland, Shrubland, and Tree Plantations were stratified into three Moisture Zones - Dry, Mesic, and Wet (Figure S1). These three zones were based on a moisture availability index (MAI), calculated as mean annual precipitation (MAP) minus potential evapotranspiration (PET; Price *et al* 2012). Areas where the MAI was less than zero, i.e. where PET exceeded MAP, were classified in the Dry Moisture Zone. Areas with MAI values between zero and 1,661 were classified in the Mesic Moisture Zone. The MAI value of 1,661 is roughly equivalent to areas at 1,000m elevation that receive 2,500 mm of annual rainfall (Price *et al* 2012). Areas with MAI values that exceeded 1,661 were classified in the Wet Mesic Zone.




Figure 1: Moisture zones of the seven main Hawaiian Islands, adapted from Jacobi et al (2017).

## Wildland Fire

This study used a new spatial database of wildland fire perimeters in the Hawaiian Islands for the years 1999-2019 compiled using data from the Monitoring Trends in Burn Severity (MTBS) program, the Hawai'i Wildfire Management Organization, the U.S. National Park Service, the O'ahu Army Natural Resources Program, and independent remote sensing analyses using publicly available Landsat data. This new wildland fire spatial database was used to calculate annual area burned (Figure S2), wildland fire probabilities, and wildland fire size distributions.




Figure 2: Annual area burned by wildland fire in the State of Hawai'i from 1999-2019, summed by year across the seven main Hawaiian Islands (A) and within each of the four largest islands (B). The dashed horizontal line in (A) represents the median area burned from 1999-2019.

## Climate

Spatially explicit mean annual data on contemporary temperature and rainfall for the main Hawaiian Islands at 250-m resolution (Figure S3) are from Giambelluca *et al* (2013) and Giambelluca *et al* (2014). Spatially explicit projections of changes in annual temperature and rainfall by the year 2100 were developed using statistically downscaled CMIP5 climate projections for the Hawaiian Islands from Timm *et al* (2015) and Timm (2017) under RCPs 4.5 and 8.5 (Figure S5).




Figure 3: Mean annual temperature (top panel) and mean annual rainfall (bottom panel) for the seven main Hawaiian Islands.




Figure 4: Projected change in mean annual temperature (top panels) and mean annual rainfall (bottom panels) by end of century under RCPs 4.5 and 8.5 (top panels) based on statistical downscaling of an ensemble of climate models (CMIP5).

## Change in State Class Area

Projections of total land area covered by each State Class were summed by year across the seven main Hawaiian Islands from LUCAS model output data.




Figure 5: Projections of total land area by year for each of five State Classes in the seven main Hawaiian Islands (Agriculture, Developed, Forest, Grassland, and Shrubland) under low and high land use change scenarios for the period 2010-2100. Solid lines represent the mean of 30 Monte Carlo realizations and shaded areas represent minimum and maximum Monte Carlo values.

## References

- Giambelluca T W, Chen Q, Frazier A G, Price J P, Chen Y-L, Chu P-S, Eischeid J K and Delparte D M 2013 Online Rainfall Atlas of Hawai‘i *Bull. Amer. Meteor. Soc.* **94** 313–6
- Giambelluca T W, Shuai X, Barnes M L, Alliss R J, Longman R J, Miura T, Chen Q, Frazier A G, Mudd R G, Cuo L and Businger A D 2014 *Evapotranspiration of Hawai‘i* Online: <http://evapotranspiration.geography.hawaii.edu/downloads.html>

50 Jacobi J, Price J, Gon III S and Berkowitz P 2017 Hawaii Land Cover and Habitat Status: U.S.  
 51 Geological Survey data release Online: <https://doi.org/10.5066/F7DB80B9>

52 Price J, Jacobi J, Gon III S, Matsuwaki D, Mehrhoff L, Wagner W, Lucas M and Rowe B 2012  
 53 *Mapping plant species ranges in the Hawaiian Islands - developing a methodology and*  
 54 *associated GIS layers* (U.S. Geological Survey) Online: <https://pubs.usgs.gov/of/2012/1192/>

55 Timm O E 2017 Future warming rates over the Hawaiian Islands based on elevation-dependent  
 56 scaling factors *International Journal of Climatology* **37** 1093–104 Online:  
 57 <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.5065>

58 Timm O E, Giambelluca T W and Diaz H F 2015 Statistical downscaling of rainfall changes in  
 59 Hawai‘i based on the CMIP5 global model projections *Journal of Geophysical Research:*  
 60 *Atmospheres* **120** 92–112