

# MELISSA

## The Melittologist's Newsletter



Ronald J. McGinley, Bryan N. Danforth, Maureen J. Mello  
Department of Entomology  
Smithsonian Institution, NHB-105  
Washington, DC 20560

Charles D. Michener  
Entomological Museum, Snow Hall  
University of Kansas  
Lawrence, KA 66045

NUMBER — 4

Summer, 1991

### EDITORIAL COMMENTS

## "Melissa is Back"

The editors of *Melissa* apologize for the lengthy delay in the publication of *Melissa 4*. Thanks to the help of Maureen Mello and Bryan Danforth, *Melissa* is back on track, and will continue as a regular newsletter for the dissemination of information on the study of solitary and social bees world-wide. Please send us articles on collecting trips, museum resources, past, present and future research, and anything which would be of general interest to bee researchers. We plan to have the 1992 issue out by the beginning of that year. We would like to thank George L. Venable for his expert assistance in the preparation of this newsletter.

In this issue we are pleased to announce that the United States National Science Foundation has funded a two year cooperative study of the bees of Mexico. The Principal Investigators on the grant are Wallace LaBerge and Ronald McGinley (see article below).

La Programa Cooperativo sobre la Apifauna Mexicana, or PCAM, traces its beginning to a meeting held in Chamela, Jalisco in 1985. At that meeting PCAM was formed as an informal organization to discuss and organize long term cooperative studies of the Mexican bee fauna. Because of the devastating earthquake which occurred a few weeks prior to the meeting, many interested Mexican scientists were unable to attend. A second workshop was held in Quintana Roo in October, 1986. A third meeting was held at Chamela in July 1989 and a fourth meeting at Hermosillo, Sonora, in July, 1990. The objectives of PCAM, established at the first two meetings, are described in the accompanying article. Requests for information regarding PCAM should be sent to George Eickwort (Cornell University) or Jerome Rozen, Jr. (American Museum of Natural History).

\* \* \*

### THE BEES OF MEXICO

## Cooperative Survey of the Bees of Mexico Funded

By

Wallace E. LaBerge  
Faunistics Section, State Natural History  
Survey Division, 607 E. Peabody,  
Champaign, Illinois, 61820, USA

In the fall of 1986, during the second PCAM meeting in the state of Quintana Roo, Mexico, it was decided that W.E. LaBerge and R.J. McGinley would prepare a proposal requesting funds from the National Science Foundation to support a survey for the native bees of Mexico. This proposal was submitted in 1987 and was not funded. A revised proposal was submitted in 1988 and was again turned down. A third proposal was prepared and submitted in 1990 and has been funded starting in the summer of 1991 for two years. The proposal was submitted through the University of Illinois in Urbana and will be administered at the Natural History Survey located on the University campus by W.E. LaBerge. This grant will provide funds to do several things listed and briefly discussed below.

1. First, travel money will be available to collect bees and associated data. This, of course, is the heart of the proposal. We hope to be able to strengthen several of the North American collections in their holding of native Mexican bees. Approximately half of all collections will eventually be returned to Mexico (to the Universidad Nacional Autonoma de Mexico or to other national insect collections). Scientists leading the native bee surveys need to agree to this stipulation, in order to receive funds from the grant. In addition, the proposal specifies that holotypes of species described from material resulting from this survey should be deposited in the collection of the Instituto de Biologia, UNAM, in Mexico City.

2. Computer data management of all collections will be centered at the Natural History Survey. Data (on discs, on paper, etc.) from all collections will be sent to LaBerge to be entered into the PCAM data bank on Mexican bees. These data will be available to all PCAM members, and other scientists interested in bee research.

3. The grant has some funds for purchase of a vehicle that will be used for field trips to collect bees. This vehicle, after the two year grant or any renewals has expired, will belong to UNAM and remain in Mexico.

4. Funds are available to support a graduate student from Mexico at the University of Illinois for two years. We hope that a suitable student will apply soon.

5. Some funds will be used to support hourly workers and a data entry person at the Natural History Survey. These workers will process bees (pin, label, etc.) for collections from trips led by investigators who do not have facilities or the resources to do this work, or who want help in this regard.

6. The grant states that all faunal studies, species lists, or keys to Mexican bees resulting from this study will be published in Spanish with English summary and that other papers in which species or nesting biologies are described will be published in either Spanish or English, depending upon nationality of the author, and summaries provided in the alternate language.

Four objectives of PCAM were agreed to by participants of the first two PCAM meetings. These are: (1) investigation of the taxonomic diversity and distribution of the bees of Mexico, (2) establishment and enhancement of permanent collections of bees in Mexico, (3) investigation of the biology and ecology of Mexican bees, and (4) establishment of a training program designed to strengthen the Mexican base of expertise on bees.

The last objective has been supported by the Smithsonian International Exchanges Program (grants to Ronald McGinley) which provided support to teach and train Mexican students in bee systematics and in field techniques at meetings held in Chamela and Hermosillo. It is hoped that the present NSF grant will advance this objective by providing the possibility for a Mexican student to earn an advanced degree. Also, Mexican students will be supported by funds from the grant to participate in the field trips. They are encouraged to take part and will be welcome participants.

The first two objectives will be directly supported by this NSF grant. Collections of Mexican bees will be enhanced, both in the United States and in Mexico. The third objective, although not directly addressed by our grant proposal, will also be advanced. In collecting native bees in Mexico, it is expected that floral data will be collected with all specimens. If the plants being used by the bees are unknown, specimens of the plants will be collected for identification later. It is hoped that a botanist may be induced to accompany the group for part or all of each trip. Also, any nesting bees will be noted, perhaps studied at the time (depending upon the discretion of the group leaders), and data entered into our data bank so that the site may be revisited.

A protocol has been designed to collect and record data from every major collecting site. This includes check-off items such as a list of plant communities, geographic location of the site, date, etc. There is space on the protocol for recording specific locations of such items as nesting sites, specific plant dominants at the site, and other pertinent information. This protocol is essential, we believe, for data management, labelling of specimens and relocation of sites.

W.E. LaBerge, R.J. McGinley (Smithsonian Institution), J.G. Rozen (American Museum of Natural History) and George Eickwort (Cornell University) will serve as an

informal committee to advise on all policy matters involving this grant, and additional Mexican committee members will be added after consultation with our Mexican counterparts. Dr. Rozen will lead the first survey trip under the auspices of this grant in August of this year to the arid northern regions of Mexico.

\* \* \*

## COLLECTION AND COLLECTING NEWS

### Catalog of Type Depositories of Bees in America North of Mexico

by

Sandra Shanks  
Department of Biology,  
Harney Science Center,  
University of San Francisco,  
San Francisco, CA, 94117-1080, USA

The updating of the Catalog of Type Depositories of Bees in America North of Mexico is essentially complete. Ron McGinley is assisting with arrangements for the Smithsonian to publish the catalog. To make sure that types of your newly described species or lectotype designations published since 1982 are included, please send a reprint or reference as soon as possible.

My sincere thanks go to all who responded to my letter of last summer by sending reprints and information to me. Library facilities here at "Pseudo-U" are woefully lacking, and time to visit other libraries is virtually non-existent. Your help made the updating considerably easier for me.

### The Brethes Types of Apoidea in the MACN Collection

by

Jorge F. Genise  
Museo Argentino de Ciencias  
Naturales, Av. Angel Gallardo 470,  
Casa Correo 220, Sucursal 5,  
1405 Buenos Aires, Argentina

Between 1901 and 1927, Juan Brethes published 77 papers in which he described 893 new species of Hymenoptera. Most of the types were deposited in the Museo Argentino de Ciencias Naturales (MACN) where he worked, building up one of the most important collections of Hymenoptera from South America.

When Brethes died, nobody paid attention to his collection for about fifty years, until M.A. Fritz began to arrange it and made the first alphabetical files.

Brethes used to put a small label with a number on each insect in his collection. He registered each number in a notebook with a determination and a copy of this notebook is kept at MACN. The number label is critical for determining the "type" status of Brethes material because he rarely identifies specimens with name labels, and his locality labels were less complete than information in the original descriptions. Definitive decisions on all "types" will have to be made by specialists after comparisons with the original descriptions.

In the following list species names are given in their original combination without considering subsequent nomenclatorial changes and synonyms. A plus sign (+) at the end of an entry indicates that there is typical material in the MACN collection. The absence of a sign means that the type material has not been found. In some cases such material may simply repose in our collection undetected, or it may have been borrowed without a record.

A corresponding list of Brethes aculeate wasp types and a complete list of Brethes papers on Hymenoptera are given in SPHECOS 20 (September, 1990), available on request from Arnold Menke (SEL/USDA, c/o National Museum of Natural History, Smithsonian Institution, Washington, DC, 20560, USA).

## APOIDEA

<i>Ancyloscelis videlai</i> Brethes, 1914	(+)
<i>Ancyloscelis bonariensis</i> Brethes, 1910	
<i>Anthidium caroliameghinoi</i> Brethes, 1903	
<i>Anthophora escomeli</i> Brethes, 1920	(+)
<i>Anthophora arequipensis</i> Brethes, 1920	(+)
<i>Arhyssusage johnsoni</i> Brethes, 1922	
<i>Augochlora (Tetraechlora) porteri</i> Brethes, 1914	(+)
<i>Augochlora proserpina</i> Brethes, 1909	(+)
<i>Augochlora clonia</i> Brethes, 1909	(+)
<i>Augochlora phylacis</i> Brethes, 1909	(+)
<i>Augochlora videlai</i> Brethes, 1914	
<i>Bombus peruvianus</i> Brethes, 1920	(+)
<i>Callonychium argentinum</i> Brethes, 1922	(+)
<i>Coelioxys fiebrigi</i> Brethes, 1909	
<i>Epeorus arechavaletai</i> Brethes, 1909	(+)
<i>Epeorus merus</i> Brethes, 1909	(+)
<i>Eulonchopria psaenythioides</i> Brethes, 1909	(+)
<i>Exomalopsis ascendens</i> Brethes, 1910	
<i>Exomalopsis latifasciata</i> Brethes, 1909	(+)
<i>Exomalopsis latifasciata sororcula</i> Brethes, 1909	(+)
<i>Exomalopsis pampeana</i> Brethes, 1910	(+)
<i>Exomalopsis solitaria</i> Brethes, 1910	(+)
<i>Exomalopsis spedazzinii</i> Brethes, 1910	(+)
<i>Exomalopsis testaceinervis</i> Brethes, 1910	(+)
<i>Exomalopsis trifasciata</i> Brethes, 1910	(+)
<i>Leptergatis fiebrigi</i> Brethes, 1909	(+)
<i>Leptometria gigantea</i> Brethes, 1920	(+)
<i>Leptometria lynchii</i> Brethes, 1910	(+)
<i>Leptometria mendoza</i> Brethes, 1910	(+)
<i>Leptometria patagonica</i> Brethes, 1910	(+)
<i>Leptometria singularis</i> Brethes, 1910	(+)
<i>Leptometria taricensis</i> Brethes, 1910	(+)
<i>Leptometria tucumana</i> Brethes, 1910	(+)
<i>Macroglossapis buccosa nigriventris</i> Brethes, 1910	(+)
<i>Megachile verrocosa</i> Brethes, 1909	(+)
<i>Melissodes gratiosus</i> Brethes, 1910	(+)

<i>Melissodes sobrinus</i> Brethes, 1910	(+)
<i>Melissodes sparsus</i> Brethes, 1910	(+)
<i>Melissodes rufithorax</i> Brethes, 1910	(+)
<i>Melissoptila argentina</i> Brethes, 1910	(+)
<i>Melissoptila fiebrigi</i> Brethes, 1910	(+)
<i>Melissoptila inducens</i> Brethes, 1910	(+)
<i>Melissoptila inducens</i> Brethes, 1910	(+)
<i>Melissoptila malvacearum</i> Brethes, 1910	(+)
<i>Melitoma nigrotarsalis</i> Brethes, 1910	(+)
<i>Nomada costalis</i> Brethes, 1909	(+)
<i>Parapsaenythia paraguaya</i> Brethes, 1909	
<i>Protodiscelis fiebrigi</i> Brethes, 1909	(+)
<i>Ptilothrix albido'hirta</i> Brethes, 1910	(+)
<i>Ptilothrix chacoensis</i> Brethes, 1910	
<i>Ptilothrix lynchii</i> Brethes, 1910	(+)
<i>Ptilothrix megasoma</i> Brethes, 1910	(+)
<i>Svastra ameghinoi</i> Brethes, 1910	(+)
<i>Svastra apicalis</i> Brethes, 1909	(+)
<i>Svastra basirufa</i> Brethes, 1910	(+)
<i>Svastra brachyura</i> Brethes, 1910	(+)
<i>Svastra corduvensis</i> Brethes, 1910	(+)
<i>Svastra mimetica</i> Brethes, 1910	(+)
<i>Svastra patagonica</i> Brethes, 1910	(+)
<i>Svastra segmentaria</i> Brethes, 1910	
<i>Svastra subapicalis</i> Brethes, 1910	(+)
<i>Svastra tucumana</i> Brethes, 1910	(+)
<i>Tetralonia chubutina</i> Brethes, 1910	(+)
<i>Tetralonia fulva</i> Brethes, 1910	(+)
<i>Tetralonia saltensis</i> Brethes, 1910	(+)
<i>Tetralonia sobria</i> Brethes, 1910	(+)
<i>Tetralonia spegazzini</i> Brethes, 1910	(+)
<i>Tetralonia squalida</i> Brethes, 1910	(+)
<i>Tetralonia tornowii</i> Brethes, 1910	(+)
<i>Tetralonia vara</i> Brethes, 1910	(+)
<i>Tetrapedia taricensis</i> Brethes, 1910	
<i>Trigona bipunctata limae</i> Brethes, 1910	(+)
<i>Trigona martinezii</i> Brethes, 1920	(+)
<i>Tyreothremma paraguayanum</i> Brethes, 1909	(+)
<i>Xylocopa paraguayensis</i> Brethes, 1909	(+)
<i>Xylocopa bellula</i> Brethes, 1916	(+)
<i>Xylocopa boliviiana</i> Brethes, 1916	(+)
<i>Xylocopa guaranica</i> Brethes, 1916	(+)
<i>Xylocopa martinezii</i> Brethes, 1916	(+)
<i>Xylocopa rotundisouta</i> Brethes, 1916	(+)

## Field Trip to Argentina: The Pleasures of the Panurginae

by

Jerome G. Rozen, Jr.  
Department of Entomology,  
The American Museum of Natural History,  
Central Park West at 79th Street,  
New York, New York 10024-5192

At the end of March 1990, I returned from a second field trip, of a month's duration, to the northwestern provinces of Argentina. This expedition followed a visit there in October–December 1989. On both trips I was accompanied by Arturo Roig-Alsina (Department of Entomology, Snow Entomological Museum, University of Kansas, Lawrence, KS, 66045), who is a specialist on

parasitic bees. The purpose of both trips was to carry out studies dealing with the comparative nesting biology, mating behavior, floral preferences and immature stages of panurgine bees. The investigations were sponsored by the National Geographic Society. Luisa Ruz (Laboratorio de Zoología, Universidad Católica de Valparaíso, Chile) joined the expedition for about a week, and we were briefly accompanied on a side trip to the high Andes by Manfredo Fritz (Ministerio de Educación, Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto de Investigaciones Entomológicas Salta).

The bee fauna of northwestern Argentina is surprisingly rich in species and genera. As an example, only four species of *Callonychium* had been described from Argentina, one of which is assignable to the subgenus *Paranychium*. As the result of these trips we identified 8 or 9 new species of *Callonychium* and 6 of these belong to the subgenus *Paranychium*. Specimens of 3 undescribed panurgine genera were also collected. This wealth of new taxa indicates that previous researchers in Argentina concentrated on large bees and overlooked smaller species. The moral of the story: if we are going to understand the diversity of the bee fauna of the world, we must not overlook the smaller species.

## Bees at Canberra

by

R. P. Mactarlane

New Zealand Department of Scientific and Industrial Research, Plant Protection, Private Bag, Christchurch, New Zealand

In the spring of 1988, I visited Canberra to attend the general assembly of the International Union of Biological Sciences as the delegate of the International Commission of Bee Plant Relationships. On October 15–16, modest numbers of 2 medium-sized halictid species were collected from a range of flowers, but only mint bushes *Prostanthera* supported more than one species. No colletids or other native bees were active. One or two hover fly species visited the various flowering *Eucalyptus*, *Grevillea*, *Telopea*, *Indigofera*, *Phebellium*, and apple flowers. In October in New Zealand, *Lasiglossum* would have been foraging on at least some of these flowers.

At the Australian National Insect Collection, Entomology Division, CSIRO, Canberra is completing a catalogue of Australian Apoidea. Most of the 10 drawers of non-Australian bees are from Papua New Guinea, Indonesia, the Solomon Islands and Malaysia. There was one drawer of bumble bees, mainly with older exchange specimens. The drawer with carpenter bees *Xylocopa* had about 200 bee specimens. Any lack of bees from the Cocos, Christmas and Norfolk Islands should not be due to an insufficient collecting effort as there were extensive collections of robber flies from these islands, but no specimens of Asilidae from Lord Howe Island.

## Bee Research in India

by

Rajiv K. Gupta

Lecturer, P.G. Department of Zoology

Raj Rishi (Autonomous) College, Alwar 301001, India

I have been collecting and identifying non-apoid bees, especially Megachilidae, from throughout India since 1980. Thanks to all apoid workers for their cooperation in this effort. I can now identify any megachilid collected in India, and shall be happy to do so. Recently, I received financial assistance for the continuation of this research. I would be grateful to anyone sending recent reprints on bee research, or apoid catalogues.

## *Perdita* Nesting Sites

by

Bryan Danforth

Department of Entomology,  
Smithsonian Institution, 105 NHB,  
Washington, DC 20560

Finding nests of solitary or communal ground nesting bees can be very frustrating business. This frustration can be greatly magnified when the bees are minute and the areas available for nesting are vast.

It is becoming increasingly evident that the nesting biology of bees in the genus *Perdita* shows great variability with some species exhibiting flightless, macrocephalic males which remain in the natal nest, and considerable interspecific variation in the numbers of females inhabiting nests. In order to find out more about the nesting biology of this genus of bees, I would like to know of localities where nests have been found, or sites which support large *Perdita* populations. Please send information to the above address.

I am particularly interested in the subgenus *Pseudomacroterea* (*Perdita turqiceps*) which occurs in Inyo, San Bernardino and Riverside counties, California. Any help would be gratefully received.

## List of Endangered Species

by

Ronald J. McGinley

The Department of the Interior has published a list of endangered vertebrates and invertebrates which includes bees such as *Perdita hirticeps*, *P. scitula antiochensis*, *Bombus franklini* and fifty-seven species of *Nesopropis* from Hawaii. For further information contact Mr. William Knapp, Chief, Division of Endangered Species and Habitat Conservation, U.S. Fish and Wildlife Service, Washington, D.C. 20240. 703-235-2771 or FTS 235-2771.

## RESEARCH NEWS

**Molecular Genetics of *Apis mellifera*:  
Genome Mapping**

by  
**Winfred W. Doane**  
 Department of Zoology,  
 University of Arizona, Tucson, AZ 85721

My laboratory has become actively involved in research on the molecular genetics of the honey bee, *Apis mellifera* L., in recent years, and I am now planning a major program to physically map the genome of the honey bee. This will be in collaboration with Dr. Charles P. Milne, Jr., at Washington State University and with the cooperation of the Carl Hayden Bee Research Laboratory in Tucson, Arizona, and the nearby Center for Insect Science at the University of Arizona. We would like the cooperation and input of other honey bee researchers interested in bee genetics for this program to be a success. As a newcomer to bee genetics, I find it an exciting challenge and believe that the time is ripe to prepare molecular linkage maps for the 16 chromosomes that make up the honey bee genome. I have been developing strategies on the best way to do this and, with the help of Dr. Milne and others, I believe we can begin to make headway in honey bee genetics. Having spent a long career in the developmental and molecular genetics of *Drosophila melanogaster*, I believe that I shall be able to contribute significantly to the success of this project.

I would appreciate input and comment from anyone interested in bee genetics.

**Research at the USDA Bee Lab,  
Weslaco, TX**

by  
**Evan A. Sugden**  
 USDA-ARS Honey Bee Research,  
 2413 E. Hwy. 83,  
 Weslaco, TX 78596

Currently I am a Research Associate at USDA-ARS Honey Bee Lab, Weslaco, TX. My responsibilities include maintenance of the Africanized honey bee trap line along the Texas-Mexico border, feral honey bee studies, and contributing to the African Honey Bee selective abatement Pilot Project. My other interests include feral honey bee ecology, theoretical and applied pollination, non-*Apis mellifera* bee biology, especially alodapine species (in particular the Australian genus *Exoneura* and allies) and Megachilidae. My previous appointments include Exotic Pest Analysis Staff, California Department of Food and Agriculture 1987-88, post-doctoral studies (honey bee-native and bee competition), Australian Museum, Sydney, New South Wales, 1985-87, and private apicultural development, Dubai, United Arab Emirates 1984-85.

**Sizing It Up: Recent Perspectives on  
Colony Growth and  
Reproduction in Bumble Bees**

by  
**Richard M. Fisher<sup>1</sup> & Nelson Pomeroy<sup>2</sup>**  
 Department of Botany and Zoology,  
 Massey University,  
 Palmerston North, New Zealand  
<sup>1</sup> Present address: Department of Biology, Acadia University, Wolfville,  
 Nova Scotia, CANADA BOP 1X0  
<sup>2</sup> Email address: N.Pomeroy@Massey.AC.NZ

We welcome this opportunity to inform *Melissa* readers of collaborative work we have conducted recently in our ongoing bumble bee domestication research. While we appear to have solved many of the intransigent problems which have previously limited attempts to mass-rear bumble bees for applied pollination purposes, there is one particular aspect of *Bombus* biology to which we keep returning: that of worker numbers. Like colonies of other annual social insects, those of *Bombus* vary tremendously in terms of the number of workers each produces, and our domesticated species, *B. terrestris*, is no exception. While we are now quite proficient at starting and maintaining colonies on a large scale, we still lack control over how large a given colony will become (i.e., how many workers it will have produced before the onset of male and queen production). We believe the following influences are important and need further investigation:

1) The role of competition in limiting colony productivity. Here we define productivity as the total biomass of workers and reproductives produced in a given nest. While much of our research has been directed towards greenhouse pollination, we have placed bumble bee colonies in the field for kiwifruit pollination trials, and to minimize labour costs have allowed the bees to maintain themselves outdoors during the long New Zealand spring. These colonies show considerable variance in colony productivity even though they have all been treated the same way prior to placement outside. We consider the role of intra- and interspecific competition to be very important in mediating these size differences and at the same time acknowledge competition as perhaps the thorniest influence to examine experimentally in the field. Bumble bees do not appear to be very bold practitioners of interference competition, although references to interference effects on their foraging behaviour appear from time to time in the literature (e.g. Thomson 1989). Thus, competition effects on colony size are likely to be subtle and hard to prove. Observations of differences in colony productivity (Fisher & Pomeroy 1989) or foraging efficiency (Plowright et al. 1978) between areas of apparently different colony densities suggest a food supply limitation effect on colony size, although we are still unable to demonstrate that competition is the underlying process which limits food supply. David Woodward, a PhD student at Massey, has recently studied the possible displacement of *Bombus terrestris* from *Borago officinalis* flowers by honey bees as the temperature increases during the day.

**2) The timing of reproduction in limiting worker numbers.** A large problem that we and others working in this area have faced is that bumble bee colonies in many cases do not follow the 'bang-bang' strategy of optimal reproduction, first espoused by Oster and Wilson (1978). In contrast to their view that it is in the colony's best interest to rear the worker 'factory' before using it to rear reproductives, we find that reproductives, particularly males, often trickle out of a colony's reproductive assembly line at irregular intervals. Some species seem to be more prone to doing this than others. Tantalizingly, other *Bombus* spp. (and *Vespula* spp.; Albert Greene, personal communication) DO seem to follow the Oster and Wilson, 'workers-first' reproductive strategy.

What influences this early pattern of reproduction, and is there anything we can do to control it, in order to produce colonies which are rich in workers? Early, or staggered reproduction may be a consequence of the battle between the queen and her workers for the control of reproductive investment in a bumble bee colony. If larval ejection, oophagy and out-and-out fisticuffs among the bees are any indication of this battle, it must surely be a ferocious and potentially debilitating one in many *Bombus* species (and again, tantalizingly, not such a source of contention in others). In one approach, cuckoo bumble bees (*Psithyrus* spp.) were used to assay possible limits on worker numbers, based on the supposition that these parasites, which live in prolonged intimacy with their hosts, would likely reflect the ways in which reproductive conflict limits colony size and reproduction. The results indicate that it is difficult for social parasites to effectively police a large colony of host bees, and that the smaller size of bumble bee colonies parasitized by *Psithyrus*, which has often been reported in the literature, may be a compromise between decreased reproductive output and better overall control of host bees by parasites.

Large colonies are unwieldy replicator vehicles for any individual, particularly the queen, which would attempt to monopolize brood rearing for its own reproductive interests in the face of large numbers of discriminating and often insubordinate nestmates. Although in New Zealand the very largest *terrestris* colonies usually produce many hundreds of queens, and the smallest produce none, there is great variation in sex ratio in the intermediate sizes. Unlike honeybees, which use worker police to maintain a type of 'community conscience' about male production (Ratnieks and Visscher 1989), anarchy appears to be the name of the game in *Bombus* societies once the queen has lost her ability to dominate workers. What promotes long-term dominance by the queen and relative stability in bumble bee social structure? We are very keen to obtain an answer to this question.

The ability to discriminate the sex and parentage of eggs and larvae, representing the offspring of many possible reproductive rivals within the nest, is an expected consequence of conflict and has been our second approach in exploring colony size. We have tried to meddle with colony ontogeny by adding male and female larvae to

pre-worker colonies and observing the effects of our 'insubordination' on queen behaviour and reproduction. Our results (Fisher & Pomeroy 1990) indicate that *B. terrestris* queens are capable of sexing larvae, and that the presence of reproductive brood at times when it doesn't normally occur produces anomalies in brood rearing. While experiments like these don't tell us how to stop colonies from reproducing early, they do provide information about some of the underlying proximate mechanisms which may influence the cessation of worker production in favour of males and queens. Larval cuticular hydrocarbon profiles may provide the basis on which discrimination by adult bees is made; profiles differ significantly in terms of the types of compounds produced by each sex. Dr. David Greenwood (Department of Scientific and Industrial Research, Biotech Division, Palmerston North, New Zealand) is currently assisting us with chemical analyses.

Another approach we are employing to study temporal trends in bumble bee reproduction is a consideration of heritable factors. Such factors are often overlooked in short-term studies, but are more accessible to us now that better methodologies for long-term *Bombus* culture are available. We hope that selective mating experiments will demonstrate to what extent colony size is controlled by these factors, and whether artificial selection among queens is a viable and desired option.

**3) Environmental influences.** It is in this area that we feel the greatest immediate gains in knowledge regarding variable colony size can be made. Proximate external factors affecting colony growth are rather malleable experimentally. Bumble bees, for example, harbour a veritable witch's brew of internal and external parasites which may act alone or in concert to debilitate the health of bees and hence their brood-rearing capabilities. We have concentrated on studies of *Nosema bombi*, a cosmopolitan microsporidian which is of special interest because it appears to weaken bumble bees but not necessarily to kill them outright. Further, its effects can be controlled through antibiotic treatment, allowing us to manipulate its epidemiology to some extent. Bumble bee tracheal mite (*Bombycarus buchneri*) is another parasite which we view with suspicion in terms of possible effects on colony size. External agents which affect 'vigour' have long been thought to influence colony growth, but to our knowledge remain unquantified parameters in social insect population biology. The parasites we have mentioned appear to rely on transmission via fall queens, and as such are ideal candidates for a sublethal infection which may affect colony strength without killing all the nest members.

So far, our attempts to inoculate bees with *Nosema* spores have met with mixed results. Like other intimate host-parasite relationships, heritable resistance may complicate inoculation procedures and success. Is the disease absent in many springqueens, for instance, because they have not had contact with it, or because they

are resistant to it? As a consequence, much of our *Nosema* research has involved post hoc analysis to compare observed incidences of infection with colony performance in the field (Fisher & Pomeroy 1989). Catherine McIvor is continuing this work as an MSc thesis at Massey University.

Our results so far suggest that it may be difficult to tease apart the various, possibly interactive factors affecting bumble bee colony size. Nonetheless, investigations in this area must continue if we are to succeed in controlling bumble bee colony growth in the following important areas: reliability and predictability.

## References

- Fisher, R.M. and N. Pomeroy. 1989. Incipient colony manipulation, *Nosema* incidence and colony productivity of the bumble bee *Bombus terrestris*. J. Kansas Entomol. Soc. 62:581-589.
- Fisher, R.M. And N. Pomeroy. 1990. Sex discrimination and infanticide by queens of *Bombus terrestris* (Hymenoptera: Apidae). Anim. Behav. 39:801-802.
- Oster, G.F. and E.O. Wilson. 1978. Caste and ecology in the social insects. Princeton Univ. Press, New Jersey.
- Plowright, R.C., B.A. Pendrel and I.A. McLaren. 1978. The impact of aerial fenitrothion spraying upon the population biology of bumble bees (*Bombus Latr.:Hym.*) in south-western New Brunswick. Can. Ent. 110:1145-1156.
- Ratnieks, F.L.W. and P. Kirk Visscher. 1989. Worker policing in the honeybee. Nature 342:796-797.
- Thomson, J.D. 1989. Reversal of apparent feeding preferences of bumble bees by aggression from *Vespula* wasps. Can. J. Zool. 67:2588-2591.

## Future Directions of Bee Research

by  
The Editors

In the previous section we presented short accounts of research in progress. In the spirit of R.P. Macfarlane's article on important and unsolved problems in bee research (*Melissa* 3), we again encourage people to send us their opinions on the future directions of bee research. What are the most important, and critical areas of bee research? Given finite resources, what questions demand the attention of researchers, and what areas of research are the most likely to yield significant advances in our understanding of bee biology and evolution? Is the ultimate question in bee biology, "why do some bees collect a small subsample of all available pollens available (i.e., oligoleptic species) while others collect a wide variety (polylectic species)?" Or are the phylogenetic placement of the family Colletidae, or the shortage of parasitic species in Australia, the crucial, unanswered questions. We hope this section of the newsletter will provide a forum for people to discuss future research projects and potential areas of collaboration.

## ADVANCES IN ELECTRONIC COMMUNICATION

### Rapid Social Communication among bee researchers via worldwide computerized networking

by  
E.E. Southwick  
Department of Biology  
State University of New York  
College at Brockport, NY 14420 USA

There is now the availability of an electronic communication network to help bee researchers to keep abreast of developing events in research and to keep in better contact with each other. This electronic network allows for very rapid spread of information and access to electronic file transfer among professional electronic mail addresses. This system is now up and running... worldwide. I am writing to encourage all researchers to "subscribe" to the network (which means get yourself into the system). A brief description follows along with the directions on how to get on (for BITNET users, or EARN users in European countries). If you are unsure of your electronic mail status (most university campuses in the United States are on BITNET), check with your campus computer people and show them this publication. You must have access to a terminal or a modem on a micro-PC. We might be able to give access in the future to other systems including public or commercial bulletin boards.

The communication System: BEE-L is a network file list dedicated to research and information concerning the biology of BEES! This includes solitary and even social bees (and maybe even wasps). We communicate about sociobiology, behavior, ecology, adaptation/evolution, genetics, taxonomy, physiology, pollination and nectar/pollen biology. Also, it is a great way to announce meetings and professional opportunities, seek help in research work or research funding, and discuss ideas.

#### How to subscribe:

You need to send a message via BITNET to :  
type in >> LISTSERV@ALBNYVM1  
then type >> SUB BEE-L your full name

Then you are on! It is that simple. Sit back and enjoy receiving the messages from BEE-L!

#### How to unsubscribe:

you need to send a message via BITNET to:  
type in >> LISTSERV@ALBNYVM1

then type >> SIGNOFF BEE-L

#### How to send a message to somebody else:

You need to send a mail via BITNET to:

type in >> BEE-L@ALBNYVM1 (note: not LISTSERV)  
the type in your message>>

and it goes out to everyone. You may send an entire text file (ASCII) instead of, or in addition to, a message.

How to get professional addresses so you can send a message to just one specific person:

You need to send a message via BITNET to:  
type in>> LISTSERV2ALBNYVM1  
then type>> REVIEW BEE-L

and you get back the list of subscribers' addresses and names. Contributions sent to this list are automatically archived.

You can obtain a list of the available archive files by sending:

INDEX BEE-L command to LISTSERV2ALBNYVM1.  
These files can then be retrieved by means of a command: GET BEE-L filename

More information may be obtained from the snail-mail address above.

Evolution and behavior of macrocephalic males in *Perdita*. B.N. Danforth, Dept. Entomol., Smithsonian Inst., NHB-105, Washington, DC 20560

Multimodal size distributions of bee faunas of North America and elsewhere. J. Seger, & J. Burner, Dept. Biol., Univ. Utah, Salt Lake City, UT 84112, J. W. Stubblefield, CERA, Charles Square, 20 Univ. Dr., Cambridge, MA 02138 & V. J. Tepedino, USDA Wild Bee Labs, UMC 53, Utah State Univ., Logan, UT 84322.

Modelling parental investment: cell construction and sex ratios. R.W. Rust, Dept. Biol. Univ. Nevada, Reno, NV 89557-0015.

Trap-nesting bees: when are the ecological interpretations qualitative, quantitative or artificial? P.F. Torchio, USDA Wild Bee Labs, UMC 53, Utah State Univ., Logan, UT 84322.

Ecology and origins of mites in solitary bee nests. G.C. Eickwort, Dept. Entomol., Cornell Univ., Ithaca, New York 14853.

Evolution of cleptoparasitism in bees. B.A. Alexander, Dept. Entomol., Univ. Kansas, Lawrence, KS 66045.

\* \* \*

## MEETING ANNOUNCEMENTS

### Ecology of Non-Social Bees

Reno, Nevada in December, 1991  
(Symposium for Section C, 1991  
ESA meetings)

Organizers:

Jim Cane, Dept. Entomol.  
Auburn Univ., Auburn, AL 36849-5413  
Tele: 205 844-2559

Rich Rust, Dept. Biol.  
Univ. of Nevada, Reno, NV 89557-0015  
Tele: 702 784-6188

Pollen-foraging of specialist and generalist bees at *Solanum*. S.L. Buchmann, USDA ARS, Carl Hayden Bee Research Labs, 2000 E. Allen Rd., Tucson, AZ 85719 & J. H. Cane., Dept. Entomol., Auburn Univ., Auburn, AL 36849-5413.

Bees as herbivores: reproductive consequences of pollen specialization. J.H. Cane, Dept. Entomol., Auburn Univ., Auburn, AL 36849-5413 & J. A. Payne, S' east Fruit & Tree Nut Research Labs, P.O. Box 87 Byron, GA 31008.

Native bees as pollinators of endangered plants. V.J. Tepedino, USDA Wild Bee Labs, UMC 53, Utah State Univ., Logan, UT 84322.

### Non-*Apis* Bees and Their Role as Crop Pollinators Logan, Utah in August, 1992

by

John D. Vandenberg  
Bee Biology and Systematics Lab,  
Utah State University, Logan, Utah, 84322-5310

An International Workshop on Non-*Apis* Bees and Their role as Crop Pollinators will be held in Logan, Utah, USA in August, 1992. Its purpose will be to facilitate exchange of current information on all aspects of bee biology and to improve prospects for establishing non-*Apis* bees as crop pollinators. The four-day workshop will include both invited symposia and contributed papers. Informal meetings and small workshops are encouraged and will be facilitated. English will be the language of the workshop.

The meeting will be hosted by the Bee Biology and Systematics Laboratory (USDA Agricultural Research Service) and by Utah State University, Logan, Utah. The conference will take place on the University campus where food services and living accommodations will be available. Off campus housing can also be obtained.

DAY 1 will be devoted to the biology, nest associates and management of leafcutting bees (*Megachile*). DAY 2 will be set aside for similar discussions of other pollinators, including *Bombus* and *Osmia*. DAYS 3 and 4 will be reserved for bee ecology, behavior, evolution, biosystematics and other basic research topics.

Logan is situated in a beautiful valley in the Wasatch Mountains of northern Utah. Typical August weather is dry with warm days and cool nights. Commercial alfalfa seed fields pollinated by the alfalfa leafcutting bee (*M. rotundata*) can be visited within a half-day drive of Logan. Many outdoor activities are available, and seven U.S. National Parks are within a one-day drive.

To be put on the mailing list for future announcements and for further information, please contact: Dr. John D. Vandenberg, USDA-ARS Bee Biology and Systematics Laboratory, Utah State University, Logan, Utah, USA, 84322-5310.

## Social Insect Studies in the Soviet Union Rybnoe (Ryazan District) in September, 1992

by

Vladilen E. Kipyatkov  
Department of Entomology, Faculty of  
Biology, Leningrad State University,  
Universetskaya nab., 7/9, Leningrad, 199034.  
U.S.S.R.

The All-Union Entomological Society Section for the Study of Social Insects was organized in 1989. The First Colloquium of the Section was held in Leningrad 2-8 October, 1990. The participants of the Colloquium have elected the Coordination Council of the Section (9 members; chairman - Dr. Vladilen E. Kipyatkov, secretary - Elena B. Lopatina) and discussed the main directions of further work. 31 papers on ants, bumble-bees, honey-bees, termites and wasps presented to Colloquium are printed now in the Proceedings of the First Colloquium (Leningrad, 1991, 212 pp., in Russian with English summaries for the most papers; available from V. E. Kipyatkov). We plan to establish a newsletter of the Section (two issues in a year, in two languages), to organize the Colloquia of the Section every two years and to publish their Proceedings in Russian and in English. The Second Colloquium of the Section will be organized in September, 1992 in Rybnoe (Ryazan District, about 160 km from Moscow). We would be happy to see our colleagues from other countries among the participants of the Second Colloquium. Interested persons and institutions may contact with Dr. Vladilen E. Kipyatkov at the address given above.



## BOOK REVIEWS

### New From Cambridge Press

The Edinburgh Building, Shaftesbury Road, Cambridge  
CB2 2RU  
Telephone (0223) 312393

Published June 1987:

### BUMBLEBEES

Oliver Prys-Jones and Sarah A. Corbet.  
Department of Applied Biology, University of Cambridge

Bumblebees are attractive and conspicuous insects and can often be identified without being caught. Their behaviour is readily observed and they form ideal subjects for field studies. This book introduces bumblebee natural history, emphasizing topics that offer scope for further research. The keys, supported by illustrations in colour and black and white, are designed to enable the reader to name the British species easily and reliably. By including keys, techniques and references as well as ideas, the book aims to encourage and facilitate individual studies of bumblebee biology by students at schools and universities and others interested in natural history.

**READERSHIP:** school biologists, entomologists, beekeepers, amateur and professional ecologists, university students for projects, general readers.

### NATURALISTS' HANDBOOK 6

210 x 148mm c. 96 pp. c. 80 line diagrams 4 colour plates

### CONTENTS:

- Editors' preface
- Acknowledgements
- 1 Introduction
- 2 Distribution and recognition
- 3 The natural history of true bumblebee (*Bombus*)
- 4 Nests and their establishment in captivity
- 5 Cuckoo bumblebees (*Psithyrus*), parasites and nest associates
- 6 Foraging behaviour
- 7 Identification
- 8 Techniques and approaches to original work
- Further reading
- Synonymy
- Index



## Recent Apoid Literature: 1987-1990

by The Editors

The following list of references has been compiled primarily from the National Agricultural Library's computer-based bibliographic retrieval system, as well as from reprints or lists of papers kindly sent to us by subscribers. Papers dealing specifically with the commercial aspects of *Apis mellifera* or *Megachile rotundata* have not been included, but papers of general biological interest have been included. Key words included most bee genera, familial and subfamilial names as well as pollination biology and bee/plant interactions. We admit in advance that these citations have not been checked for accuracy.

At present the roughly 800 references shown below are included in a single file created by v. 1.4 of the bibliographic software package Pro-Cite (Address: P.O. Box 4250, Ann Arbor, Michigan, 48106; 313-996-1580). We would like to make this and expanded lists available to other Pro-Cite users, or users of other, compatible, bibliographic programs. We hope in the future to send out this list on disk to readers who agree to add references or check those already input. Additionally, it is planned to supplement this list with references gathered from Biological Abstracts or Zoological Record, which should extend the list back to the late 1970's.

As a preliminary step towards making this reference list available to *Melissa* readers on disk, we would be willing to send copies in the form of a WordPerfect 5.0 text file to anyone who sends us a request and a blank 3.5 or 5.25 inch diskette.

Please send us comments on the utility of the reference list and how we might make it more generally available to researchers. And please continue to send reprints and reference lists to Ronald McGinley for inclusion in upcoming lists.

Abrol, D. P. (1987). Analysis of environmental factors affecting foraging behaviours of *Megachile bicolor* F. on *Crotalaria juncea* L. Proc. Indian Natl. Sci. Acad. Part B Biol. Sci., 53(1), 21-26.

Abrol, D. P. (1988). Ecology and behavior of three bee species pollinating loquat *Eriobotrya japonica* Lindley. Proc. Indian Natl. Sci. Acad. Part B Biol. Sci., 54(2-3), 161-164.

Abrol, D. P. (1988). Foraging range of subtropical bees *Megachile flavipes*, *Megachile nana* (Hymenoptera, Megachilidae) and *Apis florea* (Hymenoptera, Apidae). J. Indian Inst. Sci., 68(1-2), 43-48.

Abrol, D. P. (1989). Foraging range of alfalfa pollinating *Megachile flavipes* Spinola and *Megachile nana* Bingham (Megachilidae: Hymenoptera). J. Anim. Morphol. Physiol., 35(1), 11-16.

Abrol, D. P. (1987). Influence of thermal and energetic constraints on the pollination activity of carpenter bee *Xylocopa pubescens*. Environ. Ecol. (Kalyani), 5(1), 90-93.

- Abrol, D. P. (1988). Pollination studies in almond (*Prunus amygdalus* L.). Current Sci., India, 57(7), 397-398.
- Abrol, D. P. (1988). Significance of thermal and visual constraints on the pollination activity of *Andrena ilerda* Cameron. J. Anim. Morphol. Physiol., 35(1), 7-10.
- Abrol, D. P. (1989). Studies on ecology and behavior of insect pollinators frequenting strawberry blossoms and their impact on yield and fruit quality. Trop. Ecol., 30(1), 96-100.
- Abrol, D. P., & Bhat, A. A. (1989). *Xylocopa valga* new record Gerstacker (Hymenoptera, Anthophoridae) from India. Curr. Sci., 58(1), 41.
- Abrol, D. P., & Kapil, R. P. (1987). Factors affecting pollination activity of *Megachile lanata* Lepel. Proc. Indian Acad. Sci. Anim. Sci., 95(6), 757-770.
- Abrol, D. P., & Kapil, R. P. (1988). Pollination studies in *Crotalaria juncea* L. Sci. Cult., 54(7), 243-244.
- Ackerman, J. D. (1989). Geographic and seasonal variation in fragrance choices and preferences of male euglossine bees. Biotropica, 21(4), 340-347.
- Alcock, J. (1989). Size variation in the anthophorid bee *Centris pallida*: new evidence on its long-term maintenance. J. Kans. Entomol. Soc., 62(4), 484-489.
- Alcock, J., & Houston, T. F. (1987). Resource defense and alternative mating tactics in the Banksia bee, *Hylobaeus alcyoneus* (Erichson). Ethol., 76(3), 177-188.
- Alcock, J., & Smith, A. P. (1987). Hilltopping, leks and female choice in the carpenter bee *Xylocopa* (*Neoxylocopa*) *varipuncta*. J. Zool., 211, 1-10.
- Alexander, B., & Rozen Jr., J. G. (1987). Ovaries, ovarioles, and oocytes in parasitic bees (Hymenoptera: Apoidea). Pan-Pac. Entomol., 63(2), 155-164.
- Ali, T., & Ali, S. I. (1989). Pollination biology of *Calotropis procera* subsp. *hamiltoni* (Asclepiadaceae). Phyton, 29(2), 175-188.
- Alzuet, A. D. B. De, & Abrahamovich, A. H. (1987). Deutonymphs (Hypopi) of the genera *Sennertia oudemans*, 1905 and *Horstia oudemans*, 1905 (Acari: Astigmata) on *Xylocopa* (*S.*) *splendidula* Lepeletier, 1841 Hymenoptera: Apoidea). Rev. Soc. Entomol. Argentina, 44(3-4), 345-351.
- Anasiewicz, A. (1986)(1988). Insects pollinating flowers of celery, parsley and fennel II. Hymenoptera with special reference to the superfamily Apoidea. Biul. Lubel. Tow. Nauk. Biol., 28(1), 39-44.
- Andersen, J. F., Buchmann, S. L., Weisleder, D., Plattner, R. D., & Minckley, R. L. (1988). Identification of thoracic gland constituents from male *Xylocopa* spp. Latreille (Hymenoptera, Anthophoridae) from Arizona USA. J. Chem. Ecol., 14(4), 1153-1162.
- Anderson, G. J., & Symon D. (1988). Insect foragers on *Solanum* flowers in Australia. Ann. Mo. Bot. Gard., 75(3), 842-852.
- Andersson, S. (1988). Size-dependent pollination efficiency in *Anchusa officinalis* (Boraginaceae): causes and consequences. Oecologia, 76(1), 125-130.

- Andreev, A. V., Stratan, V. S., & Ursu, E. N. (1988). Bees (Hymenoptera, Apoidea) - pollinators of Rosaceae in Moldavia. *Izv. Akad. Nauk. Mold. SSR Ser. Biol. Khim. Nauk.*, (6), 41-45.
- Andreev, A. V., & Stratan, V. S. (1989). Ecological features of bees (Hymenoptera, Apoidea) - pollinators of fruit bearing plants and conditions of their existence in man made habitats. *Izv. Akad. Nauk. Mold. SSR Biol. Khim. Nauk.*, (1), 43-47.
- Andreev, A. V., Stratan, V. S., & Ursu, E. N. (1988). Fauna of bees (Hymenoptera, Apoidea) pollinators of fruit and berry plants belonging to rosales. *Izv. Akad. Nauk. Mold. SSR Ser. Biol. Khim. Nauk.*, (6), 41-45.
- Archer, M. E. (1988). The aculeate wasps and bees (Hymenoptera, Aculeata) of my local patch Strensall Common the first 70 visits. *Naturalist (Leeds)*, 113 (984), 25-30.
- Archer, M. E. (1988). The aculeate wasp and bee assemblage (Hymenoptera: Aculeata) of a woodland: Bernwood forest in the English midlands. *Entomol.*, 107(1), 24-33.
- Archer, M. E. (1989). The wasps and bees (Hymenoptera, Aculeata) of Allerthorpe Common England UK before and after coniferization. *Naturalist*, 114(991), 129-136.
- Arduser, M. S., & Michener, C. D. (1987). An African genus of cleptoparasitic halictid bees (Hymenoptera: Halictidae). *J. Kans. Entomol. Soc.*, 60(2), 324-329.
- Armbruster, W. S., & Guinn, D. A. (1989). The solitary bee fauna (Hymenoptera: Apoidea) of interior and arctic Alaska: flower associations, habitat use, and phenology. *J. Kans. Entomol. Soc.*, 62(4), 468-483.
- Armbruster, W. S., Keller, S., Matsuki, M., & Clausen, T. P. (1989). Pollination of *Dalechampia magnifolia* (Euphorbiaceae) by male euglossine bees. *Am. J. Bot.*, 76(9), 1279-1285.
- Armbruster, W. S., & Mziray, W. R. (1987). Pollination and herbivore ecology of an African *Dalechampia* (Euphorbiaceae): comparisons with new world species. *Biotropica*, 19(1), 64-73.
- Arrez, P., De Dantis, L., Angelica, G. M., & Lamborot, L. (1989). Presence in Chile of *Aprostocetus daira* new record Walker (Hymenoptera, Eulophidae). *Acta Entomol. Chil.*, 15(0), 275-276.
- Aston, T. J. (1987). Plant-pollinator interactions: a rich area for study. *J. Biol. Educ.*, 21(4), 167-174.
- Ayasse, M. (1987). The function of volatiles in mate choice of sweat bees (Hymenoptera: Halictidae). *Apidologie*, 18(4), 371-373.
- Babu, G. V. S., Bhatt, S. C., & Pandeya, S. C. (1987). Pollination ecology of *Solanum sisymbifolium* Lamk. *Proc. Indian Acad. Sci. Plant Sci.*, 97(4), 295-300.
- Baldovski, G. (1987). Neuere Funde von Wildbienen des Genus *Hylaeus* F. im Gebiet der DDR (Hymenoptera, Apoidea) und einige Bemerkungen zu aktuellen Problemen des Artenschutzes. *Entomol. Nachr. Ber.*, 31(1), 11-18.
- Banaszak, J. (1989). Bees (Apoidea) of moist meadows on the Mazovian Lowland. *Memorabilia Zool.*, 43, 279-287.
- Banaszak, J. (1987). Bees (Hymenoptera: Apoidea) of selected plant associations at Wielkopolski National Park (WNP). *Badania Fizjogr. Polska Zachodnia Ser. C: Zool.*, (35), 5-23.
- Banaszak, J. (1988). The collection of Aculeata (Hymenoptera) of Prof. Edward Lubicz-Niezbątowski. *Acta Univ. Lodz. Folia Zool. Anthr.*, 6, 73-81.
- Banaszak, J. (1987). Fauna of wild bees (Hymenoptera: Apoidea) of Wielkopolsko-Kujawska lowland during the last fifty years. *Badania Fizjogr. Polska Zachodnia Ser. C: Zool.*, (36), 67-77.
- Banaszak, J. (1989). Phenology of bees (Apoidea) of central Poland lowlands. *Badan. Fizjogr. Pol. Zachod. Ser. C: Zool.*, 38, 123-136.
- Banaszak, J., & Manole, T. (1987(1988)). Diversity and density of pollinating insects (Apoidea) in the agricultural landscape of Rumania. *Pol. Pismo. Entomol.*, 57(4), 747-766.
- Bastos, M. D. N. D. C. (1988). Contributions to the systematic study of some species of the genus *Machaerium* Persoon Leguminosae Papilionoideae occurring in Brazilian Amazonia. *Bol. Mus. Para. Emilio. Goeldi. Ser. Bot.*, 3(2), 183-278.
- Batra, S. W. T. (1987). Ethology of the vernal eusocial bee, *Dialictus laevissimus* (Hymenoptera: Halictidae). *J. Kans. Entomol. Soc.*, 60(1), 26-34.
- Baumgartner, D. L., & Roubik, D. W. (1989). Ecology of *Necrophilous* and filth-gathering stingless bees (Apidae: Meliponinae) of Peru. *J. Kans. Entomol. Soc.*, 62(1), 11-22.
- Beard, D. A. (1987) (September). Are bees pests? Watch over Essex, 8.
- Bego, L. R. (1989). Behavioral interactions among queens of the polygynic stingless bee *Melipona bicolor bicolor* Lepeletier (Hymenoptera, Apidae). *Braz. J. Med. Biol. Res.*, 22(5), 587-596.
- Bejko, V. B. (1987). Nest orientation in higher Hymenoptera. In: Ushatinskaya, R.S. (Ed.) *Problems of ecological physiology in insects*. Nauka, Moscow.,
- Belakova, A. (1986(1988)). Bees (Hymenoptera, Apoidea) in the vicinity of Bratislava, Czechoslovakia. *Acta Fac. Rerum. Nat. Univ. Comeniane. Zool.*, (31), 21-30.
- Belles, X., Galore, A., & Ginebreda, A. (1987). Taxonomic potential of the chemical constituents in the cephalic marking secretions of *Bombus* and *Psithyrus* species (Hymenoptera, Apidae); a numerical taxonomic study. *Apidologie*, 18(3), 231-242.
- Berezin, M. V., Berezina, N. V., & Zakhvatkin, Y. A. (1987). Species composition of the bumble bee population pollinating clover on the Mikhailovskoe training and experimental farm. *Izv. Timiryazev. Sel'Skokhozyaistvennoi Akademii*, (1), 185-191.

- Berger, L. A., Moffett, J. O., & Rummel, D. R. (1988). The foraging of *Agapostemon angelicus* Cockerell relative to hybrid cottonseed production in Texas. Southwest. Entomol., 13(1), 47-54.
- Berger, L. A., Vassiere, B. E., Moffett, J. O., & Merritt, S. (1988). *Bombus* spp. (Hymenoptera: Apidae) as pollinators of male sterile upland cotton on the Texas High Plains. Environ. Entomol., 17(5), 789-794.
- Bergstrom, G. (1987). On the role of volatile chemical signals in the evolution and speciation of plants and insects: why do they smell and why do they smell differently? Ser. Entomol., 41, 321-327.
- Bernhardt, P. (1987). A comparison of the diversity, density, and foraging behavior of bees and wasps on Australian Acacia. Ann. Mo. Bot. Gard., 74(1), 42-50.
- Bernhardt, P., & Thein, L. B. (1987). Self-isolation and insect pollination in the primitive angiosperms: new evaluations of older hypotheses. Plant. Syst. Evol., 156(3-4), 159-176.
- Berry, I. (1987). Dead bees don't pollinate. Orchardist of New Zealand, 60(8), 287.
- Bertsch, A. (1987). Flowers as food sources and the cost of outcrossing. Ecol. Stud. Anal. Synth., 61, 277-293.
- Bhat, A. A. (1987). Studies on queen rearing and breeding of *Apis cerana indica* F. (Apidae, Hymenoptera). Haryana Agric. Univ. Thesis Abstr., 12(4), 299-300.
- Bhatnagar, S., & Mishra, R. M. (1987). Pollinia loads on the sexes of the two carpenter bees *Xylocopa spinola* and *Xylocopa fenestrata* Fabr. visiting flowers of *Calotropis procera* R. Br. Asclepiadaceae. Proc. Natl. Acad. Sci. India Sect. B: (Biol. Sci.), 57(2), 171-174.
- Bichee, S. L. (1988). Effect of different modes of pollination in sunflower *Helianthus annus* L. (Compositae) sown on different days by *Trigona iridipennis* Smith. Apidacta, 23(3), 65-68.
- Billen, J. P. J. (1987). New structural aspects of the Dufour's and venom glands in social insects. Naturwiss., 3 (7), 340-341.
- Birhman, R. K. (1988). Pollination mechanisms in *Solanum chacoense* Bitt. Acta Bot. Indica., 16(1), 89-91.
- Blochtein, B., & Wittmann, D. (1988). Mating site specificity, reproduction and vector selection in *Nemognatha nigrotarsata* (Col., Meloidae), a nest parasite of leaf-cutter bees and other pollinators of crops in Rio Grande Do Sul. J. Appl. Entomol., 105, 414-419.
- Blom, J. v. der. (1989). Mutual benefit for reproducing and non-reproducing females in nests of the carpenter bee *Xylocopa pubescens* Spinola. Actes. Colloq. Insectes Soc., 5, 131-136.
- Blom, J. v. der., & Velthuis, H. H. W. (1988). Social behaviour of the carpenter bee *Xylocopa pubescens* (Spinola). Ethology, 79(4), 281-294.
- Bodnarchuk, L. I., Radchenko, V. G., & Vladimirkii, A. A. (1988). *Megachile willughbiella* - a promising alfalfa pollinator in the south-east Ukraine. Vestn. Zool., (2), 38-40.
- Bodnarchuk, L. I., Shalimov, I. I., & Olifir, A. A. (1989). Incubation of leaf-cutting bees with automatic removal of parasitic Hymenoptera. Vestn. Zool., (1), 79-81.
- Bohart, G. E., & Griswold, T. L. (1987). A revision of the Dufoureinae genus *Micralictoides* Timberlake (Hymenoptera: Halictidae). Pan-Pac. Entomol., 63(2), 178-193.
- Bohrer, K. (1987). Do sweat bees also have a trophogenic basis of caste development? Apidologie, 18(4), 369-371.
- Bohrer, K. (1987). Larval Food and Caste Differentiation in *Lasioglossum malachurum*. (Larvennahrung und Kastendifferenzierung bei *Lasioglossum malachurum* (Hymenoptera: Halictidae)). Diplomarbeit, - (-), 87.
- Bohrer, K., & Smith, B. H. (1987). Analysis of free fatty acids in the larval provisions of the halictine bee *Lasioglossum (Evylaeus) malachurum*. In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.,
- Bolchi Serini, G. (1988)(1989). Sulla morfologia delle appendici genitali maschili di alcuni *Bombus* Latr. (Hymenoptera: Bombidae). [On the morphology of the male genitalia of some *Bombus* Latr. (Hymenoptera: Bombidae)]. Boll. Zool. Agar. Bachic, 20, 1-15.
- Bollingmo, T. (1989). Worker-Queen conflict and fitness consequences in a colony of *Bombus hypnorum* L. (Hymenoptera). Fauna Norv. Ser. B., 36(1), 69-73.
- Bonetti, A. M., & Kerr, W. E. (1987). Sex determination in bees XXIII. Brood of a laying queen treated with juvenile hormone III. Rev. Bras. Genet., 10(3), 593-597.
- Bootsma, M. C., Sommeijer, M. J., & Punt, W. (1988). Etude des ressources florales de *Melipona trinitatis* (Hym., Meliponinae) a quatre localites a Trinidad, Antilles. Actes Colloq. Insectes Soc., 4, 295-299.
- Borg-Karlson, A.-K. (1987). Chemical basis for the relationship between *Ophrys* orchids and their pollinators. III. Volatile compounds of species in the *Ophrys* orchids sections Fuciflorae and Bombyliflorae as insect mimetic attractants/excitants. Chem. Scripta, 27, 313-325.
- Bourke, A. F. G. (1988). Worker reproduction in the higher eusocial Hymenoptera. Q. Rev. Biol., 63(3), 291-312.
- Broemeling, D. K. (1988). A revision of the *Nomada* subgenus *Nomadita* of North America (Hymenoptera: Anthophoridae). Pan-Pac. Entomol., 64(4), 321-344.
- Broemeling, D. K., & Moalif, A. S. (1988). A revision of the *Nomada* subgenus *Pachynomada* (Hymenoptera: Anthophoridae). Pan-Pac. Entomol., 64(3), 201-227.

- Brooks, R. W. (1988). Systematics and phylogeny of the anthophorine bees (Hymenoptera: Anthophoridae; Anthophorini). *Univ. Kans. Sci. Bull.*, 53(9), 437-575.
- Brooks, R. W., & Griswold, T. L. (1988). A key to the species of *Trachusa* subgenus *Heteranthidium* with descriptions of new species from Mexico (Hymenoptera: Megachilidae; Anthidiini). *J. Kans. Entomol. Soc.*, 61(3), 332-346.
- Brooks, R. W., & Michener, C. D. (1988). The Apidae of Madagascar and nests of *Liotrigona* (Hymenoptera). *Sociobiology*, 14(2), 299-323.
- Bruneau, A., & Anderson, G. J. (1988). Reproductive biology of diploid and triploid *Apis americana* Leguminosae. *Am. J. Bot.*, 75(12), 1876-1883.
- Buchmann, S. L. (1989). Bees assess pollen returns while sonicating *Solanum* flowers. *Oecologia*, 81(3), 289-294.
- Buchmann, S. L. (1987). The ecology of oil flowers and their bees. *Ann. Rev. Ecol. Syst.*, 18, 343-369.
- Bullock, S. H., Ayala, R., Baker, I., & Baker, H. G. (1987). Reproductive biology of the tree *Ipomoea wolcottiana* (Convolvulaceae). *Madrono*, (4), 304-314.
- Bullock, S. H., Martinez Del Rio, C., & Ayala, R. (1989). Bee visitation rates to trees of *Prokia crucis* differing in flower number. *Oecologia (Berl.)*, 78(3), 389-393.
- Buren, N. W. M. van, & Sommeijer, M. J. (1988). Study of determinant factors for the reproductive dominance of the queen of *Melipona trinitatis*. *Actes Colloq. Insectes Soc.*, 4, 285-290.
- Burnistrov, A. N., & Gorin, A. D. (1987). Artificial rearing of wild pollinators. *Pchel.*, (6), 28-29.
- Camargo, J. M. F. (1988). Meliponinae (Hymenoptera, Apidae) of the collection of the Instituto di Entomologia Agraria, Portici, Italy. *Rev. Bras. Entomol.*, 32(3/4), 351-374.
- Camargo, J. M. F., Moure, J. S., & Roubik, D. W. (1988). *Melipona yucatanica* new species (Hymenoptera: Apidae: Meliponinae) stingless bee dispersal across the Caribbean Arc and post-eocene vicariance. *Pan-Pac. Entomol.*, 64(2), 147-158.
- Camargo, J. M. F., & Moure, J. S. (1988). Notes on the Meliponinae (Hymenoptera; Apidae) collected by Filippo Silverstri in the Rio de La Plata Basin Argentina, Brazil, and Paraguay. *Rev. Bras. Entomol.*, 32(2), 293-314.
- Cameron, P. J., Hill, R. L., Valentine, E. W., & Thomas, W. P. (1987). Invertebrates imported into New Zealand for biological control of invertebrate pests and weeds, for pollination, and for dung dispersal, from 1874 to 1985. *Bull. Dep. Sci. Industr. Res., New Zealand*, (242), 51pp.
- Cameron, S. A. (1989). Temporal patterns of division of labor among workers in the primitively eusocial bumble bee *Bombus griseocollis* (Hymenoptera: Apidae). *Ethology*, 80(1-4), 137-151.
- Camillo, E., & Garofalo, C. A. (1989). Analysis of the niche of two sympatric species of *Bombus* (Hymenoptera, Apidae) in southeastern Brazil. *J. Trop. Ecol.*, 5 (pt.1), 81-92.
- Camillo, E., & Garofalo, C. A. (1989). Social organization in reactivated nests of three species of *Xylocopa* (Hymenoptera, Anthophoridae) in southeastern Brazil. *Insects Soc.*, 36(2), 92-105.
- Cane, J. H. (1989). Nesting biology notes for *Perdita (Alloperdita) bradleyi* Viereck (Hymenoptera: Andrenidae). *J. Kans. Entomol. Soc.*, 62(1), 140-143.
- Cane, J. H., & Payne, J. A. (1988). Foraging ecology of the bee *Habropoda laboriosa* (Hymenoptera: Anthophoridae), an oligolege of blueberries (Ericaceae: *Vaccinium*) in the southeastern USA. *Ann. Entomol. Soc. Am.*, 81(3), 419-427.
- Caron, D. M. (1987). Stingless bees a vanishing art. *Glean. Bee Cult.*, 115(12), 711-712.
- Cartar, R. V., & Dill, L. M. (1990). Why are bumble bees risk-sensitive foragers? *Behav. Ecol. Sociobiol.*, 26 (2), 121-127.
- Castro, L. (1987). New records of 3 Bombinae (Hymenoptera, Apidae) of the Iberian Peninsula. *Bol. Asoc. Esp. Entomol.*, 11(0), 413.
- Castro, L. (1988). On *Bombus reinigiiellus* Rasmont 1983 (Hymenoptera, Apidae). *Bol. Asoc. Esp. Entomol.*, 12 (0), 281-289.
- Celary, W. (1989). Wild bees (Hymenoptera: Apoidea) of the Bedkowska Valley Poland Czestochowa Upland Part I. (Colletidae: Halictidae). *Acta Biol. Cracov. Ser. Zool.*, 30(0), 1-18.
- Celary, W., & Dylewska, M. (1988). (Colletidae: Hymenoptera; Apoidea) of Poland. *Pol. Pismo. Entomol.*, 58(2), 359-382.
- Celli, G., & Porrini, C. (1988). Death of bees and pesticide residues in bees and hives in Italy 1983-1986. *Boll. Ist Entomol. Univ. Stud. Bologna*, 42(0), 75-86.
- Charnetski, W. A. (1988). Toxicity of insecticides to the alfalfa leafcutter bee *Megachile rotundata* F. (Hymenoptera, Megachilidae). *Can. Entomol.*, 120(4), 297-306.
- Cibula, D. A., & Zimmerman, M. (1987). Bumblebee foraging behavior: changes in departure decisions as a function of experimental nectar manipulations. *Am. Midl. Nat.*, 117(2), 386-394.
- Corbet, S. (1987). More bees make better crops. *New Sci.*, 115(1570), 40-43.
- Corbet, S. A., Chapman, H., & Saville, N. (1988). Vibratory pollen collection and flower form: bumble-bee on *Actinidia*, *Symphytum*, *Borago* and *Polygonatum*. *Funct. Ecol.*, 2(2), 147-155.
- Corbet, S., & Westgarth-Smith, A. (1988). A buzz in the air. *Nat. World*, (23), 16-18.
- Costa-Leonardo, A. M., Neves, N. A., & Cruz-Landim, C. (1987). Aspectos morfológicos da diferenciação dos músculos do voo de *Scaptotrigona postica* (Hymenoptera: Apidae) durante metamorfose. *Cien. Cult.*, 39(9), 852-857.

- Costelloe, B. H. (1988). Pollination ecology of *Gentiana andrewsii*. Ohio J. Sci., 88(4), 132-138.
- Couto, R. H., Costa, J. A., & Silveira, R. C. M. D. (1988). Effect of pollination on seed production in *Crotalaria juncea*. Cientifica (Jaboticabal), 16(1), 85-88.
- Cripps, C., & Rust, R. W. (1989). Pollen preferences of seven *Osmia* species. (Hymenoptera: Megachilidae). Environ. Entomol., 18(1), 133-138.
- Cripps, C., & Rust, R. W. (1989). Pollen foraging in a community of *Osmia* bees (Hymenoptera: Megachilidae). Environ. Entomol., 18(4), 582-589.
- Crozier, R. H., Smith, B. H., & Crozier, Y. C. (1987). Relatedness and population structure of the primitively eusocial bee *Lasioglossum zephyrum* (Hymenoptera: Halictidae) in Kansas. Evol., 41(4), 902-910.
- Cruz-Landim, C. D. (1987). Morphology of pheromone glands in Meliponinae (Hymenoptera, Apidae). In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.,
- Cruz-Landim, C. D., Silva de Moraes, R. L. M., & Costa-Leonardo, A. M. (1986-1987). Ultrastructure of the hypopharyngeal glands of *Melipona quadrifasciata anthidioides*. (Hymenoptera: Apidae). Naturalia, 11-12, 89-96.
- Cruz-Landim, C. D., & Zaniboni, M. C. (1986/1987). Study of relative size of various brain structures in bees (Hymenoptera: Apoidea). Naturalia, 11-12, 23-32.
- Csillary, G., Uncini, L., & Moor, A. (1987). Cross-pollination experiments with bees in Italy and Hungary. Capsicum Newsletter, (6), 30-32.
- Cure, J. R. (1989). Revisao de *Pseudagapostemon* Schrottky e descricao de *Oragapostemon*, gen. n. (Hymenoptera, Halictidae). Revta Bras. Ent., 33(2), 229-335.
- Current Research Information System. (1987). Bees and other pollinating insects research: a directory of USDA and state projects in CRIS. Cooperative State Research Service, Information Systems Division, Beneficial Insects Laboratory, Washington, D.C., U.S. Department of Agriculture.,
- Dafni, A., Bernadt, P., Schrimda, A., Ivri, Y., Greenbaum, S., O'Toole, C., & Losito, L. (1990). Red bowl-shaped flowers convergence for beetle pollination in the Mediterranean region. Isr. J. Bot. Basic Appl. Plant Sci., 39(1-2), 81-92.
- Dafni, A., & Calder, D. M. (1987). Pollination by deceit and floral mimesis in *Thelymitra antennifera* Orchidaceae. Plant Syst. Evol., 158(1), 11-22.
- Daly, H. V. (1988). Bees of the new genus *Ctenoceratina* in Africa south of the Sahara. (Hymenoptera: Apoidea). Univ. Cal. Pub. Entomol., 108, 1-69.
- Daly, H. V. (1988). Lectotype designations for African bees of the Genus *Ceratina* described by T. D. A. Cockerell (Hymenoptera, Apoidea). Pan-Pac. Entomol., 64(1), 87-92.
- Daly, H. V., Michener, C. D., Moure, J. S., & Sakagami, S. F. (1987). The relictual bee Genus *Manuelia* and its relation to other Xylocopinae (Hymenoptera: Apoidea). Pan-Pac. Entomol., 63(2), 102-124.
- Danforth, B. N. (1989). The evolution of hymenopteran wings: the importance of size. J. Zool., Lond., 218 (2), 247-276.
- Danforth, B. N. (1989). Nesting behavior of four species of *Perdita* (Hymenoptera: Andrenidae). J. Kans. Entomol. Soc., 62(1), 59-79.
- Danforth, B. N., & Michener, C. D. (1988). Wing folding in Hymenoptera. Ann. Entomol. Soc. Am., 81(2), 342-349.
- Danforth, B. N. (1990). Provisioning behavior and the estimation of investment ratios in a solitary bee, *Calloopsis (Hypomactotera) persimilis* (Cockerell) (Hymenoptera: Andrenidae). Behav. Ecol. Sociobiol., 27, 159-168.
- De Brujin, L. M. M., Sommeijer, M. J., & Dijkstra, E. (1989). Behavior of workers on waste dumps in the nests *Melipona favosa* (Apidae, Meliponini). Actes. Colloq. Insectes. Soc., 5, 31-37.
- De Brujin, L. M. M., Sommeijer, M. J., & Leys, R. (1989). Uptake and distribution of nectar and the storage of honey in *Melipona favosa* (Apidae, Meliponini). Actes. Colloq. Insectes. Soc., 5, 39-43.
- De Korte, M., Weissenbacher, B. K. H., & Crewe, R. M. (1988). Chemical signals in a stingless bee *Trigona (Meliplebeia) denoiti* Vachal (Hymenoptera: Apidae: Meliponinae). J. Entomol. Soc. South Afr., 51 (1), 9-16.
- Delfinado-Baker, M., & Aggarwal, K. (1987). A new *Varnoa* (Acari: Varroidae) from the nest of *Apis cerana* (Apidae). Internat. J. Acarol., 13(4), 233-237.
- Delfinado-Baker, M., & Baker, E. W. (1988). New mites (Acari, Laelapidae) from the nests of stingless bees (Apidae, Meliponinae) from Asia. Int. J. Acarol., 14(3), 127-136.
- Devall, M. S., & Thein L.B. (1989). Factors influencing the reproductive success of *Ipomoea pes-caprae* Convolvulaceae around the gulf of Mexico. Am. J. Bot., 76(12), 1821-1831.
- Devi K. R., Atluri J. B., & Reddi C. S. (1989). Pollination ecology of *Zizyphus mauritiana* Rhamnaceae. Proc. Indian Acad. Sci. Plant Sci., 99(3), 223-240.
- Dickman, G. (1987). Backyard superbees. Rodale Org. Gard., 34(11), 69-71.
- Dimitrov, P. (1987). Studies of species composition and numbers of bee pollinators (Apoidea) of lucerne during successive flowerings. Rastenievudni Nauki, 24(11), 75-80.
- Dimitrov, P. (1987). Studies on the diurnal flight activity of wild bees (Apoidea), pollinators of lucerne. Svremenno Postizheniya na Bulgarskata Zoologiya., 213-217.
- Dimitrov, P., Dimitrova, Z., Atanasov, N., & Vasileva, E. (1987). Specific population and number of bees (Apoidea) pollinating lucerne in the Rousse district (Bulgaria). Rastenievudni Nauki, 24(3), 58-62.

- Dobson, H. E. M. (1987). Role of flower and pollen aromas in host-plant recognition by solitary bees. *Oecologia* (Berl), 72(4), 618-623.
- Dobson, H. E. M. (1987). Role of pollenkit on pollen host-plant recognition by male and female solitary bees. In Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Dochkova, B. (1987). The toxic effect of some insecticides on the lucerne leaf-cutting bee - *Megachile rotundata* F. (Hymenoptera; Megachilidae). *Poch.-vozn. Agrokhim. Rositit. Zash.*, 22(5), 103-107.
- Dochkova, B. K., Atanasov, N., & Vasileva, E. I. (1987). Species composition and number of the wild bees (Hymenoptera, Apoidea) - pollinating on the forage loquats from the region of Pleven. *Acta Zool. Bulg.*, (35), 34-43.
- Dodson, C. A., Williams, N. H., & Whitten, W. M. (1989). A new species of *Gonora* (Orchidaceae) from Ecuador. *Lindleyana*, 4(1), 30-32.
- Donath, H. (1989). Considerable losses in bumble bees and other flower visiting insects due to exotic species of lime (*Tilia tomentosa* Moench, *Tilia euchlora* c. Koch). *Archiv. fur Naturschutz und Landschaftsforschung*, 29(2), 117-120.
- Doorn, A. V. (1989). Factors influencing dominance behaviour in queenless bumblebee workers (*Bombus terrestris*). *Physiol. Entomol.*, 14(2), 211-221.
- Dozhdikov, A. A. (1988). Domestication of bumblebees under conditions of Mari Autonomous Republic. Sov. Agric. Sci., (3), 72-75.
- Dreisig, H. (1989). Nectar distribution assessment by bumblebees foraging at vertical inflorescences. *Oikos*, 55 (2), 239-249.
- Duchateau, M. J., & Velthuis, H. H. W. (1989). Ovarian development and egg laying in workers of *Bombus terrestris*. *Entomol. Exp. Appl.*, 51(3), 199-213.
- Dulta, P. C., & Verma, L. R. (1987). Role of insect pollinators on yield and quality of apple fruit. *Indian J. Hortic.*, 44(3-4), 274-279.
- Dylewska, M. D. (1989). Apoidea of the Ojcow National Park, Poland, Part I. *Colletidae, Halictidae, Andrenidae, Melittidae, Megachilidae, Anthophoridae*. *Acta Biol. Cracov. Ser. Zool.*, 30(0), 19-72.
- Dylewska, M. D. (1987). The Genus *Andrena* Fabricius (Andrenidae, Apoidea) in northern and central Europe. *Acta Zool. Cracov.*, 30(12), 359-708.
- Dylewska, M. D. (1987). The genus *Andrena* in Poland. *Polskie Pismo Entomologiczne*, 57(3), 495-518.
- Eardley, C. D. (1987). Catalogue of Apoidea (Hymenoptera) in Africa south of the Sahara. Part I. The Genus *Xylocopa* Latreille (Anthophoridae). *Entomol. Mem.*, Dep. Agric. Wat. Supply Repub. S. Afr., (70), 1-20.
- Eardley, C. D. (1988). A revision of the genus *Lithurge* Latreille (Hymenoptera: Megachilidae) of sub-saharan Africa. *J. Ent. Soc. Sth. Afr.*, 51(2), 251-263.
- Eberhard, W. G. (1988). Group nesting in two species of *Euglossa* bees (Hymenoptera: Apidae). *J. Kans. Entomol. Soc.*, 61(4), 406-411.
- Ebmer, P. A. W. (1988). Critical list of non-parasitic Austrian Halictidae with regard to all Central European species. (Insecta: Hymenoptera: Apoidea: Halictidae). *Linzer Biol. Beitr.*, 20(2), 527-711.
- Ebmer, P. A. W. (1987). Die europäischen Arten der Gattungen *Halictus* Latreille 1804 und *Lasioglossum* Curtis 1833 Mit. Illustrierten Bestimmungstabellen (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). *Senckenb. Biol.*, 68(1-3), 59-148.
- Ebmer, P. A. W. (1988). Die Europäischen Arten der Gattungen *Halictus* Latreille 1804 und *Lasioglossum* Curtis 1833 mit Illustrierten Bestimmungstabellen. (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae) 2. Die Untergattung *Seladonia* Robertson 1918. *Senckenb. Biol.*, 68(4-6), 323-375.
- Efremova, Z. A. (1988). Fauna and ecology of cuckoo bumblebees (Hymenoptera, Apidae) of the Volga region U.S.S.R. *Entomol. Obozr.*, 67(3), 523-529.
- Efremova, Z. A. (1989). *Psithynus* bees and their ecology in the Volga Region U.S.S.R. (Hymenoptera, Apidae). *Entomol. Rev.*, 68(2), 64-71.
- Eickwort, G. C. (1990). Associations of mites with social insects. *Annu. Rev. Entomol.*, 35, 469-488.
- Eickwort, G. C. (1988). Distribution patterns and biology of west Indian sweat bees (Hymenoptera: Halictidae). In: Liebherr, J.K. (Ed). *Zoography of Caribbean insects*. Cornell University Press, Ithaca & London.
- Eickwort, G. C., & Kukuk, P. F. (1987). Reproductive castes in primitively eusocial Halictid bees. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Eickwort, G. C. (1988). The origin of mites associated with honey bees. Offprints from Needham et al (Eds.) *Africanised honey bees and bee mites*, Ellis Horwood Ltd., Chichester, England.
- Eisikowitch, D. (1987). *Calotropis procera* (Ait.) Ait. F. (Asclepiadaceae) and *Xylocopa* spp.: a study of inter-relationships. *Ser. Entomol.*, 41, 342-345.
- Eisikowitch, D., & Lupo, A. (1989). Wild flowers as competitors for pollinators in almond orchards. *Alon. Hanotea.*, 43(12), 1307-1312.
- Elisens, W. J. (1989). Genetic variation and evolution of the Galapagos Shrub Snapdragon. *Natl. Geogr. Res.*, 5(1), 98-110.
- Elisens, W. J., & Freeman, C. E. (1988). Floral nectar sugar composition and pollinator type among new world genera in tribe Antirrhineae (Scrophulariaceae). *Am. J. Bot.*, 75(7), 971-978.
- Elmqvist, T., Agren, J., & Tunlid, A. (1988). Sexual dimorphism and between-year variation in flowering, fruit set and pollinator behaviour in a boreal willow. *Oikos*, 53(1), 58-66.
- Engels, E., & Engels, W. (1988). Age-dependent queen attractiveness for drones and mating in the stingless bee *Scaptotrigona postica*. *J. Apic. Res.*, 27(1), 3-8.

- Engels, E., Engels, W., & Francke, W. (1987). Do stingless bees have two complexes of queen substances? In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.
- Engels, E., Engels, W., Schroder, W., & Francke, W. (1987). Intranidal worker reactions to volatile compounds identified from cephalic secretions in the stingless bee, *Scaptotrigona postica* (Hymenoptera, Meliponinae). *J. Chem. Ecol.*, 13(2), 371-386.
- Engels, W. (1987). Pheromones reproduction in Brazilian stingless bees. *Memorias Instit. Oswaldo Cruz*, 82 (Suppl. III), 35-45.
- Engels, W. (1987). Reproduction and caste development in social bees. In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.
- Engels, W. (1988). Reproductive strategies in bees. *Verehandl. Deut. Zool. Gesellschaft*, 81, 155-167.
- Equiarte, L., Martinez del Rio, C., & Arita, H. (1987). El nectar y el polen como recursos: el papel ecologico de los visitantes a las flores de *Pseudobombax ellipticum* (H.B.K.) Dugand. *Biotropica*, 19(1), 74-82.
- Eremina O. (1989). Effect of insecticides on bees and chinese silkworms. *Argokhimiya*, 0(5), 131-136.
- Erlandsson, S., Janzon, L.-A., & Svensson, B. G. (1988). Catalogus insectorum Sueciae (Hymenoptera, Apoidea) 1. Colletidae and Melittidae. *Entomol. Tidskr.*, 109(3-4), 161-163.
- Es'kov, E. K., & Markova, V. E. (1987). Effectiveness of alfalfa pollination by melliferous and other types of bees. *Dokl. Vses. Ordena. Lenina. Ordena. Trud. Krasnago. Znameni. Akad. S-Kh. Nauk. Im. V. I. Lenina.*, (5), 35-37.
- Evers, A. M. J. (1987). *Ebaeus nageli* Nov. Spec. aus Kamerun. 65. Beitrag zur Kenntnis der Malachiidae. *Entomol. Bl. Biol. Syst. Kaeffer.*, 83(2-3), 169-172.
- Fain, A., & Heard, T. A. (1987). Description and life cycle of *Cerophagus trigona* new species (Acar: Acaridae) associated with the stingless bee *Trigona carbonaria* Smith in Australia. *Bull. Inst. R. Sci. Nat. Belg. Entomol.*, 57(0), 197-202.
- Fairey, D. T., Lefkovitch, L. P., & Nelson, D. L. (1987). Viability of prepupae of the leafcutting bee *Megachile rotundata* Fab. during extended storage. *Can. J. Zool.*, 65(7), 1853-1856.
- Farinha, E. K., Torneiros, L. M., Silva de Moraes, R. L. M., & Cruz-Landim, C. d. (1988). Hormonal influence on the DNA content of salivary gland cells in larvae of *Melipona quadrifasciata anthidioides* Lep. (Apidae, Meliponinae). I. Juvenile hormone. *Naturalia*, 13, 75-83.
- Ferrazzi, P. (1989). Mallow. *Apic. Mod.*, 80(2), 89-93.
- Field, J. P., & Foster, W. A. (1988). The bees and wasps of Scott Head Island National Nature Reserve, Norfolk. *British J. Entomol. Nat. Hist.*, 1(2), 79-83.
- Fisher, R. M. (1987). Nectar and pollen collection by bumble bees: methodology for a colony-level approach. *J. N. Y. Entomol. Soc.*, 95(4), 504-508.
- Fisher, R. M. (1988). Observations on the behaviours of three European cuckoo bumble bee species (*Psithyrus*). *Insectes Soc.*, 35(4), 341-354.
- Fisher, R. M. (1987). Temporal dynamics of facultative social parasitism in bumble bees (Hymenoptera: Apidae). *Anim. Behav.*, 35(6), 1628-1636.
- Fisher, R. M., & Pomeroy, N. (1989). Incipient colony manipulation, nosema incidence and colony productivity of the bumble bee *Bombus terrestris* (Hymenoptera: Apidae). *J. Kans. Entomol. Soc.*, 62(4), 581-589.
- Fisher, R. M., & Pomeroy, N. (1989). Pollination of greenhouse muskmelons by bumble bees (Hymenoptera: Apidae). *J. Econ. Entomol.*, 82(4), 1061-1066.
- Flanigan, T. C. (1989). Protecting bees and saving bear with predator platforms. *Am. Bee J.*, 129(11), 721-722.
- Fonta, C., Masson, C., Genin, E., & Jullien, R. (1987). Comparative neurobiological and chemical data on communication in bumblebee males and females. In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.
- Fordham, R. A. (1989). Further New Zealand records of the new immigrant Australian bee *Hyleoides concinna* (Hymenoptera, Colletidae, Hylaeinae). *N. Z. Entomol.*, 12(0), 65-67.
- Foster, R. L., & Gamboa, G. J. (1989). Nest entrance marking with colony specific odors by bumble bee *Bombus occidentalis* (Hymenoptera: Apidae). *Ethol.*, 81 (4), 273-278.
- Francisco, E. R. N., & Felice, A. (1988). The Meliponinae and the American Indians. *Apicultura No. Brasil*, (28), 24-25.
- Frankie, G. W., & Vinson, S. B. (1987). Ecology of twelve sympatric *Centris* bee species in the Costa Rican dry forest. *Am. J. Bot.*, 74(5), 662.
- Frankie, G. W., Vinson, S. B., Newstrom, L. E., & Barthell, J. F. (1988). Nest site and habitat preference of *Centris* bees in the Costa Rican dry forest. *Biotropica*, 20 (4), 301-310.
- Franks, N. R. (1987). The parasitic strategies of a cuckoo bee. *Trends. Ecol. Evol.*, 2(11), 324-326.
- Free, J. B. (1987). Pheromones of social bees. Chapman & Hall, London.
- Frohlich, D. R., Clark, W. H., Parker, F. D., & Griswold, T. L. (1988). The xylophilous bees and wasps of a high cold desert Leslie Gulch, Oregon USA. (Hymenoptera: Apoidea; Vespoidea). *Pan-Pac. Entomol.*, 64(3), 266-269.
- Froneman, I. J. (1988). Subtropical crops. Pollination by bees. *Inform. Bull. Citrus Subtrop. Fruit Res. Inst. S. Africa*, (196), 12.
- Frye, R. D., Dix, M. E., & Carey, D. R. (1988). Effect of two insecticides of abundance on insect families associated with Siberian Elm Windbreaks. *J. Kans. Entomol. Soc.*, 61(3), 278-284.
- Gagnon, F. (1987). Construction of hives. *Sante Abeille*, (102), 315-317.

- Galen, C. (1989). Measuring pollinator-mediated selection on morphometric floral traits: bumblebees and the alpine sky pilot, *Polemonium viscosum*. *Evol.*, 43(4), 882-890.
- Galen, C., & Blau, S. (1988). Caste-specific patterns of flower visitation in bumble bees (*Bombus kirbyellus*) collecting nectar from *Polemonium viscosum*. *Ecol. Entomol.*, 13(1), 11-17.
- Galen, C., & Gregory, T. (1989). Interspecific pollen transfer as a mechanism of competition: consequences of foreign pollen contamination for seed set in the Alpine Wildflower, *Polemonium viscosum*. *Oecologia*, 81(1), 120-123.
- Galen, C., & Newport, M. E. A. (1987). Bumble bee behavior and selection of flower size in the sky pilot, *Polemonium viscosum*. *Oecologia*, 74(1), 20-23.
- Galen, C., & Newport, M. E. A. (1988). Pollination quality, seed set, and flower traits in *Polemonium viscosum*: complementary effects of variation in flower scent and size. *Am. J. Bot.*, 75(6), 900-905.
- Galen, C., & Stanton, M. L. (1989). Bumble bee pollination and floral morphology factors influencing pollen dispersal in the alpine sky pilot *Polemonium viscosum* Polemoniaceae. *Am. J. Bot.*, 76(3), 419-426.
- Gambino, P., Medeiros, A. C., & Loope, L. L. (1987). Introduced vespids *Paravespula pensylvanica* prey on Maui's endemic arthropod fauna. *J. Trop. Ecol.*, 3(2), 169-170.
- Gamboa, G. J., Foster, R. L., & Richards, K. W. (1987). Intraspecific nest and brood recognition by queens of the bumble bee, *Bombus occidentalis* (Hymenoptera: Apidae). *Can. J. Zool.*, 65(12), 2893-2897.
- Gangadhar, A., Subbaro, R., & Lakshminarayana, G. (1987). Synthesis of 18-octadecanolide, a major constituent of Dufour's gland secretion of *Colletes* bees. *J. Am. Oil Chem. Soc.*, 67(8), 1135-1137.
- Gardiner, B. (1989). Early nesters. *Bull. Amat. Entomol. Soc.*, 48(364), 110.
- Gebhardt, M., & Rohr, G. (1987). Zur Bionomie die Sandbienen *Andrena clarkella* (Kirby), *A. cinerea* (L.), *A. fuscipes* (Kirby) und ihrer Kuckuckbienen (Hymenoptera: Apoidea). *Drosera*, (2), 89-114.
- Gerling, D., Velthuis, H. H. W., & Hefetz, A. (1989). Bionomics of the large carpenter bees of the genus *Xylocopa*. *Annu. Rev. Entomol.*, 34, 163-190.
- Giblin-Davis, R. M., Norden, B. B., Batra, S. W. T., & Eickwort, G. C. (1990). Commensal nematodes in the glands, genitalia, and brood cells of bees (Apoidae). *J. Nemotol.*, 22(2), 150-161.
- Godzinska, E. J. (1988). Digging as tactics of escape in two bumblebee species with different nesting ecology *Bombus terrestris* L. and *Bombus pascuorum* Scopoli. *Acta Neurobiol. Exp.*, 48(5), 251-258.
- Godzinska, E. J. (1989). Extra-colony altruism in the bumblebees: Misbehavior in bumblebees. *Actes. Colloq. Insectes Soc.*, 5, 161-167.
- Goebel, R. L. (1988). Australian Native Bees. *Austr. Bee J.*, 69(2), 5-7.
- Goldblatt, J. W., & Fell, R. D. (1987). Adult longevity of workers of the bumble bees *Bombus fervidus* F. and *Bombus pennsylvanicus* (de Geer) (Hymenoptera: Apidae). *Can. J. Zool.*, 65(10), 2349-2353.
- Goldblatt, P., & Bernhardt, P. (1990). Pollination biology of *Nivenia* (Iridaceae) and the presence of heterostylous self-compatibility. *Isr. J. Bot. Basic Appl. Plant Sci.*, 39((1-2), 93-111.
- Golikov, V. I. (1987). Local wild bee species are the main alfalfa pollinators in the steppe zone of the north-western Transcaucasus USSR. *Nauchn. Dokl. Vyssh. Sch. Biol. Nauki. (Mosc.)*, (7), 49-53.
- Golikov, V. I., & Shabolta, O. M. (1988). Wild bees (Apoidae) as pollinators of Lucerne in Krasnodar region. *Zash. Semenov. Posev. Lyutserny Kompl. Vrednykh Organiz. Stepnoi Zone Sever. Kavk. Rostov-on-don, USSR: USSR Acad. Agric.*, 72-78.
- Goltz, L. (1988). Honey and pollen plants part X. Miscellaneous honey plants. *Am. Bee J.*, 128(2), 97-100.
- Gonzalez, J. M. (1989). Lista preliminar de los Euglossini (Hymenoptera: Apidae) del Marahuaka, Parque Nacional Duida, T.F.A., Venezuela. *Acta Terramaris*, (1), 81-83.
- Gonzalez, J. M. (1990). Comentarios bionómicos sobre dos nidos de *Euglossa cordata* L. (Hymenoptera: Apidae: Euglossini). *Bol. Entomol. Venez.*, 5(18), 141-143.
- Gorbunov, P. S. (1987). Endoparasitic flagellates of the genus *Critidilia* (Trypanosomatidae, Zoo-mastigophorea) from alimentary canal of bumblebees. *Zool. Zh.*, 66(12), 1775-1780.
- Gottsberger, G., Camargo, J. M. F., & Silberbauer-Gottsberger, I. (1988). A bee pollinated tropical community, the beach dune vegetation of Ilha De Sao Luis, Maranhao, Brazil. *Bot. Jahrb. Syst. Pflanzengesch. Pflanzengeogr.*, 109(4), 469-500.
- Goukon, K., Maeta, Y., & Sakagami, S. F. (1987). Seasonal changes in ovarian state in a eusocial halictine bee, *LasioGLOSSUM duplex*, based on stages of the oldest oocytes in each ovariole (Hymenoptera: Halictidae). *Res. Pop. Ecol.*, 29(2), 255-269.
- Greathead, D. J. (1989). A new species of *Anthrax* (Diptera, Bombyliidae) from *Megachile* spp. (Hymenoptera; Megachilidae) in the United Arab Emirates. *Entomol. Gaz.*, 40(1), 67-70.
- Greenberg, L. (1988). Kin recognition in the sweat bee, *Lasioglossum zephyrum*. *Behav. Genet.*, 18(4), 425-437.
- Gregg, K. B. (1989). Reproductive biology of the orchid *Cleistes divaricata* (L.) Ames var. *Bifaria* Fernald growing in a West Virginia meadow. *Castanea*, 54 (57-78).
- Griswold, T. L., & Michener, C. D. (1988). Taxonomic observations on Anthidiini of the western hemisphere (Hymenoptera: Megachilidae). *J. Kans. Entomol. Soc.*, 61(1), 22-45.

- Griswold, T. L., & Parker, F. D. (1988). New *Perdita* (*Perdita*) oligoleges of *Mentzelia* with notes on related species of the *Ventralis* group (Hymenoptera: Andrenidae). *Pan-Pac. Entomol.*, 64(1), 43-52.
- Griswold, T., & Parker, F. D. (1987). A new species of *Protosmia* Ducke from Spain with notes on related species (Hymenoptera: Megachilidae). *Psyche*, 94(1-2), 51-56.
- Guibu, L. S., Ramalho, M., Kleineri-Giovannini, A., & Imperatriz-Fonseca, V. L. (1988). Exploracao dos recursos florais por colonias de *Melipona quadrifasciata* (Apidae, Meliponinae). *Rev. Bras. Biol.*, 48(2), 299-305.
- Gupta, R. K. (1988). A new species of genus *Megachile* Latr. from Ludhiana, India (Insecta, Hymenoptera, Apoidea, Megachilidae). *Reichenbachia*, 26(11), 65-66.
- Gupta, R. K. (1988). A new species of genus *Cressoniella* Mitchell from South India (Insecta, Hymenoptera, Apoidea: Megachilidae). *Reichenbachia*, 26(12), 67-69.
- Gupta, R. K. (1987). On a new subgenus *Orientoheriades* and two new species of genus *Heriades* Spinola from India (Insecta, Hymenoptera, Apoidea, Megachilidae). *Reichenbachia*, 25((1-2)), 67-71.
- Gupta, R. K. (1988). Two new species of genus *Eumegachile* Friese from India (Insecta, Hymenoptera, Apoidea: Megachilidae). *Reichenbachia*, 26(13), 71-74.
- Gupta, R. K., & Tewari, V. K. (1987). A new species of genus *Lithurgus* Latreille (Hymenoptera: Apoidea: Megachilidae) from India. *J. Bombay Nat. Hist. Soc.*, 84(2), 405-407.
- Gupta, R. K., & Tewari, V. K. (1987). A new species of genus *Heriades* Spinola from India (Insects, Hymenoptera, Apoidea, Megachilidae). *Reichenbachia*, 25(1), 65-66.
- Gupta, U. S., & Banerjee, S. (1988). Sexual dimorphism in the hemolymph proteins of insects. *Proc. Natl. Acad. Sci. India Sect. B.*, 58(1), 17-22.
- Gyan, K. Y., & Woodell, S. R. J. (1987). Analysis of insect pollen loads and pollination efficiency of some common insect visitors of four species of woody Rosaceae. *Functional Ecol.*, 1(3), 269-274.
- Haeseler, V. (1987). Ants, wasps and bees of the Ipweger moor near Oldenburg I.O. Braunschweig Naturk. Schr., 2(4), 663-683.
- Haeseler, V. (1988). Colonization success of ants, wasps, and bees on young dune islands of the southern North Sea (Hymenoptera, Aculeata). *Drosera*, 88 (1-2), 207-236.
- Hagen, E. V. (1988). Bumble bees: identification, establishing colonies, rearing conservation. Melsungen, German Federal Republic; Verlag J. Neumann-Neudamm.
- Hallmann, M., Heller, C., & Hosenfeld, A. (1989). Die Ansiedlung und Haltung von Hummeln an Schulen. *Luscinia*, 46(3/4), 205-214.
- Hallmann, M. (1988). Die Besiedlung unterschiedlicher künstlicher Nisthilfen durch *Osmia rufa* L. (Hymenoptera: Megachilidae). *Nachr. Entomol. Ver. Apollo*, 9(3), 199-212.
- Hallmann, M. (1989). Ein Nistkasten zur künstlichen Ansiedlung von Hummeln der Gattung *Bombus* mit einem neuartigen Vorbau für einfache Stockeingangsdressuren zum Einsatz im aktiven Naturschutz. *Mitt. Int. Ent. Ver.*, 14((1/2), 49-60.
- Hallmann, M. (1989). Einige Beobachtungen zur Biologie der Solitarbiene *Osmia rufa* L. (Hymenoptera: Megachilidae). *Nachr. Entomol. Ver. Apollo*, 10 (2), 159-166.
- Hallmann, M., & Wilhelm, B. (1989). Einfache Versuche mit *Osmia rufa* L. als Motivation zum Artenschutz (Hymenoptera: Megachilidae). *Mitt. Int. Ent. Ver.*, 14(1/2), 39-48.
- Harder, L. D. (1988). Choice of individual flowers by bumble bees: interaction of morphology, time and energy. *Behaviour*, 104 (1-2), 60-77.
- Harder, L. D., & Real, L. A. (1987). Why do bumble bees risk adverse? *Ecol.*, 68(4), 1104-1108.
- Hirashima, Y. (1989). A synopsis of the bee genus *Palaeorhiza* perkins (Hymenoptera, Colletidae) of New Guinea part VII. Subgenus *Callorhiza* H. Esakia, (28), 1-10.
- Hoffmann, M., & Wittmann, D. (1987). Wild bee community in an agricultural area of Rio Grande do Sul, southern Brazil, and its impact on pollination of beans and sunflowers. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Horton, D., Payne, J., Amis, A., & Warnock, D. (1987). Insect management in Georgia rabbiteye blueberries. *Repr. U.S. Dep. Agric. Res. Ser.*, (80), 67-76.
- Hoshiba, H. (1988). Karyological analysis of a stingless bee, *Melipona favosa* (Apidae, Hymenoptera). *Cytologia*, 53(1), 153-156.
- Houston, T. F. (1987). Fossil brood cells of stenotritid bees (Hymenoptera: Apoidea) from the Pleistocene of South Australia. *Trans. R. Soc. S. Aust.*, 111(1-2), 93-97.
- Houston, T. F. (1989). *Leioproctus* bees associated with western Australian smoke bushes *Conospermum* spp. and their adaptations for foraging concealment (Hymenoptera: Colletidae: Paracolletini.). *Rec. West. Aust. Mus.*, 14(3), 275-292.
- Houston, T. F. (1987). A second contribution to the biology of *Ctenocolletes* bees (Hymenoptera: Apoidea: Stenotritidae). *Rec. West. Aust. Mus.*, 13(2), 189-201.
- Houston, T. F. (1987). The symbiosis of Acarid mites, genus *Ctenocolletacarus* (Acarina: Acariformes), and stenotritid bees, genus *Ctenocolletes* (Insecta: Hymenoptera). *Aust. J. Zool.*, 35(5), 459-468.
- Hung, A. C. F., & Norden, B. B. (1987). Biochemical systematics of bees in the *Ceratina calcarata dupla* complex. *Biochem. Syst. Ecol.*, 15(6), 691-693.

- Imperatriz, V. L. (1987). Swarm production by stingless bees. *Apic. No. Brasil*, (19), 26-27.
- Imperatriz-Fonseca, V. L., & Kleinert-Giovannini, A. (1987). The role of queens in stingless bee colonies. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social insects*. Verlag J. Peperny, Munich.,
- Imperatriz-Fonseca, V. L. (1988). How to use nucleus hives for the study (of stingless bees). *Apic. No. Brasil*, (25), 24-25.
- Inouye, D. W., & Pyke, G. H. (1988). Pollination biology in the snowy mountains of Australia: comparisons with Montane Colorado USA. *Austr. J. Ecol.*, 13(2), 191-210.
- Int. Com. Zool. Nomenc. (1988). *Halictus costulatus* Kriechbaumer 1873 currently *Lasiglossum costulatum* (Insecta, Hymenoptera) specific name conserved. *Bull. Zool. Nomenc.*, 45(3), 241.
- Ito, M. (1987). Geographic variation of an east Asian bumblebee *Bombus diversus* in some morphometric characters (Hymenoptera, Apidae). *Kontyu*, 55(2), 188-201.
- Jacob-Ramide, A. (1987). Des insectes peu connus: les abeilles solitaires. *Reserves Nat.* (Brussels), (4), 81-84.
- Jain, K. L. (1987). Analysis of nectar as a factor determining bee-flower relationship. *Haryana Agric. Univ. Thesis Abstr.*, 12(4), 305-306.
- James, D., Ariyanagam, R. P., & Duncan, E. J. (1989). Effects of tripping flowers of four early varieties of Pisum (Cicer arietinum L. Millsp.). *J. Hortic. Sci.*, 64(2), 243-247.
- Janzon, L. A., & Svensson, B. G. (1988). One hundred ten years with the carpenter bee *Ceratina cyanea* (Hymenoptera, Anthophoridae) in Sweden. *Entomol. Tidkr.*, 109(1), 19-23.
- Jedruszuk, A. (1987). Bees and their products as indicators of environmental pollution. *Med. Weter.*, 43(6), 352-356.
- Jennersten, O. (1988). Insect dispersal of fungal disease: effects of Ustilago infection on pollinator attraction in *Viscaria vulgaris*. *Oikos*, 51(2), 163-170.
- Jennersten, O. (1988). Pollination of *Viscaria vulgaris* (Carophyllaceae): the contributions of diurnal and nocturnal insects to seed set and seed predation. *Oikos*, 52, 319-327.
- Jennersten, O., Berg, L., & Lehman, C. (1988). Phenological differences in pollinator visitation, pollen deposition and seed set in the sticky catchfly, *Viscaria vulgaris*. *J. Ecol.*, 76(4), 1111-1132.
- Johnson, L. K. (1987). The pyrrhic victory of nest-robbing bees: did they use the wrong pheromone? *Biotropica*, 19(2), 188-189.
- Johnson, L. K., & Howard, J. J. (1987). Olfactory disc number in bees of different sizes and ways of life (Apidae: Meliponinae). *J. Kans. Entomol. Soc.*, 60(3), 380-388.
- Johnson, M. D. (1988). The relationship of provision weight to adult weight and sex ratio in the solitary bee, *Ceratina calcarata*. *Ecol. Entomol.*, 13(2), 165-170.
- Karron, J. D. (1987). The pollination ecology of co-occurring geographically restricted and widespread species of *Astragalus* (Fabaceae). *Biol. Conserv.*, 39 (3), 179-193.
- Kartsev, V. M. (1988). How do solitary bees locate flowers for the first time? *Priroda*, (5), 109.
- Katayama, E. (1988). Worker-like new queens in a colony of *Bombus ardens* (Hymenoptera, Apidae). *Kontyu*, 56(4), 879-891.
- Kato, M. (1988). Bumblebee visits to *Impatiens* spp.: pattern and efficiency. *Oecologia*, 76(3), 364-370.
- Kerr, W. E. (1987). Genetics of sex and caste determination. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Kerr, W. E. (1987). Genetik des Polymorphismus bei Biene. In: Schmidt, G.H. (Ed.) Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich. Wissenschaftliche Verlagsgesellschaft, Stuttgart.,
- Kerr, W. E. (1987). Geschlechts - und Kastendetermination bei Stachellosen Bienen. In: Schmidt, G.H. (Ed.) Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich, Wissenschaftliche Verlagsgesellschaft, Stuttgart.,
- Kerr, W. E. (1987). Sex determination in bees. XXI. Number of XO-Heteroalleles in a natural population of *Melipona compressipes fasciculata* (Apidae). *Insectes Soc.*, 34(4), 274-279.
- Kerr, W. E. (1987). Sex determination in bees. XVII. Systems of caste determination in the Apinae, Meliponinae and Bombinae and their phylogenetic implications. *Rev. Bras. Genet.*, 10(4), 685-694.
- Kerr, W. E., Absy, M. L., & Souza, J. M. (1986-1987). Espécies nectaríferas e polinizadoras utilizadas pela abelha *Melipona compressipes fasciculata* (Meliponinae, Apidae), no maranhão. *Acta Amazonica*, 16-17, 145-155.
- Kerr, W. E., & Rocha, R. (1988). Comunicacao em *Melipona rufiventris* e *Melipona compressipes*. *Cienc. Cult. (Sao Paulo)*, 40(12), 1200-1202.
- Kevan, P. G. (1987). Alternative pollinators for Ontario's crops: prefatory remarks to papers presented at a workshop held at the University of Guelph, 12 April, 1986. *Proc. Ent. Soc. Ont.*, 118, 109-110.
- Kevan, P. G., Mohr, N. A., Offer, M. D., & Kemp, J. R. (1988)(1989). The squash and gourd bee *Peponapis pruinosa* (Hymenoptera, Anthophoridae) in Ontario Canada. *Proc. Entomol. Soc. Ont.*, 199, 9-16.
- Khoo, S. G., & Yong, H. S. (1987). Nest structure and colony defence in the stingless bee, *Trigona terminata* Smith. *Nature Malaysiana*, 12(3), 4-15.
- Kim, C. W., & Ito, M. (1987). On the bumblebees (*Bombus*, *Psithyrus*) from the Korean Peninsula (Hymenoptera, Bombidae). *Entomol. Res. Bull.*, 13, 1-42.

- Kim, M. L., & Kim, C. W. (1989). Systematic study of Andrenidae from Korea (Hym.; Apoidea) (on the three new species and one new subspecies). *Korean J. Entomol.*, 19(3), 199-206.
- Kimsey, L. S. (1987). Generic relationships within the Eullossini (Hymenoptera: Apidae). *Syst. Entomol.*, 12(1), 63-72.
- Kirkpatrick, K. J., & Wilson, H. D. (1988). Interspecific gene flow in cucurbita: *C. texana* vs. *C. pepo*. *Am. J. Bot.*, 75(4), 519-527.
- Kleinert-Giovannini, A., & Imperatriz-Fonseca, V. L. (1987). Aspects of the trophic niche of *Melipona marginata* Lepeletier (Apidae, Meliponinae). *Apidologie*, 18 (1), 69-100.
- Kleinert-Giovannini, A., Imperatriz-Fonseca, V. L., & Rimalho, M. (1987). Exploitation of floral resources by *Plebeia saiqui* (Apidae, Meliponinae). In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Knerer, G. (1987). Photoperiod as cue for voltinism and caste regulation in halictine bees. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Knoll, F. R. N., Bego, L. R., & Imperatriz-Fonseca, V. L. (1987). Relative abundance and phenology of bees (Hymenoptera, Apoidea) in Sao Paulo, Brazil. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Knoll, F. R. N., & Imperatriz-Fonseca, V. L. (1987). Abundance and resources visited by individuals of *Tetragonisca angustula*. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Kolesnik, M. G. (1987). The role of useful entomofauna in Producing high carrot seed yields. *Nauchno-Tekhnicheskii Byull. Vsesoyuznogo Ordena Lenina, (175)*, 62-63.
- Korall, H., Leucht, T., & Martin, H. (1988). Bursts of magnetic fields induce jumps of misdirection in bees by a mechanism of magnetic resonance. *J. Comp. Physiol. A: Sens. Neural Behav. Physiol.*, 162(3), 279-284.
- Kosior, A. (1987). Impact of economic activity upon bumblebee *Bombus* Latr. Population in the west Bi-eszczady Mountains, Poland. *Ochr. Przyr.*, 45(0), 239-262.
- Kowalczyk J.K. (1988). Interesting Aculeata Hymenoptera in the Augustow forest, Poland. *Przegl. Zool.*, 32(4), 555-560.
- Kratochwil, A. (1988). Co-phenology of plants and anthophilous insects: a historical area-geographical interpretation. *Entomol. Gen.*, 13((1-2), 67-80.
- Kukal, O., & Pattie, D. L. (1988). Colonization of snow bunting, *Plectrophenax nivalis*, nests by bumblebees, *Bombus polaris*, in the high artic. *Can. Field Nat.*, 102(3), 544.
- Kukuk, P. F. (1989). Dispersal of males of the primitively eusocial sweat bee *Dialictus zephyrus* within a small nest aggregation. *Sociobiol.*, 15(1), 1-9.
- Kukuk, P. K., Crozier, R. H., Smith, D. R. R., Eickwort, G. C., & May, B. (1987). Macrogeographic genetic variation in the sweat bee *Lasioglossum (Dialictus) zephyrus*. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Kukuk, P. F., & Eickwort, G. C. (1987). Alternative social structures in halictine bees. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Kukuk, P. F., Eickwort, G. C., Raveret-Richter, M., Alexander, B., Gibson, R., Morse, R. A., & Ratnieks, F. (1989). Importance of the sting in the evolution of sociality in the Hymenoptera. *Ann. Entomol. Soc. Am.*, 82(1), 1-5.
- Kukuk, P. F., Eickwort, G. C., & May, B. (1987). Multiple maternity and multiple paternity in first generation brood from single foundress colonies of the sweat bee *Dialictus zephyrus* (Hymenoptera: Halictidae). *Soc. Insects*, 34(2), 131-135.
- Kukuk, P. F., & May, B. P. (1988). Dominance hierarchy in the primitively eusocial bee *Lasioglossum (Dialictus) zephyrus*: is genealogical relationship important? *Anim. Behav.*, 36(6), 1848-1850.
- Kukuk, P. F., & Schwarz, M. (1987). Intranest behavior of the communal sweat bee *Lasioglossum (Chilalictus) erythrurus* (Hymenoptera: Halictidae). *J. Kans. Entomol. Soc.*, 60(1), 59-64.
- Kukuk, P. F., & Schwarz, M. (1988). Macrocephalic male bees as functional reproductives and probable guards. *Pan-Pac. Entomol.*, 64(2), 131-137.
- Kulincevic, J. M., & Rothenbuhler, W. C. (1989). Variation in food consumed by caged worker bees from different colonies. *J. Apic. Res.*, 28(1), 26-29.
- Kunesch, G., & Zagatti, P. (1987). A fungal metabolite as the male wing gland pheromone of the bumblebee wax moth, *Aphomia sociella* L. Z. *Naturforsch.*, 42, 657-659.
- Kurennoi, N. M., & Mantoptin, A. I. (1988). Quality of apples in relation to bee pollination and flower thinning. *Sadovodstvo I Vinogradarstvo*, (5), 14-16.
- Kurosa, K. (1987). Two new *Chaetodactylus* (Acari, Chaetodactylidae) associated with *Osmia* (Hymenoptera, Megachilidae) in Japan. *Kontyu*, 55(2), 373-381.
- Kusumoto, K., Otsuka, Y., & Yahata, D. (1987). Wild bees (Hymenoptera: Apoidea (Apoidea) survey in the central part of Fukuoka prefecture. 2. Studies on the pollination of pear by *Osmia excavata* Alfken. *Proc. Assoc. Plant Prot. Kyushu*, 33, 199-203.
- Kwak, M. M. (1987). Pollination and pollen flow disturbed by honeybees in bumblebee-pollinated *Rhinanthus* populations? J. van Andel et al., (Eds.). *Disturbance in grasslands*. Dr. W. Junk Publishers, Dordrecht, Netherlands.,
- Kwak, M. M. (1988). Pollination ecology and seed-set in the rare annual species *Melampyrum arvense* L. (Scrophulariaceae). *Acta Bot. Neerl.*, 37(2), 153-163.

- LaBerge, W. E. (1989). A review of the bees of the genus *Pectinapis* (Hymenoptera: Anthophoridae). *J. Kans. Entomol. Soc.*, 62(4), 524-527.
- LaBerge, W. E. (1989). A revision of the bees of the genus *Andrena* of the western hemisphere. XIII. Subgenera *Simandrena* and *Taeniandrena*. *Trans. Am. Entomol. Soc.*, 115(1), 1-56.
- Lackie, P. M., Thomas, C. D., Brisco, M. J., & Hepper, D. N. (1986)(1988). On the pollination ecology of *Hamelia patens* (Rubiaceae) at Monteverde, Costa Rica. *Brenesia*, (25-26), 203-213.
- Lanham, U. N. (1987). *Andrena montrosensis* Viereck and Cockerell: evolutionary and nomenclatorial notes (Hymenoptera: Apoidea). *J. Kans. Entomol. Soc.*, 60(4), 576-577.
- Lanham, U. N. (1987). A new species of *Andrena* at the *Micrandrena*-*Scaphandrena* boundary (Hymenoptera, Apoidea). *Pan. Pac. Entomol.*, 63(4), 325-327.
- Lanham, U. N., & Weissmann, M. J. (1988). *Scaphandrena* and *Elandrena* (Hymenoptera: Andrenidae). *Pan-Pac. Entomol.*, 64(2), 183-184.
- Lanne, B. S., Bergstrom, G., Wassgren, A. B., & Tornback, B. (1987). Biogenetic pattern of straight chain marking compounds in male bumble bees. *Comp. Biochem. Physiol. B: Comp. Biochem.*, 88(2), 631-636.
- Laroca, S., Michener, C. D., & Hofmeister, R. M. (1989). Long mouthparts among 'short-tongued' bees and the fine structure of the labium in *Nittonia* (Hymenoptera, Colletidae). *J. Kans. Entomol. Soc.*, 62(3), 400-410.
- Laroca, S., Schwartz Filho, D. L., & Zanella, F. C. V. (1987). Nest of *Megachile habilis* subgenus *Austromegachile* and notes on *Megachile* (Apoidea: Megachilidae) species richness in neotropical sites. *Acta Biol. Parana*, 16(1-4), 93-106.
- Larsson, F. K., & Tengo, J. (1989). The effects of temperature and body size on the mating pattern of a gregariously nesting bee *Colletes cunicularis* (Hymenoptera: Colletidae). *Ecol. Entomol.*, 14(3), 279-286.
- Lavery, T. M., & Harder, L. D. (1988). The bumble bees of eastern Canada. *Can. Entomol.*, 120(11), 965-987.
- Lavery, T. M., & Plowright, R. C. (1988). Flower handling by bumblebees: a comparison of specialists and generalists. *Anim. Behav.*, 36(3), 733-740.
- Leclercq, J. (1989). Abeilles solitaires (Hymenoptera: Apoidea) trouvées par Fredi Darimont à la Montagne Saint-Pierre en 1933-1937. *Natura Mosana*, 42, 90-102.
- Leite, A., Neto, J. A., Leyton, J. T., Crivellaro, O., & El-Dorry, H. A. (1988). Phosphofructokinase from bumblebee flight muscle, molecular and catalytic properties and role of the enzyme in regulation of the fructose 6-phosphate/fructose 1,6-biphosphate cycle. *J. Biol. Chem.*, 263(33), 17527-17533.
- Lenardo, A. M. C., Neves, N. A., & Landim, C. D. C. (1987). Histological features of the differentiation of flight muscles of *Scaptotrigona postica* (Hymenoptera, Apidae) during metamorphosis. *Cienc. Cult. (Sao Paulo)*, 39(9), 852-857.
- Lenoir, A. (1987). Factors determining polyethism on social insects. *Exper. Suppl. (Basel)*, 54, 219-240.
- Leong, C. Y. (1986)(1987). Apiculture. *Ann. Report Agric. Commun. Dev., Dep. Agric.*, 160.
- Linsley, E. G. (1988). Further notes on halictine bees (Hymenoptera: Apoidea) visiting *Isomeris arborea* in southern California. *Pan-Pac. Entomol.*, 64(3), 275-279.
- Lipa, J. J., & Triggiani, O. (1988). *Critidia bombi* New species. A flagellated parasite of a bumble bee *Bombus terrestris* L. (Hymenoptera: Apidae). *Acta Protozool.*, 27(3-4), 287-290.
- Litt, R. (1988). Observations sur *Andrena fulva* Schrk. (Apidae, Hymenoptera). *Rev. Vervietoise Hist. Nat.*, 22-28.
- Lobreau-Callen, D., & Coutin, R. (1987). Floral resources exploited by some Apoidea in cultivated areas of arboreal savanna in Senegal during the rainy season. *Agron.*, 7(4), 231-246.
- Lobreau-Callen, D., & Suarez-Cervera, M. (1987). Relationship between the ultrastructure of the pollen exine, the pollination of flowers and the ecology of *Gaudichaudia cynanchoides* (Malpighiaceae). *Trav. Doc. Geogr. Trop.*, 59(74).
- Lombert, H. A. P. M., O'Connor, B. M., Lukoschus, F. S., & Whitaker Jr., J. O. (1987). Ontogeny, systematics and ecology of *Senertia (Ansennertia) americana* Delfinado & Baker, 1976 (Acari, Chaetodactylidae) from the nest of the carpenter bee *Xylocopa virginica* (Hymenoptera: Anthophoridae). *Internat. J. Acarol.*, 13(2), 113-129.
- Lomholdt, O. (1989). *Hylaeus pfankuchi* new record (Alfken, 1919) for Finland (Hymenoptera, Apidae). *Not. Entomol.*, 69(2), 57-58.
- Macfarlane, R. P., & Palma, R. L. (1987). The first record for *Melittobia australica* Girault in New Zealand and new host records for *Melittobia* (Eulophidae). *New Zealand J. Zool.*, 14, 423-425.
- MacGillivray, D. B. (1987). A centrifuging method for the removal of insect pollen loads. *J. Entomol. Soc. South Afr.*, 50(2), 522-523.
- Machado, M. F. P. S., & Contel, E. P. B. (1989). Glycerol-3-phosphate dehydrogenase (G-3-PDH; EC 1.1.1.8) variation in adult *Plebeia droryana* bees (Apidae, Hymenoptera). *Biochem. Genet.*, 27(7/8), 481-485.
- Maciera, O. J. D., & Hebling-Berlado, M. J. A. (1989). Laboratory toxicity of insecticides to workers of *Trigona spinipes* F. 1973 (Hymenoptera, Apidae). *J. Apic. Res.*, 28(1), 3-6.
- Macior, L. W. (1988). A preliminary study of the pollination ecology of *Pedicularis* Scrophulariaceae in Japan. *Plant Species Biol.*, 3(1), 61-66.

- Maddocks, R., & Paulus, H. F. (1987). Quantitative aspects of the breeding biology of *Osmia rufa* Latr. and *O. comuta* L. (Hymenoptera, Megachilidae): a comparative study of competition-reducing mechanisms in two closely related species. Zool. Jahr., Abteilung Syst., Okol. Geog. Tiere, 114, 15-44.
- Maeta, Y., Kubota, N., & Sakagami, S. F. (1987). *Nomada japonica* as a thelytokous cleptoparasitic bee, with notes on egg size and egg complement in some cleptoparasitic bees. Kontyu, 55(1), 21-31.
- Mahnert, V. (1987). New or badly known pseudoscorpion species mainly associated with insects from South America Arachnida. Mitt. Schweiz. Entomol. Ges., 60(3-4), 403-416.
- Malaspina, O., & Stort, A. C. (1987). Sucrose syrup-collecting behavior in Africanized and Caucasian bees and in the descendants of their crossings. Rev. Bras. Genet., 10(3), 459-470.
- Malysheva, N. Y. (1987). Description of the male of *Icteranthidium croceum* Mor. (Hymenoptera, Megachilidae). Entomol. Rev., 66(3), 131-133.
- Mann, G. S., & Sagar, P. (1987). Activity and abundance of flower visiting insects of loquat, *Eriobotrya japonica* (Thunb.) Lindl. Indian J. Hortic., 44(1-2), 123-125.
- Manson, D. C. M. (1988). The occurrence of the carpenter bee, *Xylocopa sonorina* in New Zealand. Weta., 11 (1), 4-5.
- Marikovskaya, T. P. (1989). Nesting of 2 Anthophorine species Hymenoptera. Izv. Akad. Nauk. Kaz. SSR Ser. Biol., (4), 67-73.
- Marikovskaya, T. P., & Shcherbakova, T. I. (1988). (Domestication of *Megachile centuncularis* L. in the foothills of Zailiiskii Alatau.). Vestn. S-Kh. Nauki (Kaz.), (9), 32-36.
- Marletto, F., Manino, A., & Saglia, A. (1988). Alfalfa-pollinating insects in the Susa Valley Italy. Apic. Mod., 79(4), 153-158.
- Marletto, F., & Manino, A. (1988). Insetti impollinatori di *Lupinella* coltivata in Piemonte per la produzione di seme. Apic. Mod., 79(2), 59-66.
- Marletto, F., Manino A., & Negri, J. M. (1988). Rearing of *Bombus* spp. in captivity. Apic. Mod., 79(5), 179-188.
- Martinez del Rio, C., & Eguiarte, L. E. (1987). The effect of nectar availability on the foraging behaviour of the stingless bee *Trigona testacea*. Southwest Nat., 32(3), 313-319.
- Mason, C. A. (1987). Division of labour in nestmate interactions in two species of allodapine bees. In: J. Eder & H. Remold (Eds.) Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.
- Mason, C. A. (1988). Division of labor and adult interactions in eusocial colonies of two allodapine bee species (Hymenoptera: Anthophoridae). J. Kans. Entomol. Soc., 61(4), 477-491.
- Mathew, G., Koshy, M. P., & Mohandas, K. (1987). Preliminary studies on insect visitors to teak (*Tectona grandis* Linn. F.) inflorescence in Kerala, India. Indian Forster, 113(1), 61-64.
- Matsuura, M. (1988). The life of social wasps and bees. Sapporo, Japan; Hokkaido University Press.
- May, P. G., & Spears Jr., E. E. (1988). Andromonoecy and variation in phenotypic gender of *Passiflora incarnata* Passifloraceae. Am. J. Bot., 75(12), 1830-1841.
- Mayer, D., Rathbone, L., & Miliczky, E. M. (1987). New ideas in cherry pollination. Proc. Wash. State Hortic. Assoc., 228-229.
- McFarland, J. D., Kevan, P. G., & Lane, M. A. (1989). Pollination biology of *Opuntia imbricata* Cactaceae in southern Colorado USA. Can. J. Bot., 67(1), 24-28.
- McGinley, R. J., & Rozen Jr., J. G. (1987). Nesting biology, immature stages, and phylogenetic placement of the Palaearctic bee *Parahophites* (Hymenoptera: Apoidea). Am. Mus. Novit., (2903), 1-21.
- McGinley, R. J. (1989). A catalog and review of immature Apoidea (Hymenoptera). Smithsonian Contributions to Zoology, (494), 24p.
- McMasters, J. H. (1989). The flight of the bumblebee and related myths of entomological engineering. Am. Sci., 77(2), 164-168.
- McMullen, C. K. (1989). The Galapagos Pacific Ocean carpenter bee: just how important is it? Not. Galapagos, (48), 16-18.
- Mello, M. L. S., Carcalho, H. F., & Garofalo, C. A. (1987). Polarization microscopy and topochemistry of the cocoon of *Lithurge huberi* (Hymenoptera, Megachilidae). Zool. Anz., 219(5-6), 369-376.
- Menzel, R., Steinmann, E., Souza J. de, & Backhaus, W. (1988). Spectral sensitivity of photoreceptors and colour vision in the solitary bee *Osmia rufa*. J. Exp. Biol., 136, 35-52.
- Menzel, R., Ventura, D. F., Werner, A., Joaquim, L. C. M., & Backhaus, W. (1989). Spectral sensitivity of single photoreceptors and color vision in the stingless bee, *Melipona quadrifasciata*. J. Comp. Physiol. A: Neural Behav. Physiol., 166(2), 151-164.
- Metwali, S. H., & Asmed, A. A. (1986)(1987). Mites accompanying ants and bumble bees in Egypt. Bull. Soc. Entomol. Egypte, (66), 267-277.
- Michener, C. D. (1989). Classification of American Colletinae (Hymenoptera, Apoidea). Univ. Kans. Sci. Bull., 53(11), 622-703.
- Michener, C. D. (1988). The genus *Lithurge* in the Antilles West Indies (Hymenoptera: Megachilidae). Folia. Entomol. Mex., (76), 159-164.
- Michener, C. D. (1988). The parasitic Anthophorid genus *Xeromelecta* in Cuba (Hymenoptera, Apoidea). Ann. Entomol. Soc. Am., 81(3), 377-379.
- Michener, C. D. (1987). Polymorphismus bei allodapinen Bienen. In: Schmidt, G.H. (Ed.) Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich, Wissenschaftliche Verlagsgesellschaft, Stuttgart.
- Michener, C. D., & Brooks, R. W. (1987). The family Melittidae in Madagascar (Hymenoptera, Apoidea). Ann. Soc. Entomol. France, 23(1), 99-103.

- Michener, C. D., & Grimaldi, D. A. (1988). The oldest fossil bee: Apoid history, evolutionary stasis, and antiquity of social behavior. Proc. Nat. Acad. Sci. USA, 85(17), 6424-6426.
- Michener, C. D., & Grimaldi, D. A. (1988). A *Trigona* from late cretaceous amber of New Jersey (Hymenoptera: Apidae: Meliponinae). Am. Mus. Novit., (2917), 10.
- Michener, C. D., & Smith, B. H. (1987). Kin recognition in primitively eusocial insects. In: Fletcher, D.J.C.; Michener, C.D. (Eds.) Kin recognition in animals. John Wiley & Sons, Chichester., 209-242.
- Milborrow, B. V., Kennedy, J. M., & Dollin, A. (1987). Composition of wax made by the Australian stingless bee *Trigona australis*. Austr. J. Biol. Sci., 40, 15-25.
- Miliczky, E. R. (1988). Observations on the bionomics of the bee *Andrena (Tylandrena) erythrogaster* Ashmead (Hymenoptera: Andrenidae) with notes on *A. (Micrandrena) personata* Robertson and *A. (Holandrena) c. cressonii* Roberston. Ill. Nat. Hist. Surv. Biol. Notes, (130), 1- 18.
- Minckley, R. L. (1989). Host records and biological notes for two *Anthrax* spp. in Arizona USA (Diptera, Bombyliidae). J. Kans. Entomol., 62(2), 274-278.
- Mirabzadeh, A., & Tigrari, S. (1987). The establishment of the first population of gregarious *Megachile rotundata* (Fabricius), (Hymenoptera, Megachilidae). In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.,
- Mohana Rao, G. (1987(1988). Scented Methi - a promising bee forage crop. Indian Bee J., 49(1-4), 14-15.
- Mohana Rao, G., & Suryanarayana, M. C. (1988). Studies on pollination of watermelon (*Citrullus lanatus* (Thunb.) Mansf.). Indian Bee J., 50(1), 5-8.
- Monsyavichyus, V. S. (1989). Trophic links between bees in Dainava Sandy Plain Lithuanian SSR USSR. Liet. TSR. Moskslu. Akad. Darb. Ser. C: Biol. Mokslai., (1), 61-74.
- Morris, B., Southcott, R. V., & Gale, A. E. (1988). Effects of stings of Australian native bees. Med. J. Aust., 149 (11-12), 707-709.
- Moure, J. S. (1987). Contribution to the knowledge of the genus *Ptiloglossa* (Hymenoptera: Colletidae). Acta Biol. Parana, 16, 1-4.
- Moure, J. S., & Camargo, J. M. F. (1988). A new species of *Xylocopa (Neoxylocopa)* from Brazil (Hymenoptera; Apoidea). Rev. Bras. Entomol., 32(2), 209-214.
- Moure, J. S., & Camargo, J. M. F. (1989). *Plebeia wittmanni* new-species a new Meliponinae (Hymenoptera, Apidae) from Southern Brazil. Stud. Neotrop. Fauna Environ., 24(1), 15-16.
- Mourikis, P. A., Argyriou, L. C., & Tsourgianni, A. (1988). Crumbling of the masonry of ancient settlements on Santorini caused by the Hymenoptera *Anthophora crinipes* and *Osmia latreillei*. Entomol. Hell., 6, 59-61.
- Nates Parra, G., Vergara, C., & Villa A. (In Press). Ciclo de desarrollo en *Trigona (Tetragonisca) angustula* (Hymenoptera: Apidae). Bol. Dep. Biol. U.N. Bogota, Colombia,
- Neff, J. L., & Simpson, B. B. (1988). Vibratile pollen-harvesting by *Megachile mendica* Cresson (Hymenoptera, Megachilidae). J. Kansas. Entomol. Soc., 61(2), 242-244.
- Nilsson, G. E. (1987). A gyandromorphic specimen of *Eulaeus albipes* Fabricius (Hymenoptera, Halictidae) and a discussion of possible causes of gyandromorphism in haplo-diploid insects. Not. Entomol., 67(3), 157-162.
- Nilsson, G. E. (1989). New records of Hymenoptera Aculeata from central and northern Sweden. Entomol. Tidskr., 110(1-2), 47-49.
- Nilsson, G. E. (1988). Swedish provincial records of Hymenoptera, Aculeata with *Evgates subnudus* new record for northern Europe and *Sphecodes albifrons* rediscovered from Scania. Entomol. Tidskr., 109(2), 97-100.
- Nilsson, L. A., & Rabakonandrianina, E. (1988). Chemical signalling and monopolization of nectar resources by territorial *Pachymelus limbatus* (Hymenoptera, Anthophoridae) male bees in Madagascar. J. Zool. (Lond.), 215((Pt.3)), 475-489.
- O'Connor, B. M. (1988). Coevolution in astigmatid mite-bee associations. In: Needham, G.R., Page, R.E., Jr., Delfino-Baker, M. & Bowman, C.E. (Eds). African Honey Bees and Bee Mites. Ellis Horwood LTD & John Riley & Son, Chichester, New York, Brisbane, etc.,
- O'Connor, B. M., & Eickwort, G. C. (1988). Morphology, ontogeny, biology and systematics of the genus *Vidua* (Acari, Winterschmiditiidae). Acarologia (Paris), 29(2), 147-174.
- O'Neill, K. M., & Bjostad, L. (1987). The male mating strategy of the bee *Nomia nevadensis* (Hymenoptera: Halictidae): egg structure and mate guarding. Pan-Pac. Entomol., 63(3), 207-217.
- Obrecht, E., & Scholl, A. (1987). Biochemical systematics in bumblebees: *Megabombus* S.S. (Hymenoptera, Apidae). In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.,
- Obrecht, E., & Scholl, A. (1987). Gene geographic studies in *Megabombus pascuorum* Scopoli (Hymenoptera, Apidae). In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.,
- Okazaki, K. (1987). Life cycle of a subtropical xylocopine bee, *Ceratina okinawana*, with some related problems. Kontyu, 55(1), 1-8.
- Olesen, J. M. (1988(1989)). Nest structure of a new species of *Euglossa* in a fruit of *Theobroma subincanum* from Ecuadorian Amazonas. Acta Amazonica, 18(3-4), 327-330.
- Olifer, V. N. (1989). Temperature and vitality in *Osmia rufa*. Vestn. Zool., (3), 52-55.

- Olsen, J. M. (1989). Behavior and nest structure of the Amazonian *Bombus transversalis* in Ecuador. *J. Trop. Ecol.*, 5(2), 243-246.
- Ordway, E. (1987). The Life History of *Diadasia rinconis* Cockerell (Hymenoptera: Anthophoridae). *J. Kans. Entomol. Soc.*, 60(1), 15-24.
- Ordway, E., Buchmann, S. L., Kuehl, R. O., & Shipman, C. W. (1987). Pollen dispersal in *Curbita foetidissima* (Cucurbitaceae) by bees of the genera *Apis*, *Pepponapis* and *Xenoglossa* (Hymenoptera: Apidae, Anthophoridae). *J. Kans. Entomol. Soc.*, 60(4), 489-503.
- Ornosa Gallego, C. (1987). New data on *Bombus lapidarius decipiens* Perez 1879 (Hymenoptera, Apidae). *Bol. Asoc. Esp. Entomol.*, 11(0), 415.
- Osborn, M. M., Kevan, P. G., & Lane, M. A. (1988). Pollination biology of *Opuntia polyacantha* and *Opuntia phaeacantha* Cactaceae in southern Colorado USA. *Plant Syst. Evol.*, 159(1-2), 85-94.
- Osgood, E. A. (1989). Biology of *Andrena crataegi* Robertson (Hymenoptera: Andrenidae), a communally nesting bee. *J. N. Y. Entomol. Soc.*, 97(1), 56-64.
- Ospina, T. R., Lievano, A., & Nates, G. (1987)(1988). Color pattern of the social bumblebees *Bombus atratus* Franklin in Cundinamarca Colombia a differential population. *Rev. Biol. Trop.*, 35(2), 317-324.
- Osten, T. (1989). Comparative studies on the mating behavior of *Platynopoda* and *Mesotrichia* Hymenoptera Xylocopini a study of form and function. *Stuttg. Beitr. Naturkd. Ser. A: (Biol.)*, (433), 1-18.
- Ottosen, C.-O. (1987). Male bumblebees (*Bombus hortorum* L.) as pollinators of *Lonicera periclymenum* L. in N.E.-Zealand, Denmark. *Flora (Jena)*, 179(2), 155-161.
- Owen, R. E. (1988). Body size variation and optimal body size of bumble bee queens (Hymenoptera: Apidae). *Can. Entomol.*, 120(1), 19-27.
- Owen, R. E. (1989). Differential size variation of male and female bumblebees. *J. Hered.*, 80(1), 39-43.
- Owen, R. E., & Plowright, R. C. (1988). Inheritance of metasomal pile color variation in the bumble bee *Bombus rufocinctus* Cresson (Hymenoptera, Apidae). *Can. J. Zool.*, 66(5), 1172-1178.
- Pacheco, R. D. L., & Kerr, W. E. (1989). Temperature in bees of *Melipona compressipes fasciculata*. *Cienc. Cult. (Sao Paulo)*, 41(5), 490-495.
- Packer, L. D. M. (1987). Aspects of the social biology of *Halictus ligatus* Say (Hymenoptera: Halictidae). *Diss. Abstr. Int. B: Sci. Eng.*, 47(10), 4049.
- Packer, L. (1987). Competition over oviposition in subtropical and temperate populations of the social sweat bee *Halictus ligatus*. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Packer, L. (1987). A description of the mature larva and cooon of the bee *Thygater* (Hymenoptera: Anthophoridae). *J. N. Y. Entomol. Soc.*, 95(1), 23-27.
- Packer, L. (1987). The triungulin larva of *Nemognatha (Pauronemognatha) punctulata* Leconte (Coleoptera: Meloidae) with a description of the nest of its host - *Megachile brevis pseudobrevis* Say (Hymenoptera: Megachilidae). *J. Kans. Entomol. Soc.*, 60(2), 280-287.
- Packer, L. (1987). An unexpectedly high female bias to the sex ratio in a temperate population of *Halictus ligatus* - a primitively social sweat bee. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Packer, L. (1988). The effect of *Bombylius pulchellus* (Diptera, Bombyliidae) and other mortality factors upon the biology of *Halictus ligatus* (Hymenoptera; Halictidae) in southern Ontario. *Can. J. Zool.*, 66(3), 611-616.
- Packer, L. (1990). Solitary and eusocial nests in a population of *Augochlorella striata* (Provancher) (Hymenoptera; Halictidae) at the northern edge of its range. *Behav. Ecol. Sociobiol.*, 27, 339-344.
- Packer, L., Jessome, V., Lockerbie, C., & Sampson, B. (1989). The phenology and social biology of four sweat bees in a marginal environment: Cape Breton Island. *Can. J. Zool.*, 67(12), 2871-2877.
- Packer, L., & Knerer, G. (1987). The biology of a subtropical population of *Halictus ligatus* Say (Hymenoptera; Halictidae). III. The transition between annual and continuously brooded colony cycles. *J. Kans. Entomol. Soc.*, 60(4), 510-516.
- Packer, L., & Owen, R. E. (1989). Allozyme variation in *Halictus rubicundus* (Christ): a primitively social halictine bee (Hymenoptera: Halictidae). *Can. Entomol.*, 121(12), 1049-1057.
- Packer, L., & Owen, R. E. (1989). Notes on the biology of *Lasiosiglossum (Evylaeus) cooleyi* (Crawford), an eusocial halictine bee (Hymenoptera: Halictidae). *Can. Entomol.*, 121(6), 431-438.
- Packer, L., Sampson, B., Lockerbie, C., & Jessome, V. (1989). Nest architecture and brood mortality in four species of sweat bee (Hymenoptera, Halictidae) from Cape Breton Island Nova Scotia, Canada. *Can. J. Zool.*, 67(12), 2864-2870.
- Pagliano, G. (1987)(1988). Checklist of Italian Hymenoptera. I. Halictidae. *Boll. Mus. Civ. St. Nat. Venezia*, 38, 85-128.
- Pagliano, G. (1988). Prospetto sistematico degli apoidea Italiani. *Ann. Fac. Agr. Univ. Torino*, 15, 97-128.
- Parnilo, P., Pekkarinen, A., & Vario, S. L. (1987). Clustering of bumble bee subgenera based on interspecific genetic relationships (Hymenoptera, Apidae: *Bombus* and *Psithyrus*). *Ann. Zool. Fenn.*, 24(1), 19-27.
- Parker, F. D. (1988). Influence of wood, paper and plastic nesting units on efficacy of three candidate fungicides for control of chalkbrood in the alfalfa leaf-cutting bee (Hymenoptera, Megachilidae). *J. Econ. Entomol.*, 81(3), 789-795.
- Parker, F. D. (1989). Nest clustering as a means of managing *Osmia sanrafaelae* (Hymenoptera: Megachilidae). *J. Econ. Entomol.*, 82(2), 401-403.

- Parker, F. D. (1988). Nesting biology of two North American species of *Chelostoma* (Hymenoptera: Megachilidae). *Pan-Pac. Entomol.*, 64(1), 1-7.
- Parker, F. D. (1987). Nests of *Callanthidium* from block traps (Hymenoptera: Megachilidae). *Pan-Pac. Entomol.*, 63(2), 125-129.
- Parker, G. D., Cane, J. H., Frankie, G. W., & Vinson, S. B. (1987). Host records and nest entry by *Dolichostelis*, a kleptoparasitic anthidine bee (Hymenoptera: Megachilidae). *Pan-Pac. Entomol.*, 63(2), 172-177.
- Parra, G. N., & Roubik, D. W. (1990). Sympatry among subspecies of *Melipona favosa* in Colombia and a taxonomic revision. *J. Kans. Entomol. Soc.*, 63(1), 200-203.
- Pasotti, L. (1988). Technique for raising queen bees. *Sante Abeille*, (104), 52-58.
- Pasteels, J. J. (1987). Megachilidae (Hymenoptera, Apoidea) collected by Dr. K. V. Krombein and his team in Sri Lanka. *Bull. Ann. Soc. R. Belge. Entomol.*, 123((7-9), 225-236.
- Patt, J. M., Merchant, M. W., Williams, D. R. E., & Meeuse, B. J. D. (1989). Pollination biology of *Plantathera stricta* Orchidaceae in Olympic National Park Washington USA. *Am. J. Bot.*, 76(8), 1097-1106.
- Pauly, A. (1989). Afro-tropical *Pachyhalictus* species of the subgenus *Dictyohalictus* Michener (Hymenoptera: Apoidea: Halictidae). *Rev. Zool. Afr.*, 103(1), 41-49.
- Pawlowski, T. (1987). Influence of farming systems on wild bees (Hymenoptera, Apoidea) in agricultural landscape. *Acta Univ. Nicolai Copernici. Biol.*, 35, 153-167.
- Payne, J. A., Amis, A. A., Cane, J. H., & Iyrene, P. M. (1989). Fruit size, seed size, seed variability and pollination of rabbiteye blueberries (*Vaccinium ashei* Reade). *Acta Hortic.*, (241), 38-43.
- Payne, R. M. (1987). Avon and district entomological report, 1985. Sawflies, bees, wasps, ants (Hymenoptera). *Proc. Bristol Nat. Soc.*, (45), 51-52.
- Peakall, R. (1989). A new technique for monitoring pollen flow in orchids. *Oecologia*, 79(3), 361-365.
- Pekkarinen, A., Teras, I., & Wuorenrinne, H. (1987). Endangered bees, wasps and ants in Finland. *Luonnon Tutkija*, 91, 124-129.
- Pekkarinen, A., & Teras, I. (1987). Observations on *Bombus humilis* (Hymenoptera: Apidae) in Finland. *Not. Entomol.*, 67(4), 208-209.
- Pellmyr, O. (1988). Bumble bees (Hymenoptera: Apidae) assess pollen availability in *Anemonopsis macrophylla* (Ranunculaceae) through floral shape. *Ann. Entomol. Soc. Am.*, 81(5), 792-797.
- Pellmyr, O., & Karkkainen, K. (1987). *Salix phylicifolia* as exclusive food source for all nectarivorous insects in a Taiga community, Oulanka, Finland. *Sven. Bot. Tidskr.*, 81(1), 43-46.
- Pesenko, Y. A. (1987). *Halictus costulatus* Krichbaumer 1873 currently *Lasioglossum costulatum* (Insecta, Hymenoptera) proposed conservation of specific name. *Bull. Zool. Nomencl.*, 44(1), 17-18.
- Pesenko, Y. A. (1988). Researches in systematics of hymenopterous insects. *Proc. Zool. Inst.*, Leningrad, 175, 75-101.
- Pesenko, Y. A., & Belokobylskij, S. A. (1989). Materials on systematics of hymenopterous insects. *Proc. Zool. Inst.*, Leningrad, 188, 114-129.
- Pesenko, Y. A., & Warncke, K. (1987). Beitrag zur Biogenenfauna des Iran 22. Die Tribe Nomioioidini (Hymenoptera: Halictidae). *Boll. Mus. Civ. St. Nat. Venezia*, 36, 109-115.
- Peters, C. M., & Vasquez, A. (1986-1987). Ecological studies of *Myciaria dubia*. I. Fruit production in natural populations. *Acta Amazonica*, 16-17(1), 161-174.
- Petkov, G., & Nenchev, P. (1988). A method for determination of gas exchange and heat production in bees. *Zhivotnov'd. Nauki.*, 25(3), 82-85.
- Pettersson, B. (1989). Pollination in the African species of *Nervilia* Orchidaceae. *Lindleyana*, 4(1), 33-41.
- Pham-deleuge, M. H., Etievant, P., & Masson, C. (1987). Molecular parameters involved in bee-plant relationships: a biological and chemical approach. *Biochimie*, 69, 661-670.
- Pinheiro, M. C. B., Ormond, W. T., Leite, C. D. O., & Lima, H. A. D. (1988). Pollination ecology of *Chamaecrista ramosa* Var. *ramosa*. *Rev. Bras. Biol.*, 48 (4), 665-672.
- Pinzauti, M. (1987). The pollination of strawberries in Verisia. *Notiziario di Ortoflorofrutticoltura*, 13(2), 78-84.
- Pinzauti, M., & Frediani, D. (1987). Factors determining nectars secretion from *Helianthus annuus* L. together with observation about the behavior of prounibal insects. *Apiacta*, 22(2), 45-46.
- Plant, J. D. (1987). Did eusocial behavior arise only in Apidae? In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology in Social Insects*. Verlag J. Peperny, Munich.,
- Plant, J. D., & Paulus, H. F. (1987). Comparative morphology of the postmentum of bees (Hymenoptera: Apoidea) with special remarks on the evolution of the lorum. *Z. zool. Syst. Evolut.-forsch.*, 25(2), 81-103.
- Plateaux-Quenu, C. (1989). Premières observations sur le caractère social d'*Evyiaeus albipes* F. (Hymenoptera, Halictinae). *Actes Colloq. Insectes Soc.*, 5, 335-344.
- Plateaux-Quenu, C. (1988). Réalisation expérimentale de grands individus de première couvée chez *Evyiaeus calceatus* (Scop.) (Hymenoptera, Halictinae): Biométrie et caste. *Ann. Sci. Nat. Zool. Biol. Anim.*, 9 (4), 263-270.
- Plateaux-Quenu, C., Plateaux, L., & Packer, L. (1989). Biological notes on *Evyiaeus villosulus* K. (Hymenoptera, Halictidae) a bivoltine largely solitary halictine bee. *Insectes Soc.*, 36(4), 245-263.
- Plateaux-Quenu, C., & Plateaux, L. (1986-1987). Occurrence of giant foundresses within two populations of *Evyiaeus villosulus* K. (Hymenoptera, Halictinae). *Am. Sci. Nat. Zool. Biol. Anim.*, 8(4), 259-266.

- Plateaux-Quenu, C., Plateaux, L., & Packer, L. (1987). Predaptation to sociality in the solitary, bivoltine, halictine bee *Eylaeus villosulus* (K.). In: Eder, J. & Remold, H. (Eds). Chemistry and Biology of Social Insects. Verlag J. Peperny, Munich.
- Plowright, R. C., & Fuller, G. A. (1988). The role of intruder behavior in nestmate recognition by bumble bees (Hymenoptera: Apidae). Can. J. Zool., 66(12), 2847-2851.
- Plowright, R. C., & Plowright, C. M. S. (1988). Elitism in social insects: a positive feedback model. In: Interindividual Behavioral Variability in Social Insects. Robert L. Jeanne (ed.). Boulder, Westview Press.
- Pouvreau, A. (1987(1988). Biological cycle of *Aphomia sociella* L. (Lepidoptera, Pyralidae, Gallerinae), a pest of bumble bee nests (Hymenoptera, Apoidea, Bombinae). Colloques De L'Inra. Parasitoid Insects. France., (48), 161-162.
- Pouvreau, A. (1988). Contribution to the study of the preimaginal development of *Aphomia sociella* L. (Lepidoptera, Pyralidae, Gallerinae), a pest of bumblebee nests. Entomophaga, 33(4), 467-480.
- Pouvreau, A. (1989). Contribution to the study of polyethism in bumblebees *Bombus* Latr. (Hymenoptera, Apidae). Apidologie, 20(3), 229-244.
- Pouvreau, A. (1988). Preliminary obervations on the method used by *Psithrus Rupestris* Fab. (Hymenoptera, Apoidea, Bombinae) females for the recognition of nests of specific bumble bees (*Bombus* sp.). Actes Colloq. Insectes Soc., 4, 301-305.
- Pouvreau, A., Coutin, R., & Descoins, C. (1988). Pollination of *Ophrys orchidaceae* and pseudocopulation of Hymenoptera males. Boll. Zool. Agrar. Bachic., 20(0), 37-60.
- Powell, A. H., & Powell, G. V. N. (1987). Population dynamics of male euglossine bees in Amazonian forest fragments. Biotropica, 19(2), 176-179.
- Praagh, J. P. van. (1988). Pollination of blackberry cv. thornless evergreen *Rubus laciniatus* in relation to fruit growth. Gartenbauwissenschaft, 53(1), 15-17.
- Prabucki, J., Chuda-Mickiewicz, B., & Woyke, H. W. (1987). Effect of insect pollinators on cauliflower seed yield. Biuletyn Warzywniczy, 30, 71-91.
- Priore, R. (1987). The collection of (Apoidea; Hymenoptera) of the Institute of Agarian Entomology of Portici Italy IX. *Chelostoma* Latr. *Creightonella* Cock. *Dioxys* Lepel. and Serv. *Heriades* Spin. *Osmia* Pz. *Protosmia* Duck. *Megachile* Latr. Megachilidae. Boll. Lab. Entomol. Agrar. Filippo. Silvestri., 44(0), 7-24.
- Prys-Jones, O. E., & Corbet, S. A. (1987). Bumblebees. Naturalists' Handb., (6), 1-86.
- Ptacek, V. (1987). Technical isolation in research and breeding of legumes. Sb. Ved. Pr. Vyzk. Slechtitskoho Ustavu Picninarskoho Troubsku Brna., 10(0), 117-128.
- Radchenko, V. (1989). The biology of the bumblebee family. Prepr. Inst. Zool. Acad. Sci. Ukr. SSR-Kiev, 55pp.
- Radchenko, V. G. (1987). Nesting of *Dasypoda braccata* Ev. etrsm. (Hymenoptera, Melittidae) in the southwestern Ukraine USSR. Entomol. Rev., 116(2), 299-301.
- Raeder, K., Wildfeuer, A., Wintersberger, F., Bossinger, P., & Muecke, H. W. (1987). Characterization of bee venom and its main components by high-performance liquid chromatography. J. Chromatogr., 408(0), 341-348.
- Rahman, A. (1988(1989). Taxonomic studies on megachilid bees of Haryana. Thesis Abstr. Univ., 14(4), 353-354.
- Ramalho, M., Kleinert-Giovannini, A., & Imperatriz-Fonseca, V. L. (1989). Utilization of floral resources by species of *Melipona* (Apidae, Meliponinae): floral preferences. Apidologie, 20(3), 185-195.
- Rank, G. H., & Rank, F. P. (1989). Diapause intensity in a french univoltine and a Saskatchewan commercial strain of *Megachile rotundata* (Fab.). Can. Entomol., 121(2), 141-148.
- Rasmont, P. (1988). Ecological and zoogeographical monograph on the bumble bees of France and Belgium. Dissertation, Docteur en Sciences, Faculte des Sciences Agronomiques de L'Etat, Gembloux, Belgium.,
- Rasmont, P., Delmas, R., Lecant, F., & Yarrow, I. H. H. (1987). *Megabombus (Rhodobombus) mesomelas*. (Gerstaeker, 1869) (Hymenoptera, Apidae, Bombini). Documents pour un Atlas Zoogeographique du Languedoc-Roussillon, (32), 4.
- Rasmont, P., Delmas, R., Lecant, F., & Yarrow, I. H. H. (1987). *Megabombus (Rhodobombus) pomorum* (distribution in France). Documents pour un Atlas Zoogeographique du Languedoc-Roussillon, (4), 31.
- Rasmont, P., & Mersch, P. (1988). First estimation of faunistic drift in the bumblebees of Belgium (Hymenoptera, Apidae). Ann. Soc. R. Zool. Belg., 118 (2), 141-148.
- Rasulov, M. (1989). Apidae from the Talass Valley Uzbek SSR USSR Uzb. Biol. Zh., (4), 42-43.
- Rasulov, M. (1989). Bees in the lower course of the river Syr Darya Kazakh USSR Uzb. Biol. Zh., (1), 56-58.
- Rathcke, B. (1988). Flowering phenologies in a shrub community: competition and constraints. J. Ecol., 76 (4), 975-994.
- Rathcke, B. (1988). Interactions for pollination among co-flowering shrubs. Ecol. Publ. Ecol. Soc. Am., 69(2), 446-457.
- Raw, A. (1989). The dispersal of euglossine bees between isolated patches of eastern Brazilian wet forest (Hymenoptera, Apidae). Rev. Bras. Entomol., 33(1), 103-107.
- Raw, A. (1988). Nesting biology of the leaf-cutter bee *Megachile centuncularis* (L.) (Hymenoptera: Megachilidae) in Britain. Entomologist, 107(1), 52-56.
- Read, P. E. C., Donavan, B. J., & Griffin, R. P. (1989). Use of bumble bees *Bombus terrestris* as pollinators of kiwifruit and lucerne in New Zealand. N. Z. Entomol., 12(0), 19-23.

- Reid, W. H., Sensiba, P., & Freeman, C. E. (1988). A mixed pollination system in *Penstemon pseudospectabilis* M.E. Jones (Scrophulariaceae). Great Basin Nat., 48(4), 489-494.
- Reimann, H. (1987). Bees, wasps and ants as colonizers of the scarps of low-lying drains. Abhandlungen Naturwissenschaftlichen Verein Bremen, 40(4), 333-346.
- Reinig, W. F., & Rasmont, P. (1988). Contribution to the knowledge of the bumble bee *Alpigenobombus wulfsteini* (Radoszkowski, 1959) (Hymenoptera, Apidae, Bombinae). Spixiana (Muench), 11(1), 37-67.
- Renner, S. S. (1986-1987). Reproductive biology of *Bellucia* Melastomataceae. Acta Amazonica, 16-17(1), 197-208.
- Renner, S. S. (1989). A survey of reproductive biology in neotropical Melasomataceae and Memecylaceae. Ann. Mo. Bot. Gard., 76(2), 496-518.
- Reyes, S. G., & S.F. Sakagami. (1990). A new socially parasitic *Braunsapis* from India with notes on the synonymy of *Braunsapis mixta* (Hymenoptera: Xylocopinae: Allodapini). J. Kans. Entomol. Soc., 63 (3), 458-461.
- Ribeiro, M. d. F. (1988). Knowledge of indigenous (to Brazil) stingless bees. 1. The black stingless bee (*Friesornelitta silvestrii*). Apicultura No. Brasil, (27), 34-35.
- Ricciardelli D'Albore, G. C. (1988). Daily foraging activity of *Bombus* Latr. and *Psithrus* Lep. (Hymenoptera, Apidae) in different altitudinal bands of Umbria and the Marche Italy. Redia, 71(1), 99-114.
- Riemann, H. (1988). Contribution to the fauna of Aculeata Hymenoptera in some sand pits in lower Saxony West Germany. Braunscheiger Naturkd. Schr., 3 (1), 213-242.
- Rio, C. M. del, & Eguiarte, L. E. (1987). The effect of nectar availability on the foraging behavior of the stingless bee *Trigona testacea*. Southwest. Nat., 32(3), 313-319.
- Roberts, R. B., & Brooks, R. W. (1987). Agapostemonine bees of Mesoamerica. (Hymenoptera: Halictidae). Univ. Kans. Sci. Bull., 53(7), 357-392.
- Roig-Alsina, A. (1987). The classification of the Cae-noprosopidini (Hymenoptera, Anthophoridae). J. Kans. Entomol. Soc., 60(2), 305-315.
- Roig-Alsina, A. (1989). A revision of the bee genus *Doe-ringiella* (Hymenoptera: Anthophoridae: Nomadinae). Univ. Kans. Sci. Bull., 53(10), 577-621.
- Roig-Alsina, A. (1989). The tribe Osirini, its scope, classification, and revisions of the genera *Parepolus* and *Osirinus* (Hymenoptera, Apoidea, Anthophoridae). Univ. Kans. Sci. Bull., 54(1), 1-23.
- Roman'Kova, T. G. (1989). New data on *Ctenoplectra davidi* (Hymenoptera, Apoidea, Ctenoplectridae) in the Primorye area. Vestn. Zool., (4), 60-63.
- Roman'Kova, T. G. (1988). A new genus of far eastern Russian SFSR USSR bees of the tribe Anthidiini (Hymenoptera, Apoidea, Megachilidae). Vestn. Zool., (4), 25-30.
- Romasenko, L. P. (1989). New data on the biology of *Megachile analis* (Hymenoptera: Megachilidae). Vestn. Zool., (6), 71-73.
- Romasenko, L. P. (1989). Prepupae of some megachilid bees (Hymenoptera). Vestn. Zool., (1), 75-78.
- Roseler, P. F., & Roseler, I. (1988). Influence of juvenile hormone on fat body metabolism in ovarioectomized queens of the bumblebee, *Bombus terrestris*. Insect Biochem., 18(6), 557-563.
- Roubik, D. W. (1987). Notes on the biology of Anthophorid bee *Tetrapedia* and the mite *Chaetodactylus panamensis* Baker, Roubik and Delfino-Baker (Acar: Chaetodactylidae). Int. J. Acarol., 13(1), 75-76.
- Roubik, D. W., & Ackerman, J. D. (1987). Long term ecology of euglossine orchid-bees (Apidae: Euglossini) in Panama. Oecologia (Berl.), 73(3), 321-333.
- Roubik, D. W. (1989). Ecology and natural history of tropical bees. Cambridge Tropical Biology Series, Cambridge University Press, United Kingdom.,
- Roubik, D. W. (1988). An overview of Africanized honey-bee populations: reproduction, diet, and competition. In: Needham et al (Eds.) Africanised Honey Bees and Bee Mites, Ellis Horwwod Ltd., Chichester, England.,
- Rourke, J. P. (1988). A new species of *Mimetes* Proteaceae from the southern cape South Africa. S. Afr. J. Bot., 54(6), 636-639.
- Rozen Jr., J. G. (1988). Ecology, behavior, and mature larva of a new species of the old world bee genus *Camptopoeum* (Andrenidae: Panurginae). Am. Mus. Novit., (2925), 10 pp.
- Rozen Jr., J. G. (1989). Life history studies of the 'primitive' panurgine bees (Hymenoptera: Andrenidae: Panurginae). Am. Mus. Novit., (2962), 27pp.
- Rozen Jr., J. G. (1989). Morphology and systematic significance of first instars of the cleptoparasitic bee tribe Epeolini (Anthophoridae: Nomadinae). Am. Mus. Novit., (2957), 1-19.
- Rozen Jr., J. G. (1987). Nesting biology and immature stages of a new species in the bee genus *Hesperapis* (Hymenoptera: Apoidea: Melittidae: Dasypodinae). Am. Mus. Novit., (2887), 13pp.
- Rozen Jr., J. G. (1987). Nesting biology of the bee *Asmeadiella holti* and its cleptoparasite, a new species of *Stelis* (Apoidea: Megachilidae). Am. Mus. Novit., (2900), 1-10.
- Rozen Jr., J. G. (1989). Two new species and redescription of another species of the cleptoparasite bee genus *Tripeolus* with notes on their immature stages (Anthophoridae, Nomadinae). Am. Mus. Novit., (2956), 1-18.
- Rozen Jr., J. G., & Ayala, R. (1987). Nesting biology of the squash bee *Peponapis utahensis* (Hymenoptera: Anthophoridae; Eucerini). J. N. Y. Entomol. Soc., 95 (1), 28-33.
- Rozen Jr., J. G., & Michener, C. D. (1988). Nests and immature stages of the bee *Paratetrapedia swainsonae* (Hymenoptera: Anthophoridae). Am. Mus. Novit., (2909), 1-13.

- Ruijter, A. D., Eijunde, J. V. Den, & Steen, J. V. Der. (1988). Bees improve setting of tomatoes. *Groenten en Fruit*, 43(34), 46-47.
- Ruijter, A. D., & Smeekens, C. C. (1987). Pollination by bees attractive for the culture of courgettes! *Groenten En Fruit*, 42(47), 34-35.
- Rust, R. W. (1988). Biology of *Nomadopsis larreae* (Hymenoptera: Andrenidae), with an analysis of yearly appearance. *Ann. Entomol. Soc. Am.*, 81(1), 99-104.
- Rust, R. W. (1987). Collecting of *Pinus* (Pinaceae) pollen by *Osmia* bees (Hymenoptera; Megachilidae). *Environ. Entomol.*, 16, 668-671.
- Sakagami, S. F. (1988). Bionomics of the halictine bees in northern Japan. IV. *Lasioglossum (Evylaeus) nupricola* sp. nov., a climatic relic. *Kontyu*, 56(2), 337-353.
- Sakagami, S. F. (1987). Sozialstruktur und Polyethismus bei Prachtbienen. In: Schmidt, G.H. (Ed.) *Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich*. Wissenschaftliche Verlagsgesellschaft, Stuttgart.
- Sakagami, S. F. (1989). Taxonomic notes on a Malesian bee *Lasioglossum carinatum*, the type species of the subgenus *Ctenonomia*, and its allies (Hymenoptera: Halictidae). *J. Kans. Entomol. Soc.*, 62(4), 496-510.
- Sakagami, S. F., & Ebmer, P. A. W. (1987). Taxonomic notes on oriental halictine bees of the genus *Halictus* (subgen. *Seladonia*) (Hymenoptera; Apoidea). *Linzer. Biol. Beitr.*, 19(2), 301-357.
- Sakagami, S. F., Inoue, T., Yamane, S., & Salmah, S. (1989). Nests of the myrmecophilous stingless bee, *Trigona moorei*: how do bees initiate their nest within an arboreal ant nest? *Biotropica*, 21(3), 265-274.
- Sakagami, S. F., & Inoue, T. (1987). Stingless bees of the genus *Trigona* subgenus *Trigonella* with notes on the reduction of spatha in male genitalia of the subgenus *Tetragonula* (Hymenoptera, Apidae). *Kontyu*, 55(4), 610-627.
- Sakagami, S. F., & Khoo, S. G. (1987). Taxonomic status of the Malesian stingless bee *Trigona reepenia* with discovery of *Trigona pagdeni* from northern Malaya. *Kontyu*, 55(2), 207-214.
- Sakagami, S. F., & Lacroca, S. (1988). Nests of an exomalopsine bee *Lanthanomelissa goeldiana* (Hymenoptera: Anthophoridae). *J. Kans. Entomol. Soc.*, 61(3), 347-349.
- Sakagami, S. F., & Maeta, Y. (1987). Multifemale nests and rudimentary castes of an 'almost' solitary bee *Ceratina flavipes*, with additional observations on multifemale nests of *Ceratina japonica* (Hymenoptera, Apoidea). *Kontyu*, 55(3), 391-409.
- Sakagami, S. F., & Maeta, Y. (1987). Sociality, induced and/or natural, in the basically solitary small carpenter bees (*Ceratina*). In: Ito, Y. Brown, J.L. & Kikawa, J. (Eds.). *Animal Societies, Theories and Facts*. Japan Scientific Press, Tokyo.,
- Sakagami, S. F., & Michener, C. D. (1987). Tribes of Xylocopinae and the origin of the Apidae (Hymenoptera: Apoidea). *Ann. Entomol. Soc. Am.*, 80(3), 439-450.
- Sakagami, S. F., & Yamane, S. (1987). Oviposition behavior and related notes of the Taiwanese stingless bee *Trigona (Lepidotrigona) ventris* Hoozana. *J. Ethol.*, 5(1), 17-27.
- Sakagami, S. F. (1989). Compatibility and incompatibility of solitary life with eusociality in two normally solitary bees *Ceratina japonica* and *Ceratina okinawa* (Hymenoptera, Apoidea), with notes on the incipient phase of eusociality. *Japan. J. Ecol.*, 57(2), 417-439.
- Sakagami, S. F., & Fukuda, H. (1989). Nest founding and nest survival in a eusocial halictine bee, *Lasioglossum duplex*: additional observations. *Res. Pop. Ecol.*, 31(1139-151).
- Saleh-mghir, E. (1989). Inter-individual recognition in bees *Apis mellifera*. *Annee. Biol.*, 28(3), 187-200.
- Salmah, S., Inoue, T., Mardius, P., & Shoichi, S. F. (1987). Incubation period and post-emergence pigmentation in the Sumatran stingless bee, *Trigona (Trigonella) moorei*. *Kontyu*, 55(3), 383-390.
- Sanchez, D. (1989). Ground-nesting bee best for blueberries. *Agric. Res. (Wash., D.C.)*, 37(6), 16-17.
- Santiago-Blay, J. A., & Daly, H. V. (1988). The Apoidea collection of the California insect survey (C.I.S.) at the EssigMuseum of Entomology, University of California, Berkeley. *Pan. Pac. Entomol.*, 64(3), 281-284.
- Saraiva, L. C., Cesar, O., & Monteiro, R. (1988). Pollination biology and breeding system of *Stryax camporum* Pohl and *Stryax ferrugineus neeset* Mart. Styracaceae. *Rev. Bras. Bot.*, 11((1-2), 71-80.
- Sarat Babu, G. V., Bhatt, S. C., & Pandeya, S. C. (1987). Pollination ecology of *Solanum sisymbriifolium* Lamk. *Proc. Indian Acad. Sci. Plant Sci.*, 97(4), 295-300.
- Sarycheva, Z. N., & Tarasenko, N. D. (1989). Phytospecialization of *Anthidium florentinum* F. (Apoidea, Hymenoptera, Insecta) in Novosibirsk Oblast Russian SFSR USSR. *Zh. Obshch. Biol.*, 50(5), 655-663.
- Sazima, I., & Sazima, M. (1989). *Manangavas e Irapuas* (Hymenoptera, Apoidea): vistos, interacoes e consequencias para polinizacao do Maracuja (Passifloraceae). *Rev. Bras. Entomol.*, 33(1), 109-118.
- Schmid-Hempel, R., & Schmid-Hempel, P. (1989). Superparasitism and larval competition in conopid flies (Diptera, Conopidae) parasitizing bumblebees (Hymenoptera, Apidae). *Mitt. Schweiz. Entomol.*, 62(3-4), 279-289.
- Schmidt, G. H. (1987). Mechanismen der Kastenbildung und Steuerung des Geschlechtsverhaltnisses. In: Schmidt, G.H. (Ed.) *Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich*. Wissenschaftliche Verlagsgesellschaft, Stuttgart.,

- Schmidt, G. H. (1987). Polymorphismus, Arbeitung, Kastenbildung. In: Schmidt, G.H. (Ed.) Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich. Wissenschaftliche Verlagsgesellschaft, Stuttgart.
- Schmidt, G. H. (1987). Soziogenese und Evolution des Sozialmorphismus. In: Schmidt, G.H. (Ed.) Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich. Wissenschaftliche Verlagsgesellschaft, Stuttgart,
- Schmitt, U., & Bertsch, A. (1990). Do foraging bumblebees scent-mark food sources and does it matter? *Oecologia*, 82(1), 137-144.
- Schneider, E. L., Fitch, G. K., Barrientos, R., & Koke, J. R. (1987). Contributions to the floral ecology of Texas and Chihuahuan *Argemone* (Papaveraceae). *Southwest. Natural.*, 32(1), 150-152.
- Schneider, P., & Schubert, M. (1987). Behaviour of insects on flowers. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie*, 5(1-4), 27-30.
- Schonitzer, K. (1988). Curious builders. One of the 700 European bee species- The mortar bee (*Chalicodoma*). *Reise-Kurier*, 11(125), 8.
- Schousboe, C. (1987). Deutonymphs of *Parasitellus* phoretic on Danish bumblebees (Parasitidae, Mesostigmata; Apidae, Hymenoptera). *Acarologie*, 28(1), 37-41.
- Schwarz, M. P. (1987). Intra-colony relatedness and sociality in the allodapine bee *Exoneura bicolor*. *Behav. Ecol. Sociobiol.*, 21(6), 387-392.
- Schwarz, M. P. (1988). Intra-specific mutualism and kin-association of cofoundresses in allodapine bees (Hymenoptera, Anthophoridae). *Monit. Zool. Ital.*, 22(3), 245-254.
- Schwarz, M. P. (1988). Local resource enhancement and sex ratios in a primitively social bee. *Nature*, 331 (6154), 346-348.
- Schwarz, M. P. (1988). Notes on cofounded nests in three species of social bees in the genus *Exoneura* (Hymenoptera: Anthophoridae). *Victorian Nat.*, 105 (1), 212-215.
- Schwarz, M. P., Scholz, O., & Jensen, G. (1987). Ovarian inhibition among nestmates of *Exoneura bicolor* Smith (Hymenoptera: Xylocopinae). *J. Aust. Entomol. Soc.*, 26(pt.4), 355-359.
- Schwenniger, H. R. (1987). Contribution to the bee fauna (Hymenoptera, Apoidea) in moist fallow lands of southern Pfälzerwald Palatinae West Germany. *Pollicchia O.*, (12), 241-260.
- Scott-Dupree, C. D., & Winston, M. L. (1987). Wild bee pollinator diversity and abundance in orchid and uncultivated habitats in the Okanagan Valley, British Columbia. *Can. Entomol.*, 119(7-8), 735-745.
- Shanks, S. S. (1987). Two new species of *Osiris* with a key to the species from Mexico (Hymenoptera, Anthophoridae). *Wasmann. J. Biol.*, 45(1-2), 1-5.
- Sherman, H. (1989). Polyester repair man is really a bee. *Agric. Res. (Washington D.C.)*, 37(4), 18.
- Shmida, A., & Ducas, R. (1988). A bee to suit every flower? *Israel - Land. Nat.*, 13(3), 106-109.
- Sihag, R. C. (1988). Characterization of the pollinators of cultivated cruciferous and leguminous crops of sub-tropical Hissar, India. *Bee World*, 69(4), 153-158.
- Simpson, B. B., & Neff, J. L. (1987). Pollination ecology in the arid southwest USA. *Aliso.*, 11(4), 417-440.
- Singh, A., & Tiwari, V. K. (1989). Natural enemies and nest structure of *Halictus lucidipennis* Smith (Hymenoptera, Halictidae) from Bharatpur Rajasthan India. *J. Adv. Zool.*, 10(1), 70-72.
- Singh, A., & Tiwari, V. K. (1988). A new species of genus *Nomia* Latreille from India (Hymenoptera: Apoidea: Halictidae). *Entomon.*, 13(3-4), 327-330.
- Sitdikov, A. A. (1987). Nesting of the bee *Lasioglossum (Evykæus) quadrinotatum* (Hymenoptera, Halictidae). *Vestn. Leningr. Univ.*, (1), 107-109.
- Sitdikov, A. A. (1988). Nesting of the bee *Halictus quadricinctus* (F.) (Hymenoptera, Halictidae) in the Udmurt ASSR. *Entomol. Rev.*, 67(2), 66-77.
- Sitdikov, A. A., & Pesenko, Y. A. (1988). A subgeneric classification of bees of the genus *Eucera scopoli* (Hymenoptera, Anthophoridae) with a scheme of phylogenetic relationships between the genera. *Tr. Zool. Inst. Akad. Nauk SSSR*, 175, 74-81.
- Skhirtladze, I. A. (1987). Material on the Apoidea (Hymenoptera) fauna in Talysh. *Vestn. Gos. Muz. Gruz.*, 34(4), 133-154.
- Skinner, J. A. (1988). Pollination of male-sterile sunflower by bees in California. *Dissertation Abstr. Internat. B (Sci. & Engin.)*, 48(8), 120pp.
- Skou, J. P. (1988). Japanese species of *Ascospaera*. *Mycotaxon.*, 31(1), 173-190.
- Skou, J. P., & Holm, S. N. (1989). *Ascospaera tenax* new species and a variant of *Ascospaera aggregata*. *Mycotaxon.*, 35(2), 211-218.
- Smith, B. H. (1987). Effects of genealogical relationship and colony age on the dominance hierarchy in the primitively eusocial bee *Lasioglossum zephyrum*. *Anim. Behav.*, 35(1), 211-217.
- Smith, B. H. (1988). Genealogical relationship and social dominance in bees: a reply to Kukuk and May. *Anim. Behav.*, 36(6), 1850-1851.
- Smith, B. H., & Ayasse, M. (1987). Kin-based male mating preferences in two species of halictine bee. *Behav. Ecol. Sociobiol.*, 20(5), 313-318.
- Smith, B. H., & Ayasse, M. (1987). The role of kin recognition in the mating preferences of male halictine bees (Hymenoptera: Halictidae). In: Eder, J. & Remold, H. (Eds.) *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Smith, B. H., & Weller, C. (1987). What information do social halictine bees encode in the amount of the macrocyclic lactone pheromones? In: Eder, J. & Remold, H. (Eds.) *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.

- Smith, B. H., & Wenzel, J. W. (1988). Pheromonal co-variation and kinship in social bee *Lasioglossum zephyrum* (Hymenoptera: Halictidae). *J. Chem. Ecol.*, 14(1), 87-94.
- Snelling, R. R. (1988). The generic placement of *Prosopis? Allodape? mustela* Vachal, 1895 (Hymenoptera: Apoidea). *Entomol. News*, 99(1), 10-12.
- Snelling, R. R. (1988). A new species of *Centris (Melanocentris)* from Cerro de la Neblina Venezuela (Hymenoptera: Anthophoridae). *Entomol. News*, 99(1), 13-16.
- Snelling, R. R. (1987). A revision of the bee genus *Aztecancanthidium* (Hymenoptera: Megachilidae). *Pan-Pac. Entomol.*, 63(2), 165-171.
- Snelling, R. R. (1987). Geographical inexactitude. *Pan-Pac. Entomol.*, 63(4), 339-340.
- Snelling, R. R., & Rozen Jr., J. G. (1987). Contributions toward a revision of the New World Nomadine bees. 2. The genus *Melanomada* (Hymenoptera: Anthophoridae). *Contr. Sci.*, (384), 12pp.
- Snow, A. A., & Roubik, D. W. (1987). Pollen deposition and removal by bees visiting two tree species in Panama. *Biotropica*, 19(1), 57-63.
- Soares, A. A., Campos, L. A. D. O., Vieirea, M. F., & Melo, G. A. R. D. (1989). Relationship between *Euglossa mandibularis* Freise 1899 (Hymenoptera, Apidae, Euglossini) and *Cyphomandra calycina* Solanaceae. *Cienc. Cult. (Sao Paulo)*, 41(9), 903-905.
- Soltz, R. L. (1987). Interspecific competition and resource utilization between bumblebees. *Southwest. Nat.*, 32(1), 30-52.
- Sommeijer, M. J. (1987). Age-polyethism in stingless bees and evidence of flexible individual ontogenetic sequences. In: J. Eder & H. Remold. *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Sommeijer, M. J. (1987). An ethological analysis of intra-colonial competition in relation to reproductive dominance in the stingless bee *Melipona favosa*. In: J. Eder & H. Remold. (Eds.) *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.
- Sommeijer, M. J. (1987). Social control in Meliponinae. *Jaarb. Uitg. Ned. Entomol. Ver* 1986, 24-27.
- Sommeijer, M. J., & Bootsma, M. C. (1988). *Melipona trinitatis* as a producer of honey in Trinidad, West Indies. *Actes Colloq. Insectes Soc.*, 4, 291-294.
- Sonet, M., & Jacob-Remade, A. (1987). Pollination de la legumineuse fourragere *Hedysarum coronarium* L. en Tunisie. *Bull. Rech. Agron. Gembloux*, 22(1), 19-32.
- Soomro, A. H., Khuhro, R. D., & Ansari, H. U. (1986/1987). Status of insects associated with mango blossom. *Proc. 5th Pakistan Congress Zool.*, Univ. Karachi, 123-125.
- Souza, J. M., & Ventura, D. F. (1987). Temporal summation and critical duration in bee photoreceptors: a comparison of measurement methods. *Braz. J. Med. Biol. Res.*, 20(5), 649-652.
- Sowig, P. (1989). Effects of flowering plant's patch size on species composition of pollinator communities foraging strategies and resource partitioning in bumblebees (Hymenoptera: Apidae). *Oecologia*, 78(4), 550-558.
- Spofford, M. G., Kurczewski, F. E., & Downes Jr., W. L. (1989). Nearctic species of Miltogrammini (Diptera, Sarcophagidae) associated with species of Aculeata Hymenoptera (Vespidae, Pompiloidea, Sphecoidea, Apoidea). *J. Kans. Entomol. Soc.*, 62(2), 254-267.
- Standley, L. A., Kim, S. S. H., & Hjersted, I. M. (1988). Reproductive biology of two sympatric species of *Chimaphila*. *Rhodora*, 90(863), 233-244.
- Starr, C. K. (1989). *Bombus folsomi* and the origin of Philippine bumble bees (Hymenoptera: Apidae). *Syst. Entomol.*, 14(4), 411-415.
- Starr, C. K. (1989). In reply, is the sting the thing? *Ann. Entomol. Soc. Am.*, 82(1), 6-8.
- Starr, C. K., & Sakagami, S. F. (1987). An extraordinary concentration of stingless bee colonies in the Philippines, with notes on nest structure (Hymenoptera: Apidae: *Trigona spp.*). *Insectes Sociaux*, 34(2), 96-107.
- Starr, C. K., & Jacobson, R. S. (1990). Nest structure in Philippine hornets (Hymenoptera, Vespidae, *Vespa spp.*). *Jpn. J. Ent.*, 58(1), 125-143.
- Staurengo da Cunha, M. A. (1988). Ultrastructure of the oocytes of *Scaptotrigona postica* workers (Hymenoptera: Apidae). *Naturalia*, 13, 103-115.
- Staurengo da Cunha, M. A., Cruz-Landim, C. d., & Gomes, G. M. (1988). Glandular dimorphism in female castes of *Frieseomelitta sivestri* Languida: wax glands and glands associated with the sting. *Naturalia*, 13, 37-43.
- Staurengo da Cunha, M. A., Cruz-Landim, C. D., & Gomes, G. M. (1988). Sexual dimorphism in the female castes of *Frieseomelitta silvestrilinguis* (Hymenoptera, Apidae) wax glands and glands associated with the sting. *Naturalia*, 13(0), 37-44.
- Steiner, K. E. (1990). The *Diascia* (Scrophulariaceae) window: an orientation cue for oil-collecting bees. *Bot. J. Linn. Soc.*, 102(2), 175-195.
- Steiner, K. E. (1989). The pollination of *Disperis* (Orchidaceae) by oil-collecting bees in southern Africa. *Lindleyana*, 4(4), 164-183.
- Steiner, K. E., & Whitehead, V. B. (1988). The association between oil-producing flowers and oil-collecting bees in the Drakensberg of southern Africa. *Monogr. Syst. Bot. Missouri Bot. Gard.*, 25, 249-277.
- Steinmann, E. (1986-1987). Hosenbienen in Rhazuns. *Jahresber Naturforsch Ges Graubuenden Neue Folge*, 104, 169-170.
- Stephen, W. P. (1987). *Megachile (Eutricharea) apicalis*, an introduced bee with potential as a domesticatable alfalfa pollinator. *J. Kans. Entomol. Soc.*, 60(4), 583-584.

- Stoddard, F. L., & Bond, D. A. (1987). The pollination requirements of the faba bean. *Bee World*, 68(3), 144-152.
- Stokes, J. (1988). Pollination of the wild gladiolus (*Gladiolus illyricus*) in the new forest, Hampshire by butterflies. *News Br. Butterfly Conserv. Soc.*, (41), 38-39.
- Stone, G. N., & Willmer, P. G. (1989). Endothermy and temperature regulation in bees: a critique of 'grab and stab' measurement of body temperature. *J. Exp. Biol.*, 143, 211-223.
- Stone, G. N., & Willmer, P. G. (1989). Warm-up rates and body temperatures in bees the importance of body size thermal regime and phylogeny. *J. Exp. Biol.*, 147, 303-328.
- Stort, A. C., Bueno de Moraes, M. M., & Barelli, N. (1987). Scanning electron microscope observations on the head of *Scaptotrigona postica* workers (Hymenoptera: Apidae). *J. Kans. Entomol. Soc.*, 60(1), 162-166.
- Stort, A. C., Bueno de Moraes, M. M., & Barelli, N. (1988). Scanning electron microscope observations on the abdominal terga of *Scaptotrigona postica* workers (Hymenoptera, Apidae). *Rev. Bras. Entomol.*, 32 (1), 63-68.
- Storti, E. F. (1988)(1989). Floral biology of *Solanum sessiliflorum* var. *sessiliflorum* in the Manaus region Amazonas Brazil. *Acta Amazonica*, 18(3-4), 55-66.
- Strojny, W. (1987). The sole polish colony of *Chalicodoma muraria* (F.) (Hymenoptera, Apidae) extinct in the reserve Skalki Stoeckie, near Zabkowice Slaskie. *Przegl. Zool.*, 31(3), 341-348.
- Sugden, E. A. (1989). A semi-natural, manipular observation nest for *Exoneura* spp. and other allodapine bees (Hymenoptera: Anthophoridae). *Pan-Pac. Entomol.*, 65(1), 17-24.
- Surholt, B., Grieve, H., Hommel, C., & Bertsch, A. (1988). Fuel uptake, storage and use in male bumble bees *Bombus terrestris* L. *J. Comp. Physiol. B: Biochem. Syst. Environ. Physiol.*, 158(2), 263-269.
- Sutcliffe, G. H., & Plowright, R. C. (1988). The effects of food supply on adult size in the bumble bee *Bombus terricola* Kirby (Hymenoptera: Apidae). *Can. Entomol.*, 120(12), 1051-1058.
- Suzuki, K. (1989). Nectar sucking pollinators in hybrid derivatives of *Epinedium* Berberidaceae. *Plant Species Biol.*, 2(1-2), 137-140.
- Taber, S. (1989). How to breed resistant bees to any disease. *Am. Bee J.*, 129(6), 400-401.
- Taber, S. (1987). The scientific method when applied to bees. *Am. Bee J.*, 127(11), 783-784.
- Tadauchi, O., Hirashima, Y., & Matsumura, T. (1987). Synopsis of *Andrena* (*Andrena*) of Japan (Hymenoptera: Andrenidae). Part 1. *J. Fac. Agric. Kyushu Univ.*, 31(1-2), 11-35.
- Tadauchi, O., Hirashima, Y., & Matsumura, T. (1987). Synopsis of *Andrena* (*Andrena*) of Japan (Hymenoptera: Andrenidae). Part 2. *J. Fac. Agric. Kyushu Univ.*, 31(1-2), 37-54.
- Tadauchi, O., & Hirashima, Y. (1988). Synopsis of *Andrena* (*Stenomelissa*) with a new species from Japan (Hymenoptera, Andrenidae). *J. Fac. Agric. Kyushu Univ.*, 33(1-2), 67-76.
- Takahashi, H. (1987). A comparative floral and pollination biology of *Tricyrtis flava maxim*, *T. nana yatabe* and *T. ohsumiensis masamune*. *Bot. Magaz.*, 100, 185-203.
- Tanacs, L. (1987). Biogeographic ecofaunistic and flight-dynamic composition of the Apoidea fauna at the Tisza Valley Hungary. *Acta Univ. Szeged. Acta Biol.*, 33(1-4), 105-114.
- Tanacs, L. (1987). The trend of the composition of the wild-bee population under the influence of anthropogenic effects in the biotopes of the Kiskore storage-tank region at the Tisza Valley. *Tiscia. (Szeged.)*, 22(0), 109-120.
- Tasei, J. N., Carre, S., Bosio, P. G., Debray, P., & Hariot, J. (1987). Effects of the pyrethroid insecticide, W185871 and phosalone on adults and progeny of the leaf-cutting alfalfa bee *Megachile rotundata* F. pollinator of lucerne. *Pestic. Sci.*, 21(2), 119-128.
- Tengo, J., Eriksson, J., Borg-Karlson, A. K., Smith, B. H., & Dobson, H. (1988). Mate-locating strategies and multimodal communication in male behavior of *Panurgus banksianus* and *P. calcaratus* (Apoidea, Andrenidae). *J. Kans. Entomol. Soc.*, 61(4), 388-395.
- Tepedino, V. J. (1988). Host discrimination in *Mondontomerus obsoletus* Fabricius (Hymenoptera: Torymidae), a parasite of the alfalfa leafcutting bee *Megachile rotundata* (Fabricius) (Hymenoptera: Megachilidae). *J. N. Y. Entomol. Soc.*, 96(1), 113-118.
- Tepedino, V. J., Knapp, A. K., Eickwort, G. C., & Ferguson, D. C. (1989). Death camas (*Zigadenus nuttallii*) in Kansas: pollen collectors and a florivore. *J. Kans. Entomol. Soc.*, 62(3), 411-412.
- Tepedino, V. J., & Parker, F. D. (1988). Alteration of sex ratio in a partially bivoltine bee *Megachile rotundata* (Hymenoptera, Megachilidae). *Ann. Entomol. Soc. Am.*, 81(3), 467-476.
- Tepedino, V. J., & Stackhouse, M. (1987). Bee visitors of sweetvetch, *Hedysarum boreale boreale* (Leguminosae), and their pollen-collecting activities. *Great Basin Nat.*, 47(2), 314-318.
- Tepedino, V. J., & Torchio, P. F. (1989). Influence of nest hole selection on sex ratio and progeny size in *Osmia lignaria propingua* (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.*, 82(3), 355-360.
- Teppner, H. (1988). *Lathyrus grandiflorus* (Fabaceae-vicieae): flower structure, function and *Xylocopa violacea*. *Phyton.*, 28(2), 321-336.
- Thomson, J. D. (1988). Effects of variation in inflorescence size and floral rewards on the visitation rates of traplining pollinators of *Aralia hispida*. *Evol. Ecol.*, 2(65-76),

- Thomson, J. D. (1989). Reversal of apparent feeding preferences of bumble bees by aggression from *Vespa* wasps. *Can. J. Zool.*, 67(10), 2588- 2591.
- Thomson, J. D., McKenna, M. A., & Cruzan, M. B. (1989). Temporal patterns of nectar and pollen production in *Aralia hispida*: implications for reproductive success. *Ecol. Publ. Ecol. Soc. Am.*, 70(4), 1061-1068.
- Thomson, J. D., Peterson, S. C., & Harder, L. D. (1988). Response of traplining bumble bees to competition experiments: shifts in feeding location and efficiency. *Oecologia*, 71, 295-300.
- Thorp, R. W. (1987). A new species of *Andrena* (*Onagrandrena*) from Utah's San Rafael desert (Hymenoptera: Andrenidae). *Pan-Pac. Entomol.*, 63 (2), 194-198.
- Thorp, R. W., & Sugden, E. A. (1990). Extrafloral nectaries producing rewards for pollinator attraction in *Acacia longiflora* Andr. Willd. *Isr. J. Bot. Basic Appl. Plant Sci.*, 3 (1-2), 177-186.
- Tkalcu, B. (1987). Die polytypie zweier westpalaarktischen arten der gattung *Melitta* (Hymenoptera, Apoidea). *Cas. Slezskeho. Muz. Ser. A Vedy. Prir.*, 36(2), 177-182.
- Tkalcu, B. (1987). Field observation on the *Anthocopa mocsaryi* Friese imagoes in South Slovakia Czechoslovakia (Hymenoptera: Apoidea; Megachilidae). *Biol. (Bratislava)*, 42(10), 1005-1010.
- Tkalcu, B. (1988). The Lectotype of *Chalicodoma heinii* Kohl 1906 (Hymenoptera: Apoidea: Megachilidae). *Z. Arbeitsgem. Oesterr. Entomol.*, 40(1-2), 51-52.
- Tkalcu, B. (1988). New palearctic species and subspecies of genera *Chalicodoma* and *Megachile* (Hymenoptera: Apoidea; Megachilidae). *Vestn. Cesk. Spol. Zool.*, 52(1), 48-62.
- Tkalcu, B. (1989). New taxa of Asiatic bumblebees (Hymenoptera, Apoidea). *Acta. Entomol. Bohemoslov.*, 86(1), 39-60.
- Torchio, P. F. (1989). Biology, immature development, and adaptive behavior of *Stelis montana*, a cleptoparasite of *Osmia* (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.*, 82(5), 616-632.
- Torchio, P. (1988). The blue orchard bee: an alternative pollinator of apples. *Utah Sci. Agric. Exp. Stn.*, 49 (1), 2-9.
- Torchio, P. F. (1989). In-nest biologies and development of immature stages of three *Osmia* species (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.*, 82(5), 599-615.
- Torchio, P. F., Asensio, E., & Thorp, R. W. (1987). Introduction of the European bee, *Osmia cornuta*, into California almond orchards (Hymenoptera: Megachilidae). *Environ. Entomol.*, 16, 664-667.
- Torchio, P. F., & Burdick, D. J. (1988). Comparative notes on the biology and development of *Epeorus compactus* Cresson, a cleptoparasite of *Colletes kincaidii* Cockerell. (Hymenoptera: Anthophoridae: Colletidae). *Ann. Entomol. Soc. Am.*, 81(4), 626-636.
- Torchio, P. F., & Burwell, B. (1987). Notes on the biology of *Cardequela occidentalis* (Hymenoptera: Colletidae), and a review of colletid pupae. *Ann. Entomol. Soc. Am.*, 80(6), 781-789.
- Torchio, P. F., Trostle, D. E., & Burdick, D. J. (1988). The nesting biology of *Colletes kincaidii* Cockerell (Hymenoptera: Colletidae) and development of its immature forms. *Ann. Entomol. Soc. Am.*, 81(4), 605-625.
- Tomeiros, L. M., Farinha, E. K., Cruz-Landim, C. d., & Silva de Moraes, R. L. M. (1988). Hormonal influence on the DNA content of salivary gland cells in larvae of *Melipona quadrifasciata anthidioides* Lep. (Apidae, Meliponinae). 2. Ecdysine. *Naturalia*, 13, 85-90.
- Toro, H. (1989). Addenda et corrigenda. *Acta Ent. Chilena*, 15, 335.
- Toro, H. (1989). Contribution to the knowledge of Chilean Panurginae (Apoidea, Andrenidae). *Acta Ent. Chilena*, 15(0), 229-232.
- Toro, H., & Chiappa, E. (1989). New species and subspecies of *Centris* (Hymenoptera: Apoidea) associated to *Prosopis tamarugo*. *Acta Ent. Chilena*, 15(0), 243-248.
- Toro, H., Frederick, Y., & Henry, A. (1989). Hylaeinae (Hymenoptera: Colletidae), a new subfamily of bees for the Chilean Fauna. *Acta Ent. Chilena*, 15(0), 201-204.
- Toro, H., & Magunacelaya, J. C. (1987). Femoral muscular structure of Xeromelissinae (Hymenoptera, Colletidae). *Acta Ent. Chilena*, 14, 13- 24.
- Tripplett, D. C., & Gittins, A. R. (1988). Nesting, mating and foraging habits of *Melissodes* (*Melissodes*) *tepidus tepidus* Cresson in Idaho (Hymenoptera: Anthophoridae). *Proc. Entomol. Soc. Wash.*, 90(4), 462- 470.
- U. S. Dep. Agric. (1987). Researchers seek to save queen bees from wrong mates. *Am. Bee J.*, 127(12), 871.
- Uji, T. (1987). Pollination in *Nephileium lappaceum* var. *lappaceum*. *Berita. Biol.*, 32(Suppl.), 31-34.
- Urban, D. (1989). New species of and notes on the genus *Gaesischia* Michener Laberge and Moure 1955 (Hymenoptera, Apoidea). *Rev. Bras. Entomol.*, 33(1), 75-102.
- Utami, N., & Kahono, S. (1989). Pollination on the Pinang Yaki Areca- vestaria Giseke in the botanical gardens Indonesia. *Berita. Biol.*, 3(9), 470-472.
- Vaillant, J. (1987). What is the Varoutest? *Sante Abeille*, (102), 294.
- Valenta, V., & Arbaciauskas, K. (1988). Species composition and distribution of bumblebees (*Bombus* Lat.) in the Lithuanian SSR. *Acta Entomol. Lituan.*, 9, 111-115.
- van Buren, N. W. M., & Sommeijer, M. J. (1988). Etude des facteurs determinants pour la dominance reproductive de la reine de *Melipona trinitatis*. *Actes Colloq. Insectes Soc.*, 4, 285-290.

- van Doorn, A. (1989). Factors influencing dominance behavior in queenless bumblebee workers (*Bombus terrestris*). *Physiol. Entomol.*, 14(2), 211-221.
- van Doorn, A. (1987)(1988). Investigations into the regulation of dominance behaviour and of the division of labour in bumblebee colonies (*Bombus terrestris*). *Neth. J. Zool.*, 37(3-4), 255-276.
- van Doorn, A. (1987). Juvenile hormone and worker behavior in the bumblebee *Bombus terrestris*. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Velthuis, H. H. W. (1987). Caste differentiation and egg laying in the highly social bees. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Velthuis, H. H. W. (1989). The comparative study of sociality in the higher bees. *Actes Colloq. Insectes Soc.*, 5, 2-15.
- Velthuis, H. H. W. (1987). The evolution of sociality: ultimate and proximate factors leading to primitive social behaviour in carpenter bees. *Exper. Suppl.*, 54, 405-430.
- Velthuis, H. H. W. (1987). Pollen deposition and removal by bees visiting two tree species in Panama. *Biotropica*, 19(1), 57-63.
- Vergara, C. A. Villa, & Nates Parra, G. (1987). Nidificacion de Meliponinos (Hymenoptera: Apidae) de la region Central de Colombia. *Rev. Biol. Trop.*, 34(2),
- Villalobos, E. M., & Shelly, T. E. (1987). Observations on the behavior of male *Bombus sonorus* (Hymenoptera: Apidae). *J. Kans. Entomol. Soc.*, 60(4), 541-548.
- Vinson, S. B., & Frankie, G. W. (1988). A comparative study of the ground nests of *Centris flavifrons* and *Centris aethiocesta* (Hymenoptera: Anthophoridae). *Entomol. Exp. Appl.*, 49(1-2), 181-187.
- Vinson, S. B., Frankie, G. W., & Coville, R. E. (1987). Nesting habits of *Centris flavofasciata* Friese (Hymenoptera: Apoidea: Anthophoridae) in Costa Rica. *J. Kans. Entomol. Soc.*, 60(2), 249-263.
- Vinson, S. B., Williams, H. J., McAuslane, H. J., & Frankie, G. W. (1989). Chemical contents of male mandibular glands of three *Centris* species (Hymenoptera: Anthophoridae) from Costa Rica. *Comp. Biochem. Physiol. B: Comp. Biochem.*, 93 (1), 73-76.
- Vogel, S. (1988). Die Oblumensymbiosen-parallelismus und andere Aspekte ihrer Entwicklung in Raum und Zeit. *Z. Zool. Syst. Evolutionsforsch.*, 26(5), 341-362.
- Waddington, K. D. (1987). Nutritional ecology of bees. In: Slansky, F., Jr. & Rodriguez, J.G. (Eds). *Nutritional Ecology of Insects, Mites, Spiders and Related Invertebrates*. John Wiley & Sons, New York, Chichester etc.,
- Warncke, K. (1988). Isolierte bienenvorkommen auf dem olymp in griechenland (Hymenoptera, Apidae). *Proc. Natl. Acad. Sci. USA*, 85(12), 4374-4377.
- Warncke, K. (1985)(1987)). Supplemental studies on bees of the genera *Panurgus* and *Melitturga* (Andrenidae, Apidae) chiefly from the Turkish area. *Boll. Mus. Civ. Stor. Nat. Venezia*, 36(0), 75-108.
- Waser, N. M. (1988). Comparative pollen and dye transfer by pollinators of *Delphinium nelsonii*. *Funct. Ecol.*, 2 (1), 41-48.
- Waser, N. M., & Mitchell, R. J. (1990). Nectar standing crops in *Delphinium nelsonii* flowers: spatial auto-correlation among plants. *Ecol. Publ. Ecol. Soc. Am.*, 71(1), 116-123.
- Watson, M. (1987). Strangers unawares. *Edinb. Nat. Hist. Soc. J.*, 28.
- Waslo, W. T. (1987). The role of learning in the mating biology of a sweat bee *Lasioglossum zephyrum* (Hymenoptera: Halictidae). *Behav. Ecol. Sociobiol.*, 20 (3), 170-185.
- Waslo, W. T. (1987). The roles of seasonality, host synchrony, and behaviour in the evolutions and distributions of nest parasites in Hymenoptera (Insecta), with special reference to bees (Apoidea). *Biol. Rev.*, 62(4), 515-543.
- Webster, T. C., & Peng, Y.-S. (1988). The evolution of food-producing glands in eusocial bees. (Apoidea, Hymenoptera). *J. Evol. Biol.*, 2, 165-176.
- Werren, J. H. (1987). Labile sex ratios in wasps and bees. *Biosci.*, 37(7), 498-506.
- Westrich, P. (1989). Die Wildbienen Baden-Wurttembergs. Allgemeiner Teil: Lebensräume, Verhalten, Ökologie und Schutz. Verlag Eugen Ulmer, Stuttgart.,
- Westrich, P. (1989). Die Wildbienen Baden-Wurttembergs. Spezieller Teil: die Gattungen und Arten. Verlag Eugen Ulmer, Stuttgart.,
- Westrich, P., & Schmidt, K. (1987). Pollenanalyse, ein Hilfsmittel beim Studium des Sammelverhaltens von Wildbienen (Hymenoptera, Apoidea). *Apidologie*, 18(2), 199-214.
- Whitten, W. M., Hills, H. G., & Williams, N. H. (1988). Occurrence of ipsdienol in floral fragrances. *Phytochemistry*, 27(9), 2759-2760.
- Whitten, W. M., Young, A. M., & Williams, N. H. (1989). Function of glandular secretions in fragrance collection by male euglossine bees (Apidae, Eulangini). *J. Chem. Ecol.*, 15(4), 1285-1296.
- Williams, H. J., Vinson, S. B., & Frankie, G. W. (1987). Chemical content of the dorsal mesosomal gland of two *Xylocopa* species (Hymenoptera: Anthophoridae) from Costa Rica. *Comp. Biochem. Physiol.*, 86(2), 311-312.
- Williams, I. H. (1987). The pollination of lupins. *Bee World*, 68(1), 10-16.
- Williams, N. H., & Whitten, W. M. (1988). *Stanhopea panamensis* a new species from Central Panama (Orchidaceae). *Lindleyana*, 3(1), 9-11.
- Williams, P. H. (1988). Habitat use by bumble bees (*Bombus* spp.). *Ecol. Entomol.*, 13(2), 223-237.

- Williams, P. H. (1989). Why are there so many species of bumble bees at Dungeness? *Bot. J. Linn. Soc.*, 101 (1), 31-44.
- Willmen, P. G., & Stone, G. N. (1989). Incidence of entomophilous pollination of lowland coffee *Coffea canephora* the role of leaf cutter bees in Papua, New Guinea. *Entomol. Exp. Appl.*, 50(2), 113- 124.
- Willmer, P. G. (1988). The role of insect water balance in pollination ecology: *Xylocopa* and *Calotropis*. *Oecologia*, 76(3), 430-438.
- Wirtz, P., Szabados, M., Pethig, H., & Plant, J. (1988). An extreme case of interspecific territoriality: male *Anthidium manicatum* (Hymenoptera, Megachilidae) wound and kill intruders. *Ethol.*, 78(2), 159-167.
- Wittman, D. (1989). Nest architecture, nest site preferences and distribution of *Plebeia wittmanni* (Moure & Camargo, 1989) in Rio Grande do Sul, Brazil (Apidae: Meliponinae). *Stud. Neotrop. Fauna Environ.*, 24 (1), 17-23.
- Wittmann, D., Cure, J. R., & Radtke, R. (1987). Mating and pollinating behavior of *Callonychium* sp. (Andrenidae) on *Petunia Integifolia* and *Ovalofoli* (Solanaceae). In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Wittmann, D., Hoffmann, M., & Scholz, E. (1988). Southern distributional limits of euglossine bees in Brazil linked to habitats of the Atlantic and subtropical rain forest (Hymenoptera, Apidae, Euglossini). *Entomol. Gen.*, 14(1), 53-60.
- Wittmann, D., & Scholz, E. (1987). Nectar dehydration and body weight reduction: the cues of *Xylocopa nigrocincta* to prolong duration of territorial flights. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny Munich.,
- Wittmann, D., & Scholz, E. (1989). Nectar dehydration by male carpenter bees as preparation for mating flights. *Behav. Ecol. Sociobiol.*, 25(6), 387-391.
- Wittmann, D., Scholz, E., & Hoffmann, M. (1987). Occurrence and distributional limits of euglossine bees in Rio Grande do Sul, Southern Brazil. In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Wolf, H. (1988). Massenbesuch von Furchenbienen-männchen (Hymenoptera, Apoidea, Halictidae) an blutenstanden der kanadischen goldrute. *Hess. Faun. Briefe.*, 8(4), 64-65.
- Wolf, H. (1987). Zwitter von *Sphecodes geofrellus* (Kirby) und *Sphecodes rufiventris* (Panzer) (Hym., Sphecoidea, Apidae). *Linzer Biol. Beitr.*, 19(2), 299-300.
- Wolfe, L. M., & Barrett, S. C. H. (1987). Pollinator foraging behavior and pollen collection on the floral morphs of *Tristylous pontederia cordata* L. *Oecologia*, 74(3), 347-351.
- Woyke, H. (1988). The influence of weather on insect foraging intensity in seed onions. *Biuletyn Warywniczy*, 31, 43-58.
- Wright, D. H. (1988). Temporal changes in nectar availability and *Bombus appositus* (Hymenoptera: Apidae) foraging profits. *South. Nat.*, 33(2), 219-227.
- Wu, Y.-R. (1988). Four new species of bees from China (Hymenoptera, Apoidea). *Acta Zootaxon. Sin.*, 13(1), 67-71.
- Wu, Y.-R. (1988). Hymenoptera: Apoidea. *Insects of Mt. Namjagbarwa Region of Xizang.*, 549-552.
- Wu, Y.-R. (1988). A new species of *Anthophora* from China (Hymenoptera, Apoidea, Anthophoridae). *Acta Entomol. Sin.*, 31(2), 210-212.
- Wu, Y.-R. (1987). A study on Chinese Halictoides with descriptions of 3 new species (Halictidae: Dufoureinae). *Sinozool.*, (5), 187-201.
- Wu, Y.-R. (1987). A study on Chinese Hoplitis with descriptions of new species (Hymenoptera: Megachilidae). *Acta Entomol. Sin.*, 30(4), 441- 449.
- Wu, Y. R., & Kuang, B. (1987). Two species of small honeybee-a study of the genus *Micrapis*. *Bee World*, 68 (3), 153-155.
- Yamada, M., & Sakagami, F. (1988). Non-parasitic halictid bees in Tsugaru District (Anomori Prefecture), northernmost Honshu. *J. Amomori-Ken Biol. Soc.*, 25, 10-21.
- Yanega, D. (1989). Caste determination and differential diapause within the first brood of *Halictus rubicundus* in New York (Hymenoptera: Halictidae). *Behav. Ecol. Sociobiol.*, 24(2), 97-108.
- Yanega, D. (1988). Social plasticity and early-diapausing females in a primitively social bee. *Proc. Nat. Acad. Sci. USA*, 85(12), 4374-4377.
- Yanega, D. (1990). Philopatry and nesting founding in a primitively social bee, *Halictus rubicundus*. *Behav. Ecol. Sociobiol.*, 27, 37-42.
- Yavetz, I. (1988). Jean Henri Fabre and evolution: indifference or blind hatred? *Hist. Philos. Life Sci.*, 10 (1), 3-36.
- Yong, H. S. , Khoo, S. G. , Sarjan, R., & Tho, V. P. (1987). Glucose phosphate isomerase, malate dehydrogenase and isocitrate dehydrogenase allozyme variation in the stingless bee, *Trigona (Tetragonula) fuscobalteata* (Insecta: Hymenoptera: Apidae), from Peninsular Malaysia. *Comp. Biochem. Physiol. B. Comp. Biochem.*, 87(3), 465-467.
- Yoshida, A., & Maeta, Y. (1988). Utilization of *Osmia cornifrons* (Radoszkowski) as a pollinator of apples in south-western Japan. *Honeybee Sci.*, 9(1), 1.
- Young, A. M. (1987)(1989). Differential foraging by flies (Diptera) and stingless bees (Hymenoptera: Apidae: Meliponinae). *Berenesia*, 27, 69- 73.
- Young, A. M., Erickson, B. J., & Erickson Jr., E. H. (1989). Pollination biology of *Theobroma* and *Herrania* Sterculiaceae III. Steam- distilled floral oils of *Theobroma* species as attractants to flying insects in a Costa Rican cocoa plantation. *Insect Sci. Appl.*, 10 (1), 93-98.

- Youssef, N. N., & Brindley, W. A. (1989). Effectiveness of Botran and DPX 965 as growth inhibitors of *Ascosphaera aggregata* (Ascosporeaceae) in the alfalfa leafcutting bee (Hymenoptera Megachilidae). *J. Econ. Entomol.*, 82(5), 1335-1338.
- Yumoto, T. (1988). Pollination systems in the cool temperate mixed coniferous and broad-leaved forest zone of Yakushima Island Japan. *Ecol. Res.*, 3(2), 117-130.
- Zandén, G. v. der. (1988). Contribution to the systematics and nomenclature of Palearctic Osmiini, with notes on their distribution. *Zool. Mededel.*, 62(9), 113-133.
- Zandén, G. v. der. (1987). New palaeartic taxa of the family Megachilidae (Insecta, Hymenoptera, Apoidea). *Reichenbachia*, 25(16), 73-84.
- Zandén, G. v. der. (1988). Nomenclatural and taxonomic comments on some palearctic species of the family Megachilidae (Insecta: Hymenoptera: Apoidea: Megachilidae). *Reichenbachia*, 26(10), 55-64.
- Zandén, G. v. der. (1987). Some bees from South Arabia (Hymenoptera, Aculeata, Apidae). *Mitt. Zool. Mus. Berl.*, 63(2), 331-332.
- Zeil, J., & Wittmann, D. (1987). Visual orientation towards the nest in a stingless bee (*Tetragonisca angustula*). In: Eder, J. & Remold, H. (Eds). *Chemistry and Biology of Social Insects*. Verlag J. Peperny, Munich.,
- Zeil, J., & Witmann, D. (1989). Visually controlled station-keeping by hovering guard bees of *Trigona (Tetragonisca) angustula* (Apidae, Meliponinae). *J. Comp. Physiol. A: Sens. Neural Behav. Physiol.*, 165(5), 711-718.
- Zhu, T., Zhu, F., Ning, L., & Jiang, K. (1989). A preliminary study on pollinating bees for *Amomum Compactum* Soland ex Maton. *China J. Chin. Mater. Med.*, 14(1), 17-19.
- Zimmerman, J. K., & Mandarinan, R. S. (1988). Age structure of male *Euglossa imperialis* (Hymenoptera: Apidae: Euglossini) at nectar and chemical sources in Panama. *J. Trop. Ecol.*, 4(Pt. 3), 303-306.
- Zimmerman, J. K., Roubik, D. W., & Ackerman, J. D. (1989). Asynchronous phenologies of a neotropical orchid and its euglossine bee pollinator. *Ecol. Publ. Ecol. Soc. Am.*, 70(4), 1192-1195.
- Zimmerman, M. (1988). Pollination biology of montane plants: relationship between rate of nectar production and standing crop. *Am. Midland Natur.*, 120(1), 50-57.
- Zimmerman, M. (1987). Reproduction in *Polemonium*: factors influencing outbreeding potential. *Oecologia (Berlin)*, 72, 624-632.
- Zimmerman, M., Nagy, E. S., & Galloway, L. (1987). Nectar dispersion patterns in three Australian plant species. *Aust. J. Ecol.*, 12, 183-188.
- Zimmerman, M., & Pyke, G. H. (1988). Experimental manipulations of *Polemonium foliosissimum*: effects on subsequent nectar production, seed production and growth. *J. Ecol.*, 76, 777-789.
- Zimmerman, M., & Pyke, G. H. (1988). Pollination ecology of Christmas bells (*Blandfordia nobilis*): effects of pollen quantity and source on seed set. *Aust. J. Ecol.*, 13, 93-99.
- Zlobin, V. V. (1988). The connections between entomofauna of the north Europe and Siberia. *Zool. Instit. Leningrad*, 126-145.
- Zoeckler, C. (1988). Moist meadow flora and pollinating insects, their importance in grassland expansion. *Faun-Oekol. Mitt.*, 6(1-2), 5-20.