ICHNEWS

A newsletter for students of the Ichneumonidae (Hymenoptera)

Number 4

June 1978 .

Edited and published by: M.G. Fitton

and Department of Entomology British Museum (Natural History) of Entomology Cromwell Rd,,

I.D. Gauld Commonwealth Institute

A SOL THE THE .

London, SW7 5BD England

EDITORIAL

In the year that has passed since Ichnews 3 appeared we have received little support from you. We need to have more items for inclusion. Much has been happening in the last 12 months. A large number of papers have been published, including an exceptional number of major revisionary and reference works. The work we do in preparing the bibliography and directory must be balanced by notes, news and views sent in by you.

CURRENT NEWS AND NOTES

- J-F. Aubert has completed his catalogue of the Western Palaearctic Banchinae. This work will be published in the near future and an appendix will include keys to the species of European Glypta and Lissonota.
- Gerd Heinrich has completed three works, the Eastern Palaearctic Ichneumoninae, the Burmese Ichneumoninae part XI and the Ichneumoninae of Florida and the neighbouring Austroriparian zone. These works will be in Russian, German and English respectively and should be appearing in the near future.
- Mattias Idar has completed a revision of the European Hadrodactylus which should be appearing soon. Mattias also informs us that there is a list of specimens in the Thunberg collection compiled by Lars Hedström. Further information is available through Mattias.
- John Short's monograph on Ichneumonid larvae will be appearing any day now. It is being published by the American Entomological Institute. John is now starting a revision of the Australian Mesochorinae.
- Mike Fitton recently visited Sweden in connection with the Linnaean bi-centenary celebrations. Mike read a paper about the scientific importance of the Linnaean insect collection. He plans to return to Sweden later this summer to study the Thomson collection.
- Ian Gauld's revision of African Ophioninae is in proof stage and will appear later this year. It is being published by the Commonwealth Agricultural Bureaux as the first in a new series of taxonomic monographs.

The self the contribution of

REQUESTS FOR MATERIAL, INFORMATION ETC.

Mattias Idar is interested to see any material of Synomelix, Synodites and Pantorhaestes from Europe. In return Mattias would be glad to lend specimens for revisionary purposes from his own collection. The subfamilies Campopleginae, Mesochorinae and Microleptinae are the best represented (besides of course the Ctenopelmatinae!)

Ian Gauld would like any information about the location of collections of Ophioninae from S.E. Asia, particularly from the 'Indo-China' peninsula and eastern Indonesia. Ian would be pleased to try to determine specimens of Ophioninae from the Indo-Australian region.

STYLES OF SPECIES DESCRIPTIONS

by M.E.M. Idar

Inspired by some editorial objections against my own lengthy species descriptions, I would like to have a discussion in <u>Ichnews</u> about this matter. Here are some questions to start with -

How complete should a description of a new species be?

Should it include characters of less diagnostic value, with regard to the known related species because of too strong variability or because they look the same in this character?

Should a description be based on one specimen, a fixed set of specimens or all the available material of what the author thinks is one species?

Would it be advisable to make a brief description to validate the name by fulfilling the criteria of "effective publication" and, for example circulate stencilled copies of a fuller description to one's colleagues?

What need is there for redescriptions?

(Please send us your views on the points Mattias has raised. We will attempt to include a cross section of opinions in future issues. Eds.)

A NEW CHARACTER IN THE HINDWING OF ICHNEUMONIDAE

by P.M. Sanborne

While studying Sinophorus I noticed some large bristles in the mediellan cell of the hindwing. They tend to be larger and structurally different from normal wing setae. In addition to these there are often 2-4 similar bristles on the mediellan vein. These bristles are unfortunately difficult to see unless a slide is prepared of the hind wing.

I have looked for the character in other subfamilies (except Adelognathinae, Agriotypinae and Phrudinae) and found it present in all but the Ophioninae. The search was done primarily on a tribal basis on my own collection of Nearctic material. I found the character absent in some tribes; in others it was absent in some genera. Occasionally when I found it present in a genus it was absent in one or more species.

The bristles have a distinctive arrangement in the Cremastinae and Anomaloninae where they occupy positions on the subcostella on the portion nearest the point of attachment to the thorax rather than in the mediellan cell. Other subfamilies have a more or less uniform arrangement within the mediellan cell but there are differences in the number of bristles present and their spacing and position within the cell. It is possible that this character may be of some value in determining intergroup relationships but it should also prove to be of taxonomic use at generic level and below. It is my intention to look at this problem more closely. I would welcome suggestions and comments from readers.

DIRECTORY

Change of address

Shaw, M.R. Department of Zoology, University of Reading, Whiteknights, Reading, RG6 2AJ, ENGLAND.

Additions

- Brussino, Gianfranco. Ministere Agricoltura e Foreste, Osservatorie malattie piante, Via S. Secondo, 39. 10128 Torino, ITALY.

 (Biology of species parasitic on insects of economic importance; inflorescence visitings; Ichneumonidae of the Occidental Alpine arch).
- Cederberg, Björn. Department of Entomology, University of Uppsala, Box 561, S-751 22 Uppsala, SWEDEN. (Western Palaearctic Ophioninae).
- Chao, Hsiu-fu. College of Agriculture of Fukien, Fukien, PEOPLE'S REPUBLIC OF CHINA.
- Oosterbroek, Pjotr. Instituut voor Taxonomische Zoölogie, Zoologisch Museum Universitet van Amsterdam, Plantage Middenlaan 64, Amsterdam, THE NETHERLANDS. (Western Palaearctic Ophioninae).
- Sanborne, P. Mike. Lakehead University, Thunder Bay, Ontario P7B 5El, CANADA: (Nearctic Sinophorus; wing bristles of Ichneumons).

RECENT PUBLICATIONS

- AESCHLIMANN, J.-P. 1978. Recents developpements en lutte biologique l'exemple de la tordeuse grise du meleze. (1). Ann. Soc. Hort. Hist. Nat. Hérault 118: 17-20.
- AUBERT, J.-F. & SHAUMAR, N. 1978. Supplément aux Ichneumonides d'Egypte. Bull. ent. Soc. Mulhouse Avril-Juin 1978: 13-20.

- BARRON, J.R. 1977. The Nearctic species of Orthopelma. Syst. Ent. 4: 283-299.
 - " & BISDEE, H.E. 1977. The final instars of Leptocampoplex cremastoides and Macrus parvulus and notes on the classification of the genera. Ann. ent. Soc. Am. 70: 290-292.
 - enchophorus and notes on the classification of the genus. Ann. ent. Soc. Am. 70: 48-50.
 - " " Adults and larvae of a new species of Gelis parasitizing eggs of Schizocosa saltatrix (Araneida, Lycosidae).

 J. New York ent. Soc. 85: 43-48.
- BOGENSCHÜTZ, H. 1978. Fühler-Bewegungen wirtesuchender Weibcher der Schlupfwespe Coccygomimus turionellae. Ent. Germ. 4: 122-132.
- CHANDRA, G. & GUPTA, V.K. 1977. Ichneumonologia Orientalis. VII. The tribes Lissonotini and Banchini. Oriental Insects Mon. 7: 1-290.
- CHAO, H.-F. 1976. An outline on the classification of the ichneumon-flies of China. Science Press, Peking. 413pp.
- CONSTANTINEANU, M.I. & PISICA, C. 1977. Hymenoptera: Ichneumonidae Ephialtinae-Acaenitinae. Fauna Rep. Socialiste Romania 9(7): 1-296.
- DANTHANARAYANA, W., FARRUGIA, D. & GAULD, I.D. 1977. Studies on the biology and systematic position of a new species of ichneumonid parasiting the light brown apple moth, Epiphyas postvittana (Walker) (Lep: Tortricidae), in Australia. Bull. ent. Res. 67: 607-617.
- DILLER, E.H. 1977. Die in Indien vorkommenden Taxa der Gattung Diplazon. Mitt. Münch. Ent. Ges. 66: 21-28.
- FINLAYSON, T. & HAGEN, K.S. 1977. Final-Instar larvae of Parasitic Hymenoptera. Pest Management Papers 10: 1-111. Simon Fraser University.
- GAULD, I.D. 1977. A revision of the Ophioninae of Australia. Austr. J. Zool. Suppl. Ser. 49: 1-112.
- " & MITCHELL, P.A. 1977. Ichneumonidae: Orthopelmatinae & Anomaloninae. Handbk. Ident. Br. Insects VII, 2(b): 1-32.
- GRAF, V.. 1976. Pectinorex, novo gênero de Joppini do sul do Brasil.

 Dusenia 9: 71-75.
- GUPTA, V.K. 1977. Host selection in Indian Ichneumonidae in relation to ecology of host. In <u>Insects and Host Specificity</u> (T.N. Ananthakrishnan Ed.) pp.91-95. Delhi.
 - " & MAHESHWARY, S. 1977. Ichneumonologia Orientalis IV. The tribe Porizontini. Oriental Insects Mon. 5: 1-267.
 - " & TIKAR, D.T. 1977. Ichneumonologia Orientalis I. The tribe Pimplini. Oriental Insects Mon. 1: 1-312.

. 3 . 7

- HINZ, R. 1977. Uber einige Arten der Gattung <u>Dusona</u>. <u>Nachr. Bayer</u>. <u>Ent.</u> 26: 47-54.
- " Eine neue Art der Gattung <u>Dusona</u>. <u>Nachr. Bayer</u>. Ent. 26: 109-110.
- HOWELL, J. & FISHER, R.C. 1977. Food conversion efficiency of a parasitic wasp. Nemeritis canescens. Ecological Ent. 2: 143-151.
- KASPARYAN, D.R. 1976. New species of the tribe Tryphonini from the southern part of the Far East [province]. Trudy biol.-pochven. Inst. (n.s.) 43(146): 107-120.
 - " 1976. The new species of Ichneumonids of the subfamily Campopleginae from the Eastern Palaearctic. Trudy zool. Inst. Akad. Nauk SSSR 44: 68-75.
 - " 1977. Ichneumonids of the subfamilies Pimplinae and Tryphoninae new for Mongolia and Transbaikalia. Insects of Mongolia 5: 456-469.
- " 1977. Review of the European species of Ichneumonids of the genus Delomerista. New and little known species of Insects of the European part of the U.S.S.R. Academy of Sciences, All-Union Entomological Society. 69-75.
 - KUSLITZKY, W.S. 1977. New and little known species of the tribe Glyptini in the European part of the U.S.S.R. New and little known species of Insects of the European part of the U.S.S.R. Academy of Sciences, All-Union Entomological Society, 75-82.
 - LECLANT, F., IDRISSI, A.E. & MIERMONT, Y. 1977. Les Aphides et les Tortricides du Cèdre au Maroc et en France et leurs parasites. C.R. Acad. Sc. Paris 284 (Series D): 647-649.
 - MOMOI. S. 1977. Ichneumonid parasites of Pine Tip borers in Japan with descriptions of new species. Akitu 12: 1-13.
 - PORTER, C.C. 1977. Prosthoporus, a new Bolivian genus of the subtribe Lymeonina. Psyche 83: 271-276.
 - New records of Thyreodon from south Texas. Psyche 83: 304-309.
 - " " Ecology, Zoogeography and Taxonomy of lower Rio Grand Valley Mesosterines. Psyche 84: 28-91.
 - PRESS, J.W., FLAHERTY, B.R. & ARBOGAST, R.T. 1977. Interactions among Nemeritis canescens, Bracon hebetor and Ephestia cautella.

 J. Kansas ent. Soc. 50: 259-262.
 - RASKE, A.G. 1978. Parasites of Birch Casebearer larvae in Newfoundland. Entomophaga 23: 103-108.

T FF

- SALT, G. 1977. Problems of orientation associated with cocoon-spinning by Nemeritis. Ecological Ent. 2: 171-177.
 - SAWONIEWICZ, J. 1978. Zur Systematik und Faunistik der Ichneumonidae.

 Annls zool. Warsz. 34: 121-137.
 - SHAW, M.R. 1977. On the distribution of some satyrid (Lep.) larvae at a coastal site in relation to the ichneumonid parasite. Entomologist's Gaz. 28: 133-134.
 - of Benjaminia. Nouv. Rev. Ent. 7: 85-89.
 - SHORT, J.R.T. 1977. A description and classification of some final instars of the Theriini (Anomalinae). Proc. ent. Soc. Wash. 79: 463-467.
 - SIITAN, U.V. 1977. Review of Ichneumonidae of the tribe Phaeogenini of of the European part of the U.S.S.R. Ent. Obzr. 56: 843-853.
 - SLOBODCHIKOFF, C.N. 1977. Patterns of variation in wasps of the genus Therion. Univ. Calif. Publ. (Ent.) 82: 1-65.
 - TOWNES, H. & TOWNES, M. 1978. Ichneumon-flies of America north of Mexico: 7. Subfamily Banchinae, tribes Lissonotini and Banchini. Mem. Am. ent. Inst. 26: 1-614.
 - WALLEY, G.S. & BARRON, J.R. 1977. Nearctic Glypta parasites of conifer feeding Oethreutidae (Lepidoptera). Ent. News 88: 177-184.