
Prediction of AGB using RFR, SVR, FF-NN


Ritwika Mukhopadhyay

Airborne laser scanning

The scanning over an area creates a 3d point cloud




(Remote Sensing course, SLU)






(Remote Sensing course, SLU)





(Remote Sensing course, SLU)

Study Area



RFR FF-NN





glm(TotalAGB ~ crr + crrP80, family = Gamma(link = "sqrt"))

Coefficients:

Estimate Pr(>|t|)

(Intercept) 8.93286 < 2e-16 ***

crr -14.33267 6.63e-11 ***

crrP80 1.12565 < 2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

RMSE <- 63.48

R^2 <- 0.45

Mukhopadhyay et al. (in prep.)

References:


R. Mukhopadhyay, I. Indirabai, M. Nilsson, M. Egberth, E. Holmström, and M. Ekström, "Modelling of Aboveground Biomass Change Using LiDAR Metrics and NFI Field Data: A Case Study of Southern Sweden," in *SilviLaser Conference 2021*, pp. 205–209, doi: 10.34726/WIM.1982.

R. Mukhopadhyay, H. J. Persson, E. Lindberg, S. Saarela, M. Ekström, and M. Nilsson, "Estimation of Prediction Uncertainty for Forest Aboveground Biomass using Gamma Regression Analysis: A Case Study in South-Central Sweden" (in prep.).











Thank you!!