

CHE 565 -- Homework 3

Soki Sem

April 18, 2026

Problem 1

Steepest Descent

Function :

$$f(x_1, x_2) = 0.5x_1^2 + 1.0x_1x_2 - x_1 + 1.0x_2^2 - x_2$$

At $x_0 = \text{Matrix}([1, 1])$, $f(x_0) = 0.5000000000000000$

Gradient of f:

$$\nabla f = \begin{bmatrix} 1.0x_1 + 1.0x_2 - 1.0 \\ 1.0x_1 + 2.0x_2 - 1 \end{bmatrix}$$

At $x_0 = \text{Matrix}([1, 1])$, $Df(x_0) = 1.00, 2.00$

$$\begin{bmatrix} g_1(\alpha) \\ g_2(\alpha) \end{bmatrix} = \begin{bmatrix} 1 - 1.0\alpha \\ 1 - 2.0\alpha \end{bmatrix}$$

Function ϕ_k :

$$\phi(\alpha) = 6.5\alpha^2 - 5.0\alpha + 0.5$$

Derivative of ϕ_k :

$$\begin{aligned} \frac{d}{d\alpha}\phi(\alpha) &= 13.0\alpha - 5.0 \\ 0 &= 13.0\alpha - 5.0 \end{aligned}$$

Optimal alpha: 0.38

Next iterate x_1 : 0.62, x_2 : 0.23

Function value at next iterate: -0.46

Problem 2

Conjugate Gradient Method

$$\begin{bmatrix} s_1(\alpha) \\ s_2(\alpha) \end{bmatrix} = \begin{bmatrix} 0.0946745562130178 \\ -0.195266272189349 \end{bmatrix}$$

Problem 3

Constituents	Maximum Quantity (bbl/day)	Production Cost (\$/bbl)
Crude Oil	1000	50
Natural Gas	500	30
Refined Products	2000	70