

# CHE 565 -- Homework 2

Soki Sem

April 1, 2026

## Problem 1

### Setup

$$\begin{aligned}\text{Option A: } C_{0,A} &= \$3,800,000 \\ FV_A &= \$1,100,000/\text{yr} \\ \text{Option B: } C_{0,B} &= \$5,000,000 \\ FV_B &= \$1,410,000/\text{yr}\end{aligned}$$

PV = present value, FV = future value (annual cash flow), r = yearly interest rate, n = number of years  
n = 10 years, r = 0.10

### A.) Compute Net Present Value for Options A and B

$$\text{Present Value formula: } PV = \frac{F \cdot (r + 1)^{-n} \cdot ((r + 1)^n - 1)}{r}$$

$$\text{Net Present Value formula: } NPV = C_0 + \frac{F \cdot (r + 1)^{-n} \cdot ((r + 1)^n - 1)}{r}$$

$$\text{Annual payment formula: } P = \frac{C_0 \cdot r \cdot (r + 1)^n}{(r + 1)^n - 1}$$

$$\text{Annual payment for option A: } P_A = \$ - 492,117.38/\text{year}$$

$$\text{Annual payment for option B: } P_B = \$ - 647,522.87/\text{year}$$

$$NPV_A = \$2,959,023.82, \quad NPV_B = \$3,663,839.62$$

NPV is higher for option B at 10% yearly interest, so B is preferred under these assumptions.

### B.) Annual Payment for 10 Year Lifetime, No Salvage Value, 5% Interest Rate

P = annual payment, C0 = initial cost, r = yearly interest rate, n = number of years

$$P(A) = \$492,117.38/\text{year}, \quad P(B) = \$647,522.87/\text{year}$$

## Problem 2

### Setup

\$0.94/lbs is the price of powdered detergent.

0.1 % of 4,000,000 lbs of detergent per year is carried through the exhaust

Adding a second cyclone separator costs \$10,000, with \$300/year in maintenance costs will reduce carryover to 0.0%, with a 8% yearly interest rate.

Cost of carryover detergent: \$3,760.00/year

Cost of second separator: \$10,000.00 initial + \$600.00/year maintenance = \$10,600.00 total

Lost detergent value: \$3,760.00/year

### A.) Find the additional yearly income

Gross additional income (recovered detergent): \$3,760.00/year

Net additional income after maintenance: \$3,460.00/year

### B.) Find the payback period

The NPV equation for the payback period is:  $-10000 + 43250.0 \cdot 1.08^{-n} \cdot (1.08^n - 1)$

Discounted payback period (NPV = 0): 3.42 years

## Problem 3

### Setup

Determine if each function is convex

$$A : f(x) = (x_1 - x_2)^2 + x_2^2$$

$$B : f(x) = x_1^2 + x_2^2 + x_3^2$$

$$C : f(x) = e^{x_1} + e^{x_2}$$

### Results

| Function                | Hessian                                                             | Eigenvalues                                                  | Convex? | Reason                       |
|-------------------------|---------------------------------------------------------------------|--------------------------------------------------------------|---------|------------------------------|
| $x_2^2 + (x_1 - x_2)^2$ | $\begin{bmatrix} 2 & -2 \\ -2 & 4 \end{bmatrix}$                    | $\begin{bmatrix} 3 - \sqrt{5} \\ \sqrt{5} + 3 \end{bmatrix}$ | Yes     | Eigenvalues are non-negative |
| $x_1^2 + x_2^2 + x_3^2$ | $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ | $[2]$                                                        | Yes     | Eigenvalues are positive     |
| $e^{x_1} + e^{x_2}$     | $\begin{bmatrix} e^{x_1} & 0 \\ 0 & e^{x_2} \end{bmatrix}$          | $\begin{bmatrix} e^{x_1} \\ e^{x_2} \end{bmatrix}$           | Yes     | Eigenvalues are positive     |

## Problem 4

Is the region defined by the constraints convex?

The feasible region is defined by the following constraints:  $x_2 \geq 0 \wedge x_2 \geq 1 - x_1 \wedge x_1 \leq 2 \wedge x_2 \leq \frac{x_1}{2} + 1$




Figure 1: Feasible Region

The feasible region is convex because it is defined by a set of linear inequalities, which form a convex set.

## Problem 5

### Setup

Maximize the objective function:

$$f(x) = -\frac{x^6}{6} + \frac{4x^5}{5} - \frac{x^4}{4} - \frac{10x^3}{3} + 2x^2 + 8x + 1$$

Given that:

$$\frac{df}{dx} = (2 - x)^3 (x + 1)^2$$

$$\frac{d^2 f}{dx^2} = (2-x)^3 (2x+2) - 3(2-x)^2 (x+1)^2$$

### a.) Analytical Solution

Expanded form:

$$\frac{df}{dx} = -x^5 + 4x^4 - x^3 - 10x^2 + 4x + 8$$

$$\frac{d^2 f}{dx^2} = -5x^4 + 16x^3 - 3x^2 - 20x + 4$$

The roots to df/dx are  $x^* = -1, 2$

The corresponding function values are  $f(x^*) = -2.88, 9.27$

The Hessian evaluated at the critical points are  $\frac{d^2 f}{dx^2}|_{x^*} = 0, 0$ , inconclusive 2nd derivative test

### b.) Excel Solution

The Excel data was used to perform root-finding using the Newton-Raphson method:

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)}$$

| Iteration | x_n    | f(x_n)    | f'(x_n)   | f''(x_n)   | x_n+1  |
|-----------|--------|-----------|-----------|------------|--------|
| 0.0000    | 5.0000 | -586.0833 | -972.0000 | -1296.0000 | 4.2500 |
| 1.0000    | 4.2500 | -139.2208 | -313.9541 | -538.2070  | 3.6667 |
| 2.0000    | 3.6667 | -27.0988  | -100.8230 | -224.6914  | 3.2179 |
| 3.0000    | 3.2179 | 0.5555    | -32.1432  | -94.4150   | 2.8775 |
| 4.0000    | 2.8775 | 7.2322    | -10.1590  | -39.9713   | 2.6233 |
| 5.0000    | 2.6233 | 8.8043    | -3.1799   | -17.0590   | 2.4369 |
| 6.0000    | 2.4369 | 9.1644    | -0.9854   | -7.3392    | 2.3027 |
| 7.0000    | 2.3027 | 9.2446    | -0.3025   | -3.1810    | 2.2076 |
| 8.0000    | 2.2076 | 9.2620    | -0.0920   | -1.3876    | 2.1413 |
| 9.0000    | 2.1413 | 9.2657    | -0.0278   | -0.6084    | 2.0955 |
| 10.0000   | 2.0955 | 9.2665    | -0.0084   | -0.2678    | 2.0643 |
| 11.0000   | 2.0643 | 9.2666    | -0.0025   | -0.1182    | 2.0432 |
| 12.0000   | 2.0432 | 9.2667    | -0.0007   | -0.0523    | 2.0289 |
| 13.0000   | 2.0289 | 9.2667    | -0.0002   | -0.0232    | 2.0193 |
| 14.0000   | 2.0193 | 9.2667    | -0.0001   | -0.0103    | 2.0129 |
| 15.0000   | 2.0129 | 9.2667    | -0.0000   | -0.0046    | 2.0086 |
| 16.0000   | 2.0086 | 9.2667    | -0.0000   | -0.0020    | 2.0058 |
| 17.0000   | 2.0058 | 9.2667    | -0.0000   | -0.0009    | 2.0038 |
| 18.0000   | 2.0038 | 9.2667    | -0.0000   | -0.0004    | 2.0026 |
| 19.0000   | 2.0026 | 9.2667    | -0.0000   | -0.0002    | 2.0017 |
| 20.0000   | 2.0017 | 9.2667    | -0.0000   | -0.0001    | 2.0011 |

Table 1: Optimization Results from Excel starting at  $x_0=5$

| Iteration | x_n     | f(x_n)     | f'(x_n)   | f''(x_n)   | x_n+1   |
|-----------|---------|------------|-----------|------------|---------|
| 0.0000    | -5.0000 | -4832.7500 | 5488.0000 | -5096.0000 | -3.9231 |
| 1.0000    | -3.9231 | -1208.5512 | 1775.5079 | -2114.1044 | -3.0832 |
| 2.0000    | -3.0832 | -295.6352  | 570.0315  | -883.6738  | -2.4382 |
| 3.0000    | -2.4382 | -71.0801   | 180.8132  | -373.6709  | -1.9543 |
| 4.0000    | -1.9543 | -17.8527   | 56.3066   | -160.7261  | -1.6040 |
| 5.0000    | -1.6040 | -5.9169    | 17.0747   | -70.7559   | -1.3626 |
| 6.0000    | -1.3626 | -3.4409    | 5.0003    | -32.0380   | -1.2066 |
| 7.0000    | -1.2066 | -2.9756    | 1.4068    | -14.9373   | -1.1124 |
| 8.0000    | -1.1124 | -2.8972    | 0.3808    | -7.1437    | -1.0591 |
| 9.0000    | -1.0591 | -2.8853    | 0.0999    | -3.4805    | -1.0304 |
| 10.0000   | -1.0304 | -2.8836    | 0.0257    | -1.7158    | -1.0154 |
| 11.0000   | -1.0154 | -2.8834    | 0.0065    | -0.8515    | -1.0078 |
| 12.0000   | -1.0078 | -2.8833    | 0.0016    | -0.4242    | -1.0039 |
| 13.0000   | -1.0039 | -2.8833    | 0.0004    | -0.2117    | -1.0020 |
| 14.0000   | -1.0020 | -2.8833    | 0.0001    | -0.1057    | -1.0010 |
| 15.0000   | -1.0010 | -2.8833    | 0.0000    | -0.0528    | -1.0005 |
| 16.0000   | -1.0005 | -2.8833    | 0.0000    | -0.0264    | -1.0002 |
| 17.0000   | -1.0002 | -2.8833    | 0.0000    | -0.0132    | -1.0001 |
| 18.0000   | -1.0001 | -2.8833    | 0.0000    | -0.0066    | -1.0001 |
| 19.0000   | -1.0001 | -2.8833    | 0.0000    | -0.0033    | -1.0000 |
| 20.0000   | -1.0000 | -2.8833    | 0.0000    | -0.0017    | -1.0000 |

Table 2: Optimization Results from Excel starting at x0=-5

| x0      | x*      | f(x*)   |
|---------|---------|---------|
| 5.0000  | 2.0011  | 9.2667  |
| -5.0000 | -1.0000 | -2.8833 |

Table 3: Summary of Optimization Results from Excel

### c.) Numerical Solution Using Nelder-Mead Simplex algorithm which is the same as fminsearch in matlab

Using Nelder-Mead Simplex algorithm with initial guesses x0 = 5:

One of the roots of df/dx are found to be  $x^* = [1.99987793]$

The corresponding optimum function values are  $f(x^*) = 9.26666666666667$

Actual maximum of the function:

$$\begin{aligned}
 x^* &= [1.99987793] \\
 f(x^*) &= -9.26666666666667 \\
 \text{iterations} &= 20 \\
 \text{function calls} &= 41
 \end{aligned}$$

**d.) Numerical Solution Using Newton Conjugate Gradient Method which is the same as fmincon in matlab except required to give jacobian and hessian information**

Using the Newton Conjugate Gradient method with initial guesses  $x_0 = 5$ :

One of the roots of  $df/dx$  are found to be  $x^* = [2.00384098]$

The corresponding optimum function values are  $f(x^*) = -9.266666666175944$

$$\begin{aligned}x^* &= [2.00384098] \\f(x^*) &= -9.266666666175944 \\ \text{iterations} &= 19 \\ \text{function calls} &= 19\end{aligned}$$

Using the Newton Conjugate Gradient method with initial guesses  $x_0 = -5$ :

One of the roots of  $df/dx$  are found to be  $x^* = [-1.00006112]$

The corresponding optimum function values are  $f(x^*) = 2.883333333335388$

$$\begin{aligned}x^* &= [-1.00006112] \\f(x^*) &= 2.883333333335388 \\ \text{iterations} &= 20 \\ \text{function calls} &= 20\end{aligned}$$

| Method      | Initial Guess | $x^*$         | $f(x^*)$ | Iterations | Function Calls |
|-------------|---------------|---------------|----------|------------|----------------|
| Nelder-Mead | 5             | [1.99987793]  | -9.2667  | 20         | 41             |
| Excel       | 5             | 2.0011        | 9.2667   | 21         | -              |
| Excel       | -5            | -1.0000       | -2.8833  | 21         | -              |
| Newton-CG   | 5             | [2.00384098]  | -9.2667  | 19         | 19             |
| Newton-CG   | -5            | [-1.00006112] | 2.8833   | 20         | 20             |