March 31, 2023

## An Introduction to Climate Modeling

Milestone 1 - Geography and Visualization

## 1 Read-in and Plot Geography

In this milestone, the goal is to read-in the geography information of planet Earth and plot it as a two-dimensional world map. We consider an equirectangular grid of the Earth, i.e., an equidistant rectangular grid in spherical coordinates, where the grid point (i,j) has the spherical coordinates  $(\varphi_i,\theta_j)$ , where  $\varphi_i$  is the longitude between  $-90^\circ$  south and  $90^\circ$  north (including the poles) and  $\theta_j$  the latitude between  $-180^\circ$  west and  $180^\circ$  east. The basis for this is the input file The\_World128x65.dat, which describes the distribution of the different Earth surface types. This file contains a matrix  $G \in \mathbb{N}^{65 \times 128}$  with entries  $g_{ij} \in \{1,2,3,5\}$ , where the entry  $g_{ij}$  stores the Earth surface type at grid point (i,j). Here, 1 represents the Earth surface type land, 2 represents sea ice, 3 represents snow, and 5 represents ocean. The grid resolution in longitude and latitude direction is  $2.8125^\circ$ . The basis for this distribution and grid is from Zhuang et al. You can proceed as follows:

- 1. Write a function read\_geography, which reads the file The\_World128x65.dat from the folder input and outputs a matrix  $T \in \mathbb{N}^{65 \times 128}$  with the classification of the earth surface types.
- 2. Write a function robinson\_projection, which maps an equirectangular grid in spherical coordinates to the plane. For simplicity use the approximate formula by Beineke for the Robinson projection,

$$\begin{split} x\left(\varphi,\theta\right) &= \frac{\varphi}{\pi} \left( 0.0379 \, \theta^6 - 0.15 \, \theta^4 - 0.367 \, \theta^2 + 2.666 \right), \\ y\left(\varphi,\theta\right) &= 0.96047 \, \theta - 0.00857 \, \mathrm{sign} \left(\theta\right) \left|\theta\right|^{6.41}, \end{split}$$

where  $\varphi$  is the longitude and  $\theta$  the latitude in radians. This function should return two matrices  $X=x_{ij}$  and  $Y=y_{ij}$ , where  $x_{ij}=x(\varphi_i,\theta_j), y_{ij}=y(\varphi_i,\theta_j)$ .




Figure 1: Robinson projection of the world <sup>2</sup>

- 3. Write a function plot\_geo that creates a plot of the Earth surface type  $g_{ij}$  against the mapped coordinates  $(x_{ij}, y_{ij})$ .
- 4. Use these functions in a program and run it to check your results.

<sup>&</sup>lt;sup>1</sup>K. Zhuang, G.R. North, M.J. Stevens, A NetCDF version of the two-dimensional energy balance model based on the full multigrid algorithm, SoftwareX, Vol. 6, pp. 198-202, July 7, 2017.

<sup>&</sup>lt;sup>2</sup>Daniel R. Strebe, https://en.wikipedia.org/wiki/File:Robinson\_projection\_SW.jpg

## **2 Control Solutions**




Figure 2: Robinson projection output