

# Chapter 2.9: Uniformity in the underlying distribution

Amine Hadji

University of Leiden

March 2, 2019



Universiteit Leiden

# Previously in...

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

... Reading group - Mathematical statistics

# Previously in...

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

... Reading group - Mathematical statistics

- In Chapter 2.5, we have seen
  - the conditions on  $\mathcal{F}$  so that it is  $P$ -Glivenko-Cantelli
  - the conditions on  $\mathcal{F}$  so that it is  $P$ -Donsker

# Previously in...

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

... Reading group - Mathematical statistics

- In Chapter 2.5, we have seen
  - the conditions on  $\mathcal{F}$  so that it is  $P$ -Glivenko-Cantelli
  - the conditions on  $\mathcal{F}$  so that it is  $P$ -Donsker
- In Chapter 2.3, we have seen
  - the symmetrization lemma
  - Hoeffding's inequality

# Overview

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- 1 Notations & Intentions
- 2 Glivenko-Cantelli Theorem
- 3 Donsker Theorem
- 4 Central Limite Theorem (under Sequences)

# Intentions

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

Chapter 2.8 is mostly about:

# Intentions

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

Chapter 2.8 is mostly about:

- ① finding conditions on  $\mathcal{F}$  so that it is uniformly Glivenko-Cantelli

# Intentions

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

Chapter 2.8 is mostly about:

- ① finding conditions on  $\mathcal{F}$  so that it is uniformly Glivenko-Cantelli
- ② finding conditions on  $\mathcal{F}$  so that it is uniformly Donsker



# Intentions

## Chapter 2.8

Amine Hadji

### Notations & Intentions

Glivenko-Cantelli Theorem

Donsker Theorem

Central Limit Theorem (Under Sequences)

Chapter 2.8 is mostly about:

- 1 finding conditions on  $\mathcal{F}$  so that it is uniformly Glivenko-Cantelli
- 2 finding conditions on  $\mathcal{F}$  so that it is uniformly Donsker
- 3 applying these results to a (hopefully) more concrete case

# Notations

## Chapter 2.8

Amine Hadji

### Notations & Intentions

Glivenko-Cantelli Theorem

Donsker Theorem

Central Limit Theorem (Under Sequences)

- $\mathcal{F}$  is a class of  $P$ -measurable functions on a measurable space (for all  $P \in \mathcal{P}$ )
- $F$  is a measurable envelope function of  $\mathcal{F}$
- $\mathcal{Q}_n$  is the set of all discrete probability measures with atom sized  $k/n$
- $\mathbb{P}_n := 1/n \sum_{i=1}^n \delta_{X_i}$  ( $\mathbb{P}_n \in \mathcal{Q}_n$ )
- $\mathbb{G}_{n,P} := \sqrt{n}(\mathbb{P}_n - P)$  and  $\mathbb{G}_P$  Brownian bridge

# Glivenko-Cantelli

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

First, little reminder:

**What is a Glivenko-Cantelli class of functions?**

# Glivenko-Cantelli

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

First, little reminder:

**What is a Glivenko-Cantelli class of functions?**

## Definition

We say that  $\mathcal{F}$  is  $P$ -Glivenko Cantelli iff

$\limsup_{n \rightarrow +\infty} \|\mathbb{P}_n - P\|_{\mathcal{F}}^* = 0$  almost surely.

# Glivenko-Cantelli

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

First, little reminder:

**What is a Glivenko-Cantelli class of functions?**

### Definition

We say that  $\mathcal{F}$  is  $P$ -Glivenko Cantelli iff

$$\limsup_{n \rightarrow +\infty} \|\mathbb{P}_n - P\|_{\mathcal{F}}^* = 0 \text{ almost surely.}$$

### Definition

We say that  $\mathcal{F}$  is Glivenko Cantelli uniformly in  $P \in \mathcal{P}$  iff

$$\limsup_{n \rightarrow +\infty} \|\mathbb{P}_n - P\|_{\mathcal{F}}^* = 0 \text{ almost surely for all } P \in \mathcal{P}$$

# Glivenko-Cantelli

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

First, little reminder:

**What is a Glivenko-Cantelli class of functions?**

## Definition

We say that  $\mathcal{F}$  is  $P$ -Glivenko Cantelli iff

$$\limsup_{n \rightarrow +\infty} \|\mathbb{P}_n - P\|_{\mathcal{F}}^* = 0 \text{ almost surely.}$$

## Definition

We say that  $\mathcal{F}$  is Glivenko Cantelli uniformly in  $P \in \mathcal{P}$  iff

$$\limsup_{n \rightarrow +\infty} \|\mathbb{P}_n - P\|_{\mathcal{F}}^* = 0 \text{ almost surely for all } P \in \mathcal{P}$$
$$\sup_{P \in \mathcal{P}} P_P^*(\limsup_{n \rightarrow +\infty} \|\mathbb{P}_n - P\|_{\mathcal{F}}^* = 0).$$

# Glivenko-Cantelli Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

### Theorem

*Let  $\mathcal{F}$  such that:*

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$

# Glivenko-Cantelli Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

*Let  $\mathcal{F}$  such that:*

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$
- $\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n), \forall \epsilon > 0$



# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

*Let  $\mathcal{F}$  such that:*

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$
- $\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n), \forall \epsilon > 0$

*then  $\mathcal{F}$  is Glivenko-Cantelli uniformly in  $P \in \mathcal{P}$ .*

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

Since showing almost sure convergence directly is almost impossible, we have to find another way

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

Since showing almost sure convergence directly is almost impossible, we have to find another way

- $X_n \rightarrow 0$  a.s is equivalent to  $\sup_{m \geq n} |X_m| \rightarrow 0$  in probability

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

Since showing almost sure convergence directly is almost impossible, we have to find another way

- $X_n \rightarrow 0$  a.s is equivalent to  $\sup_{m \geq n} |X_m| \rightarrow 0$  in probability
- Therefore  $\mathcal{F}$  is uniformly Glivenko-Cantelli in  $P \in \mathcal{P}$   
 $\sup_{P \in \mathcal{P}} P_P^*(\sup_{m \geq n} \|\mathbb{P}_n - P\|_{\mathcal{F}} > \epsilon) \rightarrow 0$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji


Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



① Let  $\mathcal{F}_M := \{f \mathbf{1}_{f \leq M} : f \in \mathcal{F}\}$  for fixed  $M > 0$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji


Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- 1 Let  $\mathcal{F}_M := \{f \mathbf{1}_{f \leq M} : f \in \mathcal{F}\}$  for fixed  $M > 0$
- 2  $\|\mathbb{P}_n - P\|_{\mathcal{F}} \leq \|\mathbb{P}_n - P\|_{\mathcal{F}_M} + \mathbb{P}_n F \mathbf{1}_{F > M} + P F \mathbf{1}_{F > M}$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji


Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- 1 Let  $\mathcal{F}_M := \{f \mathbf{1}_{f \leq M} : f \in \mathcal{F}\}$  for fixed  $M > 0$
- 2  $\|\mathbb{P}_n - P\|_{\mathcal{F}} \leq \|\mathbb{P}_n - P\|_{\mathcal{F}_M} + \mathbb{P}_n F \mathbf{1}_{F > M} + P F \mathbf{1}_{F > M}$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions


Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$





# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions


Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- $\mathbb{P}_n F \mathbf{1}_{F > M} \rightarrow P F \mathbf{1}_{F > M}$  almost surely uniformly

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions


Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- $\mathbb{P}_n F \mathbf{1}_{F > M} \rightarrow P F \mathbf{1}_{F > M}$  almost surely uniformly
- $\lim_{M \rightarrow +\infty} P F \mathbf{1}_{F > M}$  uniformly

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji


Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- 1 Let  $\mathcal{F}_M := \{f \mathbf{1}_{f \leq M} : f \in \mathcal{F}\}$  for fixed  $M > 0$
- 2  $\|\mathbb{P}_n - P\|_{\mathcal{F}} \leq \|\mathbb{P}_n - P\|_{\mathcal{F}_M} + \mathbb{P}_n F \mathbf{1}_{F > M} + P F \mathbf{1}_{F > M}$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

① Let  $\mathcal{F}_{nX}$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ① Let  $\mathcal{F}_n$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$
- ②  $N(\eta M, \mathcal{F}_M, L_1(\mathbb{P}_n)) \leq N_n(\eta)$  with  $\log N_n(\eta) = o(n)$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ① Let  $\mathcal{F}_{nX}$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$
- ②  $N(\eta M, \mathcal{F}_M, L_1(\mathbb{P}_n)) \leq N_n(\eta)$  with  $\log N_n(\eta) = o(n)$
- ③  $\|\mathbb{P}_n^o\|_{\mathcal{F}_M} \leq \|\mathbb{P}_n^o\|_{\mathcal{F}_{nX}} + \eta M$

# Flashback

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

**Glivenko-  
Cantelli  
Theorem**

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)





# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Lemma (Symmetrization)

*For every nondecreasing, convex  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  and  $\mathcal{F}$*

$$E^* \Phi(\|\mathbb{P}_n - P\|_{\mathcal{F}}) \leq E^* \Phi(2\|\mathbb{P}_n^o\|_{\mathcal{F}}).$$

## Lemma (Hoeffding's inequality)

*Let  $a_1, \dots, a_n$  constants and  $r_1, \dots, r_n$  i.i.d Rademacher. Then*

$$P(|\sum r_i a_i| > x) \leq 2e^{-x^2/(2\|a\|^2)}.$$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ① Let  $\mathcal{F}_{nX}$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$
- ②  $N(\eta M, \mathcal{F}_M, L_1(\mathbb{P}_n)) \leq N_n(\eta)$  with  $\log N_n(\eta) = o(n)$
- ③  $\|\mathbb{P}_n^o\|_{\mathcal{F}_M} \leq \|\mathbb{P}_n^o\|_{\mathcal{F}_{nX}} + \eta M$
- ④  $P_P(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) = E_{P,X} P_R(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) \leq E_{P,X} N_n(\eta) 2e^{-n(\epsilon - \eta M)^2 / (2M^2)} \leq 2 \exp\{-n\epsilon^2 / (4M^2)\}$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ① Let  $\mathcal{F}_{nX}$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$
- ②  $N(\eta M, \mathcal{F}_M, L_1(\mathbb{P}_n)) \leq N_n(\eta)$  with  $\log N_n(\eta) = o(n)$
- ③  $\|\mathbb{P}_n^o\|_{\mathcal{F}_M} \leq \|\mathbb{P}_n^o\|_{\mathcal{F}_{nX}} + \eta M$
- ④  $P_P(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) = E_{P,X} P_R(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) \leq E_{P,X} N_n(\eta) 2e^{-n(\epsilon - \eta M)^2 / (2M^2)} \leq 2 \exp\{-n\epsilon^2 / (4M^2)\}$
- ⑤  $\sum_{m \geq n} P_P(\|\mathbb{P}_m^o\|_{\mathcal{F}_M} > \epsilon) \rightarrow 0$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ① Let  $\mathcal{F}_{nX}$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$
- ②  $N(\eta M, \mathcal{F}_M, L_1(\mathbb{P}_n)) \leq N_n(\eta)$  with  $\log N_n(\eta) = o(n)$
- ③  $\|\mathbb{P}_n^o\|_{\mathcal{F}_M} \leq \|\mathbb{P}_n^o\|_{\mathcal{F}_{nX}} + \eta M$
- ④  $P_P(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) = E_{P,X} P_R(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) \leq E_{P,X} N_n(\eta) 2e^{-n(\epsilon - \eta M)^2 / (2M^2)} \leq 2 \exp\{-n\epsilon^2 / (4M^2)\}$
- ⑤  $\sum_{m \geq n} P_P(\|\mathbb{P}_m^o\|_{\mathcal{F}_M} > \epsilon) \rightarrow 0$
- ⑥  $\sum_{m \geq n} P_P(\|\mathbb{P}_m - P\|_{\mathcal{F}_M} > \epsilon) \rightarrow 0$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ① Let  $\mathcal{F}_{nX}$  a minimal  $\eta M$ -net for  $L_1(\mathbb{P}_n)$
- ②  $N(\eta M, \mathcal{F}_M, L_1(\mathbb{P}_n)) \leq N_n(\eta)$  with  $\log N_n(\eta) = o(n)$
- ③  $\|\mathbb{P}_n^o\|_{\mathcal{F}_M} \leq \|\mathbb{P}_n^o\|_{\mathcal{F}_{nX}} + \eta M$
- ④  $P_P(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) = E_{P,X} P_R(\|\mathbb{P}_n^o\|_{\mathcal{F}_M} > \epsilon) \leq E_{P,X} N_n(\eta) 2e^{-n(\epsilon - \eta M)^2 / (2M^2)} \leq 2 \exp\{-n\epsilon^2 / (4M^2)\}$
- ⑤  $\sum_{m \geq n} P_P(\|\mathbb{P}_m^o\|_{\mathcal{F}_M} > \epsilon) \rightarrow 0$
- ⑥  $\sum_{m \geq n} P_P(\|\mathbb{P}_m - P\|_{\mathcal{F}_M} > \epsilon) \rightarrow 0$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

① Let  $Q \in \mathcal{Q}_n$  and  $|I| = k$ , where  $I := x_i : F(x_i) \leq M$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- 1 Let  $Q \in \mathcal{Q}_n$  and  $|I| = k$ , where  $I := x_i : F(x_i) \leq M$
- 2  $Q_k \in \mathcal{Q}_k$  the discrete measure on  $I$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M}$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$

$$\sup_{Q \in \mathcal{Q}_n} N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq \sup_{n \leq k \leq 2n} \sup_{Q \in \mathcal{Q}_k} N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q))$$

- ❶ Let  $Q \in \mathcal{Q}_n$  and  $|I| = k$ , where  $I := x_i : F(x_i) \leq M$
- ❷  $Q_k \in \mathcal{Q}_k$  the discrete measure on  $I$
- ❸ Then,  $Q|f \mathbf{1}_{F \leq M}| = k/n Q_k|f|$  for any  $f$



# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji


Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- 1 Thus  $N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq N(\epsilon \|F\|_{Q_k}, \mathcal{F}, L_1(Q))$
- 2 Moreover  $\mathcal{Q}_k \subseteq \mathcal{Q}_{2k} \subseteq \dots$

# Glivenko-Cantelli Theorems

Chapter 2.8

Amine Hadji


Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F \mathbf{1}_{F > M} \\ \sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n)$$



- 1 Thus  $N(\epsilon M, \mathcal{F}_M, L_1(Q)) \leq N(\epsilon \|F\|_{Q,k}, \mathcal{F}, L_1(Q))$
- 2 Moreover  $\mathcal{Q}_k \subseteq \mathcal{Q}_{2k} \subseteq \dots$

# Donsker

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

**Donsker  
Theorem**

Central Limit  
Theorem  
(Under  
Sequences)

Again, little reminder:

**What is a Donsker class of functions?**

Again, little reminder:

**What is a Donsker class of functions?**

## Definition

We say that  $\mathcal{F}$  is  $P$ -Donsker iff

$$\mathbb{G}_n \rightsquigarrow \mathbb{G} \text{ in } \ell^\infty(\mathcal{F}).$$

# Donsker

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

Again, little reminder:

**What is a Donsker class of functions?**

## Definition

We say that  $\mathcal{F}$  is  $P$ -Donsker iff

$$\mathbb{G}_n \rightsquigarrow \mathbb{G} \text{ in } \ell^\infty(\mathcal{F}).$$

## Definition

We say that  $\mathcal{F}$  is Donsker uniformly in  $P \in \mathcal{P}$  iff

$$\mathbb{G}_{n,P} \rightsquigarrow \mathbb{G}_P \text{ in } \ell^\infty(\mathcal{F}) \text{ uniformly in } P \in \mathcal{P}.$$

Again, little reminder:

**What is a Donsker class of functions?**

## Definition

We say that  $\mathcal{F}$  is  $P$ -Donsker iff

$$\mathbb{G}_n \rightsquigarrow \mathbb{G} \text{ in } \ell^\infty(\mathcal{F}).$$

## Definition

We say that  $\mathcal{F}$  is Donsker uniformly in  $P \in \mathcal{P}$  iff

$$\mathbb{G}_{n,P} \rightsquigarrow \mathbb{G}_P \text{ in } \ell^\infty(\mathcal{F}) \text{ uniformly in } P \in \mathcal{P}.$$

$$\sup_{P \in \mathcal{P}} \sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P}) - E h(\mathbb{G}_P)| \rightarrow 0.$$

$$\mathcal{H} := \{h : \ell^\infty(\mathcal{F}) \rightarrow \mathbb{R} : |h(z_1) - h(z_2)| \leq \|z_1 - z_2\|_{\mathcal{F}} \& |h(z_1)| \leq 1\}$$

# Slide-ception

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

**Donsker  
Theorem**

Central Limit  
Theorem  
(Under  
Sequences)

In order to better rembember understand the following procedure, we have to remember better Chapter 2.5

# Slide-ception

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

In order to better remember understand the following procedure, we have to remember better Chapter 2.5

2.4: Glivenko-Cantelli Theorems

## Weak convergence in $\ell^\infty(\mathcal{F})$

Recall:  $X_\alpha$  converges weakly to a tight limit taking values in  $\ell^\infty(\mathcal{F})$  if and only if  $X_\alpha$  is asymptotically tight and its maringals converge weakly.

By the CLT, we already have marginal convergence, so asymptotic tightness of  $\mathbb{G}_n$  is what we are after.

Lasse Vuursteen

Weak Convergence and Empirical Processes



# Slide-ception

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

In order to better remember understand the following procedure, we have to remember better Chapter 2.5

2.4: Glivenko-Cantelli Theorems

## Asymptotic tightness

Recall that a net  $X_\alpha$  is asymptotically tight if for every  $\varepsilon > 0$  there exists a compact set  $K$  such that

$$\liminf P_*(X_\alpha \in K^\delta) \geq 1 - \varepsilon \quad \text{for every } \delta > 0.$$

This is hard to show directly for  $\mathbb{G}_n$ , but we can use a characterization: *asymptotic equicontinuity*.

Lasse Vuursteen

Weak Convergence and Empirical Processes

# Slide-ception

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

In order to better remember understand the following procedure, we have to remember better Chapter 2.5

2.4: Glivenko-Cantelli Theorems

## Asymptotic equicontinuity

A net  $X_\alpha : \Omega \rightarrow \ell^\infty(T)$  is *asymptotically uniformly  $\rho$ -equicontinuous in probability* if for every  $\varepsilon, \eta > 0$  there exists a  $\delta > 0$  such that

$$\limsup_\alpha P^* \left( \sup_{\rho(s,t) < \delta} |X_\alpha(s) - X_\alpha(t)| > \varepsilon \right) < \eta.$$

**Theorem 1.5.7:**  $X_\alpha$  is asymptotically tight if and only if  $X_\alpha(t)$  is asymptotically tight in  $\mathbb{R}$  for every  $t$  and there exists a semimetric  $\rho$  on  $T$  such that  $(T, \rho)$  is totally bounded and  $X_\alpha$  is asymptotically uniformly  $\rho$ -equicontinuous in probability.

# Slide-ception

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

In order to better remember understand the following procedure, we have to remember better Chapter 2.5

## 2.4: Glivenko-Cantelli Theorems

The Donsker theorems come in two flavours (two types of conditions on  $\mathcal{F}$ ):

- Based on the uniform entropy condition

$$\int_0^\infty \sup_Q \sqrt{\log N(\varepsilon \|F\|_{Q,2}, \mathcal{F}, L_2(Q))} d\varepsilon < \infty.$$

- Based on bracketing entropy

$$\int_0^\infty \sqrt{\log N_{[]}(\varepsilon, \mathcal{F}, L_2(P))} d\varepsilon < \infty.$$

These conditions are generally not comparable. Examples of function classes satisfying either or both are given in Chapter 2.7.

# Uniform conditions?

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

However, a uniform version of those conditions is stronger  
**So, just for curiosity what is a pre-Gaussian class of functions?**

# Uniform conditions?

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

However, a uniform version of those conditions is stronger  
**So, just for curiosity what is a pre-Gaussian class of functions?**

## Definition

We say that  $\mathcal{F}$  is pre-Gaussian with respect to  $P$  iff

- $E\|\mathbb{G}\|_{\mathcal{F}} < +\infty$
- $\lim_{\delta \rightarrow 0} E \sup_{\rho(f,g) < \delta} |\mathbb{G}(f) - \mathbb{G}(g)| = 0$

# Uniform conditions?

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

However, a uniform version of those conditions is stronger  
**So, just for curiosity what is a pre-Gaussian class of functions?**

## Definition

We say that  $\mathcal{F}$  is pre-Gaussian with respect to  $P$  iff

- $E\|\mathbb{G}\|_{\mathcal{F}} < +\infty$
- $\lim_{\delta \rightarrow 0} E \sup_{\rho(f,g) < \delta} |\mathbb{G}(f) - \mathbb{G}(g)| = 0$

## Definition

We say that  $\mathcal{F}$  is pre-Gaussian uniformly in  $P \in \mathcal{P}$  iff

- $\sup_{P \in \mathcal{P}} E\|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
- $\lim_{\delta \rightarrow 0} \sup_{P \in \mathcal{P}} E \sup_{\rho_P(f,g) < \delta} |\mathbb{G}(f) - \mathbb{G}(g)| = 0$

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

*Let  $\mathcal{F}$  then the following statements are equivalent:*

- *$\mathcal{F}$  is Donsker and pre-Gaussian uniformly in  $P \in \mathcal{P}$*

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

*Let  $\mathcal{F}$  then the following statements are equivalent:*

- *$\mathcal{F}$  is Donsker and pre-Gaussian uniformly in  $P \in \mathcal{P}$*
- *the sequence  $\mathbb{G}_{n,P}$  is*
  - *asymptotically  $\rightarrow_P$ -equicontinuous uniformly in  $P \in \mathcal{P}$*
  - *$\sup_{P \in \mathcal{P}} N(\epsilon, \mathcal{F}, \rho_P) < +\infty$  for every  $\epsilon > 0$*



# Assertion $\Rightarrow$

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

**Donsker  
Theorem**

Central Limit  
Theorem  
(Under  
Sequences)

① Fix  $\delta > 0$  and  $P$

# Assertion $\Rightarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

① Fix  $\delta > 0$  and  $P$

②  $z \mapsto h_P(z) = \sup_{\rho_P(f,g) < \delta} |z(f) - z(g)| \wedge 1 \in 2\mathcal{H}$

# Assertion $\Rightarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- 1 Fix  $\delta > 0$  and  $P$
- 2  $z \mapsto h_P(z) = \sup_{\rho_P(f,g) < \delta} |z(f) - z(g)| \wedge 1 \in 2\mathcal{H}$
- 3 Since  $\mathcal{F}$  is Donsker,  $E_P^* h_P(\mathbb{G}_{n,P}) \rightarrow E h_P(\mathbb{G}_P)$  when  $n \rightarrow +\infty$

# Assertion $\Rightarrow$

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- ① Fix  $\delta > 0$  and  $P$
- ②  $z \mapsto h_P(z) = \sup_{\rho_P(f,g) < \delta} |z(f) - z(g)| \wedge 1 \in 2\mathcal{H}$
- ③ Since  $\mathcal{F}$  is Donsker,  $E_P^* h_P(\mathbb{G}_{n,P}) \rightarrow Eh_P(\mathbb{G}_P)$  when  $n \rightarrow +\infty$
- ④ Since  $\mathcal{F}$  is pre-Gaussian,  $Eh_P(\mathbb{G}_P) \rightarrow 0$  when  $\delta \rightarrow 0$

# Assertion $\Rightarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- ① Fix  $\delta > 0$  and  $P$
- ②  $z \mapsto h_P(z) = \sup_{\rho_P(f,g) < \delta} |z(f) - z(g)| \wedge 1 \in 2\mathcal{H}$
- ③ Since  $\mathcal{F}$  is Donsker,  $E_P^* h_P(\mathbb{G}_{n,P}) \rightarrow E h_P(\mathbb{G}_P)$  when  $n \rightarrow +\infty$
- ④ Since  $\mathcal{F}$  is pre-Gaussian,  $E h_P(\mathbb{G}_P) \rightarrow 0$  when  $\delta \rightarrow 0$
- ⑤  $\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow +\infty} E_P^* h_P(\mathbb{G}_{n,P}) = 0$

① Let  $\mathcal{G} \subseteq \mathcal{F}$  with  $\mathcal{G}$  finite

# Assertion $\Leftarrow$

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

① Let  $\mathcal{G} \subseteq \mathcal{F}$  with  $\mathcal{G}$  finite

②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}\}$

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- ① Let  $\mathcal{G} \subseteq \mathcal{F}$  with  $\mathcal{G}$  finite
- ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}\}$
- ③ in  $\mathcal{G}$ ,  $P(\sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| < \epsilon) \leq$   
 $\liminf_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^*(\sup_{\rho(f,g) < \delta} |\mathbb{G}_{n,P}(f) - \mathbb{G}_{n,P}(g)| < \epsilon)$



# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- ① Let  $\mathcal{G} \subseteq \mathcal{F}$  with  $\mathcal{G}$  finite
- ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}\}$
- ③ in  $\mathcal{G}$ ,  $P(\sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| < \epsilon) \leq \liminf_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^*(\sup_{\rho(f,g) < \delta} |\mathbb{G}_{n,P}(f) - \mathbb{G}_{n,P}(g)| < \epsilon)$
- ④ in  $\mathcal{G}$ ,  $P(\sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| < \epsilon) \rightarrow 0$

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- ① Let  $\mathcal{G} \subseteq \mathcal{F}$  with  $\mathcal{G}$  finite
- ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}\}$
- ③ in  $\mathcal{G}$ ,  $P(\sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| < \epsilon) \leq \liminf_{n \rightarrow \infty} \sup_{P \in \mathcal{P}} P_P^*(\sup_{\rho(f,g) < \delta} |\mathbb{G}_{n,P}(f) - \mathbb{G}_{n,P}(g)| < \epsilon)$
- ④ in  $\mathcal{G}$ ,  $P(\sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| < \epsilon) \rightarrow 0$
- ⑤ Letting  $\mathcal{G}$  increasing to  $\mathcal{F}$  leads to
$$\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$$

# Assertion $\Leftarrow$

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$

- 1 Fix  $\delta > 0$  and  $\mathcal{G}_P$  a  $\delta$ -net for  $\rho_P$  over  $\mathcal{F}$  and  $|\mathcal{G}_P| \leq k$  uniformly

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

$$\bullet \lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$$

- ① Fix  $\delta > 0$  and  $\mathcal{G}_P$  a  $\delta$ -net for  $\rho_P$  over  $\mathcal{F}$  and  $|\mathcal{G}_P| \leq k$  uniformly
- ②  $\|\mathbb{G}_P\|_{\mathcal{F}} \leq \|\mathbb{G}_P\|_{\mathcal{G}_P} + \sup_{\rho_P(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)|$

# Assertion $\Leftarrow$

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$

- 1 Fix  $\delta > 0$  and  $\mathcal{G}_P$  a  $\delta$ -net for  $\rho_P$  over  $\mathcal{F}$  and  $|\mathcal{G}_P| \leq k$  uniformly
- 2  $\|\mathbb{G}_P\|_{\mathcal{F}} \leq \|\mathbb{G}_P\|_{\mathcal{G}_P} + \sup_{\rho_P(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)|$
- 3 Thus  $E\|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$  uniformly in  $P \in \mathcal{P}$

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\sup_{P \in \mathcal{P}} E \|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
- $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$
- ① Let  $\Pi_P : \mathcal{F} \rightarrow \mathcal{G}_P$  a (sort of) projection

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\sup_{P \in \mathcal{P}} E \|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
  - $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$
- 1 Let  $\Pi_P : \mathcal{F} \rightarrow \mathcal{G}_P$  a (sort of) projection
  - 2  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}_P\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}_P\}$

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\sup_{P \in \mathcal{P}} E \|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
  - $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$
- ① Let  $\Pi_P : \mathcal{F} \rightarrow \mathcal{G}_P$  a (sort of) projection
  - ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}_P\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}_P\}$
  - ③  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E h(\mathbb{G}_P \circ \Pi_P)| \rightarrow 0$



# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\sup_{P \in \mathcal{P}} E \|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
  - $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$
- ① Let  $\Pi_P : \mathcal{F} \rightarrow \mathcal{G}_P$  a (sort of) projection
  - ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}_P\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}_P\}$
  - ③  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E h(\mathbb{G}_P \circ \Pi_P)| \rightarrow 0$
  - ④ Since  $h \in \mathcal{H}$ ,  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E^* h(\mathbb{G}_{n,P})| \leq \epsilon + 2P_P^*(\|\mathbb{G}_{n,P} \circ \Pi_P - \mathbb{G}_{n,P}\|_{\mathcal{F}} > \epsilon)$

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\sup_{P \in \mathcal{P}} E \|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
  - $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$
- ① Let  $\Pi_P : \mathcal{F} \rightarrow \mathcal{G}_P$  a (sort of) projection
  - ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}_P\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}_P\}$
  - ③  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E h(\mathbb{G}_P \circ \Pi_P)| \rightarrow 0$
  - ④ Since  $h \in \mathcal{H}$ ,  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E^* h(\mathbb{G}_{n,P})| \leq \epsilon + 2P_P^*(\|\mathbb{G}_{n,P} \circ \Pi_P - \mathbb{G}_{n,P}\|_{\mathcal{F}} > \epsilon)$
  - ⑤ By construction of  $\mathcal{G}_P$ ,  
 $\limsup P_P^*(\|\mathbb{G}_{n,P} \circ \Pi_P - \mathbb{G}_{n,P}\|_{\mathcal{F}} > \epsilon) \rightarrow 0$

# Assertion $\Leftarrow$

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

- $\sup_{P \in \mathcal{P}} E \|\mathbb{G}_P\|_{\mathcal{F}} < +\infty$
- $\lim_{\delta \rightarrow 0} \sup_{P \in \mathbb{P}} E \sup_{\rho(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0$
- ① Let  $\Pi_P : \mathcal{F} \rightarrow \mathcal{G}_P$  a (sort of) projection
- ②  $\{\mathbb{G}_{n,P}(g) : g \in \mathcal{G}_P\} \rightsquigarrow \{\mathbb{G}_P(g) : g \in \mathcal{G}_P\}$
- ③  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E h(\mathbb{G}_P \circ \Pi_P)| \rightarrow 0$
- ④ Since  $h \in \mathcal{H}$ ,  $\sup_{h \in \mathcal{H}} |E_P^* h(\mathbb{G}_{n,P} \circ \Pi_P) - E^* h(\mathbb{G}_{n,P})| \leq \epsilon + 2P_P^*(\|\mathbb{G}_{n,P} \circ \Pi_P - \mathbb{G}_{n,P}\|_{\mathcal{F}} > \epsilon)$
- ⑤ By construction of  $\mathcal{G}_P$ ,  
 $\limsup P_P^*(\|\mathbb{G}_{n,P} \circ \Pi_P - \mathbb{G}_{n,P}\|_{\mathcal{F}} > \epsilon) \rightarrow 0$
- ⑥ Besides,  $\sup_{h \in \mathcal{H}} |E h(\mathbb{G}_P \circ \Pi_P) - E h(\mathbb{G}_P)| \rightarrow 0$  since  $\mathcal{F}$  uniformly pre-Gaussian

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

*Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta,P} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P,2} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = \{(f - g)^2 : f, g \in \mathcal{F} \text{ are } P\text{-measurable for every } \delta > 0 \text{ and } P \in \mathcal{P}\}$ .*

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta,P} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P,2} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = (f - g)^2 : f, g \in \mathcal{F} \text{ are } P\text{-measurable for every } \delta > 0 \text{ and } P \in \mathcal{P}$ . Furthermore, suppose

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F^2 \mathbf{1}_{F > M} \rightarrow 0,$
- $\int_0^{+\infty} \sup_{Q \in \mathcal{Q}_n} \sqrt{\log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, L_2(Q))} d\epsilon < +\infty,$

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta,P} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P,2} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = (f - g)^2 : f, g \in \mathcal{F} \text{ are } P\text{-measurable for every } \delta > 0$  and  $P \in \mathcal{P}$ . Furthermore, suppose

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F^2 \mathbf{1}_{F > M} \rightarrow 0,$
- $\int_0^{+\infty} \sup_{Q \in \mathcal{Q}_n} \sqrt{\log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, L_2(Q))} d\epsilon < +\infty,$

then  $\mathcal{F}$  is Donsker and pre-Gaussian uniformly in  $P \in \mathcal{P}$

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta,P} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P,2} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = (f - g)^2 : f, g \in \mathcal{F}$  are  $P$ -measurable for every  $\delta > 0$  and  $P \in \mathcal{P}$ . Furthermore, suppose

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F^2 \mathbf{1}_{F > M} \rightarrow 0,$
- $\int_0^{+\infty} \sup_{Q \in \mathcal{Q}_n} \sqrt{\log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, L_2(Q))} d\epsilon < +\infty,$

then  $\mathcal{F}$  is Donsker and pre-Gaussian uniformly in  $P \in \mathcal{P}$

## Theorem

Let  $\mathcal{F}$  such that

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F^2 \mathbf{1}_{F > M} \rightarrow 0,$
- $\int_0^{+\infty} \sup_{P \in \mathcal{P}} \sqrt{\log N_{[]}(\epsilon \|F\|_{P,2}, \mathcal{F}, L_2(P))} d\epsilon < +\infty,$

# Donsker & pre-Gaussian Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta,P} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P,2} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = (f - g)^2 : f, g \in \mathcal{F}$  are  $P$ -measurable for every  $\delta > 0$  and  $P \in \mathcal{P}$ . Furthermore, suppose

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F^2 \mathbf{1}_{F > M} \rightarrow 0,$
- $\int_0^{+\infty} \sup_{Q \in \mathcal{Q}_n} \sqrt{\log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, L_2(Q))} d\epsilon < +\infty,$

then  $\mathcal{F}$  is Donsker and pre-Gaussian uniformly in  $P \in \mathcal{P}$

## Theorem

Let  $\mathcal{F}$  such that

- $\lim_{M \rightarrow +\infty} \sup_{P \in \mathcal{P}} P F^2 \mathbf{1}_{F > M} \rightarrow 0,$
- $\int_0^{+\infty} \sup_{P \in \mathcal{P}} \sqrt{\log N_{[]}(\epsilon \|F\|_{P,2}, \mathcal{F}, L_2(P))} d\epsilon < +\infty,$

then  $\mathcal{F}$  is Donsker and pre-Gaussian uniformly in  $P \in \mathcal{P}$



# Application of the results

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

For each  $n \in \mathbb{N}$ , let  $X_{n1}, \dots, X_{nn}$  be i.i.d according to  $P_n$  and  $\mathbb{P}_n := \frac{1}{n} \sum_{i=1}^n \delta_{X_{ni}}$ .

# Application of the results

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

For each  $n \in \mathbb{N}$ , let  $X_{n1}, \dots, X_{nn}$  be i.i.d according to  $P_n$  and

$$\mathbb{P}_n := \frac{1}{n} \sum_{i=1}^n \delta_{X_{ni}}.$$

If  $P_n \rightarrow P_0$  in a suitable sense, then hopefully

# Application of the results

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

For each  $n \in \mathbb{N}$ , let  $X_{n1}, \dots, X_{nn}$  be i.i.d according to  $P_n$  and  $\mathbb{P}_n := \frac{1}{n} \sum_{i=1}^n \delta_{X_{ni}}$ .

If  $P_n \rightarrow P_0$  in a suitable sense, then hopefully

$$\mathbb{G}_{n,P_n} := \sqrt{n}(\mathbb{P}_n - P_n) \rightsquigarrow \mathbb{G}_{P_0} \text{ in } \ell^\infty(\mathcal{F})$$

# Lemmas

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

### Lemma

Let  $\mathcal{F}$  be Donsker and pre-Gaussian uniformly in  $\{P_m, m \leq n\}$  and

- $\sup_{f,g \in \mathcal{F}} |\rho_{P_n}(f, g) - \rho_{P_0}(f, g)| \rightarrow 0,$
- $\limsup_{n \rightarrow +\infty} P_n F^2 \mathbf{1}_{F \geq \epsilon \sqrt{n}} = 0$  for all  $\epsilon > 0$ .

Then  $\mathbb{G}_{n, P_n} \rightsquigarrow$  in  $\ell^\infty(\mathcal{F})$ .

### Lemma

Let  $\mathcal{F}$  be pre-Gaussian uniformly in  $\{P_m, m \leq n\}$  and

- $\sup_{f,g \in \mathcal{F}} |\rho_{P_n}(f, g) - \rho_{P_0}(f, g)| \rightarrow 0,$

Then  $\mathbb{G}_{P_n} \rightsquigarrow$  in  $\ell^\infty(\mathcal{F})$ .

# Final Theorems

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

*Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta, P_n} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P_n, 2} < \delta\}$   
and  $\mathcal{F}_{\infty}^0 = (f - g)^2 : f, g \in \mathcal{F}$  are  $P_n$ -measurable for every  
 $\delta > 0$  and  $n$ .*

# Final Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

### Theorem

Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta, P_n} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P_n, 2} < \delta\}$  and  $\mathcal{F}_{\infty}^0 = (f - g)^2 : f, g \in \mathcal{F} \text{ are } P_n\text{-measurable for every } \delta > 0 \text{ and } n$ . Furthermore, suppose

- $P_n F^2 = O(1)$ ,
- $\int_0^{+\infty} \sup_{Q \in \mathcal{Q}_n} \sqrt{\log N(\epsilon \|F\|_{Q, 2}, \mathcal{F}, L_2(Q))} d\epsilon < +\infty$ ,
- $\sup_{f, g \in \mathcal{F}} |\rho_{P_n}(f, g) - \rho_{P_0}(f, g)| \rightarrow 0$ ,
- $\limsup_{n \rightarrow +\infty} P_n F^2 \mathbf{1}_{F \geq \epsilon \sqrt{n}} = 0 \text{ for all } \epsilon > 0$ .

# Final Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

### Theorem

Let  $\mathcal{F}$  such that  $\mathcal{F}_{\delta, P_n} = \{f - g : f, g \in \mathcal{F}, \|f - g\|_{P_n, 2} < \delta\}$  and  $\mathcal{F}_{\infty}^0 = (f - g)^2 : f, g \in \mathcal{F} \text{ are } P_n\text{-measurable for every } \delta > 0 \text{ and } n$ . Furthermore, suppose

- $P_n F^2 = O(1)$ ,
- $\int_0^{+\infty} \sup_{Q \in \mathcal{Q}_n} \sqrt{\log N(\epsilon \|F\|_{Q, 2}, \mathcal{F}, L_2(Q))} d\epsilon < +\infty$ ,
- $\sup_{f, g \in \mathcal{F}} |\rho_{P_n}(f, g) - \rho_{P_0}(f, g)| \rightarrow 0$ ,
- $\limsup_{n \rightarrow +\infty} P_n F^2 \mathbf{1}_{F \geq \epsilon \sqrt{n}} = 0 \text{ for all } \epsilon > 0$ .

Then  $\mathbb{G}_{n, P_n} \rightsquigarrow$  in  $\ell^\infty(\mathcal{F})$ .

# Final Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

### Theorem

*Let  $\mathcal{F}$  such that*



# Final Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

## Theorem

Let  $\mathcal{F}$  such that

- $\mathcal{F}$  totally bounded for  $\rho_{P_0}$
- $\int_0^{+\infty} \sup_{P \in \mathcal{P}} \sqrt{\log N_{[]}(\epsilon \|F\|_{P,2}, \mathcal{F}, L_2(P))} d\epsilon < +\infty,$
- $\sup_{f,g \in \mathcal{F}} |\rho_{P_n}(f, g) - \rho_{P_0}(f, g)| \rightarrow 0,$
- $\limsup_{n \rightarrow +\infty} P_n F^2 \mathbf{1}_{F \geq \epsilon \sqrt{n}} = 0$  for all  $\epsilon > 0.$

# Final Theorems

## Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

### Theorem

Let  $\mathcal{F}$  such that

- $\mathcal{F}$  totally bounded for  $\rho_{P_0}$
- $\int_0^{+\infty} \sup_{P \in \mathcal{P}} \sqrt{\log N_{[]}(\epsilon \|F\|_{P,2}, \mathcal{F}, L_2(P))} d\epsilon < +\infty,$
- $\sup_{f,g \in \mathcal{F}} |\rho_{P_n}(f, g) - \rho_{P_0}(f, g)| \rightarrow 0,$
- $\limsup_{n \rightarrow +\infty} P_n F^2 \mathbf{1}_{F \geq \epsilon \sqrt{n}} = 0$  for all  $\epsilon > 0$ .

Then  $\mathbb{G}_{n, P_n} \rightsquigarrow$  in  $\ell^\infty(\mathcal{F})$ .

# In the next...

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

... session of the Reading group - Mathematical statistics, we'll see

# In the next...

Chapter 2.8

Amine Hadji

Notations &  
Intentions

Glivenko-  
Cantelli  
Theorem

Donsker  
Theorem

Central Limit  
Theorem  
(Under  
Sequences)

... session of the Reading group - Mathematical statistics, we'll see

- Chapter 2.9, with
  - the multiplier central limit theorem
  - the conditional central limit theorem

Let  $Z_i := \delta_{X_i} - P$ , then we have directly

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_i \rightsquigarrow \mathbb{G},$$

but also,

Let  $Z_i := \delta_{X_i} - P$ , then we have directly

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_i \rightsquigarrow \mathbb{G},$$

but also,

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \xi_i Z_i \rightsquigarrow \mathbb{G},$$

if the  $\xi_i$ 's satisfy some condition and  $\mathcal{F}$  is Donsker (why?).

Let  $Z_i := \delta_{X_i} - P$ , then we have directly

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n Z_i \rightsquigarrow \mathbb{G},$$

but also,

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \xi_i Z_i \rightsquigarrow \mathbb{G},$$

if the  $\xi_i$ 's satisfy some condition and  $\mathcal{F}$  is Donsker (why?).  
Moreover the previous assertion is true given almost every  
sequence  $Z_1, Z_2, \dots$  (why?)