

Reading group mathematical foundations of statistics

Stefan Franssen, Msc

September 9, 2019

Goals

- To understand the current knowledge of nonparametric statistics better

Goals

- To understand the current knowledge of nonparametric statistics better
- To translate this knowledge into new results in statistics

Getting to know the Audience

Who are you?

Set up

- Held biweekly.

Set up

- Held biweekly.
- Alternates with the causality seminar.

Set up


- Held biweekly.
- Alternates with the causality seminar.
- Amine, Bart, Geerten, Lasse and I will be the main speakers.

Set up

- Held biweekly.
- Alternates with the causality seminar.
- Amine, Bart, Geerten, Lasse and I will be the main speakers.
- First two talks will be held in Room 312, the rest of the semester will be in 176.

Set up

- Held biweekly.
- Alternates with the causality seminar.
- Amine, Bart, Geerten, Lasse and I will be the main speakers.
- First two talks will be held in Room 312, the rest of the semester will be in 176.
- Any questions so far?



The book we will start with.

You can get it from Springer link, where you can also order a softcover version for 25 euro.

I also have a pdf version of the second version of the book which I can share with the people here.

Topic of the book

The main object of interest is the empirical distribution

$$\mathbb{P}_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}.$$

Topic of the book

The main object of interest is the empirical distribution

$$\mathbb{P}_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}.$$

We will study its asymptotic normality, not just for a single function f but also uniformly over a class \mathcal{F} of functions.

Topic of the book

The main object of interest is the empirical distribution

$$\mathbb{P}_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}.$$

We will study its asymptotic normality, not just for a single function f but also uniformly over a class \mathcal{F} of functions. We will develop tools for showing asymptotic normality in a uniform sense.

Outline of the book

- Part 1: Introduction of the basic definitions

Outline of the book

- Part 1: Introduction of the basic definitions
- Part 2: The theory on empirical processes

Outline of the book

- Part 1: Introduction of the basic definitions
- Part 2: The theory on empirical processes
- Part 3: Applications of the theory to statistical problems.

Part 1

Part 1 contains the basic theory of convergence of random variables in various settings.

Part 2

Part 2 contains the empirical process theory. The first 5 chapters are dealing with the main theory and the latter chapters deal mostly with examples.

Part 3

Part 3 contains a lot of statistical applications, for example the theory of M- and Z-estimators and the bootstrap is treated here.

Overview for today

We start the first two weeks with the first part of the book. Today I will do the following sections:

- Introduction
- Outer integrals and Measurable Majorants
- Weak convergence
- Product spaces

Introduction

The classical theory of weak convergence.

Introduction: weak convergence of probability measures

Definition

Let (\mathbb{D}, d) be a metric space, and let $(P_n)_{n \in \mathbb{N}}$ and P be probability measures on the borel measurable space $(\mathbb{D}, \mathcal{D})$. Then P_n converges weakly to P if and only if

$$\int f dP_n \rightarrow \int f dP, \quad \forall f \in C_b(\mathbb{D})$$

Introduction: Weak convergence of random variables

Definition

In the same notation as before, let X_n, X be random variables taking values in \mathbb{D} . Then X_n converges weakly to X if and only if

$$\mathbb{E}[f(X_n)] \rightarrow \mathbb{E}[f(X)] \quad \forall f \in C_b(\mathbb{D}).$$

Introduction: The limitation of the classical theory

In the classical theory of weak convergence, we run into measurability issues, namely all the random variables X_n are measurable maps.

Introduction: The limitation of the classical theory

In the classical theory of weak convergence, we run into measurability issues, namely all the random variables X_n are measurable maps. However, classical examples of objects of interest are not measurable.

Introduction: The limitation of the classical theory

In the classical theory of weak convergence, we run into measurability issues, namely all the random variables X_n are measurable maps. However, classical examples of objects of interest are not measurable. For example let X_i be iid uniformly distributed random variables. Then the following two maps are not measurable in the natural spaces (exercise 1.7.3):

$$\mathbb{F}_n(t) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{X_i \leq t}$$

$$\mathbb{G}_n(t) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbb{1}_{X_i \leq t}$$

Introduction: Suggested solutions

Several people have suggested solutions, but the one that can cover the most is suggested by Hoffmann-Jørgensen. We will follow this approach.

The suggested solution

The suggested solution is to drop the measurability requirement and only require asymptotic measurability.

Introduction: Key differences

The four main differences are:

- The notion of (uniform) tightness of sequences needs a modification

Introduction: Key differences

The four main differences are:

- The notion of (uniform) tightness of sequences needs a modification
- Prohorov's theorem, asymptotic tightness implies relative compactness, must be modified by the addition of the requirement of asymptotic measurability

Introduction: Key differences

The four main differences are:

- The notion of (uniform) tightness of sequences needs a modification
- Prohorov's theorem, asymptotic tightness implies relative compactness, must be modified by the addition of the requirement of asymptotic measurability
- (Almost) sure convergence is meaningless without asymptotic measurability

Introduction: Key differences

The four main differences are:

- The notion of (uniform) tightness of sequences needs a modification
- Prohorov's theorem, asymptotic tightness implies relative compactness, must be modified by the addition of the requirement of asymptotic measurability
- (Almost) sure convergence is meaningless without asymptotic measurability
- No general Fubini's theorem.

Outer integrals and Majorants: The outer integral

Definition (p. 6)

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space and let $T : \Omega \rightarrow \bar{\mathbb{R}}$ be any (possibly nonmeasurable) function. The **outer integral** of T with respect to P is defined as:

$$\mathbb{E}^*[T] = \inf \{ \mathbb{E}U : U \geq T, U : \Omega \rightarrow \bar{\mathbb{R}} \text{ integrable} \}$$

Outer integrals and Majorants: The outer probability

Definition (p. 6)

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be an probability space and let B be any (possibly nonmeasureable) subset of Ω . The **outer probability** of B with respect to P is defined as:

$$\mathbb{P}^*(B) = \inf \{ \mathbb{P}(A) : B \subset A, A \in \mathcal{A} \}$$

Outer integrals and Majorants: Inner versions

The inner integral and probability are defined by reversing the direction of the inequalities.

Outer integrals and Majorants: Existence of a Majorant

Lemma (Lem 1.2.1)

For any map $T : \Omega \rightarrow \bar{\mathbb{R}}$ there exists a measurable function $T^ : \Omega \rightarrow \bar{\mathbb{R}}$ with*

- $T^* \geq T;$

Outer integrals and Majorants: Existence of a Majorant

Lemma (Lem 1.2.1)

For any map $T : \Omega \rightarrow \bar{\mathbb{R}}$ there exists a measurable function $T^ : \Omega \rightarrow \bar{\mathbb{R}}$ with*

- $T^* \geq T$;
- $T^* \leq U$ almost surely, for any measurable $U : \Omega \rightarrow \bar{\mathbb{R}}$ with $U \geq T$ a.s.

Outer integrals and Majorants: Existence of a Majorant

Lemma (Lem 1.2.1)

For any map $T : \Omega \rightarrow \bar{\mathbb{R}}$ there exists a measurable function $T^ : \Omega \rightarrow \bar{\mathbb{R}}$ with*

- $T^* \geq T$;
- $T^* \leq U$ almost surely, for any measurable $U : \Omega \rightarrow \bar{\mathbb{R}}$ with $U \geq T$ a.s.

For any T^ satisfying these requirements, it holds that $\mathbb{E}^* T = \mathbb{E} T^*$, provided that $\mathbb{E} T^*$ exists. The latter is certainly true if $\mathbb{E}^* T < \infty$.*

Outer integrals and Majorants: Relation between outer measure and probability

Lemma (Lem 1.2.3)

For any subset B of Ω ,

- $P^*(B) = \mathbb{E}^* 1_B; \mathbb{P}_*(B) = \mathbb{E}(1_B)_*;$

Outer integrals and Majorants: Relation between outer measure and probability

Lemma (Lem 1.2.3)

For any subset B of Ω ,

- $P^*(B) = \mathbb{E}^* \mathbb{1}_B$; $\mathbb{P}_*(B) = \mathbb{E}(\mathbb{1}_B)_*$;
- *there exists a measurable set B^* containing B with $\mathbb{P}(B^*) = \mathbb{P}^*(B)$, for any such B^* it holds that $\mathbb{1}_{B^*} = (\mathbb{1}_B)^*$.*

Outer integrals and Majorants: Relation between outer measure and probability

Lemma (Lem 1.2.3)

For any subset B of Ω ,

- $P^*(B) = \mathbb{E}^* \mathbb{1}_B$; $\mathbb{P}_*(B) = \mathbb{E}(\mathbb{1}_B)_*$;
- *there exists a measurable set B^* containing B with $\mathbb{P}(B^*) = \mathbb{P}^*(B)$, for any such B^* it holds that $\mathbb{1}_{B^*} = (\mathbb{1}_B)^*$.*
- $(\mathbb{1}_B)^* + (\mathbb{1}_{\Omega-B})_* = 1$.

Outer integrals and Majorants: Fubini

Lemma (Lemma 1.2.6)

Let T be defined on a product probability space. Then

$$\mathbb{E}_* T \leq \mathbb{E}_{1*} \mathbb{E}_{2*} T \leq \mathbb{E}_1^* \mathbb{E}_2^* T \leq \mathbb{E}^* T$$

Outer integrals and Majorants: Fubini

Lemma (Lemma 1.2.6)

Let T be defined on a product probability space. Then

$$\mathbb{E}_* T \leq \mathbb{E}_{1*} \mathbb{E}_{2*} T \leq \mathbb{E}_1^* \mathbb{E}_2^* T \leq \mathbb{E}^* T$$

Lemma (Lem 1.2.7)

Let $(\Omega_1, \mathcal{A}_1)$ be a separable metric space equipped with its Borel σ -field.

Outer integrals and Majorants: Fubini

Lemma (Lemma 1.2.6)

Let T be defined on a product probability space. Then

$$\mathbb{E}_* T \leq \mathbb{E}_{1*} \mathbb{E}_{2*} T \leq \mathbb{E}_1^* \mathbb{E}_2^* T \leq \mathbb{E}^* T$$

Lemma (Lem 1.2.7)

Let $(\Omega_1, \mathcal{A}_1)$ be a separable metric space equipped with its Borel σ -field. Suppose the map $T : \Omega_1 \times \Omega_2 \rightarrow \mathbb{R}$ is defined on a product probability space

Outer integrals and Majorants: Fubini

Lemma (Lemma 1.2.6)

Let T be defined on a product probability space. Then

$$\mathbb{E}_* T \leq \mathbb{E}_{1*} \mathbb{E}_{2*} T \leq \mathbb{E}_1^* \mathbb{E}_2^* T \leq \mathbb{E}^* T$$

Lemma (Lem 1.2.7)

Let $(\Omega_1, \mathcal{A}_1)$ be a separable metric space equipped with its Borel σ -field. Suppose the map $T : \Omega_1 \times \Omega_2 \rightarrow \mathbb{R}$ is defined on a product probability space and satisfies

$$|T(\omega_1, \omega_2) - T(\omega'_1, \omega_2)| \leq d(\omega_1, \omega'_1)H(\omega_2),$$

Outer integrals and Majorants: Fubini

Lemma (Lemma 1.2.6)

Let T be defined on a product probability space. Then

$$\mathbb{E}_* T \leq \mathbb{E}_{1*} \mathbb{E}_{2*} T \leq \mathbb{E}_1^* \mathbb{E}_2^* T \leq \mathbb{E}^* T$$

Lemma (Lem 1.2.7)

Let $(\Omega_1, \mathcal{A}_1)$ be a separable metric space equipped with its Borel σ -field. Suppose the map $T : \Omega_1 \times \Omega_2 \rightarrow \mathbb{R}$ is defined on a product probability space and satisfies

$|T(\omega_1, \omega_2) - T(\omega'_1, \omega_2)| \leq d(\omega_1, \omega'_1)H(\omega_2)$, for a function H with $P_2^ H < \infty$ and for every sufficiently small $d(\omega_1, \omega'_1)$.*

Outer integrals and Majorants: Fubini

Lemma (Lemma 1.2.6)

Let T be defined on a product probability space. Then

$$\mathbb{E}_* T \leq \mathbb{E}_{1*} \mathbb{E}_{2*} T \leq \mathbb{E}_1^* \mathbb{E}_2^* T \leq \mathbb{E}^* T$$

Lemma (Lem 1.2.7)

Let $(\Omega_1, \mathcal{A}_1)$ be a separable metric space equipped with its Borel σ -field. Suppose the map $T : \Omega_1 \times \Omega_2 \rightarrow \mathbb{R}$ is defined on a product probability space and satisfies

$|T(\omega_1, \omega_2) - T(\omega'_1, \omega_2)| \leq d(\omega_1, \omega'_1)H(\omega_2)$, for a function H with $P_2^ H < \infty$ and for every sufficiently small $d(\omega_1, \omega'_1)$. Then*

$$\mathbb{E}^* T = \mathbb{E}_1 \mathbb{E}_2^* T$$

The monotone convergence theorem

Lemma (Ex. 1.2.3)

Let T_n, T be maps on a probability space with $T_n \uparrow T$ pointwise on a set of probability one.

The monotone convergence theorem

Lemma (Ex. 1.2.3)

Let T_n, T be maps on a probability space with $T_n \uparrow T$ pointwise on a set of probability one. Then $T_n^ \uparrow T^*$ almost surely.*

The monotone convergence theorem

Lemma (Ex. 1.2.3)

Let T_n, T be maps on a probability space with $T_n \uparrow T$ pointwise on a set of probability one. Then $T_n^ \uparrow T^*$ almost surely. If the maps are bounded from below, then $\mathbb{E}^* T_n \uparrow \mathbb{E}^* T$.*

The dominated convergence theorem

Lemma (Ex 1.2.4)

Let T_n, T be maps on a probability space with $|T_n - T|^$ converging to zero almost surely.*

The dominated convergence theorem

Lemma (Ex 1.2.4)

Let T_n, T be maps on a probability space with $|T_n - T|^$ converging to zero almost surely. Suppose $\mathbb{E}^*[\sup_n |T_n|] < \infty$, then $\mathbb{E}^* T_n \rightarrow \mathbb{E}^* T$.*

Polish spaces

Definition (P. 17)

*A topological space X is called **Polish** if it is separable and its topology can be generated by a complete metric.*

Tight and separable random variables

Definition (p. 16, 17)

- A Borel probability measure P is **tight** if for every $\epsilon > 0$ there exists a compact K such that $P(K) \geq 1 - \epsilon$.

Tight and separable random variables

Definition (p. 16, 17)

- A Borel probability measure P is **tight** if for every $\epsilon > 0$ there exists a compact K such that $P(K) \geq 1 - \epsilon$.
- A Borel probability measure P is **pre-tight** if for every $\epsilon > 0$ there exists a totally bounded measurable K such that $P(K) \geq 1 - \epsilon$.

Tight and separable random variables

Definition (p. 16, 17)

- A Borel probability measure P is **tight** if for every $\epsilon > 0$ there exists a compact K such that $P(K) \geq 1 - \epsilon$.
- A Borel probability measure P is **pre-tight** if for every $\epsilon > 0$ there exists a totally bounded measurable K such that $P(K) \geq 1 - \epsilon$.
- We call a Borel probability measure **separable** if there exists a separable measurable set S with $P(S) = 1$.

Tight and separable random variables

Definition (p. 16, 17)

- A Borel probability measure P is **tight** if for every $\epsilon > 0$ there exists a compact K such that $P(K) \geq 1 - \epsilon$.
- A Borel probability measure P is **pre-tight** if for every $\epsilon > 0$ there exists a totally bounded measurable K such that $P(K) \geq 1 - \epsilon$.
- We call a Borel probability measure **separable** if there exists a separable measurable set S with $P(S) = 1$.

Lemma (1.3.2)

On a complete metric space these concepts are equivalent.

Definition

Let A be a directed set with order relation \geq and X be a topological space. A map $f : A \rightarrow X$ is called a net.

Example

- Sequences

Weak convergence of random variables

Definition (1.3.3)

Let $(\Omega_\alpha, \mathcal{A}_\alpha, P_\alpha)$ be a net of probability spaces and let $X_\alpha : \Omega_\alpha \rightarrow \mathbb{D}$ be arbitrary maps. The net X_α **converges weakly** to a borel measure L if

$$\mathbb{E}^* f(X_\alpha) \rightarrow \int f dL, \quad \forall f \in C_b(\mathbb{D}).$$

We denote this convergence by $X_\alpha \rightsquigarrow L$.

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L;$

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L$;
- $\liminf P_*(X_\alpha \in G) \geq L(G)$ for every open G ;

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L$;
- $\liminf P_*(X_\alpha \in G) \geq L(G)$ for every open G ;
- $\limsup P^*(X_\alpha \in F) \leq L(F)$ for every closed F ;

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L$;
- $\liminf P_*(X_\alpha \in G) \geq L(G)$ for every open G ;
- $\limsup P^*(X_\alpha \in F) \leq L(F)$ for every closed F ;
- $\liminf \mathbb{E}_* f(X_\alpha) \geq \int f dL$ for every lower semicontinuous f that is bounded below;

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L$;
- $\liminf P_*(X_\alpha \in G) \geq L(G)$ for every open G ;
- $\limsup P^*(X_\alpha \in F) \leq L(F)$ for every closed F ;
- $\liminf \mathbb{E}_* f(X_\alpha) \geq \int f dL$ for every lower semicontinuous f that is bounded below;
- $\limsup \mathbb{E}^* f(X_\alpha) \leq \int f dL$ for every upper semicontinuous f that is bounded above;

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L$;
- $\liminf P_*(X_\alpha \in G) \geq L(G)$ for every open G ;
- $\limsup P^*(X_\alpha \in F) \leq L(F)$ for every closed F ;
- $\liminf \mathbb{E}_* f(X_\alpha) \geq \int f dL$ for every lower semicontinuous f that is bounded below;
- $\limsup \mathbb{E}^* f(X_\alpha) \leq \int f dL$ for every upper semicontinuous f that is bounded above;
- $\lim P^*(X_\alpha \in B) = \lim P_*(X_\alpha \in B) = L(B)$ for every Borel set B with $L(\delta B) = 0$;

Portmanteau lemma

Lemma (1.3.4)

The following statements are equivalent.

- $X_\alpha \rightsquigarrow L$;
- $\liminf P_*(X_\alpha \in G) \geq L(G)$ for every open G ;
- $\limsup P^*(X_\alpha \in F) \leq L(F)$ for every closed F ;
- $\liminf \mathbb{E}_* f(X_\alpha) \geq \int f dL$ for every lower semicontinuous f that is bounded below;
- $\limsup \mathbb{E}^* f(X_\alpha) \leq \int f dL$ for every upper semicontinuous f that is bounded above;
- $\lim P^*(X_\alpha \in B) = \lim P_*(X_\alpha \in B) = L(B)$ for every Borel set B with $L(\delta B) = 0$;
- $\liminf \mathbb{E}_* f(X_\alpha) \geq \int f dL$ for every bounded, Lipschitz continuous, nonnegative f .

The continuous mapping theorem for weak convergence

Lemma (1.3.6)

Let $g : \mathbb{D} \rightarrow \mathbb{E}$ be continuous at every point of set $\mathbb{D}_0 \subset \mathbb{D}$.

The continuous mapping theorem for weak convergence

Lemma (1.3.6)

Let $g : \mathbb{D} \rightarrow \mathbb{E}$ be continuous at every point of set $\mathbb{D}_0 \subset \mathbb{D}$. If $X_\alpha \rightsquigarrow X$ and X takes its values in \mathbb{D}_0 ,

The continuous mapping theorem for weak convergence

Lemma (1.3.6)

Let $g : \mathbb{D} \rightarrow \mathbb{E}$ be continuous at every point of set $\mathbb{D}_0 \subset \mathbb{D}$. If $X_\alpha \rightsquigarrow X$ and X takes its values in \mathbb{D}_0 , then $g(X_\alpha) \rightsquigarrow g(X)$.

Asymptotically measurability and asymptotically tight

Definition (1.3.7)

The net of maps X_α is:

- *asymptotically measurable iff*

$$\mathbb{E}^* f(X_\alpha) - \mathbb{E}_* f(X_\alpha) \rightarrow 0, \quad \forall f \in C_b(\mathbb{D}).$$

Asymptotically measurability and asymptotically tight

Definition (1.3.7)

The net of maps X_α is:

- *asymptotically measurable iff*

$$\mathbb{E}^* f(X_\alpha) - \mathbb{E}_* f(X_\alpha) \rightarrow 0, \quad \forall f \in C_b(\mathbb{D}).$$

- *Asymptotically tight if for every $\epsilon > 0$ there exists a compact K such that*

$$\liminf P_*(X_\alpha \in K^\delta) \geq 1 - \epsilon, \quad \forall \delta > 0.$$

Here $K^\delta = \{y \in \mathbb{D} : d(y, K) < \delta\}$ the δ -enlargement around K .

Prohorov's theorem 1: The forward statement

Lemma (1.3.8)

- *If $X_\alpha \rightsquigarrow X$, then X_α is asymptotically measurable.*

Prohorov's theorem 1: The forward statement

Lemma (1.3.8)

- If $X_\alpha \rightsquigarrow X$, then X_α is asymptotically measurable.
- If $X_\alpha \rightsquigarrow X$, then X_α is asymptotically tight if and only if X is tight.

Prohorov's theorem 2: The reverse statement

Lemma (1.3.9)

- *If the net X_α is asymptotically tight and asymptotically measurable, then it has a subnet $X_{\alpha(\beta)}$ that converges in law to a tight Borel law.*

Prohorov's theorem 2: The reverse statement

Lemma (1.3.9)

- *If the net X_α is asymptotically tight and asymptotically measurable, then it has a subnet $X_{\alpha(\beta)}$ that converges in law to a tight Borel law.*
- *If the sequence X_n is asymptotically tight and asymptotically measurable, then it has a subsequence X_{n_j} that converges weakly to a tight Borel law.*

Prohorov's theorem 2: The reverse statement

Lemma (1.3.9)

- *If the net X_α is asymptotically tight and asymptotically measurable, then it has a subnet $X_{\alpha(\beta)}$ that converges in law to a tight Borel law.*
- *If the sequence X_n is asymptotically tight and asymptotically measurable, then it has a subsequence X_{n_j} that converges weakly to a tight Borel law.*

<https://math.stackexchange.com/questions/2150490/trying-to-understand-how-a-subnet-of-a-sequence-differs-from-a-subsequence>
<https://math.stackexchange.com/questions/1209341/example-of-converging-subnet-when-there-is-no-converging-subsequence?noredirect=1&lq=1>

The end

Any questions?