

# Chapter 2.1: Introduction

**Geerten Koers**

Reading group *Weak Convergence and Empirical Processes*

**2019-10-07**

- 1 Empirical processes
- 2 Limit distributions
- 3 Entropy
- 4 Glivenko-Cantelli and Donsker classes
- 5 VC-classes
- 6 Miscellaneous
- 7 Maximal Inequalities

## Definition

The *empirical measure* is defined by

$$\mathbb{P}_n(C) := \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \in C)$$

for  $C \subset \mathcal{X}$ .

## Definition

The *empirical measure* is defined by

$$\mathbb{P}_n(C) := \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \in C)$$

for  $C \subset \mathcal{X}$ .

**This induces the map**

$$f \mapsto \mathbb{P}_n f = \frac{1}{n} \sum_{i=1}^n f(X_i),$$

**for any  $f : \mathcal{X} \rightarrow \mathbb{R}$ .**

## Definition

The *empirical process*  $\mathbb{G}_n$  is defined by

$$f \mapsto \mathbb{G}_n f = \frac{1}{\sqrt{n}} \sum_{i=1}^n (f(X_i) - Pf)$$

with  $Pf = \int f dP$ .

**For a fixed  $f : \mathcal{X} \rightarrow \mathbb{R}$  we have**

$$\mathbb{P}_n f \rightarrow Pf,$$

$$\mathbb{G}_n \rightsquigarrow N(0, P(f - Pf)^2)$$

**if  $P|f| < \infty$  and  $Pf^2 < \infty$ .**

**For a fixed  $f : \mathcal{X} \rightarrow \mathbb{R}$  we have**

$$\mathbb{P}_n f \rightarrow Pf,$$

$$\mathbb{G}_n \rightsquigarrow N(0, P(f - Pf)^2)$$

**if  $P|f| < \infty$  and  $Pf^2 < \infty$ .**

**For what  $\mathcal{F} \subset \mathbb{R}^{\mathcal{X}}$  can we make these limits uniform?**

# Uniform convergence

## Definition

For a measure  $Q$  on  $\mathcal{X}$ , define

$$\|Q\|_{\mathcal{F}} := \sup\{|Qf| : f \in \mathcal{F}\}.$$



# Uniform convergence

## Definition

For a measure  $Q$  on  $\mathcal{X}$ , define

$$\|Q\|_{\mathcal{F}} := \sup\{|Qf| : f \in \mathcal{F}\}.$$

## Definition

If  $\|\mathbb{P}_n - P\|_{\mathcal{F}} \rightarrow 0$  in outer probability, then  $\mathcal{F}$  is called *Glivenko-Cantelli*.

# Uniform convergence

## Definition

For a measure  $Q$  on  $\mathcal{X}$ , define

$$\|Q\|_{\mathcal{F}} := \sup\{|Qf| : f \in \mathcal{F}\}.$$

## Definition

If  $\|\mathbb{P}_n - P\|_{\mathcal{F}} \rightarrow 0$  in outer probability, then  $\mathcal{F}$  is called *Glivenko-Cantelli*.

$$\sup \left\{ \frac{1}{n} \left| \sum_{i=1}^n f(X_i) - Pf(X_1) \right| : f \in \mathcal{F} \right\} \rightarrow 0.$$

# Limiting theorems

**Assume**

$$\sup_{f \in \mathcal{F}} |f(x) - Pf| < \infty.$$

**Now  $\{G_n f : f \in \mathcal{F}\}$  is a map into  $\ell^\infty(\mathcal{F})$ .**

# Limiting theorems

**Assume**

$$\sup_{f \in \mathcal{F}} |f(x) - Pf| < \infty.$$

**Now  $\{\mathbb{G}_n f : f \in \mathcal{F}\}$  is a map into  $\ell^\infty(\mathcal{F})$ .**

## Definition

If

$$\mathbb{G}_n \rightsquigarrow \mathbb{G}$$

for a tight Borel measurable element  $\mathbb{G} \in \ell^\infty(\mathcal{F})$ , then  $\mathcal{F}$  is called a *P-Donsker class*.

# Limiting theorems

**Assume**

$$\sup_{f \in \mathcal{F}} |f(x) - Pf| < \infty.$$

**Now  $\{\mathbb{G}_n f : f \in \mathcal{F}\}$  is a map into  $\ell^\infty(\mathcal{F})$ .**

## Definition

If

$$\mathbb{G}_n \rightsquigarrow \mathbb{G}$$

for a tight Borel measurable element  $\mathbb{G} \in \ell^\infty(\mathcal{F})$ , then  $\mathcal{F}$  is called a *P-Donsker class*.

**$\mathbb{G}$  is a centered Gaussian process with covariance**

$$\mathbb{E} \mathbb{G} f_1 \mathbb{G} f_2 = P f_1 f_2 - P f_1 P f_2.$$

**We define the  $X_i$  on the product space  $(\mathcal{X}^\infty, \mathcal{B}^\infty, P^\infty)$ . This is possible since the projection on  $\mathcal{X}^n$  is perfect.**

# Entropy


## Definition (2.1.5)

The *covering number*  $N(\epsilon, \mathcal{F}, \|\cdot\|)$  is the minimum number of balls  $B_\epsilon(f) := \{g \in \mathcal{F} : \|g - f\| < \epsilon\}$  such that  $\mathcal{F}$  is covered. The *entropy without bracketing* is  $\log N(\epsilon, \mathcal{F}, \|\cdot\|)$ .

# Entropy

## Definition (2.1.5)

The *covering number*  $N(\epsilon, \mathcal{F}, \|\cdot\|)$  is the minimum number of balls  $B_\epsilon(f) := \{g \in \mathcal{F} : \|g - f\| < \epsilon\}$  such that  $\mathcal{F}$  is covered. The *entropy without bracketing* is  $\log N(\epsilon, \mathcal{F}, \|\cdot\|)$ .






# Entropy

## Definition (2.1.5)


The *covering number*  $N(\epsilon, \mathcal{F}, \|\cdot\|)$  is the minimum number of balls  $B_\epsilon(f) := \{g \in \mathcal{F} : \|g - f\| < \epsilon\}$  such that  $\mathcal{F}$  is covered. The *entropy without bracketing* is  $\log N(\epsilon, \mathcal{F}, \|\cdot\|)$ .



# Entropy

## Definition (2.1.5)

The *covering number*  $N(\epsilon, \mathcal{F}, \|\cdot\|)$  is the minimum number of balls  $B_\epsilon(f) := \{g \in \mathcal{F} : \|g - f\| < \epsilon\}$  such that  $\mathcal{F}$  is covered. The *entropy without bracketing* is  $\log N(\epsilon, \mathcal{F}, \|\cdot\|)$ .



## Definition (2.1.6)

The *bracketing number*  $N_{[]}(\epsilon, \mathcal{F}, \|\cdot\|)$  is the minimum number of brackets  $[l, u] := \{g \in \mathcal{F} : l \leq g \leq u\}$  with  $\|u - l\| < \epsilon$  such that  $\mathcal{F}$  is covered.


The *entropy with bracketing* is  $\log N_{[]}(\epsilon, \mathcal{F}, \|\cdot\|)$ .

# Entropy

## Definition (2.1.6)

The *bracketing number*  $N_{[]}(\epsilon, \mathcal{F}, \|\cdot\|)$  is the minimum number of brackets  $[l, u] := \{g \in \mathcal{F} : l \leq g \leq u\}$  with  $\|u - l\| < \epsilon$  such that  $\mathcal{F}$  is covered.

The *entropy with bracketing* is  $\log N_{[]}(\epsilon, \mathcal{F}, \|\cdot\|)$ .



**Figure:** Lower and upper bounds on sine functions.

**For norms such that  $|f| \leq |g|$  implies  $\|f\| \leq \|g\|$ , we have**

$$N(\epsilon, \mathcal{F}, \|\cdot\|) \leq N_{[]} (2\epsilon, \mathcal{F}, \|\cdot\|).$$

**For norms such that  $|f| \leq |g|$  implies  $\|f\| \leq \|g\|$ , we have**

$$N(\epsilon, \mathcal{F}, \|\cdot\|) \leq N_{[]} (2\epsilon, \mathcal{F}, \|\cdot\|).$$

- **There is no general reverse inequality.**

**For norms such that  $|f| \leq |g|$  implies  $\|f\| \leq \|g\|$ , we have**

$$N(\epsilon, \mathcal{F}, \|\cdot\|) \leq N_{[]} (2\epsilon, \mathcal{F}, \|\cdot\|).$$

- **There is no general reverse inequality.**
- **For the uniform norm, this is an equality.**

# Entropy and envelopes

## Definition

An *envelope function* for a class  $\mathcal{F}$  is a function  $F : \mathcal{X} \rightarrow \mathbb{R}$  with  $|f(x)| \leq F(x)$  for all  $x \in \mathcal{X}, f \in \mathcal{F}$ .



# Entropy and envelopes

## Definition

An *envelope function* for a class  $\mathcal{F}$  is a function  $F : \mathcal{X} \rightarrow \mathbb{R}$  with  $|f(x)| \leq F(x)$  for all  $x \in \mathcal{X}, f \in \mathcal{F}$ .

## Definition

The *uniform entropy numbers* are defined as

$$\sup_Q \log N(\epsilon \|F\|_{Q,r}, \mathcal{F}, L_r(Q)),$$

with the supremum over all probability measures  $Q$  on  $(\mathcal{X}, \mathcal{A})$  with  $0 < QF^r < \infty$ .

## 2.4: Glivenko-Cantelli classes

### Theorem

$\mathcal{F}$  is a  $P$ -Glivenko-Cantelli class if

$$N_{[]}(\epsilon, \mathcal{F}, L_1(P)) < \infty, \quad \text{for all } \epsilon > 0.$$

**This is a corollary of Theorem 2.4.3.**

## 2.4: Glivenko-Cantelli Theorems

### Theorem (2.4.3)

$\mathcal{F}$  is a  $P$ -Glivenko-Cantelli class if

$$\sup_Q N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) < \infty, \quad \text{for all } \epsilon > 0$$

with  $P^*F < \infty$  and

$$(X_1, \dots, X_n) \mapsto \left\| \sum_{i=1}^n e_i f(X_i) \right\|_{\mathcal{F}}$$

is measurable in the completion of  $(\mathcal{X}^n, \mathcal{A}^n, P^n)$  for every vector  $(e_1, \dots, e_n) \in \mathbb{R}^n$ .

**The measurability of**

$$(X_1, \dots, X_n) \mapsto \left\| \sum_{i=1}^n e_i f(X_i) \right\|_{\mathcal{F}}$$

**is needed for the bound**

$$\mathbb{E}^* \Phi (\| \mathbb{P}_n - P \|_{\mathcal{F}}) \leq \mathbb{E}^* \Phi (2 \| \mathbb{P}_n^o \|_{\mathcal{F}}),$$

**with  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  nondecreasing, convex,**

**The measurability of**

$$(X_1, \dots, X_n) \mapsto \left\| \sum_{i=1}^n e_i f(X_i) \right\|_{\mathcal{F}}$$

**is needed for the bound**

$$\mathbb{E}^* \Phi (\| \mathbb{P}_n - P \|_{\mathcal{F}}) \leq \mathbb{E}^* \Phi (2 \| \mathbb{P}_n^o \|_{\mathcal{F}}),$$

**with  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  nondecreasing, convex, and**

$$\mathbb{P}_n^o f = \frac{1}{n} \sum_{i=1}^n \epsilon_i f(X_i),$$

**where  $\mathbb{P}(\epsilon_1 = 1) = \mathbb{P}(\epsilon_1 = -1) = \frac{1}{2}$ .**

## 2.5: Donsker Theorems

### Theorem

$\mathcal{F}$  is a  $P$ -Donsker class if

$$\int_0^\infty \sqrt{\log N_{[]}(\epsilon, \mathcal{F}, L_2(P))} d\epsilon < \infty.$$

## 2.5: Donsker Theorems

### Theorem

$\mathcal{F}$  is a  $P$ -Donsker class if

$$\int_0^\infty \sqrt{\log N_{[]}(\epsilon, \mathcal{F}, L_2(P))} d\epsilon < \infty.$$

### Theorem (2.5.2)

$\mathcal{F}$  is a  $P$ -Donsker class if

$$\int_0^\infty \sup_Q \sqrt{\log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, L_2(Q))} d\epsilon < \infty$$

for all  $P$  such that  $P^*F^2 < \infty$ . The supremum is taken over all finitely discrete probability measures  $Q$  on  $(\mathcal{X}, \mathcal{A})$  with  $\int F^2 dQ > 0$ .

## 2.6: VC-Classes

### Definition

$\mathcal{C} \subset \mathcal{P}(\mathcal{X})$  *picks out* a subset  $A$  of  $\{x_1, \dots, x_n\} \subseteq \mathcal{X}$  if there is a  $C \in \mathcal{C}$  such that  $A = \{x_1, \dots, x_n\} \cap C$ .



## 2.6: VC-Classes

### Definition

$\mathcal{C} \subset \mathcal{P}(\mathcal{X})$  picks out a subset  $A$  of  $\{x_1, \dots, x_n\} \subseteq \mathcal{X}$  if there is a  $C \in \mathcal{C}$  such that  $A = \{x_1, \dots, x_n\} \cap C$ .

### Definition

$\mathcal{C}$  shatters  $\{x_1, \dots, x_n\}$  if  $\mathcal{C}$  picks out every  $A \subseteq \{x_1, \dots, x_n\}$ .

## 2.6: VC-Classes

### Definition

$\mathcal{C} \subset \mathcal{P}(\mathcal{X})$  picks out a subset  $A$  of  $\{x_1, \dots, x_n\} \subseteq \mathcal{X}$  if there is a  $C \in \mathcal{C}$  such that  $A = \{x_1, \dots, x_n\} \cap C$ .

### Definition

$\mathcal{C}$  shatters  $\{x_1, \dots, x_n\}$  if  $\mathcal{C}$  picks out every  $A \subseteq \{x_1, \dots, x_n\}$ .

### Definition

The VC-index  $V(\mathcal{C})$  is the smallest  $n$  such that for all  $B \subseteq \mathcal{X}$  with  $|B| \geq n$ ,  $\mathcal{C}$  does not shatter  $B$ .

## 2.6: VC-Classes

### Definition

$\mathcal{C} \subset \mathcal{P}(\mathcal{X})$  picks out a subset  $A$  of  $\{x_1, \dots, x_n\} \subseteq \mathcal{X}$  if there is a  $C \in \mathcal{C}$  such that  $A = \{x_1, \dots, x_n\} \cap C$ .

### Definition

$\mathcal{C}$  shatters  $\{x_1, \dots, x_n\}$  if  $\mathcal{C}$  picks out every  $A \subseteq \{x_1, \dots, x_n\}$ .

### Definition

The VC-index  $V(\mathcal{C})$  is the smallest  $n$  such that for all  $B \subseteq \mathcal{X}$  with  $|B| \geq n$ ,  $\mathcal{C}$  does not shatter  $B$ .

### Definition

A collection of measurable sets  $\mathcal{C}$  is a VC-class if  $V(\mathcal{C}) < \infty$ .

## 2.6: VC-Classes and Covering Numbers

**A VC-class picks out  $O(n^{V(\mathcal{C})-1})$  subsets of  $\{x_1, \dots, x_n\}$  if  $n \geq V(\mathcal{C})$ .**

## 2.6: VC-Classes and Covering Numbers

**A VC-class picks out  $O(n^{V(\mathcal{C})-1})$  subsets of  $\{x_1, \dots, x_n\}$  if  $n \geq V(\mathcal{C})$ .**

*"Some thought shows that the number of subsets picked out (...) is closely related to the covering numbers (...)"*.

## 2.6: VC-Classes and Covering Numbers

**A VC-class picks out  $O(n^{V(\mathcal{C})-1})$  subsets of  $\{x_1, \dots, x_n\}$  if  $n \geq V(\mathcal{C})$ .**

*"Some thought shows that the number of subsets picked out (...) is closely related to the covering numbers (...)"*

### Theorem (2.6.4)

*There exists a universal  $K$  such that for all VC-classes  $\mathcal{C}$  and measures  $Q$ ,*

$$N(\epsilon, \mathcal{C}, L_r(Q)) \leq KV(\mathcal{C})(4e)^{V(\mathcal{C})} \left(\frac{1}{\epsilon}\right)^{r(V(\mathcal{C})-1)}$$

*for all  $r \geq 1$  and  $0 < \epsilon < 1$ .*

## 2.6: VC-Classes and Covering Numbers

**A VC-class picks out  $O(n^{V(\mathcal{C})-1})$  subsets of  $\{x_1, \dots, x_n\}$  if  $n \geq V(\mathcal{C})$ .**

*"Some thought shows that the number of subsets picked out (...) is closely related to the covering numbers (...)"*

### Theorem (2.6.4)

*There exists a universal  $K$  such that for all VC-classes  $\mathcal{C}$  and measures  $Q$ ,*

$$N(\epsilon, \mathcal{C}, L_r(Q)) \leq KV(\mathcal{C})(4e)^{V(\mathcal{C})} \left(\frac{1}{\epsilon}\right)^{r(V(\mathcal{C})-1)}$$

*for all  $r \geq 1$  and  $0 < \epsilon < 1$ .*

**This thought requires 2.5 pages.**

## 2.6: VC-classes of Functions

### Definition

A general  $\mathcal{F}$  is called a *VC-subgraph class* if the collection of all subgraphs

$$\{(x, t) \in \mathcal{X} \times \mathbb{R} : t \leq f(x)\}$$

of  $f \in \mathcal{F}$  is a VC-class in  $\mathcal{X} \times \mathbb{R}$ .



## 2.6: VC-classes of Functions

### Definition

A general  $\mathcal{F}$  is called a *VC-subgraph class* if the collection of all subgraphs

$$\{(x, t) \in \mathcal{X} \times \mathbb{R} : t \leq f(x)\}$$

of  $f \in \mathcal{F}$  is a VC-class in  $\mathcal{X} \times \mathbb{R}$ .

**If  $\mathcal{F}$  is a VC-subgraph class, it is a Glivenko-Cantelli and Donsker class.**

## 2.7: Bracketing Numbers

**Classes of smooth functions have nice entropy properties.**

**$C_1^\alpha(\mathcal{X})$  is (roughly) the set of functions on  $\mathcal{X} \subseteq \mathbb{R}^d$  with uniformly bounded partial derivatives up to order  $\alpha$ .**

## 2.7: Bracketing Numbers

**Classes of smooth functions have nice entropy properties.**

$C_1^\alpha(\mathcal{X})$  is (roughly) the set of functions on  $\mathcal{X} \subseteq \mathbb{R}^d$  with uniformly bounded partial derivatives up to order  $\alpha$ .

**Theorem (2.7.1 and 2.7.2)**

$$\log N(\epsilon, C_1^\alpha(\mathcal{X}), \|\cdot\|_\infty) \leq K\epsilon^{-d/\alpha},$$

*and*

$$\log N_{[]}(\epsilon, C_1^\alpha(\mathcal{X}), L_r(Q)) \leq K\epsilon^{-d/\alpha}.$$

## 2.7: Bracketing Numbers

**Classes of smooth functions have nice entropy properties.**

$C_1^\alpha(\mathcal{X})$  is (roughly) the set of functions on  $\mathcal{X} \subseteq \mathbb{R}^d$  with uniformly bounded partial derivatives up to order  $\alpha$ .

**Theorem (2.7.1 and 2.7.2)**

$$\log N(\epsilon, C_1^\alpha(\mathcal{X}), \|\cdot\|_\infty) \leq K\epsilon^{-d/\alpha},$$

and

$$\log N_{[]}(\epsilon, C_1^\alpha(\mathcal{X}), L_r(Q)) \leq K\epsilon^{-d/\alpha}.$$

**Theorem (2.7.5)**

*The class  $\mathcal{F}$  of monotone functions  $f : \mathbb{R} \rightarrow [0, 1]$  satisfies*

$$\log N_{[]}(\epsilon, \mathcal{F}, L_r(Q)) \leq K\epsilon^{-1}.$$

## 2.8: Uniformity in the Underlying Distribution

**Some results can be stated uniformly in the distribution of the  $X_i$ 's.**

## 2.8: Uniformity in the Underlying Distribution

**Some results can be stated uniformly in the distribution of the  $X_i$ 's.**

### Theorem (2.8.1)

*Let  $\mathcal{F}$  be a  $P$ -measurable class of functions on a measurable space for every probability measure  $P$  in a class  $\mathcal{P}$ . Suppose that, for some measurable envelope function  $F$ ,*

$$\lim_{M \rightarrow \infty} \sup_{P \in \mathcal{P}} P\{F > M\} = 0,$$

$$\sup_{Q \in \mathcal{Q}_n} \log N(\epsilon \|F\|_{Q,1}, \mathcal{F}, L_1(Q)) = o(n), \quad \text{for every } \epsilon > 0,$$

*where the supremum is taken over the set  $\mathcal{Q}_n$  of all discrete probability measures with atoms of size integer multiples of  $1/n$ . Then  $\mathcal{F}$  is Glivenko-Cantelli uniformly in  $P \in \mathcal{P}$ .*

## 2.8: Uniformity in the Underlying Distribution

### Theorem (2.8.2)

*Let  $\mathcal{F}$  be a class of measurable functions with envelope function  $F$  that is square integrable uniformly in  $P \in \mathcal{P}$ . Then the following statements are equivalent:*

- ❶  $\mathcal{F}$  is Donsker and pre-Gaussian, both uniformly in  $P \in \mathcal{P}$ ;*
- ❷ the sequence  $\mathbb{G}_{n,P}$  is asymptotically  $\rho_P$ -equicontinuous uniformly in  $P \in \mathcal{P}$  and  $\sup_{P \in \mathcal{P}} N(\epsilon, \mathcal{F}, \rho_P) < \infty$  for every  $\epsilon > 0$ .*

## 2.9: Multiplier Central Limit Theorem

When are the following equivalent:

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n (\delta_{X_i} - P) \rightsquigarrow \mathbb{G},$$

and

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \xi_i (\delta_{X_i} - P) \rightsquigarrow \mathbb{G},$$

with convergence in  $\ell^\infty(\mathcal{F})$ .



## 2.9: Multiplier Central Limit Theorem

When are the following equivalent:

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n (\delta_{X_i} - P) \rightsquigarrow \mathbb{G},$$

and

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \xi_i (\delta_{X_i} - P) \rightsquigarrow \mathbb{G},$$

with convergence in  $\ell^\infty(\mathcal{F})$ .

### Theorem

If  $\xi_i$  are i.i.d. with zero mean and variance 1, and

$$\|\xi_1\|_{2,1} = \int_0^\infty \sqrt{P(|\xi_1| > t)} dt < \infty,$$

then it is equivalent to the  $\mathcal{F}$  being Donsker.

If  $\mathcal{F}$  and  $\mathcal{G}$  are Donsker, then so are

- 1 The convex hull of  $\mathcal{F}$ ;

If  $\mathcal{F}$  and  $\mathcal{G}$  are Donsker, then so are

- 1 The convex hull of  $\mathcal{F}$ ;
- 2  $\{\phi(f_1, \dots, f_k), f_i \in \mathcal{F}\}$  if  $\phi$  is a Lipschitz function on  $\mathbb{R}^k$ ;

If  $\mathcal{F}$  and  $\mathcal{G}$  are Donsker, then so are

- ❶ The convex hull of  $\mathcal{F}$ ;
- ❷  $\{\phi(f_1, \dots, f_k), f_i \in \mathcal{F}\}$  if  $\phi$  is a Lipschitz function on  $\mathbb{R}^k$ ;
- ❸  $\mathcal{F} + \mathcal{G}$ ,  $\mathcal{F} \wedge \mathcal{G}$ , and  $\mathcal{F} \cup \mathcal{G}$ ;

If  $\mathcal{F}$  and  $\mathcal{G}$  are Donsker, then so are

- ① The convex hull of  $\mathcal{F}$ ;
- ②  $\{\phi(f_1, \dots, f_k), f_i \in \mathcal{F}\}$  if  $\phi$  is a Lipschitz function on  $\mathbb{R}^k$ ;
- ③  $\mathcal{F} + \mathcal{G}$ ,  $\mathcal{F} \wedge \mathcal{G}$ , and  $\mathcal{F} \cup \mathcal{G}$ ;
- ④  $\mathcal{F}\mathcal{G}$  and  $\mathcal{F}/\mathcal{G}$  if they are uniformly bounded from above, or uniformly bounded away from zero.

## 2.11: Central limit theorem for Processes

**We can extend the Donsker theorem to sums of independent, but not identically distributed processes  $\sum_{i=1}^n Z_{n,i}$ .**

## 2.12: Partial-Sum Processes

**For  $s \in [0, 1]$ , consider**

$$\mathbb{Z}_n(s, f) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} (f(X_i) - Pf).$$

## 2.12: Partial-Sum Processes

**For  $s \in [0, 1]$ , consider**

$$\mathbb{Z}_n(s, f) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} (f(X_i) - Pf).$$

**It converges in  $\ell^\infty([0, 1] \times \mathcal{F})$  to a tight limit if and only if  $\mathcal{F}$  is Donsker.**



## 2.12: Partial-Sum Processes

**For  $s \in [0, 1]$ , consider**

$$\mathbb{Z}_n(s, f) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} (f(X_i) - Pf).$$

**It converges in  $\ell^\infty([0, 1] \times \mathcal{F})$  to a tight limit if and only if  $\mathcal{F}$  is Donsker.**

**It converges to the centered process  $\mathbb{Z}$  with**

$$\text{cov}(\mathbb{Z}(s, f), \mathbb{Z}(t, g)) = (s \wedge t)(Pfg - PfPg).$$

## 2.13: Other Donsker Classes

### Theorem (2.13.1)

*Any sequence  $\{f_i\}$  of square-integrable, measurable functions with  $\sum_{i=1}^{\infty} P(f_i - Pf_i)^2 < \infty$  is  $P$ -Donsker.*

## 2.13: Other Donsker Classes

### Theorem (2.13.1)

*Any sequence  $\{f_i\}$  of square-integrable, measurable functions with  $\sum_{i=1}^{\infty} P(f_i - Pf_i)^2 < \infty$  is  $P$ -Donsker.*

### Theorem (2.13.2)

*Let  $\{f_i\}$  be a sequence of measurable functions such that  $Pf_i f_j = 0$  for every  $i \neq j$  and  $\sum_{i=1}^{\infty} P f_i^2 < \infty$ . Then the class of all pointwise converging series  $\sum_{i=1}^{\infty} c_i f_i$ , such that  $\sum_{i=1}^{\infty} c_i^2 \leq 1$ , is  $P$ -Donsker.*

**By Kolmogorov's Extension Theorem, there always exists a zero-mean Gaussian process  $\{\mathbb{G}f : f \in \mathcal{F}\}$  with covariance function**

$$\mathbb{E}\mathbb{G}f_1\mathbb{G}f_2 = Pf_1f_2 - Pf_1Pf_2.$$

**By Kolmogorov's Extension Theorem, there always exists a zero-mean Gaussian process  $\{\mathbb{G}f : f \in \mathcal{F}\}$  with covariance function**

$$\mathbb{E}\mathbb{G}f_1\mathbb{G}f_2 = Pf_1f_2 - Pf_1Pf_2.$$

**$\mathcal{F}$  is pre-Gaussian if  $\mathbb{G}$  is a tight, Borel measurable map into  $\ell^\infty(\mathcal{F})$ .**

**Pre-Gaussianity is necessary for being Donsker, but not sufficient.**

**Pre-Gaussianity is necessary for being Donsker, but not sufficient.  $\mathcal{F}$  is Donsker if and only if**

$$\lim_{\delta \searrow 0} \limsup_{n \rightarrow \infty} \mathbb{P}^* \left( \sup_{\rho_P(f-g) < \delta} |\mathbb{G}_n(f-g)| > \epsilon \right) = 0,$$

**and  $\mathcal{F}$  is totally bounded in  $\mathcal{L}_2(P)$ .**

**Pre-Gaussianity is necessary for being Donsker, but not sufficient.  $\mathcal{F}$  is Donsker if and only if**

$$\lim_{\delta \searrow 0} \limsup_{n \rightarrow \infty} \mathbb{P}^* \left( \sup_{\rho_P(f-g) < \delta} |\mathbb{G}_n(f-g)| > \epsilon \right) = 0,$$

**and  $\mathcal{F}$  is totally bounded in  $\mathcal{L}_2(P)$ .**

**Here**

$$\rho_P(f) = (P(f - Pf)^2)^{1/2}.$$



**2.1.1: If  $\mathcal{F}$  is totally bounded in  $L_2(P)$ , then it is totally bounded for the seminorm  $\rho_P$ . If  $\mathcal{F}$  is totally bounded for  $\rho_P$  and  $\|P\|_{\mathcal{F}} = \sup\{|Pf| : f \in \mathcal{F}\}$  is finite, then it is totally bounded in  $L_2(P)$ .**

**2.1.2: Suppose  $\sup\{|Pf| : f \in \mathcal{F}\}$  is finite. Then a class  $\mathcal{F}$  of measurable functions is  $P$ -Donsker if and only if  $\mathcal{F}$  is totally bounded in  $L_2(P)$  and the empirical process is asymptotically equicontinuous in probability for the  $L_2(P)$ -semimetric.**

**2.1.5: If  $a_n : [0, 1] \rightarrow [0, \infty)$  is a sequence of nondecreasing functions, then there exist  $\delta_n \searrow 0$  such that  $\limsup a_n(\delta_n) = \lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} a_n(\delta)$ .**

**2.1.5: If  $a_n : [0, 1] \rightarrow [0, \infty)$  is a sequence of nondecreasing functions, then there exist  $\delta_n \searrow 0$  such that  $\limsup a_n(\delta_n) = \lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} a_n(\delta)$ .**

**Deduce that  $\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} a_n(\delta) = 0$  if and only if  $a_n(\delta_n) \rightarrow 0$  for every  $\delta_n \rightarrow 0$ .**

# Chapter 2.1: Introduction

**Geerten Koers**

Reading group *Weak Convergence and Empirical Processes*

**2019-10-07**

## Chapter 2.2: Maximal Inequalities

**Geerten Koers**

Reading group *Weak Convergence and Empirical Processes*

**2019-10-07**

## Definition

The *Orlicz norm* is defined as

$$\|X\|_{\psi} = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\},$$

with  $\psi$  non-decreasing, convex and  $\psi(0) = 0$ .

## Definition

The *Orlicz norm* is defined as

$$\|X\|_{\psi} = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\},$$

with  $\psi$  non-decreasing, convex and  $\psi(0) = 0$ .

- 1 If  $\psi(x) = x^p$ , then  $\|X\|_{\psi} = \|X\|_p = (E|X|^p)^{1/p}$ ,  $p \geq 1$ .



## Definition

The *Orlicz norm* is defined as

$$\|X\|_{\psi} = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\},$$

with  $\psi$  non-decreasing, convex and  $\psi(0) = 0$ .

- ❶ If  $\psi(x) = x^p$ , then  $\|X\|_{\psi} = \|X\|_p = (E|X|^p)^{1/p}$ ,  $p \geq 1$ .
- ❷  $\psi_p(x) = e^{x^p}$ .

$$\|X\|_\psi = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\}$$

**is indeed a norm:**

- 1  $\|X\|_\psi \geq 0;$

$$\|X\|_\psi = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\}$$

**is indeed a norm:**

- 1  $\|X\|_\psi \geq 0$ ;
- 2  $\|\lambda X\|_\psi = |\lambda| \|X\|_\psi$ ;

$$\|X\|_\psi = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\}$$

**is indeed a norm:**

- 1  $\|X\|_\psi \geq 0$ ;
- 2  $\|\lambda X\|_\psi = |\lambda| \|X\|_\psi$ ;
- 3  $\|X\|_\psi = 0$  **if and only if**  $X = 0$  **a.s.;**

$$\|X\|_\psi = \inf \left\{ C > 0 : \mathbb{E} \psi \left( \frac{|X|}{C} \right) \leq 1 \right\}$$

**is indeed a norm:**

- 1  $\|X\|_\psi \geq 0$ ;
- 2  $\|\lambda X\|_\psi = |\lambda| \|X\|_\psi$ ;
- 3  $\|X\|_\psi = 0$  **if and only if**  $X = 0$  **a.s.**;
- 4  $\|X + Y\|_\psi \leq \|X\|_\psi + \|Y\|_\psi$ .

**Exponential norms give better bounds than  $L_p$ -norms:**

$$\bullet \quad \|X\|_{\psi_p} \leq \|X\|_{\psi_q} (\log 2)^{p/q}, \quad p \leq q,$$

**Exponential norms give better bounds than  $L_p$ -norms:**

- $\|X\|_{\psi_p} \leq \|X\|_{\psi_q} (\log 2)^{p/q}, \quad p \leq q,$
- $\|X\|_p \leq p! \|X\|_{\psi_1}, \quad p \geq 1.$

## Lemma (2.2.1)

*If for all  $x \geq 0$ ,  $\mathbb{P}(|X| > x) \leq Ke^{-Cx^p}$  for some constants  $K$  and  $C$ , then*

$$\|X\|_{\psi_p} \leq ((1 + K)/C)^{1/p}.$$



## Lemma (2.2.1)

*If for all  $x \geq 0$ ,  $\mathbb{P}(|X| > x) \leq Ke^{-Cx^p}$  for some constants  $K$  and  $C$ , then*

$$\|X\|_{\psi_p} \leq ((1 + K)/C)^{1/p}.$$

**This follows from**

$$\mathbb{E}(e^{D|X|^p} - 1) = \mathbb{E} \int_0^{|X|^p} De^{Ds} ds = \int_0^\infty \mathbb{P}(|X| > s^{1/p}) De^{Ds} ds.$$

## Theorem (2.2.2)

Assume  $\psi$  is a convex, nondecreasing, nonzero function, with  $\psi(0) = 0$  and such that  $\limsup_{x,y \rightarrow \infty} \psi(x)\psi(y) / \psi(cxy) < \infty$  for some  $c \in \mathbb{R}$ . Then we have

$$\left\| \max_{1 \leq i \leq m} X_i \right\|_{\psi} \leq K \psi^{-1}(m) \max_{1 \leq i \leq m} \|X_i\|_{\psi},$$

for  $K$  depending only on  $\psi$ .

## Theorem (2.2.4)

*Assume that  $\psi$  satisfies the previous conditions. Then for any  $\eta, \delta > 0$ , we have*

$$\left\| \sup_{d(s,t) \leq \delta} |X_s - X_t| \right\|_{\psi} \leq K \left[ \int_0^{\eta} \psi^{-1}(N(\epsilon, d)) d\epsilon + \delta \psi^{-1}(N^2(\eta, d)) \right],$$

*with  $d(s, t) = \|X_s - X_t\|_{\psi}$ .*

## Definition

If the  $\psi_2$ -norms of  $X_s - X_t$  are finite, then  $\{X_t : t \in T\}$  is *sub-Gaussian*.

## Definition

If the  $\psi_2$ -norms of  $X_s - X_t$  are finite, then  $\{X_t : t \in T\}$  is *sub-Gaussian*.

**This happens if and only if**

$$\mathbb{P}(|X_s - X_t| > x) \leq 2e^{-\frac{1}{2}x^2/d^2(s,t)}, \quad x > 0.$$

## Definition

If the  $\psi_2$ -norms of  $X_s - X_t$  are finite, then  $\{X_t : t \in T\}$  is *sub-Gaussian*.

**This happens if and only if**

$$\mathbb{P}(|X_s - X_t| > x) \leq 2e^{-\frac{1}{2}x^2/d^2(s,t)}, \quad x > 0.$$

**Now the equality in the theorem can be simplified to**

$$\mathbb{E} \sup_{d(s,t) \leq \delta} |X_s - X_t| \leq K \int_0^\delta \sqrt{\log N(\epsilon, T, d)} \, d\epsilon.$$