

Country	RMNCH					Infectious diseases		
	Family Planning	Antenatal care, 4+ visits	Child immunization (DTP3)	Care seeking behaviour for ARI	Tuberculosis treatment ⁺	HIV anti-retroviral therapy ⁺	Insecticide treated nets use ¹	Access to at least basic sanitation
Afghanistan	46	24	66	68	66	10		54
Bahrain	58	≥80	≥80	≥80	≥80	63		≥80
Djibouti	51	26	59	≥80	80	31	9	67
Egypt	≥80	≥80	≥80	68	60	40		≥80
Iran (Islamic Republic of)	76	≥80	≥80	76	60	30		≥80
Iraq	57	68	78	44	55	29		≥80
Jordan	57	≥80	77	61	47	53		≥80
Kuwait	67	≥80	≥80	≥80	≥80	58		≥80
Lebanon	62	≥80	67	74	≥80	66		≥80
Libya	38	≥80	73	≥80	49	41		≥80
Morocco	75	61	≥80	70	≥80	≥80		≥80
Oman	39	74	≥80	56	≥80	65		≥80
Pakistan	51	52	≥80	71	55	14		69
Qatar	64	≥80	≥80	≥80	≥80	58		≥80
Saudi Arabia	46	80	≥80	≥80	≥80	79		≥80
Somalia	4	24	42	22	41	50	18	40
Sudan	34	51	≥80	48	69	27	51	37
Syrian Arab Republic	61	64	48	77	≥80	38		≥80
Tunisia	70	≥80	≥80	≥80	59	29		≥80
United Arab Emirates	60	≥80	≥80	≥80	≥80	≥80		≥80
Yemen	48	25	72	34	59	31		55

Note: Values in italics correspond to imputed values. Country index values of 80 and over are reported as ≥80 for presentation purposes and to avoid comparisons that are not meaningful given the inability of the index to adequately distinguish between countries with high level of service coverage provision.

* Proxy indicators; + Values are rescaled for incorporation into the index calculations.

1 Pertains only to countries with highly endemic malaria in sub-Saharan Africa.

Legend

Very high coverage (80+)
High coverage (60-79)
Medium coverage (40-59)
Low coverage (20-39)
Very low coverage (<20)