	RMNCH				Infectious diseases			
Country	Family Planning	Antenatal care, 4+ visits	Child immunization (DTP3)	Care seeking behaviour for ARI	Tuberculosis treatment [±]	HIV anti-retroviral therapy [†]	Insecticide treated nets use1	Access to at least basic sanitation
Bangladesh	73	37	≥80	46	≥80	31		59
Bhutan	80	80	≥80	74	67	42		78
Democratic People's Republic of Korea	≥80	≥80	41	≥80	66	18		≥80
India	76	58	≥80	56	67	65		74
Indonesia	≥80	≥80	67	<i>7</i> 5	45	28		≥80
Maldives	33	≥80	≥80	74	44	23		≥80
Myanmar	78	59	37	59	33	70		74
Nepal	62	78	≥80	≥80	41	72		≥80
Sri Lanka	74	≥80	≥80	52	48	66		≥80
Thailand	≥80	≥80	≥80	80	70	≥80		≥80
Timor-Leste	52	77	≥80	70	50	55		57

Note: Values in italics correspond to imputed values. Country index values of 80 and over are reported as ≥80 for presentation purposes and to avoid comparisons that are not meaningful given the inability of the index to adequately distinguish between countries with high level of service coverage provision.

- * Proxy indicators; + Values are rescaled for incorporation into the index calculations.
- 1 Pertains only to countries with highly endemic malaria in sub-Saharan Africa.

Legend

Very high coverage (80+)

High coverage (60-79) Medium coverage (40-59) Low coverage (20-39)

Very low coverage (<20)