

Photoionization Cross Sections in EXHALE: H I, He I, He II, and the He I 2^3S Triplet, with Benchmarks against Verner+1996 and Norcross 1971

Contents

1. Overview	1
2. The four cross sections	2
2.1 Hydrogenic atoms: H I and He II	2
2.2 He I ground state (1^1S)	2
2.3 He I 2^3S metastable (two VFKY96 forms + bridge)	2
2.4 Verner+1996 single-shell fit (VFKY96)	3
3. Overview comparison	3
4. He I ground state: EXHALE two-term fit vs Verner+1996	4
5. He I 2^3S: EXHALE broken power law vs p-winds/Norcross	4
5.1 Cross-check against TOPbase / Opacity Project	5
5.2 Does the $> 59\text{eV}$ truncation matter for the rate?	6
5.3 Impact of the autoionizing resonances	7
5.4 A smooth high-energy extension ($E > 59\text{eV}$)	9
5.5 A unified full-range representation from TOPbase	9
5.6 Recommended minimal-change extension	10
5.7 He 2^3S as two VFKY96 (Verner) forms	11
6. Summary	12
7. Reproducing this comparison	14

1. Overview

EXHALE computes the photoionization rate of each species by integrating its energy-dependent cross section $\sigma(E)$ against the local stellar spectrum on a fixed energy grid (the cross sections are pre-tabulated in `set_energy_vectors.f90`). All cross-section routines live in `src/modules/functions/cross_sec.f90` (module `Cross_sections`) and return σ in units of $1\text{Mb} \equiv 10^{-18}\text{cm}^2$, with photon energy E in eV.

This memo (i) collects the four atomic cross sections relevant to the He 10830 / H α / Ly α diagnostics — H I, He I, He II, and the metastable He I 2³S triplet — and (ii) benchmarks the two *fitted* helium cross sections against independent references:

- **He I ground state:** the EXHALE two-term ATES fit (`sigma_HeI`) versus the Verner, Ferland, Korista & Yakovlev (1996; hereafter VFKY96) single-shell analytic fit, over 24.6–500 eV.
- **He I 2³S:** the EXHALE broken power-law fit (`sigma_HeI23S`) versus the tabulation of Norcross (1971) data shipped with the p-winds code, over 4.77–60 eV.

All numbers and figures below are produced by the companion script `docs/compare_photoion_cross_sections.py` (§7.), which ports the Fortran routines to Python verbatim and writes vector-PDF figures into `docs/figures/`.

2. The four cross sections

2.1 Hydrogenic atoms: H I and He II

H I ($Z = 1$) and He II ($Z = 2$) use the analytic hydrogenic cross section (`sigma(E, Z)`), with threshold $E_0 = 13.6Z^2$ eV. For $E > E_0$,

$$\sigma(E) = \frac{6.3}{Z^2} \left(\frac{E_0}{E} \right)^4 \frac{\exp[4 - 4 \arctan(\epsilon)/\epsilon]}{1 - \exp(-2\pi/\epsilon)} \text{ Mb}, \quad \epsilon = \sqrt{E/E_0 - 1}, \quad (1)$$

which is the exact non-relativistic result with the Gaunt-factor correction; at threshold $\sigma = 6.3/Z^2$ Mb (6.3 Mb for H I, 1.575 Mb for He II).

2.2 He I ground state (1¹S)

By default the He I ground state (`sigma_HeI`) uses the Verner+1996 (VFKY96) single-shell fit — the form of §2.4 with the He I parameters of Table 1:

$$\sigma_{\text{HeI}}(E) = \sigma_{\text{VFKY96}}(E; E_{\text{th}}=24.59, E_0=13.61, \sigma_0=949.2, y_a=1.469, P=3.188, y_w=2.039, y_0=0.4434, y_1=2.136). \quad (2)$$

Setting `ATES_photoionization_rate: True` in `input.inp` reverts to the legacy ATES two-term fit (valid above $E_{\text{th}} = 24.6$ eV):

$$\sigma_{\text{HeI}}^{\text{ATES}}(E) = \frac{0.6935}{(E/100)^{1.82} + (E/100)^{3.23}} \text{ Mb}. \quad (3)$$

(The 2³S metastable *triplet* is a separate species, Eq. 4 below; Eqs. 2–3 are the 1¹S *singlet* ground state only.)

2.3 He I 2³S metastable (two VFKY96 forms + bridge)

The metastable triplet — the lower level of He 10830 — (`sigma_HeI23S`) is built from *two* Verner+1996 (VFKY96, §2.4) single-shell forms joined by a log-linear (power-law) bridge, used piecewise. With $\ell \equiv \log_{10}(E/\text{eV})$ and the node

energies $E_3=34.70$, $E_4=45.59$ eV ($x_3 \equiv \log_{10} E_3$, $x_4 \equiv \log_{10} E_4$):

$$\sigma_{\text{He } 2^3\text{S}}(E) = \begin{cases} \sigma_{\text{VFKY96}}(E; \text{wing A}) & E \leq E_3 \quad (\text{threshold} \rightarrow \text{Cooper min}) \\ 10^{m_b(\ell-x_3)+y_3^b} & E_3 < E < E_4 \quad (\text{power-law bridge}) \\ \sigma_{\text{VFKY96}}(E; \text{wing B}) & E \geq E_4 \quad (\text{bump} \rightarrow \text{tail}) \end{cases} \quad (4)$$

Wing A and wing B use the parameters of Table 8 (§5.7; each a $\sim 4\%$ fit to the resonance-averaged cross section), and the bridge is the straight log–log line joining $y_3^b = \log_{10} \sigma_{\text{VFKY96}}(E_3; A)$ to $y_4^b = \log_{10} \sigma_{\text{VFKY96}}(E_4; B)$ with $m_b = (y_4^b - y_3^b) / (x_4 - x_3)$. Wing A vanishes below the 4.78 eV threshold (its E_{th}), and wing B supplies the smooth high- E tail ($\sim E^{-7/2}$), so there is no hard cutoff. The construction is continuous at both E_3 and E_4 and reproduces the former broken power law to $\sim 4\%$ (deepest at the bump, 2.57 vs 2.68 Mb).

The *former* five-segment broken power law (Norcross-1971 fit + TOPbase-tail extension of §5.6), retained commented-out in the code, was $\log_{10} \sigma = -0.8134\ell + 1.240$ (4.78–7.49), $-1.772\ell + 2.078$ (7.49–34.70), $9.098(\ell - x_3) - 0.6515$ (34.70–45.59, the bump rise), $-2.660\ell + 4.841$ (45.59–70), and $-2.996\ell + 5.462$ (> 70 eV, the TOPbase tail); $\sigma=0$ below 4.78 eV.

2.4 Verner+1996 single-shell fit (VFKY96)

This is the VFKY96 analytic fit, the form EXHALE adopts for metals, for He II¹ and — now by default — for the He I ground state (sigma_VFKY96); for $E \geq E_{\text{th}}$,

$$x = E/E_0 - y_0, \quad z = \sqrt{x^2 + y_1^2}, \quad Q = 5.5 - P/2, \quad (5)$$

$$\sigma(E) = \sigma_0 [(x-1)^2 + y_w^2] z^{-Q} (1 + \sqrt{z/y_a})^{-P} \text{ Mb}. \quad (6)$$

Verner+1996 Table 1 provides ground-shell parameters for *all* ions $Z = 1\text{--}30$, including HI, He I and He II (Table 1); the He I 2³S excited level is *not* in that table, which is why a separate Norcross-based fit is required for the triplet.

Table 1: VFKY96 ground-shell parameters (Verner+1996 Table 1) used here. σ_0 in Mb.

ion	E_{th}	E_0	σ_0	y_a	P	y_w	(y_0, y_1)
HI	13.60	0.4298	5.475×10^4	32.88	2.963	0.0	(0, 0)
He I	24.59	13.61	949.2	1.469	3.188	2.039	(0.4434, 2.136)
He II	54.42	1.720	1.369×10^4	32.88	2.963	0.0	(0, 0)

3. Overview comparison

Figure 1 shows the four EXHALE-adopted cross sections on a common log–log scale. The ordering of thresholds (He 2³S at 4.78, HI at 13.6, He I at 24.6, He II at 54.4 eV) and the peak magnitudes ($\sim 5\text{--}8$ Mb near threshold for HI, He I and He 2³S; 1.6 Mb for He II) are evident. The He 2³S curve displays the characteristic minimum near 35 eV and the rise toward 45 eV inherited from the Norcross data.

¹ via the hydrogenic routine




Figure 1: Photoionization cross sections adopted in EXHALE for H I, He I, He II and the metastable He I 2³S triplet. Dotted vertical lines mark the ionization thresholds.

4. He I ground state: EXHALE two-term fit vs Verner+1996

Figure 2 and Table 2 compare the EXHALE two-term fit with the VFKY96 He I cross section over 24.6–500 eV. The two agree to within $\sim 14\%$ everywhere: the fits match within a few percent at threshold and at high energy, with the two-term form running $\sim 12\text{--}14\%$ *below* VFKY96 in the 50–100 eV range (the band that dominates He photoionization in a typical XUV field). The median ratio over the interval is 0.94.

Because both are accurate fits to the same underlying He I data, the $\leq 14\%$ difference is well within the photoionization-rate uncertainty budget. Replacing `sigma_HeI` with the VFKY96 He I parameters of Table 1 (a one-line wrapper around the existing `sigma_VFKY96`) would unify He I with the metal and He II treatment and slightly improve the 50–100 eV band, but is not required for the ionization balance.

5. He I 2³S: EXHALE broken power law vs p-winds/Norcross

The p-winds code (dos Santos et al. 2022) ships the He 2³S photoionization cross section as a 23-point table of Norcross (1971), converting a differential oscillator strength $df/d\epsilon$ to a cross section via $\sigma[\text{cm}^2] = 8.0670 \times 10^{-18} (df/d\epsilon)$. Crucially, the EXHALE broken-power-law breakpoints coincide *exactly* with the Norcross wavelength nodes, confirming that `sigma_HeI23S` is a fit to this same data set.

Figure 3 overlays the EXHALE curve, the tabulated Norcross points, and their log–log interpolation; Table 3 lists the comparison. The agreement is excellent: the broken power law reproduces the table to within $\sim 10\%$ across 4.78–59 eV, with a median ratio of 1.00. The only localized deviation ($\sim 15\%$) occurs right at the Cooper minimum near

Table 2: He I cross section (Mb): EXHALE two-term fit vs Verner+1996.

E [eV]	EXHALE	Verner	ratio
24.6	7.821	7.430	1.053
30	5.244	5.361	0.978
50	1.779	2.022	0.880
100	0.347	0.394	0.880
200	0.054	0.055	0.973
300	0.0165	0.0160	1.028
500	0.0035	0.0032	1.089
ratio over 24.6–500 eV: min 0.861, max 1.089, median 0.939			

30 eV, where the cross section is smallest and therefore least important for the rate.

Table 3: He I 2³S cross section (Mb): EXHALE broken PL vs p-winds/Norcross. (At threshold 4.78 eV the table interpolation is undefined; EXHALE gives 4.867 Mb, matching the Norcross threshold value $8.0670 \times 0.605 = 4.881$ Mb.)

E [eV]	EXHALE	p-winds	ratio
6	4.046	4.092	0.989
8	3.006	3.110	0.966
12	1.465	1.463	1.002
20	0.593	0.546	1.085
30	0.289	0.277	1.042
45	2.382	2.442	0.975
55	1.517	1.516	1.001
59	1.226	1.251	0.980
ratio over 4.78–59 eV: min 0.853, max 1.095, median 1.000			

5.1 Cross-check against TOPbase / Opacity Project

As an independent, fully *ab initio* reference we extracted the He I 2³S ($1s2s^3S$) photoionization cross section from the **TOPbase** (Opacity Project) R-matrix database at the CDS server. The relevant level is NZ = 2, NE = 2, iSLP = 300 (³S, even parity), iLV = 1; the table has 922 points with the photon energy in Rydberg and σ in Mb (level energy -0.35032 Ry, i.e. ionization threshold 4.766 eV). The query URL and parsing are in the companion script (§7.); the raw table is saved as docs/data/topbase_HeI_2_3S_photo.dat.

Two points must be kept in mind when comparing. (i) The TOPbase threshold sits at the *theoretical* energy (4.63 eV for the first point), ~ 0.14 eV below the experimental 4.77 eV; this is the usual OP threshold offset. (ii) Above ~ 35 eV TOPbase resolves a dense forest of **autoionizing resonances** (doubly-excited states converging on the excited He II




Figure 2: He I ground-state photoionization cross section: EXHALE two-term fit (sigma_HeI, green) vs the Verner+1996 single-shell fit (black dashed). Bottom panel: ratio EXHALE/Verner.

core), with peaks reaching $\sigma \simeq 3200 \text{ Mb}$. These are real but extremely narrow, and the smooth Norcross/EXHALE fits represent only their resonance-averaged background.

Figure 4(a) shows the full range with the resonance structure; panel (b) zooms into the smooth near-threshold region (4.6–25 eV). In that smooth region the three independent descriptions agree to within $\sim 8\%$ (Table 4): EXHALE/TOPbase has min 0.92, max 1.07, median 1.00 over 5–20 eV. This confirms that the EXHALE broken power law — and the Norcross data it is fit to — reproduces the OP R-matrix background where it matters for the He 2³S population.

Two caveats remain for completeness: (a) EXHALE sets $\sigma_{\text{He } 2^3\text{S}} = 0$ above 59.2 eV, whereas TOPbase shows a smooth $\sim E^{-3}$ tail continuing to several hundred eV (small in magnitude, $\lesssim 0.1 \text{ Mb}$); (b) the broken-power-law “bump” near 45 eV is a smooth stand-in for the resonance complex and should not be read as a true cross section at any single energy there.

5.2 Does the $> 59 \text{ eV}$ truncation matter for the rate?

The relevant quantity for the He 2³S population (and hence He 10830) is the *photoionization rate*, $\Gamma_{2^3\text{S}} \propto \int \sigma_{2^3\text{S}}(E) N(E) dE$, where $N(E) = J_\nu / h\nu \propto J_{\text{inc}}(E)/E$ is the photon-number flux. In the EXHALE power-law SED, $J_{\text{inc}} \propto E^{\text{PLind}}$ over the EUV band $[e_{\text{low}}, e_{\text{mid}}] = [4.8, 124] \text{ eV}$ (normalized to L_{EUV}) and the X-ray band $[124, 1240] \text{ eV}$ (to L_X), so $N(E) \propto E^{\text{PLind}-1}$. Because both $N(E)$ and $\sigma_{2^3\text{S}}$ fall steeply, the integrand peaks at the threshold; the rate is dominated by the near-threshold UV ($\sim 4.8\text{--}15 \text{ eV}$), where σ is largest (1–5 Mb) and photons are most numerous.




Figure 3: He I 2³S photoionization cross section: EXHALE broken power law (sigma_HeI23S, red) vs the p-winds log–log interpolation (blue dashed) of the Norcross (1971) table (black points). Bottom panel: ratio EXHALE/p-winds.

Table 5 gives the relative increase of Γ_{2^3S} when σ_{2^3S} is extended above 59.18 eV with the TOPbase tail (integrated to 323 eV). The effect is only 0.1–1.2% across realistic spectral indices (PLind = −1 to −2) and X-ray/EUV ratios; for the EUV slopes typical of cool stars (PLind $\simeq$ −1.3 to −1.6) it is 0.3–0.6%. This is far below the $\sim$ 10% uncertainty of the cross-section fit itself (§5.) and of the composition/SED. The 35–59 eV resonance “bump” likewise carries only $\sim$ 2% of the current rate.

Moreover, these are *upper bounds*: the power-law SED extrapolates the EUV slope down to 4.8 eV, whereas a real star has a much larger photospheric near-UV flux (4.8–13.6 eV); with a realistic SED the near-threshold region dominates even more and the > 59 eV contribution shrinks further. (Heating is also unaffected: the 2³S number density is negligible compared with the H/He ground states, so its photoionization deposits negligible energy regardless of the high-energy tail.) We therefore retain the 59 eV cutoff; extending it is a cosmetic, sub-percent refinement.

5.3 Impact of the autoionizing resonances

Between $\sim$ 40 and 55 eV, TOPbase shows a dense forest of autoionizing resonances (doubly-excited states converging on the He II $n = 2$ threshold at 65.4 eV), with peaks reaching $\sigma \simeq 3223$ Mb, whereas the EXHALE broken power law represents this region by a smooth bump peaking at only $\sim$ 2.4 Mb (Fig. 4a). It is tempting to ask whether the resonances, being so tall, substantially raise the He 2³S photoionization rate.

They do not — but the analysis requires care, because *the raw TOPbase table must not be integrated directly*. Integrat-

Table 4: He I 2³S cross section (Mb) in the smooth near-threshold region: EXHALE vs Norcross/p-winds vs TOPbase.

E [eV]	EXHALE	p-winds	TOPbase	EX/TB	PW/TB
5.0	4.693	4.671	5.081	0.924	0.919
6.0	4.046	4.092	4.231	0.956	0.967
8.0	3.006	3.110	2.866	1.049	1.085
10.0	2.024	2.069	2.025	1.000	1.022
12.0	1.465	1.463	1.488	0.985	0.983
16.0	0.880	0.872	0.877	1.004	0.994
20.0	0.593	0.546	0.574	1.033	0.952
EXHALE/TOPbase over 5–20 eV: min 0.924, max 1.066, median 0.999					

Table 5: Relative increase (%) of the He 2³S photoionization rate when σ_{2^3S} is extended above 59 eV with the TOPbase tail, for an EXHALE power-law SED of index PLind and X-ray/EUV luminosity ratio L_X/L_{EUV} .

PLind	$L_X/L_{EUV} = 0.1$	0.3	1.0
−1.0	1.13%	1.15%	1.21%
−1.3	0.57%	0.59%	0.65%
−1.6	0.29%	0.30%	0.36%
−2.0	0.12%	0.13%	0.19%

ing both cross sections on the *same* native TOPbase energy grid (so the quadrature is identical) gives the total-rate ratio TOPbase/EXHALE in Table 6. With no clipping the resonances appear to raise the rate by $\sim 13\%$ (at PLind = −1.3); but this increase is produced almost entirely by the 2–4 tallest grid points. The narrowest resonances ($\Delta E \sim 0.01$ eV) are sampled by only one or two points, so the trapezoidal rule assigns them a triangular area fixed by the arbitrary mesh spacing rather than by the physical Lorentzian width. Clipping σ at $\gtrsim 100$ Mb removes just these few unresolved spikes and the ratio collapses to 0.99–1.02: **the robust net effect of the resonances on the rate is only ~ 1 –2%.**

Physically this is expected: although individual resonances are very tall, they are extremely narrow and therefore intercept little photon flux. The resonance-averaged cross section $\langle \sigma \rangle = \sigma_{bg} + \sum_i (\text{resonance area})_i / \Delta E$ sits only modestly above the smooth background, which the few-Mb Norcross/EXHALE values already approximate. The net resonance effect (~ 1 –2%) is comparable to the > 59 eV tail (§5.2) and well below the $\sim 10\%$ fit and composition/SED uncertainties, so it does not change the He 10830 conclusions.

Practical caveat: for percent-level work one must use a properly resonance-averaged Opacity-Project cross section (e.g. the resonance-averaged tabulations in the spirit of Bautista et al.); the raw 922-point TOPbase table should never be fed into a broadband rate integral, as the quadrature artifact in Table 6 illustrates.




Figure 4: He I 2³S photoionization: EXHALE broken power law (red), Norcross/p-winds table (blue points), and TOPbase/OP (gray). **(a)** Full range: TOPbase resolves autoionizing resonances above ~ 35 eV (peaks to ~ 3200 Mb) that the smooth fits average over; EXHALE is truncated to zero above 59 eV. **(b)** Smooth near-threshold region: the three agree to within $\sim 8\%$.

5.4 A smooth high-energy extension ($E > 59$ eV)

Although the > 59 eV truncation is dynamically harmless (§5.2), a smooth analytic extension is trivial to provide and removes the (cosmetic) hard cutoff. Above the He II $n = 2$ resonance limit ($E \gtrsim 66$ eV) the TOPbase cross section is smooth and monotonic, and a single power law fits the resonance-free background to $\sim 5\%$ over 60–320 eV (Fig. 5):

$$\sigma_{\text{He } 2^3\text{S}}(E) \simeq 0.294 \left(\frac{E}{100 \text{ eV}} \right)^{-2.99} \text{ Mb} = 10^{5.45} E^{-2.99} \text{ Mb} \quad (E > 59 \text{ eV}), \quad (7)$$

with E in eV. The index -2.99 is close to the asymptotic hydrogenic s -shell value ($-7/2$ is approached only at much higher E), and the fit residuals are ≤ 0.023 dex. This is the resonance-free background by construction: it is fit only to points above the $n = 2$ resonance limit and deliberately ignores the 40–55 eV resonance forest (§5.3).

To splice it into `sigma_HeI23S`, replace the “zero above x_5 ” branch with one more log-linear segment of slope -2.99 . Two variants: (i) *data-matched* — use the boxed normalization, which reproduces TOPbase to 5% but sits $\sim 16\%$ above the broken-PL endpoint at the 59.2 eV junction (a step at $\sigma \sim 1.4$ Mb, irrelevant to the rate); (ii) *continuity-matched* — anchor at the broken-PL value $\sigma(59.2) = 1.21$ Mb, i.e. $\sigma = 1.21 (E/59.2)^{-2.99}$ Mb, which is continuous but runs $\sim 15\%$ below TOPbase. Either is adequate; given the $< 1\%$ rate impact, the continuity-matched form is the tidier drop-in.

5.5 A unified full-range representation from TOPbase

The power-law tail of §5.4 only patches the high-energy end. A cleaner option is a single representation that follows the TOPbase *resonance-averaged* background over the whole range, 4.77–320 eV, with no ad-hoc junction. Because the cross section has a deep Cooper minimum near 30 eV, a resonance-averaged enhancement (the He II $n = 2$ channel) peaking near 50 eV, and an E^{-3} tail, no single compact analytic form captures it. Instead we tabulate a short (E, σ) node

Table 6: Convergence/spike test of the resonance contribution. Total He 2³S rate ratio TOPbase/EXHALE (PLind = −1.3), both integrated on the same native grid, as the TOPbase cross section is clipped at an upper value; “pts>cap” counts the grid points removed.

σ clip [Mb]	TOPbase/EXHALE	pts > cap
none (raw)	1.134	0
500	1.024	2
200	1.001	2
100	0.993	4
50	0.989	7

table and interpolate it **log–log with a monotone cubic (PCHIP)** — exactly the scheme EXHALE already uses for the metal cooling fits, so it is a drop-in. Below 30eV and above 66eV the nodes are the clean TOPbase background; in 30–66eV they are the *resonance-averaged* (spike-capped) values, i.e. the sharp resonances are deliberately averaged over rather than followed.

Table 7: Node table for the unified He 2³S photoionization cross section (log–log PCHIP interpolation; $\sigma = 0$ outside the range).

E [eV]	4.77	6	8	10	13.6	20	30	40	52	63	80	100	150	200	300
σ [Mb]	4.88	4.09	3.11	2.07	1.16	0.55	0.273	0.50	1.9	1.14	0.577	0.294	0.085	0.037	0.0111

This representation (Fig. 6) reproduces the TOPbase smooth background to within $\sim 10\%$ across the full range (the residual at the very threshold is the ~ 0.14 eV theoretical-threshold offset, §5.1), is C^1 -continuous by construction, and avoids the hard cutoff. Crucially, the He 2³S photoionization *rate* it produces differs from the current broken power law by only +1.7% (PLind = −1.3; consistent with §5.2 and §5.3), confirming that the upgrade is a fidelity/tidiness improvement rather than a change that moves the He 10830 prediction. The 30–66eV nodes carry the largest uncertainty ($\sim 30\%$, from the resonance averaging) but the least weight.

Recommendation. For a clean, well-connected cross section over all energies, adopt the Table 7 node set with log–log PCHIP in sigma_HeI23S. It is faithful to the OP R-matrix background, smooth everywhere, and changes the rate by $< 2\%$. If instead only the cutoff is to be removed with minimal code change, the continuity-matched power-law tail of §5.4 suffices — or, better, the minimal-change extension of §5.6 below.

5.6 Recommended minimal-change extension

The tidiest upgrade keeps essentially all of the existing broken power law and changes only its *last* segment. The procedure is:

1. Keep the broken power law unchanged up to the resonance-averaged bump peak at x_4 (45.6eV; $\sigma = 2.68$ Mb) — i.e. segments a_1 , a_2 and the rising segment are untouched.




Figure 5: Smooth high-energy extension of $\sigma_{\text{He } 2^3\text{S}}$. The power-law fit (red) to the resonance-free TOPbase background (blue points, $E \geq 66\text{ eV}$) follows $\sigma \propto E^{-2.99}$ to $\sim 5\%$ up to 320 eV . The current EXHALE broken power law (black dashed) rises to a resonance-averaged bump near 45 eV and is cut to zero above 59.2 eV (dotted line).

2. Fit the smooth TOPbase background above $E_J = 70\text{ eV}$ with a power law:

$$\sigma_{\text{tail}}(E) = 10^{5.462} E^{-2.996} \text{ Mb} \quad (E > 70\text{ eV}), \quad (8)$$

($\sigma_{\text{TOPbase}}(70\text{ eV}) = 0.858\text{ Mb}$; fit residuals $\leq 3\%$).

3. Re-aim the last power-law segment so it runs from the bump peak at x_4 to exactly $\sigma_{\text{tail}}(70\text{ eV})$ at 70 eV : its slope changes only slightly, from -3.039 to -2.658 (intercept $c_3 = 4.838$ in $\log_{10} \sigma = a'_3 \log_{10} E + c_3$). Above 70 eV use σ_{tail} ; drop the old 59 eV cutoff.

By construction the result is continuous at *both* joins — at 45.6 eV (with the kept rising segment) and at 70 eV (with the TOPbase tail) — so there is no step and no kink beyond the tiny 0.38 -in-slope change. It reproduces the TOPbase background to $\sim 3\%$ above 70 eV and tracks Norcross/p-winds throughout (Fig. 7), while changing the He 2^3S photoionization rate by only $+0.65\%$ (PLind = -1.3) — the smallest of all the options. This is the recommended drop-in: it is a one-line slope change plus a new tail branch in `sigma_HeI23S`, retains the validated threshold-to-bump shape exactly, and removes the unphysical cutoff.

5.7 He 2^3S as two VFKY96 (Verner) forms

It is natural to ask whether the He 2^3S cross section could use the same VFKY96 fits (§2.4) EXHALE already applies to He I and the metals, instead of the bespoke broken power law. Fitting a *single* VFKY96 term to each smooth part of the curve works very well (Fig. 8, Table 8): segments 1–2 (threshold $\rightarrow$ Cooper minimum, 4.85 – 34.7 eV) and segments 4–5 (resonance-averaged bump $\rightarrow$ tail, 45.7 – 320 eV) are *each* reproduced to $\sim 4\%$. This is physically sensible — the two terms are the $2s$ -shell channel and the (resonance-averaged) He II $n = 2$ channel.




Figure 6: Unified full-range representation (red) of $\sigma_{\text{HeI}2^3\text{S}}$: log–log PCHIP through the node table (black points; Table 7) following the TOPbase resonance-averaged background (gray, with resonances). The Norcross (1971) / p-winds table (green squares) and the current EXHALE broken power law (blue dashed; cut to zero above 59 eV) are overplotted. Bottom: ratio of the unified fit to the TOPbase smooth-background points (gray) and to the Norcross points (green); both within $\sim \pm 15\%$, with the larger Norcross scatter at 40–55 eV reflecting its coarse sampling of the resonance region.

The one region the two wings do not cover is the *transition* (34.7–45.6 eV, shaded in Fig. 8), where the cross section climbs from the Cooper minimum to the bump. This is bridged by a short log-linear (power-law) segment joining wing A at $E_3 = 34.70$ eV to wing B at $E_4 = 45.59$ eV. **This two-VFKY96-plus-bridge form is now the EXHALE implementation** of $\sigma_{\text{HeI}2^3\text{S}}$ (Eq. 4); it reproduces the former broken power law to $\sim 4\%$ (deepest at the bump, 2.57 vs 2.68 Mb), is continuous at E_3 and E_4 , and reuses the same σ_{VFKY96} routine as He I and the metals. The former broken power law is retained, commented out, in the code.

6. Summary

- EXHALE uses three distinct prescriptions: an exact hydrogenic form for H I and He II, Verner+1996 (VFKY96) for the He I ground state (default; legacy two-term fit behind `ATES_photoionization_rate`), and — for the He I 2³S triplet — **two VFKY96 forms joined by a power-law bridge** (§5.7; formerly a Norcross-1971 broken




Figure 7: Recommended minimal-change extension (red): the original broken power law up to the 45.6 eV bump, a re-aimed last segment (slope $-3.039 \rightarrow -2.658$) reaching the TOPbase fit at 70 eV (dotted line), and the TOPbase power-law tail above. Norcross/p-winds (green) and TOPbase (gray) are overplotted; the original broken power law (blue dashed) cuts to zero at 59 eV. The curve is continuous at both joins.

Table 8: Single-VFKY96 fits to the two smooth parts of He 2³S (form of §2.4; σ_0 in Mb, energies in eV). Each reproduces its region to $\sim 4\%$.

region	E_{th}	E_0	σ_0	y_a	P	y_w	y_0	y_1
seg 1–2 (4.85–34.7)	4.78	2.645	20.8	1.0×10^{12}	3.42	2.681	1.956	2.603
seg 4–5 (45.7–320)	45.59	49.68	1052	0.0439	2.941	1.717	5.5×10^{-5}	1.118

power law, retained commented-out in the code).

- The He I two-term fit agrees with Verner+1996 to within 14% (24.6–500 eV; median 0.94), running slightly low at 50–100 eV.
- The He I 2³S broken power law reproduces its parent Norcross/p-winds table to within $\sim 10\%$ (4.78–59 eV; median 1.00), including the Cooper minimum and the rise near 45 eV.
- An independent check against the **TOPbase / Opacity Project** R-matrix cross section (§5.1) confirms agreement to $\sim 8\%$ in the smooth near-threshold region (5–20 eV; median 1.00). TOPbase additionally shows autoionizing resonances above ~ 35 eV (averaged over by the fits) and a smooth tail above 59 eV (where EXHALE truncates to zero).
- The 40–55 eV autoionizing resonances raise the He 2³S rate by only $\sim 1\text{--}2\%$ once the few numerically under-




Figure 8: He 2³S (black: implemented broken power law + TOPbase tail) with single-VFKY96 fits to segments 1–2 (blue dashed) and 4–5 (red dash-dot), each good to ~4%. The shaded band (34.7–45.6 eV) is the Cooper-minimum/rise transition where the two must be spliced piecewise; a simple sum fails there.

resolved spikes are excluded (§5.3); the raw TOPbase table must not be integrated directly.

- A simple power law $\sigma_{\text{He } 2^3\text{S}} \simeq 0.294(E/100\text{ eV})^{-2.99}\text{ Mb}$ reproduces the resonance-free TOPbase background to ~5% and provides a clean smooth extension above the 59 eV cutoff (§5.4). For a single well-connected cross section over the *whole* range, a 15-node log–log PCHIP table (Table 7, §5.5) follows the TOPbase resonance-averaged background to ~10% and changes the rate by only +1.7%. The *recommended* drop-in (§5.6) keeps the broken power law intact up to the 45.6 eV bump and only re-aims its last segment (slope $-3.039 \rightarrow -2.658$) to meet the TOPbase tail $10^{5.462}E^{-2.996}\text{ Mb}$ at 70 eV: continuous at both joins, 3% to TOPbase, and +0.65% in rate.
- Both fitted helium cross sections are adequate for the photoionization balance. Extending the He 2³S fit above 59 eV with the TOPbase tail changes its photoionization rate by only 0.1–1.2% (§5.2), so the cutoff is harmless; the one worthwhile refinement, if any, is to migrate He I to the VFKY96 form (Table 1) for internal consistency.

7. Reproducing this comparison

All figures and numbers are generated by docs/compare_photoion_cross_sections.py:

```
cd EXHALE/docs
python3 compare_photoion_cross_sections.py
# -> figures/xsec_overview.pdf
#   figures/xsec_HeI_vs_Verner.pdf
#   figures/xsec_HeI23S_vs_pwinds.pdf
```

```
# figures/xsec_HeI23S_topbase.pdf
# figures/xsec_HeI23S_tail.pdf
# figures/xsec_HeI23S_unified.pdf
# figures/xsec_HeI23S_ext.pdf
# figures/xsec_HeI23S_vfky.pdf
# reads: data/topbase_HeI_2_3S_photo.dat
```

The TOPbase table was retrieved (once) from the CDS Opacity Project server with

```
curl -L -G "https://cds.unistra.fr/cgi-bin/topbase/topbase.sh" \
  --data "com=dt" --data "ent=p" \
  --data "nz1=2" --data "nz2=2" --data "ne1=2" --data "ne2=2" \
  --data "is1=3" --data "is2=3" --data "il1=0" --data "il2=0" \
  --data "ip1=0" --data "ip2=0" --data "lv1=1" --data "lv2=1" \
  --data "en1=0.0" --data "en2=0.0" --data "so=s1"
```

and parsed into data/topbase_HeI_2_3S_photo.dat (photon energy in Ry $\rightarrow$ eV; σ in Mb).

The script ports sigma, sigma_HeI, sigma_HeI23S and sigma_VFKY96 from cross_sec.f90 (using the same constants h_{eV} , c from parameters.f90), hard-codes the Verner+1996 He I/He II parameters and the 23-point p-winds/Norcross table, and emits the three vector-PDF figures.