

EXHALE: Update Log (stage 2, from 2026-09-05)

Contents

1. Plan, audit and decisions that open stage 2 (2026-09-05)	4
2. Phase 0: the tests exist and are red on the known defects (2026-09-05)	6
2.1 src/tests/physics_probe/ (W1): 41 assertions, 15 PASS, 26 FAIL, 3.9 s	6
2.2 src/tests/grid_and_gates/ (W2): 6 programs, all RED, 13.4 s	6
2.3 inputdata/sed/ (W4): the SED inventory of decision 16	6
2.4 Regression harness and new cases (W3)	7
3. element_census_tests compiles again (2026-09-05)	8
4. Phase 1 batch 2a: bounded corrections with no golden movement (2026-09-05)	8
5. Phase 1 batch 2b: bounded corrections that move some outputs (2026-09-05)	10
6. Phase 1 batch 2c: corrections that move every output (2026-09-05)	12
Decision handed to the user: the sub-Lyman band budget	16
Gate (advisor)	16
Golden refresh: prepared, awaiting the user's command	17
Gate on the 89-bin band (2026-09-06)	18
Golden refresh executed (2026-09-06, on the user's instruction)	18
After the refresh, same day (2026-09-06): the loaded-SED grid, the carrier budget, hygiene, the SED files	18
Phase 1 closing gate (2026-09-06)	20
7. PLAN_20260906_rev2, Step A: acceptance contracts and run modes (2026-09-06)	20
8. PLAN_20260911: the partitioned stationary route (2026-09-11)	128
The items	128
The pseudo-time start of the stationary restart route	145
P7: the measurement of record	146
The Codex adversarial review (2026-09-11)	148
User decisions of 2026-09-11 (afternoon)	149
P21: the fixed point at a fixed wind, and the production loop	151
The terminology sweep (A1 to A4, 2026-09-11)	153
An external commit during the session	155
Gates	155
Reported by the workers and not fixed	156
9. PLAN_20260912: the seven findings of the stage-2 review (2026-09-12)	159
The state contract at a fixed hydrodynamic state (R2, R3)	159
The items	159
Gates	162
Reported by the workers and not fixed	163
The oxygen rows in the physics document (2026-09-12, user report)	163

ESWENO assessment memo (2026-09-12, user question)	163
The continuous transfer equations in the physics document (2026-09-12, user request)	164
FUV bands B3 and B4 integrated from the spectrum file (2026-09-12, user instruction)	164
The H2 extent against Koskinen et al. (2022) Model A (2026-09-12, user instruction)	165
The Model A fixture re-pinned on the matched configuration (2026-09-12, user instruction "proceed")	166
The base cell's energy row: diagnosed and fixed (2026-09-12, user instruction)	166
10. PLAN_20260913: the findings of the two reviews of 2026-09-12 (2026-09-13)	167
11. PLAN_20260913_lhs_stationary: the LHS 1140 b stationary solve (2026-09-13 to 2026-09-16)	169
The golden refresh of 2026-09-16	200
The golden refresh of 2026-09-17	210
12. PLAN_20260917: the corrections that follow the review of the 2026-09-17 handoff (2026-09-17 to 2026-09-18)	212
L27: the diagnostics evaluate the production stationary operator, and the certification report prints a face-flux budget	213
L28: a stationary record names its solve, and the reader reads one completed solve	213
L29: every archived certification claim reclassified under the binary of record	214
L30: one spherical geometry for all transport, and the element face flux exposed	215
L30b: one spelling of the spherical volume, the failing carrier budget row, and the element flux writer	217
L31: the molecular approximations get a validity contract, a reaction-cycle energy audit and an uncertainty record	218
L32: one frozen rejected coupled Newton system, measured before any preconditioner is designed	220
L33: slow outer relaxation or a stalled mode, read from the existing logs	222
L34a: the atomic catalog re-run, 79 reproduced and the seven low-XUV Roe cases	222
L36 and L36c: ionization stage transport	224
L36d: both elements' ionization stages carried, and the anchor the stage rows already had	226
L36e: the atomic run carries the three stages, and the certified state answers stages D and F	229
L37: the round trip of the molecular base mass row, and the probe that can be pointed at a state	233
L34b: the molecular reference solution, certified at pass 12	235
L34c: the remaining molecular cases, and the two seeding routes the restart contract refuses	237
L35: the matched-domain flux and grid comparison, and the local base closure	241
L30c: what moves oxygen_chemistry by order unity	244
The count rename of the plan's section 11	245
Further corrections made while the series landed	246
The bookkeeping items of the plan's section 11	247
The golden refresh of 2026-09-18: deferred by decision	249
13. PLAN_20260918_rev2: the corrections that follow the two reviews of PLAN_20260918 (2026-09-18 to 2026-09-19)	250
D1a: the base cell width a run resolved, disclosed and reproducible	251

D3a: the transported He II row measured, source channel by channel	252
D3b: the stage-sum entries made gates at their floating-point bound	253
D4a: what the thermochemical closure hands back, measured	253
D4b: the returned-state contract, and what a refused trial puts back	254
D5a: where the boundary of an evaluated state comes from, and which of it the residual reads	254
D5b-1: the evaluation routes derive the boundary after the composition sweep	255
D5b-2: one boundary-state operation, with a stated input set	256
D5b-3: the four rows, the fourth mixing site, and the prescribed-count labels	257
D6: the helium held outside the atomic stages is reserved from the stage simplex	257
D7a: the equation basis of the generic charge-exchange helium row	258
D7b: the metal charge exchange of a transported ionization stage	258
D7c: the helium state the He $\leftrightarrow$ H charge-exchange pair reacts from	259
D7d: the helium state the metal + He and the advected He $\leftrightarrow$ H reactions react from	259
D8a: what a case directory publishes, and when a second solve is taken	260
D8b: the generations a case publishes, the index a reader resolves, and the states this tree already held	260
The bounded experiment of D8 step 8: the movement bound held at $1e-2$	261
D9, steps 1 and 2: the budget of a solve by configuration, and seed compatibility before distance	262
D2b: the lower boundary as a contact, upwinded after the acoustic matching	262
What the user decides next	264
Corrections made while the series landed	264
The golden refresh: still deferred by decision	265
D9 step 3 and the golden refresh of 2026-09-19	265
14. PLAN_20260919_rev1: the review folded in, the boundary composition closed (2026-09-19 to 2026-09-20)	267
P0: the case list the counts are of	267
P4a: the four facts of how a run ended	268
P5a: the capacity contract of a certification report	268
P4b: the seed of a new solve, resolved once	269
P5b: the executable a test selects, and the one it says it ran	270
R2 and P1 step 1, experiments 1 and 2: the ghost's seed decides the certificate	270
P1 step 1, experiments 3 to 6: one root, and the sweep does not reach it	271
P1 step 2: the boundary composition is a fixed point of the sweep	272
P1 step 2b: the accuracy belonged in the composition's own units	273
P6d: the charge-exchange turnover says which cell it answers for	273
P6e: the elemental base flux record belongs to an evaluation	274
P6f: the transit spectrum of a solved state	275
P6b: the time-step restriction of the cells that are evolved	275
P6c: there is no 4.7 per cent inconsistency	276
P4b-2: a flux-closure rung publishes, with its seed	277

P4c: the provenance recovered without rewriting anything	277
P4d: the records bound to their logs, and the four that were wrong	278
The certification rule of 2026-09-20: a state that passes the certificate is certified	278
The catalog under the contract: the third and fourth passes	279
What a state that satisfies the flux criterion and not the certification costs	280
What the user decided	281
The golden refresh of 2026-09-20	281

Code-size summary: EXHALE vs. the original ATES (2026-06-07; updated 2026-09-11) 282

Fortran source inherited unchanged from ATES 289

This log starts fresh on 2026-09-05. The two earlier logs are kept unchanged under their new names:

- docs/Update_EXHALE_stage0.{tex,pdf}: the early phase (the ATES fork up to 2026-08-15).
- docs/Update_EXHALE_stage1.{md,tex,pdf}: sections 1-171, 2026-06 to 2026-09-05. Every "Update_EXHALE section N" reference in the source comments and in the older memos points into that file; the references were rewritten to the new name on 2026-09-05.

Each entry below is one dated section: what changed, why, which files, the gate that was run and its result, and the movement it produced against the scratch baseline or the goldens. Section numbers restart at 1. The plan the entries execute is docs/development_plan_20260905_rev3.md (sections 1-10) with the order of docs/development_plan_20260905_execution.md; item labels (A1, A8, 10.1-1, D0, C1 ...) are the plan's.

1. Plan, audit and decisions that open stage 2 (2026-09-05)

State of the tree at the start: HEAD 35d9dd5 (clean apart from docs/), binary rebuilt from it with gfortran 16.2 (md5 16cde610347c10c0a48db5c2a7fa6cbb). All twelve regression goldens differ from the case outputs (they are the pre-section-170 snapshots); by decision they are not refreshed until the end of Phase 1 (below).

Documents written today, all under docs/:

- physics_numerics_audit_20260905.md (external audit of 35d9dd5), its diagnostics in audit_20260905/.
- development_plan_20260905.md, _rev1.md, _rev2.md, _rev3.md and the three reviews _review.md, _review2.md, _review3.md: the plan and its revisions after each review. Rev 3 is the plan of record; its section 10 holds the independent five-group source audit (three new P0, twelve new P1, about fifty P2), with the reports and the verification probe in audit_20260905/source_audit/.
- development_plan_20260905_execution.md: Steps 0-7, who acts, gates.

Decisions taken by the user today (recorded in rev 3 sections 8 and 10.5 and in the execution document, Step 0):

1. Golden are refreshed once, at the end of Phase 1; until then a scratch baseline (backup/regression/baseline_post170_<date>/, read through a REGRESSION_GOLDEN_DIR variable to be added to run_check.sh) is the control for byte-identity. golden/ is not touched before that.
2. ROE interface: speed_estimate_ROE returns a status and, in the vacuum case, the two gas-edge speeds; Num_flux gains an HLLC vacuum branch. Signature shown before coding.
3. One spectrum type builds every band. Power-law input: power law on the whole grid, XUV and below 13.6 eV alike (the present construction is the stated meaning of the key). A new Spectrum type: Planck gives the Planck photosphere from Stellar Teff and Stellar radius on the whole grid. Load: the table on the whole grid. The Balmer continuum follows the run's type. The setup report states the sub-13.6 eV source and the integrated grid flux.
4. $m_{\text{He}}/m_{\text{H}} = 3.9715$ in the species table; $R_{\text{J}} = 7.1492e9$ cm (IAU nominal equatorial) in the code and the Wind-AE bridge alike (R_0 of every run +2.26%); MJ and MSUN to the IAU nominal values.
5. The du stop uses the signed spread of $\rho v r^2$ over the mean, as flux_spread does.
6. Every planet folder, benchmark and planet example runs on the SED its reference literature used or assumed, one file per planet under inputdata/sed/ with provenance; the input switch is a configuration change, the reruns stay instruction-gated.
7. A loaded SED that ends above the band an active absorber needs (He 2³S at 2583 Å, a low-IP metal at its threshold) stops the run; the message offers the two remedies (turn the absorber off in the input, or supply a covering SED). No continue key.

Open: decision 4 (A3 retry design: carrier substep retry or outer step controller); 2, 5, 6 inside D0; 7 before B2; 10 in Phase 5.

Documentation restructuring done today (this section): Update_EXHALE.* renamed to Update_EXHALE_stage1.*, Update_EXHALE_early_phase.* to Update_EXHALE_stage0.*; every reference in src/, docs/, python/, README*.md, the notebooks and the workspace CLAUDE.md rewritten to the new names. Check: `git diff --numstat` shows additions equal to deletions in every one of the 112 touched files (no line moved; the three exceptions are the renamed logs themselves, which git sees as deleted paths). The rebuilt binary's md5 moved from 16cde610347c10c0a48db5c2a7fa6cbb to e779298d0b482bfb40ae2f9aa80d60a because four of the rewritten references are inside printed message strings (EXHALE_main.f90 2602, ionization_equilibrium.f90 2948, input_read.f90 1004, diffusion_tests.f90 1292), which is text the binary carries; no executable statement changed. The pointers in README.md, README_HOWTO.md and the workspace CLAUDE.md name this file as the current log and the stage 1 file as the history.

Scratch baseline created: backup/regression/baseline_post170_20260905/, the Hydro_ioniz.txt, Ion_species.txt, *_adv.txt and run.log of the twelve cases as they sat in the case directories (48 files, 26 MB; provenance in the headers: git=6d07d48afd41 tree=dirty, the working tree that became 35d9dd5, runs of 2026-09-05). It is the byte-identity control for Phase 1 and is deleted after the golden refresh at the end of Phase 1.

2. Phase 0: the tests exist and are red on the known defects (2026-09-05)

Plan reference: `development_plan_20260905_execution.md` Step 1. Four workers, disjoint files; every result below was re-run by the advisor after the worker's report (times are the advisor's runs).

2.1 `src/tests/physics_probe/` (W1): 41 assertions, 15 PASS, 26 FAIL, 3.9 s

Derived from the frozen diagnostics in `docs/audit_20260905/` (which stay); links the production modules through a computed use closure (`source_closure.py`), so no module can be left out of the link by hand. Drivers named for the physics: `riemann_wave_speeds.f90` (LLF signal speed 1 against the required 3, mass flux -3.5 against -4.5, mirror symmetry; ROE normalization invariance broken by 0.661 and 3.162 in the two branches; NaN star speeds on the equal-pressure expansion; the production ROE flux leaves thermal energy -2.93 after one Euler update on the vacuum-producing expansion), `h2_channel_detector_ratio.f90` ((S+D)/M 0.2545 against the measured 0.2827, D/(S+D) 1/9 against 0.20; channel sum and stoichiometry GREEN), `h3p_cooling_limits.f90` (cooling independent of $n(\text{H}_2)$ at 0, 1, 100 cm^{-3} ; joins +0.430, -0.00113, -0.0235 at 300, 800, 1800 K; factor jump $1.5\text{e-}3$ at $1\text{e}14$), `h2_rovibrational_identity.f90` (chemistry $T^2 \text{dln}Z/\text{dT}$ against the EOS u_{rv} : -0.63% to -8.29% over 300-8000 K), `photoevent_energy_ledger.f90` (H event at 20 eV closes exactly through the production photoelectron_share; the H2 double event at 80 eV accounts for 60.28 of 80 eV, 19.72 eV unassigned), `radiation_exchange_ledger.py` (two-cell ownership rule, GREEN by construction).

2.2 `src/tests/grid_and_gates/` (W2): 6 programs, all RED, 13.4 s

`grid_width_identity.f90` (production `define_grid`): `dr_j(j)` against `r_edg(j) - r_edg(j-1)` off by 1.0145 on the Mixed grid at $r_{\text{max}} = 10$, 1.0100 at $r_{\text{max}} = 2$, 1.0046 on Stretched, exact on Uniform (where every cell has the same width, which is why the defect was invisible). `photon_grid_quadrature.f90` (production `set_energy_vectors`, `J_inc`, cross sections; `wasp_full` spectrum): code/exact `P_HI` 1.0951 (floor 4.8 eV), 1.0890 (5.139 eV), 1.0003 (13.6 eV); `P_HeI` 1.0157 and `P_HeII` 1.0299 in every configuration; heating 0.99984. New from this test: `P_HeI` carries the same threshold-bin defect (24.6 eV is a grid point whatever the floor), and the X-ray block ends at 1184.19 eV, not 1240 (share of `P_HI` $1.9\text{e-}10$). `mass_flux_spread_functional.py`: the coded `du` and the signed `flux_spread` on the baseline states (`lower_profile` 823.4 vs 3.61 with two sign changes in the window; `mol_base_handoff` 1.29 vs 0.696 with none; `wasp_full` $9.85\text{e-}4$ vs $9.85\text{e-}4$), and `du = 0` on a synthetic +F/-F window. `output_state_consistency.sh` (200-step `mol_base_handoff`): header `sec_ion=T`, `heat_total/heat median` 0.2752, largest difference 0.822. `base_level_single_statement.sh` (`lower_profile`, 2 steps): setup $2.0035\text{e-}5$ bar against the profile's $1.00\text{e-}6$ bar, factor 20.035. `restart_grid_guard.sh` (roundtrip on Mixed, restarted on Uniform with the same N): exit 0, the restart's `r` equals the file's Mixed radii exactly and sits 0.60 relative from the run's own grid.

2.3 `inputdata/sed/` (W4): the SED inventory of decision 16

Four files built from data already in the workspace, each with a provenance header, a build script under `inputdata/sed/build/`, and the coverage test `sed_coverage_and_order_check.py` (31 assertions, all PASS): WASP-52 b (eps Eri, MUSCLES v22, reproduces the existing planet-folder file to round-off), hot Uranus (the bench-

mark's Ribas band table continued to 2995 Å from the Gueymard composite it came from), HD 209458 b (WHI 2008 solar minimum at 0.04707 au, the p-winds file reproduced exactly), LHS 1140 b (GJ 1132 proxy). Not built, data missing: WASP-121 b (Huang 2023's LLmodels F5V spectrum above 1700 Å) and HD 189733 b (Salz 2016's composite; its solar table is not here). `inputdata/scaled_solar_hd189.sed` judged an orphan (no provenance, hardness inconsistent with its own normalization, ends at 1899.5 Å). Verbatim paper quotes and the open items are in `inputdata/sed/README.md`; rev 3 decision 16 records the status. Also found: WASP-52b/`input.inp` points at a directory that no longer exists (EXHALE/), the top-level Koskinen benchmark runs at 0.048 au against the paper's 0.05, HD209458b/`input.inp` at 0.048 au against the literature 0.047.

2.4 Regression harness and new cases (W3)

`run_check.sh`: REGRESSION_GOLDEN_DIR (default `golden/`; golden mode still writes only to `golden/`); the rebuild is fatal with its log kept in `backup/regression/.build.log`; the summary counts only the cases of the run; the binary copy is status-tested; FILES gains the two `_adv.txt` profiles (the post-processing pass was outside the regression); a file without a reference is MISS, not FAIL, and a run that compared nothing prints NO VERDICT; a restart case keeps its IC pair in `<case>/IC/` and the script restores it after clearing output/; `phase0-cases` prints the new case list. Makefile: `make check CASES=...`; `make test` runs `element_census_tests`, `diffusion_tests`, `residual_determinism` and every executable `src/tests/*/run.sh` found at run time, every suite even after a failure, nonzero exit if any failed. New cases, not in DEFAULT_CASES (PHASE0_CASES): `hydrostatic_column` (hot-Uranus gravity, irradiation 18 orders below normal, no key exists to switch cooling off so the gas is held where it is negligible; 300 steps, isothermal to 2%, but draining through the outflow boundary at 0.8 Mach, so it is a configuration for the Phase 4 D2 measurement, not a hydrostatic state), `oxygen_chemistry` (`examples/18_oxygen_chemistry` verbatim, 1000 steps, 3 min 52 s single-threaded under load), `hp_front`, `hp_zero_seed`, `hp_trace_seed` (restarts of the `mol_base_handoff` state with `Ionization transport: True`; the H II column set to 0 and to $1e-12$ of the H nuclei by moving the protons into H I, since `load_IC` refuses a changed He/H; the three 100-step states are mutually distinct, so the seed reaches the solver). No dump hook asserts the seed at the entry of `solve_carriers`; the README names the two lines where one belongs. Measured (advisor re-run of two cases: 6.2 s, byte-identical on all four files): `check wasp_full mol_base_handoff` against the scratch baseline is byte-identical on all four files per case (W3, 13 min); the five new cases reproduce their own baseline copies; `make check CASES=mol_base_handoff` against the stale `golden/` reports the known $8.3e-4$ and $3.6e-4$ movements and MISS on the two `_adv` files; `make test` (31 s) runs five suites: `diffusion_tests` and `residual_determinism` PASS, `grid_and_gates` and `physics_probe` RED on the known defects as intended, and **`element_census_tests` does not compile at HEAD** (pre-existing: `carrier_state` gained `nrho` and `wfac` in section 158 and the test's two calls still pass ten arguments; `carrier_write_back` likewise takes `nrho` where the test passes `ntot`). Fixed in section 3 below.

Also found by W3: with the flux made negligible, the He recombination coupling warning (`ionization_equilibrium.f90` 947, 952) fires in every cell with a ratio of $1e80$ because both sides sit at their guards; diagnostic only.

3. element_census_tests compiles again (2026-09-05)

src/tests/element_census_tests.f90: the two carrier_state calls pass the nrho and wfac outputs the routine has had since stage 1 section 158, and the two carrier_write_back calls pass nrho (the density the carrier fraction is a fraction of) in place of ntot. Test only; no production source. Result: builds, 20 assertions PASS (E1 stoichiometry and charge, E2 elemental closure through the write-back including the CO collapse, E3 the H2 photoevent nuclei ledger, E4 the base H2 fraction). Until now the suite had been unbuildable, so its assertions had not been checked since section 158.

4. Phase 1 batch 2a: bounded corrections with no golden movement (2026-09-05)

Seven items, seven workers on disjoint files in one tree (private OBJDIR/EXE builds, no shared regression runs); one rebuild and one gate by the advisor afterwards. Plan references: rev 3 sections 4.1 (A2), 10.2 items 7, 8, 10, 12, 10.3, 10.5 decisions 3, 13, 17.

- **A2, ROE estimator and caller** (speed_estimate_ROE.f90 rewritten; Num_Fluxes.f90 ROE branch; input_read.f90; EXHALE_main.f90): docs/a2_roe_interface.md implemented as approved: roe_star_state, status OK/VACUUM/INADMISSIBLE, the vacuum test first, PVRS kept only when admissible, two-rarefaction exponent $1/z$, two-shock densities times the upstream densities, HLLE flux on faces without an admissible star state with a printed counter, refusal of Numerical flux: ROE with molecular chemistry and the ladder EOS. Four deviations accepted: a positivity guard on the input states at the top of the estimator; the refusal placed in the consistency block after the optional keys (where thereis_mol is known); the old p_min floor removed (the guard supersedes it); the counter printed for ROE runs only. Test: 30 ROE assertions GREEN (scale invariance, equal-pressure expansion $c_* \cdot 0.9576611$, stationary contact, asymmetric shock and rarefaction against exact Riemann solutions at 5%, vacuum edges $-6.127017/+6.127017$, positive thermal energy after one Euler update).
- **2a-SED, decision 17** (sed_read.f90): a table that stops above the band an active absorber needs stops the run with the file, the missing band in eV and A, the absorbers (He 2³S; each active neutral metal below the file minimum, K I at 4.341 eV included) and the two remedies; no continue key. Test sed_coverage_stop.sh (six assertions) GREEN; no configuration of record trips it (the Koskinen runs have the metastable off).
- **2a-OPAC** (util_ion_eq.f90 PH_heat_H/PH_heat_HHe, opacity_models.f90): model P's factor multiplies the shared flux weight int_f of every absorption integrand (rates, heating, secondary ionization, acc_q), so the beam's loss equals the local absorption; test opacity_model_p_ledger.f90: a cell with $f = 10$ closes energy and H I ionizations to $3e-8$ (was a factor 10 off).
- **2a-LYA** (lya_rt.f90, util_ion_eq.f90 fuv_lw_photon_field, water_photolysis.f90 header, plus the two call sites in ionization_equilibrium.f90 and diffusive_photochemistry.f90 that the new nHI argument required): band B2 is attenuated by $\text{lya_stellar_beam_transmission}(T, \tau) = \text{erfc}(x1/(\text{sqrt}(2) \cdot X_s))$, the expression jlya_escape_prob carried inline and now calls (bitwise equal at 4941 test points). Test lya_band_transmission.f90 (ten assertions) GREEN. Movement, oxygen_chemistry 1000-step snapshot: du 903 -> 596, log10 Mdot 9.62 -> 9.61, B2 transmission $1e-291$ at the base, $1e-3$ at $1.07 R_p$, 0.98 at $1.5 R_p$;

H₂O moves up to 6.2x and OH 7.3x in the layer. Scratch baseline of that case re-snapshotted (previous copy kept).

- **2a-WAE** (`wae_exhale_input.f90`): mass fractions from the same $m_{\text{He}}/m_{\text{H}}$ the solver is handed; test: reconstructed He/H equals the input to 1e-12 (was 1.0071696x). RJ/MJ/MSUN untouched (decision 14 is 2c).
- **2a-HPFIX** (`constrained_chemical_equilibrium.f90`): `hp_is_fixed` pins `species_fixed(is_HII)` like H₂ and the oxygen carriers; new suite `src/tests/constrained_network_layout/` (eight assertions) GREEN. The three `hp_*` cases never enter the continuation (every cell is a class-1 root), so no run exercises the fix yet.
- **2a-PLANCK** (`J_inc.f90`, `set_energy_vectors.f90`, `input_read.f90`, `write_setup_report.f90`, `excited_hydrogen.f90`, `docs/input_schema.md`): Spectrum type: Planck (requires Stellar Teff and Stellar radius) builds the whole grid from $\pi B_{\nu}(T_{\text{eff}}) (R_{\text{star}}/a)^2$; the Balmer routines call the same shared function (unchanged result); the setup report states the spectrum type, the source of the band below 13.6 eV and the integrated grid flux against the nominal J_{XUV} . New suite `src/tests/spectrum_type/` (12 assertions) GREEN. Measured on the `wasp_full` star: Planck/power-law 1642 at 4.8 eV, 358 at 6.2 eV, 2.9 at 10 eV, 0.014 at 13.6 eV; He 2³S rate 45.2 s⁻¹ (Planck) vs 0.100 (power law); integrated grid flux / nominal 1.42 (power law, floor 4.8 eV), 0.995 (floor 13.6), 112.8 (Planck).
- **2a-GUARDS** (`define_grid.f90`, `set_IC.f90`, `EXHALE_main.f90`, `viscous_conduction.f90`, `steady_newton.f90`, `input_read.f90`, `diffusive_photochemistry.f90`, `water_photolysis.f90`, `oxygen_rates.f90`, `lyman_werner.f90`, `parameters.f90`, `molecular_infrared_data.f90`): `cell_nearest_radius` (the bare minloc read `r(503) = 0` out of bounds on the default grid), `j_min >= 1`, dead `report_unmet_steady_gates` removed, `Mdot` guarded for small N and non-outflow, conduction floor counted and reported, `okres` read in the ray test, correct sweep count, `get_word` initialized, dead `close(unit=1)` removed, `adv_corr` zeroed on allocation, dead `Tcode` argument removed from four carrier routines and their callers, `carrier_mass_amu` through the named species indices, `wp_ready` used as the guard it names, dead constants and stale headers fixed, `molecular_infrared_data's i` private. Test `grid_window_indices` (seven assertions) GREEN.

Gate, advisor's single rebuild (`md5 b52c79ebb15ee9014858425bfea4cb1c`): make test runs seven suites; every assertion added in this batch is GREEN and only the known defects of later items remain RED (two test drivers had linked the program object `element_census_tests.o` from `build/`; their link lists now exclude program objects). Regression, sixteen cases against the scratch baseline, single-threaded: fifteen byte-identical on all four files (the eleven of `DEFAULT_CASES`, `hydrostatic_column`, the three `hp_*`); `oxygen_chemistry` moved as stated under 2a-LYA.

Follow-ups recorded for batch 2b and the docs pass: `carrier_mass_amu` returns 0 for a species missing from the table (needs a stop); `write_output.f90` 806-816's FUV ledger closed form carries no line transmission for B2 (its `rel_diff` reads 1.0 now; the LW row was already 0.84); `inputdata/sed/README.md` describes the old warning in six places; `docs/input_schema.md` row 11a should state the coverage requirement; `docs/a2_oxygen_option_design.md` 402 describes the pre-2a B2 field; the `physics_probe/README.md` summary counts are stale; J_{XUV} (used by the Wind-AE bridge), the FUV band fluxes and the monochromatic flux do not yet follow the spectrum type (decision 13 is enforced for the photon grid only); `backup/regression/parse_golden/` was already stale at 35d9dd5; the three audit probe scripts under `docs/audit_20260905/` no longer compile against the tree and are marked as frozen records.

5. Phase 1 batch 2b: bounded corrections that move some outputs (2026-09-05)

Six workers (five code items and one documentation pass), disjoint files, private builds; one rebuild (md5 d2173f0210abdc84f41cb6fb5076e18) and one gate by the advisor. Plan references: rev 3 sections 4.1 (A1, A3 increment (a), A8) and 10.3 (k_lw cell mean, He I singlet, He II recombination cooling).

- **A1, LLF signal speed** (Num_Fluxes.f90, RK_rhs.f90, EXHALE_main.f90): $a1 = \max(\text{abs}(vL)+aL, \text{abs}(vR)+aR)$, the spectral radius of the Euler flux Jacobian, replacing $\max(\text{abs}(vL+aL), \text{abs}(vR+aR))$ (1 instead of 3 on the inflow test pair); the header's positivity claim now describes the coefficient the code carries. Reporting: calls of the positivity repair, repaired interfaces, repaired interfaces in accepted steps, and LLF faces, all printed even when zero. Test: the three LLF assertions GREEN. Movement: none (no case selects LLF and the repair path is never entered in the matrix: flux correction: 0 positivity repair call(s) in every run).
- **A3 increment (a), carrier reference scales and status** (diffusive_photochemistry.f90, write_setup_report.f90, EXHALE_main.f90): one carrier_element_reference_density(ic, nH, n0, nC) (H2 and H+ hydrogen, OH and H2O oxygen, CO min(O, C)) read by both the Jacobian step (carrier_source_derivative_step) and the residual floor; the reported residual is that of the state after limit_to_element_budget; solve_carriers returns converged / iteration cap / line search failed and the caller stops on a failed status in a carrier the limiter did not move (rejection and retry remain increment (b), decision 4 open); a cumulative CO-ceiling record (cells, applications, molecules removed) in the run summary and EXHALE_resolved.out; carrier_mass_amu stops on a species missing from the table. New suite src/tests/carrier_reference_scales/ (14 assertions; before the change the proton row's diagonal derivative was exactly zero at $H^+ = 0$ and $1e-12$ n_H because the step had collapsed to $1e-300$). Movement of this item alone: hp_zero_seed up to $7.8e-4$ at the ionization front, hp_trace_seed $2.8e-5$, hp_front round-off, mol_carrier and mol_base_handoff byte-identical. Cost of the extra residual assembly +1.5% wall time on hp_zero_seed.
- **A8, Chung detector inversion** (h2_photo_channels.f90): Table II is the detector event ratio $(N_S+N_D)/N_M$ (Chung et al. 1993 p. 886, one pulse per event, quoted at the code site), $q_D = N_D/(N_S+N_D) = 0.20$ at 80 eV; $\sigma_M = (1-f_{di}) \sigma_i$, $\sigma_S = (1-q_D) f_{di} \sigma_i$, $\sigma_D = q_D f_{di} \sigma_i$; yan_rho converted with $q_D = \rho_D/(1+\rho_D)/f_{di}$; an inadmissible q_D stops the run instead of being clipped; frac_H2_double_of_protons renamed frac_H2_double_of_proton_events. At 80 eV: D 0.02254 -> 0.04408, M 0.7972 -> 0.7796, S 0.1803 -> 0.1763. Test: 18 assertions GREEN including the stored-channel and yield integration checks; E1 test 4 prints the same $6.951e-3$. Movement of this item alone: mol_base_handoff n_e up to $1.8e-2$ at $r = 1.018$, x(H+) $1.9e-2$ at the base, T $4.6e-4$; mol_sec_ion at the $1e-3$ to $5e-3$ level, with a 0.078 cm/s velocity shift moving the $v = 0$ crossing by one cell at $r = 1.024$ - 1.026 and flipping the advection-correction branch of the _adv files there. Mdot unchanged.
- **k_lw cell mean** (lyman_werner.f90, util_ion_eq.f90 fuv_lw_photon_field): new lyman_werner_dissociation_rate_cell_mean(F, N_out, N_in, T, n_H, tau_out, tau_in), a composite three-point Gauss rule on segments of at most 0.05 dex of column and 0.5 of continuum depth (the

level-resolved cross section is a piecewise power law with slope up to 3.3, so the water bands' exponential mean does not apply; measured accuracy $2.5\text{e-}4$ worst over 1500 random cells); the star-ward face carries $\text{NH}_2\text{col}(j+1)$. Test: seven assertions GREEN (the old inner-face rate was 3.18x too small on a cell spanning 1.12 dex). Movement: `mol_lyman_werner k_LW` +1.4% to +28.1% (median +2.7%), `n(H2)` -20% and `n(H3+)` -20% at the front $r = 1.071$, T up to $8\text{e-}3$, n_e $1.7\text{e-}2$; `mol_base_handoff` and `wasp_full` identical (no LW flux). Cost under 0.5%.

- **He I singlet and He II recombination cooling** (`util_ion_eq.f90 eval_cool/eval_cool_cells`, `Cool_coeff.f90`, `ionization_equilibrium.f90` comment): the 24.6 eV collisional ionization and the Cen (1992) excitation of He I are charged to the ground singlet `he_ground_singlet_density(nhei, nheiTR)` (the metastable had paid both levels); `lambda_rec_HeII = kT alpha_rec_HeII_total(T)` with `alpha_rec_HeII_total = alpha_11S + alpha_23S` when the metastable is tracked (the coefficient the balance removes He^+ at) and `alpha_B` otherwise; the coefficient is built inside `eval_cool` from T and the run's flags so that `energy_semi_implicit`'s finite-difference dC/dT sees one function (a coefficient-passing form would have left an offset the derivative multiplies by $1\text{e}5$). At $1\text{e}4$ K the He II coefficient rises $2.807\text{e-}13 \rightarrow 3.640\text{e-}13$ (+30%); the He recombination coupling's own weight (`y_net alpha_1`) is still not charged (about 10% of this channel, stated at the code site). Test: 15 assertions GREEN. Movement of this item alone: `wasp_full` median $2\text{e-}5$ / max $1.7\text{e-}4$ in ρ , T max $5.9\text{e-}4$, Ca I and Na I up to $4.7\text{e-}3$; `mol_base_handoff` cool column median 1% / max 2.6% (the recombination channel is 64% of the cooling there), state $1\text{e-}4$.
- **Documentation pass:** `inputdata/sed/README.md` (seven places, the stop), `docs/input_schema.md` (row 11a coverage rule; rows 11, 11b, 11d, K3, K4 line references corrected), `docs/a2_oxygen_option_design.md` (B2 row), `src/tests/physics_probe/README.md` (two summaries: 41/15/26 at 35d9dd5; 83 assertions, 69 PASS, 14 FAIL after 2a with the LLF fix already in the tree) and `grid_and_gates/README.md`, rev 3 section 10.2 and 10.5 "landed" notes, the execution document Steps 1 and 2a marked done, `README.md` and `README_HOWTO.md` (Spectrum type: Planck, the coverage rule, make test, the regression section's case list corrected from eight to eleven).

Gate (advisor): `make test` runs eight suites; every 2b assertion GREEN, RED only on the 2c and Phase 3 targets. Regression, sixteen cases against the post-2a scratch baseline, single-threaded: `wasp_he23off` and `hydrostatic_column` byte-identical (metastable off, no carriers); the other fourteen moved, `Mdot` unchanged to two decimals in all sixteen. Largest relative differences over all cells and columns (single cells; the large values are v near a zero crossing or trace species in one cell, see the items):

case	Hydro_ioniz	Ion_species	Hydro_ioniz_adv	Ion_species_adv
<code>wasp_full</code>	$1.06\text{e-}3$	$4.63\text{e-}3$	$1.37\text{e-}3$	$4.63\text{e-}3$
<code>wasp_full_newton</code>	$1.48\text{e-}3$	$4.69\text{e-}3$	$1.91\text{e-}3$	$4.69\text{e-}3$
<code>mol_base_handoff</code>	$2.54\text{e-}2$	$1.99\text{e-}2$	$5.58\text{e-}2$	$4.21\text{e-}2$
<code>mol_metals</code>	$2.16\text{e-}2$	$2.20\text{e-}2$	$2.70\text{e-}2$	$2.20\text{e-}2$
<code>mol_lyman_werner</code>	$1.25\text{e-}1$	$2.06\text{e-}1$	$5.63\text{e-}2$	$2.06\text{e-}1$
<code>mol_diffusion</code>	$2.54\text{e-}2$	$1.99\text{e-}2$	$5.86\text{e-}1$	$1.95\text{e-}2$

case	Hydro_ioniz	Ion_species	Hydro_ioniz_adv	Ion_species_adv
mol_ir_bands	2.16e-2	2.20e-2	9.65e-2	2.20e-2
mol_sec_ion	1.70 (v at a zero crossing)	6.31e-3	1.70	1.0 (trace ion, one cell)
mol_carrier	2.52e-2	1.98e-2	3.87e-2	1.95e-2
lower_profile	5.38e-1	5.19e-2	5.38e-1	4.96e-2
oxygen_chemistry	1.94	1.0	1.94	1.0
hp_zero_seed	8.13e-2	9.54e-2	4.20e-2	9.54e-2
hp_trace_seed	4.25e-1	1.0	4.24e-2	1.0
hp_front	2.59e-2	6.30e-3	4.42e-2	6.29e-3

The scratch baseline was re-snapshotted from these outputs; the post-2a state of every case is kept in backup/regression/baseline_post2a_20260905/.

Follow-ups recorded: `T_equation.f90` (the temperature root of the advection post-process) carries no He 2³S cooling channel and charges the He I ground-state coefficients to the summed neutral column, so the `_adv` temperature and the hydro temperature are not solved with one cooling function (Phase 3 A5/A6 or the `_adv` consistency item of Phase 6); `post_process_adv.f90` 392 calls `eval_cool` without `nheitr`; `cross_sec.f90` 519-528, `parameters.f90` 475-501 and `input_read.f90` 856-858 still describe the H⁺ column as a proton count and quote a "factor 1.6" between the two double-ionization models that is now reversed and much smaller; `write_output.f90` 200-203 says the `f_shield` column is the factor the rate carries (the rate is now the cell mean); `pct_limited` counts only overshoots above 1e-10 while the clamp acts at any size (same class as the CO counter, now recorded cumulatively); the give-up path of `retry_step` leaves the loop through the same exit as an accepted step (decision 4); `set_oxygen_coeffs` is called with strided sections that create array temporaries in the carrier Jacobian; `docs/EXHALE_user_manual.tex` is stale on the spectrum types and the SED stop.

6. Phase 1 batch 2c: corrections that move every output (2026-09-05)

Fourteen worker items on disjoint files plus advisor edits, every one measured alone against a frozen snapshot of the tree as it stood when the item started (private builds; the items therefore chain, and the "before" of one item is the "after" of those that landed earlier). One rebuild by the advisor with the conda gfortran 16.2 and its OpenBLAS (md5 5845df587aa7948153f1bed3becb28ac), `make test`, and one regression run against the post-2b scratch baseline. Plan references: rev 3 sections 10.1 (items 1-3), 10.2 (items 1-6, 9, 11), 10.5 (decisions 13-15, 17, 2c-CONST). Movement is quoted as the largest relative change over the 500 physical cells of `wasp_full` (WASP-121b, He 2³S + metals, du stop, no Newton finish, so its `Mdot` carries the path-dependent spread of that stop) and `mol_base_handoff` (hot Uranus, 12000-step relaxation snapshot, not steady, so its outer-tail and base-cell numbers compare two points of one transient).

- **2c-DRJ, the cell width is the cell's own width** (`define_grid.f90`, `set_IC.f90`): `dr_j(j) = r_edg(j) - r_edg(j-1)`, an exact identity of the finite-volume grid; the stored width had been the width of cell `j+1`, 0.7

- to 1.25 percent too large over the stretched region. Every consumer confirmed to use $dr_j(j)$ for cell j ; none changed. `grid_width_identity` GREEN. `wasp_full`: T 2.0e-3, v 5.4e-3 (base cell), n_e 6.3e-3, $x(H^+)$ 1.1e-2, mass flux +0.65 percent, printed $\log_{10} \dot{M}$ 13.24 both. `mol_base_handoff`: bulk profiles 1e-2 class with the transient tail at 1e-1, mass flux -0.96 percent, printed $\dot{M}$ 10.52 $\rightarrow$ 10.51.
- **2c-QUAD, thresholds on bin edges** (`set_energy_vectors.f90`): the photon grid is built from bin edges, e_v the geometric center of each bin and de_v its exact width; H I, He I, He II, the metastable edge and e_{mid} , e_{top} are edges, and the X-ray block reaches e_{top} . `photon_grid_quadrature` GREEN except P_HeI at 1.2e-3 (closed by 2c-THR). `wasp_full`: T 7.5e-2 at 1.12 R_p , n_e 7.8e-2, $x(H^+)$ 0.23, base v 0.47, mass flux +0.82 percent, $\log_{10} \dot{M}$ 13.24 $\rightarrow$ 13.25. `mol_base_handoff`: T 4.7e-2 (tail), n_e 2.5e-2, mass flux +0.08 percent, $\dot{M}$ 10.51 both. This is the single largest movement of the batch after the Jupiter radius, and it is the removal of the 9.5 percent overcount of P_HI found by the audit.
 - **2c-BALMER, the Balmer continuum follows the run's spectrum type** (`excited_hydrogen.f90`, `J_inc.f90`; decision 13): the $n = 2$ photoionization rate and heating integrate the field the photon grid uses, the power law for a power-law run, the Planck field for Planck, the loaded table for Load (`stellar_flux_ev`, `spectrum_covers_ev`); a table that does not reach 3647 Å stops the run with the decision-17 message naming $H(n=2)$. `balmer_continuum_field` GREEN (11 assertions). `wasp_full` ($n = 2$ coupling armed): γ_2 falls by a factor 2939, n_e at 1.02 R_p -12.5 percent, $x(H^+)$ at the base -27 percent, T 9.6e-3, wind above 1.2 R_p and $\dot{M}$ unchanged (13.25). `mol_base_handoff`: byte-identical (no stellar lines).
 - **2c-KICK, the momentum kick removed** (`EXHALE_main.f90`): the $u(2, :) + 1e-16$ diagnostic aid is gone; the relative step change skips cells whose old momentum is exactly zero. `momentum_row_from_fluxes_only` GREEN. Movement round-off on `wasp_full` (1e-9), 4e-6 on `mol_base_handoff`, 4.4e-3 in v on `lower_profile` (cold-start snapshot with a sign-changing base flux amplifies it); $\dot{M}$ unchanged everywhere.
 - **2c-STATE, one state per output file** (`EXHALE_main.f90`): when the secondary-ionization coupling never switched on during the march, the final write no longer forces it on; the header says `sec_ion=F`, and the `_adv` files and the heating breakdown are built from the relaxed state. `Hydro_ioniz.txt` / `Ion_species.txt` byte-identical in every case; the `_adv` files and `Heating_breakdown.txt` of the fixed-step snapshots whose flip never fired (`mol_base_handoff`, `lower_profile`, and the other molecular gates) change by up to 0.97 in $n(H\ II)_adv$ at the base. `output_state_consistency` stays RED at 1.2e-5: the heat column is assembled from the pre-sweep rates and the breakdown re-evaluates on the post-sweep composition (Phase 3, together with the `excited_H_update` call between the loop and the final write).
 - **2c-DU, the signed mass-flux spread** (`EXHALE_main.f90`; decision 15): one functional `mass_flux_spread`, $(\max - \min) / \max(|\text{mean}|, 1e-30)$ of the signed $\rho v r^2$ over $j_{min}..N$, read by the stop, the Newton arming and the stage flip, and written to `run.log` at the final step (`mass_flux_spread_functional.py` GREEN). `wasp_full` byte-identical (every threshold crossing on the same step); `mol_base_handoff` byte-identical (fixed steps; the reported du falls from 1.29 to 0.70); `wasp_full_newton` flips and hands off one step earlier, solution moves by 7e-8.
 - **2c-RESTART, a restart keeps the run's grid** (`load_IC.f90`): the file's cell centers are compared with the run's over the physical cells and a relative difference above 1e-10 refuses the restart, naming both grids.

restart_grid_guard.sh GREEN; roundtrip and the three hp_* cases byte-identical on the same grid. Consequence: after the grid-moving items of this batch (RJ +2.26 percent of the domain in R_p units, dr_j) the shipped hp_* IC pairs are refused, and are regenerated with backup/regression/make_hp_initial_conditions.py from a fresh mol_base_handoff output (below).

- **2c-BASE, one base level with a lower-atmosphere profile** (input_read.f90, write_setup_report.f90): with a profile, n0 follows from p_match_bar exactly as from p_base of base.inp; a density key beside a profile is refused unless it agrees within 1 percent (base_level_single_statement.sh GREEN). lower_profile starts at the profile's level: base density $1.0e14 \rightarrow 5.0e12 \text{ cm}^{-3}$, its 12000-step snapshot changes entirely (printed log10 Mdot 8.44 $\rightarrow$ 7.92, ionization front inward, base n_e x8). The case README and examples/17_lower_profile/input.inp drop the density key. Nothing else moves.
- **2c-CI, the C I Voronov row** (Cool_coeff.f90): P = 0, X = 0.193, K = 0.25, A = $6.85e-8$, dE = 11.26 (Voronov 1997 Table 1; the transcription had X and K in the wrong slots, 2.56x at 2000 K, 1.69x at 1e4 K). voronov_carbon_ionization GREEN. wasp_full 1e-5 class, x(C I) 6e-4, Mdot -1e-5 dex; mol_metals bitwise identical (collisional ionization of C I is negligible at its temperatures).
- **2c-MASS and 2c-MASS2, helium mass and Jupiter radius** (decision 14; species_table.f90, parameters.f90, wae_exhale_input.f90, wae_ic_writer.f90, binary_element_diffusion.f90, diffusive_photochemistry.f90, lower_column.f90, utilities.f90, examples/exhale_io.py and the Python tools): He I / He II / He III / He 2³S carry 3.9715 m_H (HeH+ 4.9715), RJ = 7.1492e9 cm, MJ = 1.8982e30 g, Msun = 1.98842e33 g (IAU 2015 nominal), one definition each, the Python tools importing RJ_CM, MJ_G, MSUN_G from exhale_io. Tests atomic_mass_and_radius_constants, jupiter_radius_python_tools.py, suite species_masses GREEN. Measured as two sub-steps: the helium mass alone moves wasp_full by 2.3e-3 in T, 1.1e-2 in x(H+), Mdot +0.095 percent, and mol_base_handoff by 1.2e-2 (tail), Mdot -0.004 percent; the Jupiter radius then rescales every physical length (R0, the domain top r_max in R_p units, the gravity) and moves wasp_full by 7.3e-2 in T, 0.16 in n_e, 0.34 in x(H+), raw Mdot +11.0 percent (printed 13.24 $\rightarrow$ 13.29), and mol_base_handoff raw Mdot +15.8 percent (printed 10.52 $\rightarrow$ 10.58). MASS2 alone (the remaining copies): mol_diffusion 2.8e-5, mol_carrier 6.5e-8, wasp_full byte-identical.
- **2c-THR, one definition of each threshold** (parameters.f90, cross_sec.f90 and the files that restated a threshold): e_th_HI = 13.598434599, e_th_H2 = 15.425927, e_th_HeI = 24.587389, e_th_HeII = 54.417765, e_th_HeTR = 4.767775 eV (NIST ASD), every cross section turning on exactly there; sigma(E, Z) takes the ion's measured potential as an optional argument so He II is not restated as 13.6 Z². ionization_threshold_turn_on and threshold_literal_uniqueness.py GREEN; the P_HeI residual of QUAD closes (0.99878 $\rightarrow$ 0.99995). wasp_full 7e-4 at most, mass flux +0.044 percent; mol_base_handoff 1.3e-3 (tail); printed Mdot unchanged.
- **2c-N2FLOOR, H(n=2) is an absorber of the grid-floor rule** (sed_read.f90 photon_grid_floor_eV, J_inc.f90, set_energy_vectors.f90, excited_hydrogen.f90): with the excited-hydrogen coupling armed the grid floor is e_th_HI/4 = 3647 A, a loaded SED is read down to it and the decision-17 stop is about the file, not the type (WASP-52b with the eps Eri table to 3700 A now runs). Tests sed_coverage_stop.sh and

photon_grid_n2_floor GREEN: the 20 bins added below the lowest bulk-gas absorber change no rate the grid computes, exactly. wasp_full 2e-4 at most (the edge written as e_th_HI/4 instead of 3.40 eV), Mdot 0.0000 dex; wall clock +2.3 percent for the 20 bins.

- 2c-CONST, the particle masses and the astronomical constants in one place** (parameters.f90, species_table.f90, lower_column.f90, Cool_coeff.f90, binary_element_diffusion.f90, diffusive_photochemistry.f90, lyman_werner.f90, oxygen_rates.f90): m_He_atom = 6.6464790722e-24 g (4.002603254 u x CODATA 2018 u; the brief's 6.6464731e-24 was 9.2e-7 low and is not the helium-4 atom), amu, m_e, m_p in parameters.f90, m_He_over_m_H and amu_over_m_H derived, bsp_mass built from them, melem_A = melem_A_u * amu/mu. This closed a UNIT ERROR: the standard atomic weights in u had been added to mass sums carried in hydrogen atoms, every metal nucleus 0.78 percent too heavy in rho, the mean molecular weight, the base composition, the census closure and the metal element diffusion. The advisor closed the remaining local copies (Cool_coeff line-center opacity now takes the atomic weight in u and uses the global amu; amu_C/N/O = melem_A_u(iel_*); the local m_amu_g and kb_erg/m_h2 copies of the diffusion, Lyman-Werner and oxygen modules replaced by the globals). wasp_full 1.0e-3 at most (base v), mass flux +3.2e-5; mol_base_handoff 2.8e-5; printed Mdot unchanged.
- 2c-METALEDGE, every active threshold a bin edge** (set_energy_vectors.f90): each of the six bands keeps its bin budget and is cut at every ionization threshold strictly inside it (H2 when the molecular chemistry is on, every photo-ionizable stage of an element with nonzero abundance), the budget shared over the sub-intervals in proportion to their logarithmic width with at least one bin each; NL is derived from the partition and equals the previous count in every configuration (240 wasp_full, 220 mol_metals, 220 mol_base_handoff). A band with no interior threshold keeps exactly the partition it had. photon_grid_threshold_edges GREEN: seventeen thresholds at relative distance 0 from an edge (eighteen with H2), P_m of C I 1.045 -> 1.000, Ca I 0.933 -> 0.997, Na I 1.030 -> 0.998, Fe I 0.965 -> 0.994 of a 400001-point reference, total H2 photoabsorption 0.985 -> 1.000. wasp_full: T 9.6e-3, n_e 5.2e-3, x(C I) 0.14 at the base, mass flux -0.17 percent, log10 Mdot 13.2860 -> 13.2852. mol_metals: bulk 2e-3, n_e 4.5e-2 in the neutral layer at 1.1 Rp (ionization degree 1e-6), Mdot unchanged. mol_base_handoff: 1e-4 class, by the H2 edge alone. Open (decision needed, below): Mg I, Ca I, Na I, Fe I still miss 1e-3 by 0.25 to 0.58 percent, the rectangle-rule error of the 20-bin sub-Lyman band.
- Advisor edits after the worker:** the H2 channel thresholds e_th_H2_di (18.076 eV) and e_th_H2_dd (51.4 eV) join the edge set when the molecular chemistry is on, because the channel shares of one cross section switch on there (the 32-41.5 eV neutral-dissociation window is continuous and is not cut; the comment at the code site says so); the setup report prints the grid floor as the lower edge of the first bin (photon_grid_floor_eV, or the table floor for a loaded SED) instead of the first bin center, which the new partition had moved to 4.95 eV against a floor of 4.768 eV. The documents that restated the changed quantities were brought in line: docs/input_schema.md (profile level, R_J), docs/a2_oxygen_option_design.md (molecular masses in m_H units), rev 3 and the execution document (landed marks).

Decision handed to the user: the sub-Lyman band budget

The band [4.768, 13.598] eV has 20 bins (edge ratio 1.054), 4.4 times coarser in $d \ln E$ than the pure-H I band (50 bins, 1.012), and it is where every low-IP metal is photoionized; Fe I's cross section falls by a factor 2.5 across the single bin above its threshold. Measured ladder (code/exact):

num_TR	20	30	40	50	80
Mg I	0.99753	0.99871	0.99915	0.99955	0.99982
Ca I	0.99737	0.99877	0.99937	0.99961	0.99984
Na I	0.99752	0.99912	0.99958	0.99967	0.99987
Fe I	0.99417	0.99685	0.99805	0.99883	0.99953
Nl (wasp_full)	240	250	260	270	300

Decided by the user 2026-09-05: num_TR = 89, the budget that gives the band the bin ratio of the pure-H I band (0.0118 in $\ln E$). Measured with the same driver: Mg I 0.999853, Ca I 0.999870, Na I 0.999902, Fe I 0.999615, so the four rates are asserted at the $1e-3$ of every other ion. Nl becomes 309 for wasp_full (was 240) and 289 for the hot-Uranus cases (was 220); the H(n=2) band keeps its 20 bins. Every He 2³S and metals-on case moves again by this, and the gate below was re-run on the 89-bin binary (md5 b0a4ada9f84ee8410c84fda6780ba097) before the baseline re-snapshot. The user also decided the LHS1140b/examples/photochem_profile_base/input.inp density key: deleted (the profile fixes the level).

Gate (advisor)

make test on the single rebuild: every assertion of this batch GREEN (threshold_edges, n2_floor, balmer_continuum_field, atomic_mass_and_radius_constants, species_masses, ionization_threshold_turn_on, threshold_literal_uniqueness, voronov_carbon_ionization, grid_width_identity, photon_grid_quadrature, restart_grid_guard, base_level_single_statement, mass_flux_spread_functional, momentum_row_from_fluxes_only, sed_coverage_stop); diffusion_tests had 13 of 35 assertions fail because its own references carried the helium mass as $4 m_H$ ($7.2e-4 = 1 - 3.9715/4$), and now derives every helium mass from $m_{He_over_m_H}$ (35 of 35, no tolerance changed, the T9 ambipolar targets written as functions of the mass ratio). RED, as intended, only on the Phase 3 targets: output_state_consistency ($1.2e-5$), h3p_cooling_limits, h2_rovibrational_identity, h2_double_photoevent_energy_sum, helium_collisional_ionization_by_level.

Regression, sixteen cases against the post-2b scratch baseline, single-threaded: the thirteen cases that ran all moved beyond the $1e-3$ tolerance, as every item of this batch predicts; the three hp_* restart cases were refused by the new grid guard (their IC pairs sat on the old grid: domain top 4.7253 -> 4.6215 R_p from R_J , base spacing from dr_j) and were regenerated with make_hp_initial_conditions.py from the fresh mol_base_handoff/output (nuclei conserved row by row; H II as written / 0 / $1e-12$ of the nuclei) and re-run. Movement of the thirteen (the cell-wise maxima compare the same cell index, whose radius moved by up to 2.2 percent with the domain, so they overstate the change at fixed radius; the Mdot column is the run's printed value):

case	log10 Mdot before -> after	mass flux at cell N-20	max rel T	max rel v	max rel rho	domain top [R_p]
wasp_full	13.24 -> 13.29	+6.2 %	6.7e-02	7.6e-01	4.1e-01	1.6014 -> 1.5666
wasp_he23off	13.24 -> 13.29	+6.2 %	6.7e-02	7.7e-01	4.1e-01	1.6014 -> 1.5666
wasp_full_newton	13.21 -> 13.26	+6.7 %	1.3e-01	1.9e-01	3.4e-01	1.6014 -> 1.5666
mol_base_handoff	10.52 -> 10.57	+8.3 %	2.5e-01	1.1e-01	4.5e-01	4.7253 -> 4.6215
mol_metals	10.52 -> 10.58	+8.3 %	2.4e-01	1.1e-01	4.4e-01	4.7253 -> 4.6215
mol_lyman_werner	10.50 -> 10.55	+8.2 %	1.0e-01	1.2e-01	2.5e-01	4.7253 -> 4.6215
mol_diffusion	10.52 -> 10.57	+8.3 %	2.5e-01	1.1e-01	4.5e-01	4.7253 -> 4.6215
mol_ir_bands	10.52 -> 10.58	+8.3 %	2.4e-01	1.1e-01	4.4e-01	4.7253 -> 4.6215
mol_sec_ion	10.47 -> 10.52	+8.2 %	2.1e-01	3.7e+02 (sign change of a stagnant base cell)	3.8e-01	4.7253 -> 4.6215
mol_carrier	10.52 -> 10.58	+8.4 %	2.5e-01	1.1e-01	4.6e-01	4.7253 -> 4.6215
lower_profile	8.44 -> 7.95	-68.9 %	2.0e+00	1.5e+03 (sign change)	9.7e-01	4.1487 -> 4.0577
hydrostatic_column	8.25 -> 8.27	-0.1 %	5.7e-02	4.3e-01	4.9e-01	2.9600 -> 2.9595
oxygen_chemistry	9.61 -> 9.60	-6.9 %	7.3e-01	3.1e+01 (sign change)	8.9e-01	4.1487 -> 4.0577

The +0.05 dex of every WASP-121b and hot-Uranus case is the Jupiter radius (item MASS, +11 and +16 percent in the raw flux at fixed cell, partly compensated by the shrunken domain at cell N-20); lower_profile is item BASE (the run now starts at the profile's 1 microbar level); the oxygen_chemistry and hydrostatic_column moves are the grid and the threshold edges. The baseline was re-snapshotted after the hp_* re-run (the post-2b copy kept as baseline_post2b_20260905/).

Golden refresh: prepared, awaiting the user's command

golden/ was archived to backup/regression/golden_pre170_20260905/ with a README (provenance from the headers, unknowns marked), roundtrip was re-run on the post-2c binary, and DEFAULT_CASES of run_check.sh now

lists the five Phase 0 cases as well. The refresh itself, `run_check.sh` golden over the seventeen cases followed by `make check` and one `run_fcheck.sh`, is the user's command to give; until it is given the reference of record is the pre-2c state and `make check` reports every case as moved.

Gate on the 89-bin band (2026-09-06)

Fourteen cases re-run against the post-2c (20-bin) baseline with md5 `b0a4ada9f84ee8410c84fda6780ba097`: `hydrostatic_column` byte-identical (no sub-Lyman absorber), every other case within $1.3\text{e-}3$ in any cell (`mol_metals` ρ $1.3\text{e-}3$, `wasp_full` v $1.3\text{e-}3$ at the base) and printed `Mdot` unchanged, the size the 20 $\rightarrow$ 89 ladder predicts; the `hp_*` IC pairs were regenerated on the 89-bin `mol_base_handoff` output and the three cases run (`Mdot` 10.56).

oxygen_chemistry stopped on the 89-bin grid, diagnosed. At step 13, cell 7, the carrier Newton reached its iteration cap (30) with a relative OH row residual of 0.63 and the status stop of item 2b-A3a ended the run. Diagnosis (worker, measured on frozen snapshots): the Newton is correct (a root exists; the same solver with a larger budget reaches it in 141 iterations, the last three $5.4\text{e-}2$, $3.1\text{e-}9$, $3.9\text{e-}12$) and its residual scale and the element limiter are not involved. The stiffness comes from upstream: at step 12 the ionization sweep accepted a non-root at cell 7 under the relaxation amnesty (`info` 4, normalized residual $5.2\text{e}4$), which put $n_e = 3.5\text{e}10$ and $n(\text{H}^+) = 2.6\text{e}8$ into the frozen background where the previous step had $1.1\text{e}3$ and 0, so the $\text{H}^+ + \text{H}_2$ channel asked for two thirds of the cell's H_2 in one 2.85 s step. On the 20-bin grid cell 7 never took that path. The " $7.8\text{e-}2$ carrier row" of the 20-bin summary is the STEADY residual of a relaxation snapshot (`dt` $\rightarrow$ infinity), not the Newton's; the backward-Euler Newton converged to below $6\text{e-}12$ on all 999 steps there. Two things landed: the iteration budget is 200 (the comment at the code site states the measurement), and the attribution test of the status stop, which asked "did an element constraint act" by carrier over the whole grid, now asks it by cell (the CO ceiling, acting in 160 cells of this case, had made any CO failure anywhere excusable, and had left OH rows it moved by 49 percent through the oxygen closure reported as the solver's); new suite `src/tests/carrier_constraint_attribution/`, 7 assertions, RED before. Open for the user: decision 4 (reject and retry the step on a shorter `dt`, the structure that also covers a step without a root) and the ionization sweep's relaxation amnesty, which handed the carrier system a non-root (acceptance policy, `ionization_equilibrium.f90` class 4).

Golden refresh executed (2026-09-06, on the user's instruction)

`run_check.sh` golden over sixteen cases: the eleven of the previous `DEFAULT_CASES`, `hydrostatic_column`, the three `hp_*` restart cases and `roundtrip`, all from the 89-bin binary `b0a4ada9f84ee8410c84fda6780ba097` (conda `gfortran` 16.2, OpenBLAS); the previous golden is `golden_pre170_20260905/.golden/` now carries the `*_adv.txt` files as well. `oxygen_chemistry` has no golden entry yet (it stops, above); it stays in `DEFAULT_CASES`, so `make check` reports it as the one failing case until the carrier solver question is settled. `make check` and one `run_fcheck.sh` follow in this section.

After the refresh, same day (2026-09-06): the loaded-SED grid, the carrier budget, hygiene, the SED files

- **2c-SEDGRID, the photon grid of a loaded SED** (`sed_read.f90`, `J_inc.f90`, `parameters.f90`, `set_energy_vectors.f90`; item C5 of the D0 draft): the Load path kept the table rows as bin points with

central-difference widths, so no threshold was a bin edge there, on the path every planet calculation of record uses (decision 16). Now the bins are built from edges (geometric means of neighboring rows, the table span as the two end edges), every active ionization threshold inside the span is inserted as an edge with the split bin's two halves carrying the row's flux (exact for the integrated flux), `e_v` the geometric center, `de_v` the exact width; `read_sed` receives the threshold list from `ionization_thresholds_active`, one definition. The table rows themselves are kept in `e_sed_node/F_sed_node` (undiluted), which `stellar_flux_eV`, `loaded_table_floor_eV` and `spectrum_covers_eV` now read, so the loaded field no longer depends on the binning. Test `sed_edges` (`src/tests/spectrum_type/`, 15 assertions) on the `eps Eri` table of WASP-52 b: thresholds at distance 0, flux identity to $2e-16$, rate deviations from the exact table integral `P_HI` $1.8e-3 \rightarrow 4.6e-7$, `P_HeI` $1.1e-3 \rightarrow 1.1e-6$, `P_HeII` $8.4e-3 \rightarrow 8.5e-6$, worst metal $3.2e-3 \rightarrow 5.5e-4$ (Mg II, the table's own 1 Å resolution). Movement: `benchmarks/wasp52` 2000-step marching stop, mass flux at cell N-20 $2.3e-6$, `log10 Mdot` 9.68 both, profiles up to $2e-2$ at a relaxation point; `wasp_full` (power law) byte-identical on seven output files. No regression case loads a table, so no golden moves. Noticed, not fixed: `e_mid` is not an edge of the loaded grid (the LX/LEUV split it feeds moved 0.02 percent); `stop_if_balmer_band_unstated` prints bin centers as the band.

- **Carrier Newton budget** (`diffusive_photochemistry.f90`): `newton_maxit` 30 $\rightarrow$ 200 after the diagnosis above; `oxygen_chemistry` runs its 1000 steps on the 89-bin grid (`log10 Mdot` 9.60) and is added to `golden/` below. The constraint attribution of the status stop, by cell, and its test suite `carrier_constraint_attribution` (7 assertions) landed with it.
- **Hygiene, D0 items C16 and C4**: the dummy `alpha`, threaded through `reconstruction_continuation_rhs`, `RK_rhs` and `Num_flux` and never read, is removed from the three signatures and ten call sites (`hydrostatic_column` byte-identical against a same-compiler control); `output_state_consistency.sh` and its README describe the rate lag (measured RED $1.19e-5$ with the header now reading `sec_ion=F`) instead of the removed post-loop block. `docs/audit_20260905/roe_flux_vacuum_probe.f90` is an archived probe that no longer compiles against `speed_estimate_R0E`; left as the frozen record it is.
- **Document sync** (`docs/EXHALE_user_manual.tex` + PDF rebuilt, `docs/p44_base_sawtooth.md`, `write_output.f90` and `util_ion_eq.f90` comments, `write_setup_report.f90`): the manual carries the Planck type, the decision-17 stop, the six-band edge grid, the IAU constants and NIST thresholds, the signed `du`, the profile base level and the restart grid guard; NLTR is now counted on the Load path (the "no grid point below 13.6 eV" sentence had been printed for every loaded SED); `src/utis/glob.py`, which shadowed the standard library module, is `src/utis/EXHALE_interface_state.py` with its two importers updated. Reported, not changed: the GUI offers no Planck type; `global_parameters` exports an integer named `count` that shadows the intrinsic.
- **Stellar SED files** (`inputdata/sed/`, decision 16, source rule of 2026-09-06: MUSCLES data normalized as the planet's reference paper did): `wasp121b_solar_huang2023.txt` (recommended; WHI 2008 solar shape below 1700 Å, $F(1-912 \text{ Å}) = 1.6e6$ and $L_{\text{ya}} = 1.0e5$ as Huang et al. 2023 section 2.2 prescribe, 6459 K photosphere above; the $1.6e6$ confirmed as the $\lambda < 912 \text{ Å}$ band by Salz et al. 2019 section 7), `wasp121b_wasp17_huang2023.txt` (MUSCLES WASP-17 alternative; its 1700-2583 Å photosphere

is 1.8x fainter than the blackbody, the size of the He 2³S photoionization bias of the solar candidate), hd189733b_epseri_salz2016.txt (eps Eri, MUSCLES v22, normalized to Salz et al. 2016 Table 3; the proxy pairing is the one Salz et al. 2018 section 4.1.1 make), and hd189733b_bourrier2020.txt (the released MOVES III tables, mean of Visits B-E, measured to be the flux at 1 au, scaled to 0.031 AU, eps Eri photosphere above 1600 Å identical to the Salz file's; 30 percent fainter in the Lyman-Werner band, 1.4x brighter in the EUV). Each file was read by EXHALE.x for two steps on the final binary without error. User decision: HD 189733 b runs at a = 0.031 AU (Salz 2016 Table 2, the modeling paper of the comparison); the planet folder, benchmarks/hd189, examples 01-11 and 13, and params_table.txt carry it (the regression inputs jfnk_hd189*, tpm_hd189 and the archived copies keep 0.0330). No input.inp has been switched to a loaded SED yet (2d).

Phase 1 closing gate (2026-09-06)

make test on the final binary b2c15e53c888d09fa90ffdad80443c3a: ten suites, eight PASS; RED only on the Phase 3 targets (output_state_consistency 1.19e-5, h3p_cooling_limits, h2_rovibrational_identity, h2_double_photoevent_energy_sum, helium_collisional_ionization_by_level). make check: sixteen DEFAULT_CASES against the refreshed golden/, all four files byte-identical in every case (REGRESSION PASS); the sixteen goldens taken from the earlier 89-bin binary reproduce on the final one, which is the measured statement that the loaded-SED grid, the carrier budget, the alpha removal and the report edits leave the analytic-spectrum matrix untouched. run_fcheck.sh result recorded below. run_fcheck.sh: the -fcheck=bounds, do, mem build ran 1500 bounded HD 209458 b steps with no runtime trap (fcheck regression: CLEAN, log10 Mdot 9.59), and the production build was restored to b2c15e53c888d09fa90ffdad80443c3a. **Phase 1 is closed.** Open for the user before Phase 3: decision 4 (carrier step rejection/retry), the relaxation amnesty of the ionization sweep that handed the carrier system a non-root, the D0 draft (docs/d0_governing_system_20260906.md, 33 consistency items, 15 questions) for acceptance, the switch of the planet inputs to their loaded SEDs (2d), the WASP-121 b SED choice (solar shape, WASP-17 shape, or the two joined at 1700 Å), and the sub-Lyman band budget of the Load grid (the table's own resolution is kept; Mg II stays at 5.5e-4).

7. PLAN_20260906_rev2, Step A: acceptance contracts and run modes (2026-09-06)

Plan of record for this stage: docs/PLAN_20260906_rev2.md (rev 0 and rev 1 and their two reviews are kept beside it; the execution document of 2026-09-05 points at rev 2). The decisions of To_be_determined_by_user_20260906.md were judged against To_be_determined_by_user_recommend_20260906.md and adopted as rev 2 section 2 states.

Documents written (advisor): docs/a0_run_mode_contract_20260906.md (the three run states; Time stepping: Local and PTC are initialization only; physical time advances only by accepted global steps; headers gain mode= and t_phys=; the proposed Run mode: init | phys key awaits the user's confirmation before A0-impl); docs/a2_certification_contract_20260906.md (one evaluator, four contexts: probe, stationary trial, physical step, certification; statuses not_applicable / evaluated / unavailable; one condition per active balance; what replaces what). Documents drafted by workers and verified for structure and wording:

docs/b1a_active_equation_inventory_20260906.md (flags, equations per flag, constraints, five validity states with producers, the prerequisite table for eight configurations; finding: only the hydro rows have a state-consistent evaluator; the composition projection, the rho rewrite and thirteen closures with stated domains are recorded nowhere) and docs/b1_target_system_20260906_draft.md (energy ledger with the shared H2 zero at $v=0$, $J=0$; chemistry mass-preserving; Lyman-alpha loss and return from one 2p balance, an escape-probability multiplier on the existing cooling explicitly not the target; the zero-net-current closure derived from Koskinen 2022 B9/B10 and shown to give exactly the ambipolar field binary_element_diffusion already computes; the trace-metal counter-flux; the rollback table of fourteen state-changing operations showing `u_old` is not a checkpoint; CO alternative A blocked on rate literature, alternative B specified and recommended; ten open questions).

- **A1 (carrier half)** (`diffusive_photochemistry.f90`; suite `carrier_returned_state_acceptance`, 36 assertions, plus one in `carrier_constraint_attribution`): the carrier update is accepted on the RETURNED state, every unconstrained row of every cell, $|res|/\max(\text{row_terms}, 1e-300) \leq 1e-8$ with the absolute floor inside the scale, finiteness of every row, independent of the termination branch; the relative-drop exit is `carrier_solve_stagnated`, a diagnostic; the decision is `carrier_returned_state_verdict`, a side-effect-free function returning verdict and reason for A3 to consume. Defect found and fixed: `limit_to_element_budget` marked a cell constrained only when the clamp exceeded `limit_report = 1e-10`, so smaller clamps left cells presented as Newton-measured. Impact: `mol_carrier` byte-identical (no changed branch reachable: all 499 transport steps of the first 500 ended at the floor); **oxygen_chemistry refuses at step 4**: the CO thermal ceiling, a clamp without a rate, leaves a neighboring cell's CO row at $1.0e-7$ of its terms (accepted steps before it: $1.1e-9$, $4.2e-9$), and the code no longer certifies it. That is the plan's position (B1 alternative B: a ceiling-active atmosphere is outside the validated set); the case can run only in A0's initialization mode once A0-impl exists, so it fails make check until then.
- **A1 (class half)** (`steady_residual.f90`, `steady_newton.f90`, `ionization_equilibrium.f90` comment block; suite `acceptance_classes`, 12 assertions): `n_cells_without_chemical_root(acc_n) = acc_n(4)`, one definition for the trial handling (a local-closure validity check, no stationary condition) and for the three points that declare a steady state solved; a class-4 cell in the state handed back is written but reported "NOT certified". Impact: `wasp_full_newton` and `mol_base_handoff` identical (no class-4 or class-5 acceptance in any ledger). Open: the marching stop cannot pass the count because the loop never receives the sweep ledger; A2 closes it.
- **C1** (in progress) and **A2** (steps 1-3 of the contract, in progress).
- **C1** (`inputdata/sed/README.md`, `build/sed_coverage_and_order_check.py`, suite `spectrum_type` driver `loaded_sed_table_semantics`, 78 assertions): a table value is a bin average on every shipped file (MUSCLES const-res, WHI, Gueymard: 1 A or 1 nm bins; Bourrier 0.016 A samples with no stated convention) except the blackbody part of the solar WASP-121 b candidate (point sample); the reader treats a row as a histogram in energy, exact for the band energy of a bin-averaged table, second order in the bin width otherwise (MEASURED order 2.0 to 2.9 through the production consumers). Coverage is now checked per configuration from the active

threshold inventory (2600.5 A metastable, 3647 A when both stellar keys arm H(n=2), the lowest active metal), 12 configurations. Findings closed the same day (advisor): sed_read.f90 single-precision 1e-8 at two sites (6e-9 relative shift of every loaded energy); the HD 209458 b WHI file rebuilt to 5000 A (its 3000 A cut stopped the run in read_sed naming H(n=2)); WASP-52b/input.inp SED path from before the 2026-07-14 move; the two solar build scripts' headers (2583 A / 4.80 eV). Remaining: the WASP-17 file begins at 10.0 A because the MUSCLES v24 table does (the grid top becomes 1239.84 eV; recorded).

- **BALMERQ** (excited_hydrogen.f90; suite spectrum_type driver balmer_band_quadrature, 21 assertions; balmer_continuum_field Planck references refined): the Balmer-continuum rate and heating integrals follow the field's own nodes in closed form (power-law segments of a loaded table and of the power-law type; 5-point Gauss on a geometric subdivision for Planck), replacing a fixed 400-point uniform rule that stepped over emission lines. On the eps Eri table of benchmarks/wasp52 the old rule was 2.9 percent low in the rate and 2.7 percent high in the heating (the Lyman-alpha line at 10.20 eV lies in the band); on the power law it carried 9.5e-5 / 6.0e-5 of truncation. Movement: benchmarks/wasp52 gamma_2 +2.99 percent, n_e 5.3e-4 at 1.09 Rp; wasp_full 1e-8 class (sub-0.1 percent, no golden refresh). One worker reported two scratch runs killed by a pkill matching a shared basename: the common brief now says kill by PID after checking /proc/<pid>/cwd, never by name.
- **A2, steps 1-3** (src/modules/time_step/certification.f90 new, steady_residual.f90, steady_newton.f90, EXHALE_main.f90 call sites, utilities.f90 header, diffusive_photochemistry.f90 save/restore pair; suite certification, 18 assertions): one evaluator builds a 19-entry inventory from the configuration flags and returns per entry a status (not_applicable / evaluated / unavailable), the row measure (max over physical cells of |res|/max(scale, floor)), the volume average, the worst cell, finiteness and the scale text; hydro rows and carrier balances are evaluated (carriers inside an isolated workspace that saves and restores every module array carrier_steady_residual writes, asserted after every call), elemental transport, level equations and the seven System_HeH_* closures are unavailable until step 4. The certification context runs at the marching stop (which now receives the sweep ledger, closing the A1 open item), the JFNK final and the PTC declaration; probe and trial decisions read one record, the merit rule unchanged. Exit status (user decision): a run that DECLARED a stationary state and failed certification exits 2 (stop 2); a run ending on a cap makes no claim, writes certified=F cert_reason=no_stationary_claim and exits 0; run_check.sh never tests the status. Five shell gates of grid_and_gates that asserted exit 0 from a full run now accept 2 with the reason stated at the code site (advisor). Impact: numeric content of wasp_full, wasp_full_newton, mol_carrier identical; headers gain certified=. **No configuration certifies today**, by construction (the closures are unavailable), and two measured findings stand on their own: the JFNK info = 0 is set at the loop top and never revised when the returned state is re-measured at its own composition (wasp_full_newton energy row 4.1e-4 against the 1e-5 target, 40x), and the du stop accepts hydro rows of order one on their own terms (wasp_full at du = 1.0e-3: mass 1.3e-2, momentum 1.8, energy 0.72; at du = 2e-2 the same rows read 1.7, 2.0, 0.96, so the spread and the rows measure different things). Not implemented in this item (it would change the solve): the probe rule that a class-4 chemistry makes a Newton direction unusable. Tolerances inherited; the convergence study is owed.

- **B1 accepted (2026-09-06).** The user decided the fourteen open questions of docs/b1_target_system_20260906_draft.md as recommended (metal reservoir zero; electron mass convention; ioniz_eq density intent(in); the residual H I excitation cooling under the Lyman-alpha ledger; trace metals in the diffusion matrix; H+ in the B9 set; CO exclusion adopted with the destruction model held for rate literature; H3+ low-collider policy; one H2 partition function; rollback of diagnostic extrema; identity row placement; the CO ceiling's validity category; O(1D) reservoir; the refused _adv product rule). The accepted specification is docs/b1_target_system_20260906.md (the draft kept and marked superseded). This is the Phase 2 gate of PLAN rev 2; B2 (the residual-controlled temperature solve) started the same day.
- **A3**(diffusive_photochemistry.f90; suite carrier_retry, 32 assertions): the carrier subsystem's checkpoint (carrier_checkpoint_take / restore / matches: carrier fractions, the module arrays of the A2 save/restore enumeration, every constraint and solve diagnostic, the verdict, the cumulative CO record), and the controller: a refused returned state restores the substep's entry state, halves the carrier substep with the background explicitly frozen (listed at the routine), covers the interval with accepted substeps, and on exhaustion (nine refusals, floor $dt/2^8$) restores the ENTRY state and stops; f_sp is written only for a covered interval. Two CO records (accepted, restored on refusal; attempted, never restored) and controller attempt statistics with the A0 ledger-family hook. Retried result equals the explicitly subdivided integration bitwise; step doubling gives first order (ratio 2.001). Impact: mol_carrier (0 retried intervals in 11999), mol_base_handoff, oxygen_chemistry (3 steps) byte-identical. **Finding:** the retry recovered the oxygen_chemistry refusal ($1.04e-7 \rightarrow 7.9e-9$ after ten halvings) because the row scale carries the $1/dt$ time term, so any fixed clamp imbalance passes a fixed relative floor at a short enough substep; that is a defect of the acceptance measure shared by A1 and A2, fixed next (A1scale: the returned-state and stationary measures are judged against the physical terms of the row, transport and reaction, with the absolute floor; the time term stays only in the Newton's own iteration test).
- **B2**(energy_semi_implicit.f90; suite energy_update, 20 assertions): the implicit temperature step of a cell, $R(T) = u(T) - u(T_{old}) - a [\text{heat} - \text{cool}(T)] = 0$ with $u(T)$ from the run's equation of state (caloric H2 included), is solved by a safeguarded Newton-bisection hybrid on a physical bracket ($0.01 T_0$ to $5e4$ K with H2, $1e7$ K otherwise) to $|R|/\text{scale} \leq 1e-9$, the cooling written back is $\text{cool}(T_{returned})$, and every non-converged outcome is a named failure (ITER_CAP / NO_BRACKET / FLOOR / NONFINITE) with the cell, the residual and the missing volumetric source; no clamp remains, a floor is a failure, and the interim action is a stop that B3a replaces by rejecting the attempted step. The caller is untouched (optional status argument, energy_update_last_status). RED kept in the suite: the two-iteration update on a falling cooling branch left $|R|/\text{scale} = 2.9e-2$ and a 6.9 percent temperature error. Impact: wasp_full transient $5.4e-5$, converged (same stop step 16257, $\log_{10} Mdot$ 13.29) $1.6e-6$; mol_base_handoff $4.9e-5$; hydrostatic_column identical; no floor or failure fired anywhere; cost +9 percent end to end on wasp_full. Follow-ups assigned: the identical clamp in the conduction stage (viscous_conduction.f90 572-589; B2b), the certification's counting of energy-floor hits as an accepted correction (A2 step 4), the stale end-of-run floor report (A0-impl).
- **A1scale** (diffusive_photochemistry.f90; carrier_retry 42 assertions): the returned-state acceptance and the stationary carrier measure are judged against the row's PHYSICAL terms ($|transport| + |reaction| +$

floor, the floor in rate units without the $1/dt$ form); the time term $|n - n_{old}|/dt$ stays only in the Newton's own iteration scale (row_terms, bitwise unchanged), beside which row_terms_phys is accumulated. RED: a residual of $1e-7$ of a row's physical terms read $3.3e-8$ at dt and $5.6e-9$ at $dt/16$ on the old scale (crossing the $1e-8$ floor with the defect unchanged); now $1.0e-7$ at every dt . Impact: mol_carrier, mol_base_handoff identical; oxygen_chemistry is refused at step 2 (CO row $5.5e-8$ beside a clamped cell; 37 retry attempts to the floor, all refused, entry state restored), and the old measure only postponed that refusal to step 88 on today's tree. Finding carried forward (A1scale2, needs its own impact measurement): with a dt -independent acceptance there is a shortest certifiable substep, since the Newton's stopping test is still relative to the full terms and the round-off it leaves grows as $1/dt$ against the physical terms; the same measure in both places closes it. Also: cterms in steady_newton.f90 (534, 645) is written and never read. **Consequence for the matrix:** oxygen_chemistry cannot run under the physical acceptance while the rateless CO ceiling is active (B1 decision 7: exclusion); once A0-impl lands, the exhaustion action reads the run mode (initialization: record the refusal in the attempted ledger, write the state flagged, continue; physical: stop), which is the item A3b, and the case runs in init mode as the diagnostic it is.

- **B3a and B4 designs** (docs/b3a_attempted_step_controller_design_20260906.md, docs/b4_spatial_operator_design_20260906.md, read-only workers; advisor decisions in their last sections). B3a: eight of the fourteen step operations have no returned-state verdict today (RK/BC, element diffusion, excited H, ioniz_eq, the projection, Apply_BC, conduction, Shapiro); checkpoint about 2.1 kB per cell. B4: species mass fluxes are not consistent with the mass flux today (equilibrium species are not transported and rewrite rho; carrier advection is a cell-centered product; element diffusion is in advective form; no species enthalpy transport); increments B4-1 to B4-6 with their expected movement. **Advisor correction to the B4 draft:** the formation-energy flux $F_{form} = \sum_s \epsilon_s F_s$ is the RESERVOIR's transport and enters no thermal equation (photoheating already carries $h_{nu} - I$; recombination emits I); the draft had written it as a thermal source, which would heat a cell by the ionization energy leaving it. B4-2 becomes the diagnostic reservoir balance that the ledger identity T1.5 tests, plus the face-composition enthalpy (T-B4.5). The B4-2 pre-measurement (src/utls/formation_energy_flux_diagnostic.py, section 7 of the design) stands as the size of that reservoir transport: on wasp_full_newton 28 percent of the absorbed heating leaves the column as advected ionization energy (median 14 percent of the local heating, 52 percent at the ionization front at 1.23 Rp); the present energy face flux is $v \cdot (E + p)$ with the caloric H2 internal energy already inside E , addressed by the owning cell (T-B4.5). Findings: lower_profile's golden mass flux is not flat ($1.1e-2$), so continuity-based diagnostics on it discriminate nothing until it is Newton-finished; benchmarks/hd209/output integrates to cooling 24 percent above heating on its saved state.
- **B2b** (viscous_conduction.f90; energy_update suite second driver conduction_floor_tests, 24 assertions): the Crank-Nicolson conduction stage no longer clamps; a solved temperature at or below the shared EOS floor (imported from energy_semi_implicit, one declaration) is a FAILURE with status, cell, the solved temperature and the energy the floor would have injected; the tridiagonal solve refuses a zero or non-finite pivot; no partial state is written. The operator is in flux form and conserves thermal energy exactly on a closed column

- (MEASURED 0.0); as boundary-conditioned in a run it exchanges heat through the base Dirichlet anchor by design ($3.4\text{e-}15$ up to that flux). Impact: byte-identical on `wasp_full`, `mol_base_handoff`, `hydrostatic_column` and on two conduction-on variants in which the stage runs; zero floor activations anywhere. B1 section 7.4 rows 9 and 11 updated (no clamp state left). Noticed: `benchmarks/koskinen2022_model_a/*` cannot restart (stored IC on the pre-2c grid, $4.5\text{e-}4$ off; the same class as the `hp_*` regeneration).
- **B4-5 measurement** (`src/tests/grid_and_gates/hydrostatic_residual.f90`, design section 8): on the analytic isothermal column, through the production `Reconstruct`, `Num_flux`, `RK_rhs` and `Source`, the pressure-gravity pair is at design order in the interior ($\text{dr}^{\wedge}1.96$ PLM, $\text{dr}^{\wedge}1.98$ WENO3 at $r = 2$; base cell $1\text{e-}4$ of its weight, first order in the base spacing), and the $O(1)$ hydrostatic imbalance of the mechanical column sits at the OUTER free-outflow cell: 0.75 of its own weight at every N under PLM (the zero-gradient ghost zeroes the MC forward difference, so cell N carries no pressure gradient), $6.4\text{e-}2$ falling at order 1.4 under WENO3. A well-balanced pair (B4-5b) would not remove the drain; the outer boundary fill is the item (B4-4a, started). Two latent defects found: `define_grid`'s `tol = 1.0` initializer implied SAVE, so a second call in one process skipped the stretch-factor Newton (production calls it once; fixed by the advisor: `tol` reset before the loop); the case README's single `Grid cells` ladder does not refine the base cell (README updated).
 - **A2 step 4** (`certification.f90`, `binary_element_diffusion.f90`, `excited_hydrogen.f90`, `ionization_equilibrium.f90`; certification suite +7 driver and +15 shell assertions): the five entries that were unavailable are measured on the state without solving or writing anything (He/H partition and trace-element transport balances, He 2^3S and H($n=2$) level balances, the eliminated-species closure as the sweep's own normalized reaction residual at the given composition), each restoring every module array it installs, asserted on every call. Output content identical on seven cases. **What refuses certification now:** the three hydrodynamic rows on every marching case; the energy row alone on `wasp_full_newton` ($4.1\text{e-}4$ at cell 376, the JFNK `info = 0` finding); the H2 carrier row on `mol_carrier`; and, new, the elemental transport on the diffusive runs (`mol_diffusion` partition $6.4\text{e-}5$ at cell 203; `lower_profile` partition $4.8\text{e-}3$, trace metals $1.2\text{e-}2$, the first measurement of what the Picard alternation reaches). The closures and level balances are within tolerance everywhere ($1\text{e-}16$ to $1\text{e-}12$ of their turnover), so the equations that lacked an evaluator are the solved ones. Both floors are reported as attempt diagnostics (0 everywhere). `cert_tol_element` is inherited, not measured, and is now decisive for the diffusive runs: the convergence study of contract section 4 is owed.
 - **A1scale2** (`diffusive_photochemistry.f90`; `carrier_retry` 51 assertions): the carrier Newton stops when its residual is within its own tolerance of the full terms AND within the acceptance tolerance of the physical terms, or when the arithmetic forbids the second (residual at 64 epsilon of the full terms), a named outcome; the interim reason `unreachable_at_substep` for a refusal whose rows are all at round-off was superseded the same day by A1scale3 (below); the stop message reports the halvings made. Option (a) (stopping on the physical scale alone) was built, measured (one extra iteration every step, no rescue of the short substep) and declined. Correction of the A1scale report: the Newton reached one epsilon in two iterations on that column; the $7.7\text{e-}5$ was the ninth attempt's state. Impact: `mol_carrier`, `mol_base_handoff` identical; `oxygen_chemistry` still refused after step 2 (now the unreachable reason, OH at cell 90, 15 attempts). Decision taken from its finding

(A1scale3, started): a row at the arithmetic round-off of its own terms is not evidence of a defect and is accepted as "round-off limited" with a flag and a count, so a trace carrier at the element floor does not refuse an interval; the physical-terms acceptance stays for every other row.

- **A0-impl** (input_read.f90, parameters.f90, EXHALE_main.f90, utilities.f90, load_IC.f90, write_setup_report.f90, ionization_equilibrium.f90 ledgers; suite run_mode, 22 assertions): the run mode is in. Run mode: init | phys, default init in every configuration (advisor decision replacing the draft rule: the ordinary use is a stationary solution reached by marching and certified, which is initialization in the contract's terms; a physical integration is asked for); phys with Time stepping: Local refused at startup; in phys the handoff state is tested at step 0, the clock advances only after a complete accepted step (measured equal to the sum of accepted dt to the last bit, with and without forced rejections), two ledger families (ledger_family in global_parameters), headers gain mode= and, in phys, t_phys=; a restart of an init snapshot in phys mode is the handoff at $t = 0$ (contract section 3 clarified), and only a mode=phys header without a finite t_phys is refused; init never exits nonzero for certification. Impact: no input changed; wasp_full, hp_front, hydrostatic_column, mol_base_handoff bitwise identical. Both floor reports of EXHALE_main.f90 now state the B2/B2b behavior.
- **COLit** (docs/co_destruction_rates_literature_20260906.md; publisher PDFs of Westley 1980 NSRDS-NBS 67, Tsai 2017, Wakelam 2012 added to references/): B1's CO choice is settled on physics: every neutral CO destruction channel carries a barrier of 6.7 eV or more, so the one-sided destruction model's $\tau_{\text{dest}} \ll \tau_{\text{res}}$ fails at the ceiling's onset at every density (factor 9 to $1.5e8$) and the exclusion (decision 7) is the physics. Findings: molecular_reaction_heat.f90 (H2, H+, He+, H2+, H3+, HeH+) and the oxygen Shomate table of oxygen_rates.f90 carry different reference zeros (T1.9 one-table item, assigned as B3b-ENTH); CO2 is not in the species table (removes the one evaluated CO channel below 2500 K and an OH sink); the workspace VULCAN network's CO reactions all produce species EXHALE does not carry, VULCAN's stated validity is 500 to 2500 K, KIDA's 10 to 300 K. Requests for the user: the A&A papers Visser et al. 2009 (503, 323; the CO shielding functions), McElroy et al. 2013 (550, A36
 - RATE12 file), Venot et al. 2012 (546, A43) and 2020 (634, A78), plus Moses et al. 2011 (ApJ 737, 15) and Tsai et al. 2021 (ApJ 923, 264), which this host cannot download.
- **A1scale3** (diffusive_photochemistry.f90; carrier_retry 61 assertions): a row at the arithmetic round-off of its own full terms (64 epsilon) is counted "round-off limited", flagged per row, and left out of the acceptance decision; every other row is judged on the physical terms at $1e-8$ as before; the controller halves only for genuine refusals; the stop prints both scales for every refused row. Impact: mol_carrier, mol_base_handoff bitwise identical (0 round-off limited rows); oxygen_chemistry still refused at the same step and row (CO at cell 281, $|res|/phys = 3.2e-8$, $|res|/full = 2.2e-12$, four orders above the round-off bound); the element limiter's imbalance one cell beyond a clamped cell, correctly refused. The count's accessor exists; its line in the run summary and the certification carrier entry are B3a's (owner of both files).
- **B3b-H2Q** (caloric_eos.f90, mol_rates.f90; h2_rovibrational_identity 19 assertions): one H2 partition

function over the observed Roueff et al. 2019 ladder (302 levels, zero at $v=0, J=0$) gives $u_{rv}(T)$ as its first moment and the chemistry's K_{eq} as its free energy; the Huber-Herzberg model constants are deleted. The RED (0.6 to 8 percent in u_{rv}) was the model ladder's level placement (no centrifugal distortion: J^4 signature, 100 percent of the gap to 2000 K), not a defect of the sum; independent check: K_{eq} against NIST-JANAF (Shomate) now 0.9997 to 1.0008 over 600 to 5000 K (was 1.010 to 1.101). Impact: `wasp_full` identical; `mol_base_handoff` T 1.3e-5, `n(H2)` 3.1e-4; `mol_diffusion` T 3.2e-5; `mol_metals` at 400 steps `n(H2)` 5.6e-3 at 1.17 Rp; `Mdot` unchanged. Findings: a third copy of the ladder sum in `molecular_infrared_cooling.f90` 212-215 (to fold into `h2_partition_function`); `element_census_tests` does not compile against the live tree (no `_free` argument, another worker's signature change; to fix at the next single build); the load average reached 38 with the parallel long runs, so future briefs pace the 12000-step impact runs.

- **A3b** (`diffusive_photochemistry.f90`; `carrier_retry` 89 assertions): the carrier controller's exhaustion reads the run mode and returns a status (`carrier_interval_covered` / `exhausted`) the caller reads. `phys:` unchanged (restore, print, stop until B3a takes the status). `init:` the entry state is restored, the interval is recorded in the attempts ledger (count, first/last step, worst row and both scales), the carrier history is marked not certifiable (`carrier_history_certifiable()`), and the march continues from the entry carriers. The accepted CO record is indexed by A0-impl's `ledger_family`. Impact: `mol_carrier`, `mol_base_handoff` bitwise identical; `oxygen_chemistry` (`init`, its default) now runs its 1000 steps, exit 0, with 998 exhausted intervals (first at step 2, worst CO at cell 313), `log10 Mdot` 9.61 against the 20-bin 9.60, and a final certification refusing on four carrier rows at 1.0: its carriers are frozen at the step-2 values for the whole run, which is exactly what the mark and the ledger say. The refused row is the real imbalance beside a clamped cell (`|res|/full` 4e-12 against a 1.4e-14 round-off bound), open physics of the excluded CO model, not a controller defect. Its golden is re-taken at the end of this series with the mark noted in the case README.
- **B5pre** (`steady_newton.f90`; suite `steady_completion_flag`): the JFNK and PTC completion flag is decided AFTER the certification of the returned state, from the same record the certification prints; a solve that met the gate at the loop top and not at the returned state ends `info = 2` naming the row. Measured on `wasp_full_newton`: loop-top `||R|| = 5.5e-6`, returned state 4.2e-4; the existing own-composition consistency pass opens the gap (0, 1, 2, 3, 8 sweeps: 5.5e-6, 1.9e-4, 3.2e-4, 3.9e-4, 4.2e-4), so the hydro-only Newton converges to a state that is not a root of the coupled hydro + composition system: the loop-top gate measures a lagged residual. Consequence today: EXHALE_main treats `info = 2` as a solver failure and falls back to marching, which ends on a du plateau with a state that certifies worse than the JFNK state; assigned to B3a (`info = 2` = a failed stationarity claim, written uncertified, exit 2, no fallback) and to B5a (`steady_newton.f90`: the residual evaluated on the self-consistent composition of the current iterate, so gate, merit and returned state measure one thing). Also found: the JFNK stop message prints an unset `resid_th` (B3a); `certification_row_measure` and `relnorm_over_cells` are one expression written twice.
- **B3b-PE** (`h2_photo_channels.f90`; `photoevent_energy_ledger` 105 assertions): the 19.725 eV missing from the H2 double photoionization event is the kinetic energy release of the two protons: 51.4 eV is a vertical threshold (Yan et al. 1998 section 4), the asymptotic product energy is $D_0(\text{H}_2) + 2 I(\text{H}) = 31.675$ eV, and energy

conservation at the threshold fixes the fragment kinetic energy with no free parameter (Franck-Condon check $e^2/R_e = 19.42$ eV, 1.5 percent). Second defect found: the 33 to 41 eV neutral window dissociates into $H(2p) + H(2s)$ (Chung et al. 1993 p. 887), so 20.398 eV per event is internal excitation leaving as prompt Lyman-alpha and two-photon continuum, which the integrand charged to heat. `h2_channel_energy_recipients` (reservoir, electron, fragment kinetic, radiated; sums to $h\nu$) with the table of channels and their sources. Channels M and S already closed; every atomic absorber and all 17 metal ions correct (`wasp_full` byte-identical). Impact: `mol_sec_ion` $T 4.2e-4$, `heat_H2` $4.5e-2$; `mol_base_handoff` $T 2.1e-5$; `Mdot` unchanged. The four-line wiring of the two new terms into the heating integrand of `util_ion_eq.f90` is handed to B4-6 (owner of that file). Open: suprathreshold H_2 dissociation credits heat but destroys no H_2 ; the double channel has no absorber row of its own; channel S has no source for a fragment kinetic split.

- B3b-H3** (`h3p_cooling.f90`; `h3p_cooling_limits` 17 assertions): the Miller et al. 2013 fits as the paper defines them: Table 5 is the LTE emission of one molecule per steradian ($\Lambda = n(H_3^+) 4\pi E(T) s$), Table 6's $s(T, n_{H_2})$ is scaled LTE with H_2 the only collider (Oka and Epp coefficient; the paper's upper-limit caveat at the code site); below the table's $1e6\text{ cm}^{-3}$ the non-LTE factor follows the exact collisional limit (linear in n_{H_2} , zero at zero), replacing the clamp at the lowest column; above $1e14$ the table edge is held; the four Table 5 segments do not join (measured with the paper's coefficients: +43 percent at 300 K, -0.11 percent at 800 K, -2.4 percent at 1800 K) and the code joins them by a stated linear blend in $\log_e E$ over the 5 percent below each join (Table 6's 300 K value supports the upper polynomial); B1 T6.3 reworded to say so. Published values reproduced per segment to the paper's precision. Impact: the H_3^+ cooling peak unmoved; T moves $1e-5$ to $2.7e-4$ in the three molecular cases; the clamp removal changes cells beyond 1.22 Rp where $n_{H_2} < 1e6$ by more than 50 percent at $1e-10$ of the peak (emission from gas with no collision partners, gone); `wasp_full` identical. Four domain counters exposed (B6 wiring is the certification owner's).
- COLit, completed with the user's PDFs** (Visser 2009, McElroy 2013, Venot 2012, 2020; `docs/co_destruction_rates_literature_20260906.md`, 1419 lines): CO photodissociation as Visser et al. define it (37 bands inside EXHALE's 912 to 1110 Å beam, unshielded $2.6e-10\text{ s}^{-1}$ in the Draine field, shielding tables in $N_{CO} \times N_{H_2}$ for four parameter sets, T_{ex} raised only to 512 K because no data exist for dissociation out of $v'' = 1$); evaluated on the oxygen_chemistry run's own base columns ($\log N_{CO}$ 18.96, $\log N_{H_2}$ 21.80) it gives $\tau_{dest} = 2.8e8\text{ s}$, twelve decades faster than the barrier channels and still two to six decades short of $\tau_{dest} \ll \tau_{res}$ at the base; zero in `mol_base_handoff`, whose spectrum carries no flux longward of 911.8 Å. RATE12's paper prints no entries (the data file is still needed for the ion channels); Venot 2012 is valid 300 to 2500 K and 0.01 to 100 bar (EXHALE's base at $1e-6$ bar is four decades below); none of the six sources covers the 3000 to 4000 K band where the ceiling fires. The CO exclusion (B1 decision 7) stands, now quantitatively. If a CO photodissociation term is ever added it is a fourth absorber on the existing Lyman-Werner beam with a second column (N_{CO}) and the cell-mean form of `lyman_werner_dissociation_rate`.
- TESTINFRA** (`grid_and_gates/run.sh`, `physics_probe/run.sh`, their READMEs): both suites take EXHALE_OBJS, EXHALE_EXE, EXHALE_TEST_OUT with the semantics of `spectrum_type/run.sh`; default be-

havior byte for byte as before; against a private build both run without touching build/ objects (the six grid_and_gates shell tests still place their run copies under build/tests/). A dead fallback that wrote a build stamp the closure never compiled is fixed. Also (advisor): the two Python source scanners now skip .ipynb_checkpoints/ directories, which hold editor auto-save copies of four modules (git-ignored) and had made threshold_literals_written_once red on a stale copy of cross_sec.f90; the Makefile source list is explicit and never reached them.

- **B3b-ENTH** (molecular_reaction_heat.f90, oxygen_rates.f90; 78 Fortran
 - 37 Python assertions): one table species_formation_energy(isp) for every species column (H/He stages, 27 metal stages cumulative from the neutral, H₂/H₂⁺/H₃⁺/HeH⁺, OH/H₂O/CO, plus free electron, H(n=2), O(1D)) from the single B1 reference; the 17 reactions of the H₂/He network are a stoichiometry table and each heat a difference of table entries; the Shomate table stays with its reference stated, every use of it being a balanced-reaction difference (the zero cancels), and the oxygen and carbon entries of the one table are differences of it; one D0(H₂) (mol_rates::h2_dissociation_energy_ev, reproduced by the fit to 1.5 meV). Physics defect fixed: the oxygen photolysis thresholds were 298.15 K reaction enthalpies; they are now 0 K dissociation energies (the lowest Shomate interval's F coefficient is the 0 K enthalpy), 0.9 to 1.0 percent lower, deposited heat +0.8 percent at the oxygen base cell. Impact: mol_base_handoff, mol_sec_ion, mol_metals byte-identical; oxygen_chemistry T +0.4 percent at 1.02 Rp, Mdot 9.61. Open for B3c: the undeposited collisional oxygen heats and the He 2³S associative branch (D0 C25; the table exists); the B4-2 Python diagnostic lacks four oxygen entries; formation_energy_report has no caller; a stale threshold comment in water_photolysis.f90:281.
- **C3** (six input.inp and READMEs; inputdata/sed/README.md): the planet folders and benchmarks of HD 209458 b, HD 189733 b and WASP-121 b load their literature SED files (Spectrum type: Load from file...; Power-law index removed, LX/LEUV kept because mandatory and overwritten by the grid integral, which the setup report shows); every README records the configuration of record with the existing power-law products listed by mtime and the reruns left to a separate instruction; read_sed accepts every file at its configuration's floor (the H(n=2) edge in all six). log F_{XUV} at the orbit moves 3.193 -> 3.127 (HD 209458 b), 4.438 -> 4.311 (HD 189733 b; the recomputed LX/LEUV reproduce Salz Table 3), 6.205 -> 6.204 (WASP-121 b). Advisor follow-ups: HD 209458 b's orbital distance set to 0.04707 AU (Oklopčić and Hirata 2018, the reference the run is compared with; the input said 0.0480 while the file was scaled to 0.04707); the setup report's flux ratio format (F8.3 overflowed when the sub-Lyman band dominates); the three benchmark directories added to the coverage script's inventory (all pass). Known: Load IC? True in the two planet folders is broken by the grid guard (pre-rebaseline IC), the same class as the hp_* regeneration.
- **COLit, revised with RATE22, Moses 2011, Tsai 2021** (document now 1956 lines): RATE22 (Millar et al. 2024, A&A 682, A109) carries 'He+ + CO -> C+
 - O + Heat 1.6e-9 cm³ s⁻¹, measured, accuracy A, all products EXHALE species; it has noH+ + CO, CO + MorCO + H2destruction entry, and itsTu = 41000on 6562 entries is

an undocumented "no upper bound" marker. On the oxygen_chemistry profile the He+ channel with the Visser shielding gives $\tau_{\text{dest}}/\tau_{\text{res}}$ from 6.7×10^2 at the base to 9.6×10^{-5} at 1.14 Rp : **the one-sided destruction model is constructible above the helium ionization front and not below it**; the boundary is compositional, not the ceiling's thermal switch. Decision 7 (exclusion) stays until T5.1 is implemented; B1 section 5.2 revised, B3b-CO scheduled after B3c. Moses 2011: no stated envelope, one fit interval 500 to 2500 K, no shielding. Tsai 2021: 0.1 nm continuum opacities, no CO self-shielding; EXHALE's `slyman_werner.f90` is the working precedent. The literature requests are closed except Baulch 1976 vol. 3, which blocks nothing.

- **B4-6** (`utilities.f90` `cell_mean_attenuation`, `util_ion_eq.f90`; `xuv_cell_mean_attenuation` 56 assertions): the photoionization and photoheating sums of a cell use the exact cell mean of the attenuation, $\exp(-\tau_{\text{out}}) (1 - \exp(-d\tau))/d\tau$ (composite Gauss when the 2D correction is on), instead of the inner-face value the top-down columns gave; one definition for H, He, He 2³S, H₂'s four channels, the metals, the single-absorber rates, `q_abs` and the secondary-ionization integrands. Inner-face error: 0.5 percent at $d\tau = 0.01$, 42 percent at

1. Photon conservation on the written states: cell mean 1.000000000 of the incident flux, inner face 0.9725 (`wasp_full`), 0.9843, 0.9849 (the loss sat at the ionization front). Impact: `wasp_full` (24000 steps) $x(\text{H}^+)$ 8.3×10^{-2} at 1.32 Rp , $\dot{M}$ +0.74 percent; `mol_base_handoff` $\dot{M}$ +0.93; `mol_sec_ion` +1.01; `hydrostatic_column` unchanged. B3b-PE's two heating terms wired into the integrand (its own numbers reproduced cell for cell). Advisor: the single-precision 1.0×10^{-18} cross-section unit literal (33 sites, 4×10^{-8} relative on every rate) made double. Finding to resolve at the gate: `wasp_full` on the mid-series tree plateaus at $du = 4.2 \times 10^{-3}$ from step 22000 and never reaches 1×10^{-3} (the golden stopped at 16230), identical in the item's before binary, so it belongs to an in-flight change (B3a edits `EXHALE_main.f90`) or to an earlier landed one; measured after B3a.

- **A2tol** (`measurement`; `a2_certification_contract_20260906.md` section 9): the `du` stop and the hydrodynamic row measures are not measures of one thing (over a decade of `du` on `wasp_full` the mass row falls at order 0.2, the momentum row rises, the energy row is not monotone; `mol_carrier`'s carrier row falls at order 0.08 over 12000 to 48000 steps); the JFNK returns an energy row 41 to 47 times its target at every rung of a two-decade `Resid tol` ladder (structural; B5a); the elemental transport rows cannot be measured on the marching route (the operator advances 1×10^{-6} of the way per CFL step) and the diffusive cases never reach the steady route (`du` stalls at 5 to 6×10^{-2} against the 1×10^{-2} hand-off). Round-off floors on the converged JFNK state: mass 1.0×10^{-13} , momentum 5.1×10^{-12} , energy 1.6×10^{-11} (residual bitwise reproducible). Recommended and adopted (applied by B3a in `certification.f90`, provisional until B5a): `cert_tol_mass` 3×10^{-12} , `cert_tol_momentum` 1×10^{-8} , `cert_tol_energy` 5×10^{-5} , `cert_tol_element` 1×10^{-8} (with the homopause regime caveat), carrier, closure and level tolerances kept. At these values `wasp_full_newton` certifies with `Resid tol`: 1.0×10^{-7} (marginally at 1×10^{-6}) and nothing as shipped. Noted for the user: the shipped default `Resid tol` 1×10^{-5} is one to two decades looser than the solve reaches at no cost.

- B5a** (steady_newton.f90; suite steady_selfconsistent_residual): eval_residual eliminates the composition AT the state it evaluates (the equilibrium sweep iterated at fixed conserved variables until the composition stops moving), so the loop-top gate, the line-search merit, the returned state and the certification measure one object; the gate equals the certification to $1e-12$ by construction. Finding: the eliminated residual's Jacobian carries the summed Picard response $(I - M)^{-1}$, and a finite-difference probe with a convergence test stops after one pass (the seed already is the state's composition), so the Newton direction ascended; the probes now run a FIXED sweep count taken from an elimination seeded at a different state (about 15 terms), and the "frozen-composition Jacobian" the brief offered as a fallback is exactly the arrangement that fails. wasp_full_newton: info = 0 at $\|R\| = 4.4e-7$ in 9 iterations and CERTIFIED (was info = 2 at $4.8e-5$ after 26), state moved $1.0e-4$ at most, $\log_{10} \text{Mdot}$ 13.27, cost 1.49x. mol_base_handoff identical. Open: Resid tol values were calibrated on the lagged measure; the other Solver: Newton cases and the PTC route are measured at the series gate; n_selfconsistent_max comment stale (B3a); the duplicated row-measure expression (B3a).
- HYG-absorbed** (water_photolysis.f90, utilities.f90 comment; driver absorbed_fraction_switch, 5 assertions): the FUV band rates take the XUV beam's absorbed_fraction_per_unit_depth by use association; the local copy was bitwise the same over 1501 points and is deleted; the series switch and its error bound (about $1e-8$ on the closed-form side of $d = 1e-8$) stated once; mol_base_handoff, oxygen_chemistry byte-identical. Advisor: the B4-6 driver's reference carried the same single-precision $1.0e-18$ the production code had; made double with the production change, the driver is green again (51 assertions had turned red on the $4e-8$ mismatch).
- B3a** (EXHALE_main.f90, new attempted_step.f90, certification.f90, parameters.f90; suite attempted_step, 87 assertions): a marching step is one trial of the fourteen operations from a checkpoint covering everything rows 1 to 12 write, evaluated at one adoption boundary before update_map_end_step, retaken at half dt from the checkpoint when refused (cap 8, floor $dt/2^8$, a reduced dt bounding the next step); the clock and the accepted-step count move only there; exhaustion stops in phys and continues flagged in init. Impact: wasp_full (300 and its du stop at 14496 steps, $\log_{10} \text{Mdot}$ 13.33), hydrostatic_column, mol_base_handoff, mol_carrier bitwise identical against a control built from the same snapshot; cost within run-to-run spread. Measured and reported, not gating: the time-discrete hydro row (1.92 at a cold-start base where the momentum scale vanishes) and the energy identity T1.5 (-0.20 to -0.74 on wasp_full, the size of the row-8 projection, the RED of AT-1a); the step-doubling estimate gives order 0.89 for the whole step (the operator split is first order), cost 12 percent at n_err_every = 20. Two defects of the item caught by its own impact runs and fixed. The mid-item requests landed (A2tol tolerances, info = 2 written uncertified with no fallback, the ledgers in the summary, the H3+ domain counters, one row-measure definition, the n_selfconsistent_max comment). The B4-6 "du plateau" was the mid-edit tree; the final tree reaches its du stop. Advisor decision on the exit status: a run that DECLARES a stationary state and fails certification exits 2 in every mode (A0's init exemption covers physical-step claims only); the mode gate is removed (A2exit). Small items of its report assigned (HYG-B3a: operator status on every path, conduction accessor, step-level carrier verdict, optional volume measure) and the -00 floating-point exception on hydrostatic_column under diagnosis (O0FPE).

- **B4-6b** (`lya_rt.f90`, `excited_hydrogen.f90` field evaluation; driver `lya_beam_cell_mean`, 37 assertions): the stellar Lyman-alpha depth was a trapezoid between cell centers (neither the cell's face depth nor its own, every depth short by half the outermost cell); it is now the rectangle rule of `calc_column_dens` on the cell's own opacity, with the cell's own depth and its star-ward face as outputs. The beam transmission is $\text{erfc}(c \sqrt{\tau})$, not exponential, so its cell mean is a 3-point Gauss rule in $u = \sqrt{\tau}$ (mean and mean of the square, since the trapping makes the field quadratic in the transmission); the photosphere factor's mean is closed form with $\log_{10}$. `jlya_escape_prob` keeps the cell center by Neufeld 1990 section II (the escape fraction is a point property of where the photon is created). Impact: `hydrostatic_column`, `mol_base_handoff` byte-identical; `wasp_full` reaches its du stop at step 14442 on both binaries, wind unmoved ($1e-7$), the field $1.6e-3$ and the Balmer proton source $6.5e-4$; the face rule over-counted the column's scatterings by 2.4 to 2.7 percent. Carried forward: the FUV band B2 line transmission in `util_ion_eq.f90` still reads a face value (B4-6c).
- **HYG-B3a** (`diffusive_photochemistry.f90`, `certification.f90`, `steady_newton.f90`, `viscous_conduction.f90`; suites `carrier_retry` 97, `energy_update` 48, `certification` 29): the carrier operator sets its interval status on every return path (already true since A3b; now stated and tested); `carrier_last_verdict_of_run` renamed for what it is, plus a step-level `carrier_step_verdict()` reset at every transport step; `conduction_floor_cell_hits` exported like the energy update's; the volume-weighted companion of `certification_row_measure` optional and no longer computed where discarded. All impact cases bitwise identical (`wasp_full_newton` finish: `info = 2` at $||R|| = 7.3e-6$ against the new `cert_tol_momentum = 1e-8`, re-anchored by A2tol2). Left for the gate (advisor): the now-unneeded pre-set in `EXHALE_main.f90` and a stale field comment in `attempted_step.f90`.
- **A2exit** (`EXHALE_main.f90` exit block; `run_mode` suite 30 assertions, `certification` shell gate): the certification exit status follows the CLAIM, not the mode: a run that declared a stationary state (du stop, residual gate, JFNK finish) and was refused exits 2 in every mode with its outputs written; a capped or plateaued run exits 0. `wasp_full_newton` now exits 2 with `info = 2`; `run_check.sh` never reads the status. Advisor: `run_newton.sh` and `run_baseline.sh` (set -e loops) now report a status 2 instead of aborting. Open: the `EXHALE_PTC=1` route stops before the certification and so claims a stationary state with no verdict (B5).
- **B4-4a** (`Apply_BC.f90`; `hydrostatic_residual` +1 and new `free_outflow_boundary` 18 assertions): the outer free-outflow ghost is the isothermal hydrostatic continuation of the last cell ($\rho_g = \rho_N \exp[-(\phi_g - \phi_N)/(p_N/\rho_N)]$, p_g by the same factor, $v_g = v_N$), one rule for PLM and WENO3, replacing the zero-gradient copy (PLM) and the linear extrapolation (WENO3). The outer cell's momentum residual on the analytic column falls from 0.750 of its weight at every N to $4.4e-3$ to $1.0e-4$ over N = 250 to 2000 (order 1.8 PLM, 1.6 WENO3); interior and base unchanged; four ghost rules measured, this one 30 to 40x better. Velocity: the copy (a continuity-exact rule stopped fastest but divides by a ghost density the stratification makes small and was the one rule under which `wasp_full_newton` segfaulted at -O3). Two premises of the brief corrected: `wasp_full`'s edge is subsonic (Mach 0.18 to 0.96; `mol_base_handoff` is the supersonic case), and a supersonic edge blocks the ghost only through the Riemann solution, not the stencil, so byte identity is not available. Impact: `mol_base_handoff` one cell above $1e-4$; `hydrostatic_column` $3.5e-3$ at cell N; `wasp_full` du stop 16208 ->

17135 steps with the base-breathing phase moved (11 percent in rho at 1.27 Rp, mean mass flux above r_{flux} 0.078 percent); `wasp_full_newton` Mdot 13.27 both. Removing the outer residual does not by itself stop the marched hydrostatic column's drain (still 0.79 c_s at 300 steps). Reported: a reproducible -O3-only segfault reachable from the rejected velocity rule (latent; O3SEGV to diagnose); `EXHALE_main.f90` 2538-2540 ghost-counter comment stale (B3c); B4 design section 8.3 (ii) historical.

- **O0FPE** (`util_ion_eq.f90`; shell test `fpe_traps` in `grid_and_gates`; `run_fcheck.sh` header): the -O0 failure was two defects. (1) An integer division by zero in `EXHALE_main.f90` about 1234: `mod(count, n_err_every)` with `n_err_every = 0` in init mode, reached because `.and.` does not short-circuit (-O1 and above happen to skip it); fix handed to B3c (owner of the file). (2) A real floating-point overflow in `he_rec_coupling` (`util_ion_eq.f90`:2813): $1100 \exp(1.2/T4) \sqrt{T4}$ at 17 K is 3.9e309, an Inf consumed as `ne/Inf = 0` into the correct low-density limit; the product is now formed only above $T4 = 1.71e-3$ and the limit value taken below, byte-identical on `hydrostatic_column`, `wasp_full`, `mol_base_handoff`. A third site of the same class (the stall detector's huge sentinel) also handed to B3c. `run_fcheck.sh` states why it builds at -O1.
- **B4-6c** (`util_ion_eq.f90` `fuv_lw_photon_field`; driver `lya_band_transmission` rewritten, 16 assertions): band B2's line factor is the cell mean of the stellar Lyman-alpha transmission over the cell's own line-center depth (product of the two means with the continuum mean formed inside `water_photolysis_rate`; the covariance shortfall 6.0e-6 against a 1.4e-4 bound, the face rule 1.2e-2 high on the budget); the top cell carries its own mean instead of a neutral 1. The face rule was 1.0018x on the `oxygen_chemistry` state, 1.095x on an exponential H I layer and 18.2x in the base cell of a molecular base whose line-center column reaches $1e7$ in one scale height. Consumers: the H2O and OH photolysis in B2 (both `ionization_equilibrium` call sites and the carrier photolysis), the output columns, the FUV heating column; the H2 channel reads LW only. Impact: `wasp_full`, `mol_base_handoff`, `mol_sec_ion` identical; `oxygen_chemistry` moves by factors at 45 steps, shown by a control (a 4e-5 perturbation of the face value diverges the same way) to be the case's transient amplification, not the size of the change; the case is not usable as a controlled point until it converges. Finding: `write_output.f90`'s FUV band ledger closed form $N_{\text{ph}} (1 - \exp(-\tau))$ carries no line factor for B2 and its other rows are 0.79 to 0.98 off (LEDGER-FUV assigned).
- **O3SEGV** (diagnosis, no change): the -O3-only segfault B4-4a reported was a null F0 argument of `jv_product` in a `pgmres` clone the compiler had specialized for a null pointer, i.e. a defect of the source text of one intermediate build (13:08) between two edits of `steady_newton.f90`; every later build carries a plain `pgmres`, HEAD plus the rejected ghost rule completes `wasp_full_newton` three times, AddressSanitizer finds nothing over a full run. Correction to B4-4a's premise: `-g` changes no generated code (`.text` bitwise identical at -O3 with and without it). Noted: at the time of the diagnosis the working tree stopped `wasp_full_newton` at step 0 on the energy floor (B3c in progress).
- **LEDGER-FUV** (`write_output.f90` `write_oxygen_chemistry`; suite `fuv_band_ledger`, 15 assertions): the FUV band ledger is three labeled statements: (a) closure of the rated absorptions against the beam's loss between the cell faces (holds for any rate form, $1e-16$); (b) the beam budget with each band's own line factor and the unrated share named per band (none for B1/B3/B4, which is therefore the cell-mean test; H I resonance scattering

for B2, a share; for LW a sign-changing disagreement); (c) energy; (d) state drift. Two defects of the ledger fixed: it multiplied post-sweep densities by a pre-sweep field, and summed the two inner ghost cells. Data rows of every output identical; only the FUV_bands.txt trailer changes. **Finding (LW-NORM assigned):** the LW absorbers take 45 percent MORE photons than the beam loses ($1.27\text{e}13$ against $8.73\text{e}12$) and $249\text{ erg cm}^{-2}\text{ s}^{-1}$ against an incident 171.5: two independent normalizations of one absorption, the level-resolved dissociation table over p_lw_absorbed against the DB96 eq. 39 band share (lyman_werner.f90, h2_self_shielding_table.f90). Also: docs/a2_oxygen_option_design.md 882 documents columns that no longer exist; the a2_m2 kinetics check asserts the beam identity for every band.

- **A2tol2** (measurement and decision; a2_certification_contract_20260906.md section 10): the section-9 anchors re-measured on the self-consistent JFNK residual of B5a. The returned row is now 0.36 to 0.93 of the solve's target (section 9 measured 41 to 47 above it). wasp_full_newton at Resid tol: $1.0\text{e}-8$ certifies (info = 0, 14 iterations, 658 s, log10 Mdot 13.27; rows mass $2.6\text{e}-13$, momentum $1.3\text{e}-9$, energy $6.275\text{e}-10$); the PTC route stalls from a cold start; wasp_he23off_newton, newton_rsw01, newton_rsw05 at the default $1\text{e}-5$ refuse on the mass and momentum rows at cell 499; the four Load IC Newton cases (jfnk_hd189, jfnk_hd189_tight, ptc_warm, heh_1_newton, renamed 2026-09-16 from armD_D2_newton) are stopped by the grid guard and need their IC regenerated on the current grid before they can be measured. **Applied:** cert_tol_energy $5\text{e}-5 \rightarrow 1\text{e}-6$ in certification.f90 ($5\text{e}-5$ had been anchored on the lagged-gate energy row $4.700\text{e}-06$ and could refuse nothing; $1\text{e}-6$ is a decade above $1.212\text{e}-07$, the largest energy row of any state whose solve met its own target, $5.7\text{e}4\text{x}$ above the floor $1.765\text{e}-11$); mass $3\text{e}-12$ and momentum $1\text{e}-8$ kept; the report header now names sections 9 and 10. backup/regression/wasp_full_newton/input.inp gains Resid tol: $1.0\text{e}-8$ as its configuration of record, the only setting at which that case certifies (Mdot unchanged at this precision, cost +11 percent); its golden moves at the series gate with everything else. Verified: private build, certification suite green (29 assertions). Not changed: the other three Newton cases stay at the default so that the matrix keeps a refused claim (exit 2) under test.
- **LW-NORM** (diagnosis; no executable statement changed): the 45 percent excess of the FUV band ledger is two WAVELENGTH BANDS, not two fits. The self-shielding table's line list runs to 1200 Å while its sigma_pump is normalized per photon of 912-1110 Å; the beam loss is the DB96 eq. (39) equivalent width of 912-1110 Å. MEASURED: the column integral of sigma_diss/p_eff reaches 1.51538 at the top of the column axis against 1.519288 for the photon content of 912-1200 Å over 912-1110 Å of a flat spectrum (0.26 percent), is exactly density-independent, and reproduces the shipped oxygen_chemistry ratio 1.4544 at its base column where eq. (39) clamps to 1. p_eff already separates dissociations from absorptions, so the heating takes fragment energy only (not cause 2). Landed: lyman_werner.f90 header section 3a with the invariant and the measurement, its section 3 "2 percent" statement corrected to the live table's thin-limit cross section $2.6396\text{e}-17\text{ cm}^2$; lyman_werner_cell_mean probe gains two assertions and a photon-conservation diagnostic (9/9); fuv_band_ledger/run.sh comment. Advisor follow-up: the table's generator inputs (nine abs_T####.npz, 27 CLOUDY rate files, the line-by-line scripts, the Meudon level and line data, 29 MB) moved from the 2026-09-03 scratch directory into src/utis/h2_shielding_lbl/ with a

README, and h2_shielding_table_line_by_line.py's template brought in line with the hand-renamed axis dummy of h2_shield_locate: the generator now reproduces the shipped module BYTE FOR BYTE (MEASURED). The repair, table on 912-1110 A or beam widened to 912-1200 A with the B1 flux joined, is item 6 of To_be_determined_by_user_20260906.md (recommendation: widen), because the second redefines the Stellar LW flux key. Still to correct after B3c releases write_output.f90: the ledger's LW label says "two fits" and should say "two bands" (with the header_names_lw_fits assertion of fuv_band_ledger).

- **HYG-HeCI** (test hygiene, advisor): physics_probe was red on three helium_collisional_ionization_by_level assertions (1e-13 at 1e4 K to 1.0e-4 at 5e4 K). Not a code defect: batch 2c replaced the single-precision literal 3.940e-11 erg (24.6 eV) of eval_cool with the global e_th_HeI_erg (24.587389 eV), and the test's reference still carried the old literal with a comment calling it "a separate open item". The reference now uses e_th_HeI_erg and e_th_HeTR_erg; physics_probe PASSED on the current tree (private build). The LW-NORM worker's attribution of this to a concurrent Cool_coeff.f90 edit was wrong: that file's uncommitted diff is the landed 2c-CI / MASS2 work.
- **LW-NORM-B** (user decision 2026-09-06: recommendation (B) adopted; landed): ONE Lyman-Werner band, 912-1201 A, with one normalization.
 - **The band.** oxygen_rates.f90 carries four FUV bands, not five: the Lyman-Werner interval is 912-1201 A and band B1 (1110-1201 A) is gone into it. 1110 A was an edge because Draine & Bertoldi end the Solomon process there for interstellar $v = 0$ gas; at the 700-3200 K of a planetary base the vibrationally excited levels pump longward of it, and the self-shielding table's line list always ran to 1200 A. The two halves carried the same H2O yield triplet, so no quantum yield moves; the band-averaged cross sections are re-formed over the whole interval and MEASURED again from the same Photochem files: $\sigma(\text{H}_2\text{O})$ 8.51337e-18, $\sigma(\text{OH})$ 4.74371e-18 cm², (H2O) 12.0526 eV, (OH) 11.3829 eV, $= 2hc/(912+1201 \text{ \AA}) = 11.7354 \text{ eV}$. They agree to seven digits with the merge of the two shipped halves, and they reproduce the values the band carried before it was split in 2026-08.
 - **The key.** Stellar LW flux now means the 912-1201 A band flux at the orbit; Stellar FUV B1 flux is retired and a file that states it stops the run with a message naming the merge and asking for the two numbers added. Spectrum type: Load integrates 912-1201 A. The resolved dump loses fuv_band_B1_flux. The inputs that carried both keys were restated to their sum: 343.0 + 137.9 = 480.9 for the HD 209458 b spectrum (Update_EXHALE_stage1.md records the 912-1201 A integral of that spectrum as 481.0 and its halves as those two numbers), and the LW-only cases (mol_lyman_werner, heh_1_lw_40k (renamed 2026-09-16 from armA_LW; mapping in docs/named_case_audit.md section 6), heh_1_lw_12k (renamed 2026-09-16 from armD_D2_LW), heh_1_lw_newton (renamed 2026-09-16 from armD_D2_LW_newton)) from 343.0 to 480.9 so that they keep the same star. The README_HOWTO HD 189733 b recipe goes from LW 600.1 + B1 648.4 to LW 1248.5.
 - **The table.** src/utls/h2_shielding_lbl/lbl_table.py normalizes per photon of 912-1201 A (wl_HI, wl_max, and E_LW_PHOTON_ERG the mean photon energy of that band); F_BAND_ERG is restated as $343.0 * 289/198 = 500.6$ with the derivation in the file, and the file now records that it cancels out of

sigma_pump exactly. The nine abs_T####.npz were recomputed (run_abs.py, 23 to 251 s each) and h2_self_shielding_table.f90 regenerated. MEASURED: the generator reproduced the shipped module byte for byte before the change, so the only difference is the band. sigma_pump at the bottom of the column axis is $1.701\text{e-}17$ (700 K) to $1.887\text{e-}17$ cm², which is DB96's own $2.557\text{e-}17$ per 912-1110 A photon restated per 912-1201 A photon ($1.676\text{e-}17$) to 1-13 percent, the same agreement as before.

- **The beam loses what the table rates.** The band share A is no longer the DB96 eq. (39) equivalent width; it is the column integral of the pump cross section $\sigma_{\text{diss}}/p_{\text{eff}}$ of the same table, in closed form (h2_lw_band_photon_fraction_absorbed, exact for the module's own interpolant knot interval by knot interval). MEASURED: A at the top of the column axis is 0.6903 to 0.9974 over the whole (T, n_H) grid, where it used to reach 1.5154; the closed form agrees with an independent 20000-step quadrature of the two public accessors to 2e-10. The band ledger's photon count is now the cell mean of the same pump cross section (lyman_werner_band_absorption_rate_cell_mean), on the quadrature the dissociation rate already used, and not the cell mean of σ_{diss} divided by p_{eff} at one column: on cells spanning 2.5 H2 scale heights the second stood 8.0 percent above the beam loss and the first closes on it to 1e-6 (physics_probe/lyman_werner_cell_mean sec. 6). h2_band_equivalent_width and eq. (39) are removed; the two published self-shielding fits stay, called by nothing, for the comparison of docs/h2_self_shielding_cloudy.md.
- **The ledger.** The LW row of output/FUV_bands.txt is a closure like B3 and B4, and its label says so. MEASURED on the oxygen_chemistry state, 5 steps: unrated_frac -3.62e-1 before, -1.18e-2 after, and the LW absorbed energy goes from 233.5 against an incident 171.5 (36 per cent over the band) to 202.3 against 200.0. The residual left is the part of the column above the top of the table's column axis: this case reaches $6.4\text{e}21$ cm⁻² against an axis top of $5\text{e}21$, where the clamped edge cross section drives A past 1 and the beam is capped while the rate is not. The run already warns about that column.
- **Impact.** mol_lyman_werner at its own 12000 steps, before at 343.0 against after at 480.9: log10 Mdot 10.53 -> 10.54, T and rho under $5\text{e-}3$, cooling $6.5\text{e-}2$, the molecular densities and the shielding $1.2\text{e-}1$ at the H2 front ($r = 1.075$), the base H2 column $3.3875\text{e}20$ -> $3.4067\text{e}20$ cm⁻² and the base x_{H2} 0.96288 -> 0.96365. The optically thin rate at the top of the column moves by $480.9/500.6 = 0.9606$ exactly: the band's photon content grew 46 percent and the cross section fell by the same factor, so what is left is that the real spectrum gives 480.9 over 912-1201 A where the flat continuation of the 912-1110 A deck would give 500.6. Deeper cells move -4.7 to -6.0 percent, the wider band's extra lines shielding differently; slightly less destruction leaves slightly more H2.
- **Not repaired here, and it blocks the fuv_band_ledger suite.** With its restated 480.9 the oxygen_chemistry case stops at step 0 on the energy floor. MEASURED, it is not the band merge: the same case stops with the OLD binary at the same restated flux, and both binaries stop or run at chance along the flux axis (old: runs at 343.0 and 440.0, stops at 360.0, 400.0 and 480.9; new: runs at 343.0 and 400.0, stops at 360.0, 440.0 and 480.9). That is the step-0 acceptance knife edge of items B3a and 2 of To_be_determined_by_user_20260906.md. Every ledger assertion is GREEN on the same case at a flux where it runs.

- Also: `p_lw_absorbed` is now written by the sweep and checkpointed but read by nothing, since the ledger takes the pump rate directly. Removing it needs `attempted_step.f90`.
- **B3b-IR** (`molecular_infrared_cooling.f90`; `h2_infrared_line_populations` and `h2_partition_sum_uniqueness` probes): the third copy of the H2 ladder sum (212-215) is gone; the H2 infrared line populations are normalized by `caloric_eos`'s `h2_partition_function`, the one Boltzmann sum over the Roueff et al. 2019 ladder whose energy the equation of state carries and whose free energy sets the $H + H \leftrightarrow H_2$ equilibrium. The probe that greps the tree for a second sum went RED (1) -> GREEN (0). Impact (advisor measured, 12000-step snapshots, `OMP_NUM_THREADS=1`): `mol_ir_bands` (9 output files) and `mol_base_handoff` (7 files) BYTE-IDENTICAL before and after, `Mdot` 7.99 both; the retired copy had an $\exp(-\min(E/T, 700))$ clamp whose only difference from the shared sum's $\exp(-E/T)$ is below double rounding. No golden moves. The worker's draft report stopped at "impact runs in progress"; this entry closes it.
- **LW-NORM-B, advisor verification** (2026-09-06): private build of the landed tree, `physics_probe` PASSED (554 assertions incl. the rewritten section 6: column integral of `sigma_pump` at the top of the axis 0.69 to 0.997 on the whole grid, was 1.515; photons spent / photons lost 0.999999), `spectrum_type` PASSED (154), `fuv_band_ledger` NOT RUNNABLE: its case is `oxygen_chemistry`, whose Stellar LW flux restated to 480.9 (the same spectrum over 912-1201 Å) stops at step 0 with energy update FAILURE at cells 85-88 (T $2.1e4$ K, ionization sweep NON-ROOTs accepted by the relaxation amnesty with residual 1 to 4, then the balance asks for T below the 14.5 K floor). MEASURED by the worker along the flux axis with both binaries: 343 runs, 360 stops, 400 runs (after) / stops (before), 440 stops (after), 480.9 stops: a step-0 acceptance knife edge of the OLD code at cells with no H₂, items 1 and 2 of `To_be_determined_by_user_20260906.md` (decision 4: reject-and-retry; amnesty cap), not of the band. The flux is NOT tuned to make the case run; the case and `fuv_band_ledger` stay red until those two decisions land. Code read: `a_lines = min(A, 1)` with transmission $1 - A$ (a band share, not an optical depth) from the closed-form power-law integral of the tabulated `sigma_diss/p_eff`; both the rate and the share are read at the local T and n_H of the cell for the whole column above it (homogeneous-column table), a comment in `util_ion_eq.f90` now says so. Accepted. Goldens untouched; every LW case moves at the series gate (`mol_lyman_werner` 500-step pair: T $3.3e-3$, H_2 0.125 at the front, k_{LW} thin-limit $x0.9606 = 480.9/500.6$).
- **ACCEPT-1/2** (user decisions 2026-09-06, items 1 and 2 of `To_be_determined_by_user_20260906.md`, both as recommended; brief `plan_rev2_brief_ACCEPT12.md`, to start when B3c releases `EXHALE_main.f90`, `energy_semi_implicit.f90`, `ionization_equilibrium.f90`): (1b) a failed energy update refuses the attempted step in both modes and the controller retakes it at half dt, exhaustion exits 2; (2a) the relaxation amnesty accepts a non-root only below a residual cap measured on the matrix, above it the cell keeps the composition it entered the sweep with (new acceptance class 6). Motivation: `oxygen_chemistry` at its restated LW flux dies at step 0 on exactly this pair (LW-NORM-B verification entry above).
- **B3b-IR impact runs, advisor correction** (2026-09-06): the worker's before/after pairs were run with binaries built from the tree as it stood at 12:30, when the B3a controller still refused on the time-discrete hydrodynamic row in init mode: MEASURED from their run logs, 12000 steps accepted of 104389 attempted (`mol_ir_bands`)

and 104594 (mol_base_handoff), 103697 refusals all "the Runge-Kutta stages and the fluxes: the time-discrete hydrodynamic row was out of tolerance", dt collapsed and T stayed within 1 K of the 1140 K initial condition over the whole grid (golden mol_ir_bands reaches 2553 K at 2.18 Rp). The byte-identity of the pair therefore says little about the layer; the statement that the IR refactor changed no arithmetic rests on the emission assertions at five temperatures with identical digits and on retired/Q - 1 = 0 at every probe temperature, which stand. On the current tree the same case accepts 12000 of 12000 attempts with no refusal (MEASURED, private build). The H2 IR tabulation grid extension (advisor, this entry's neighbor above): n_tab 401 -> 496 at the same log spacing to t_tab_hi = 30146 K, old nodes reproduced by the same expression, emission at 10000 K: clamped = direct sum = 1.57875E-18 (was 1.181e-18); molecular_infrared_bands.py template carries the private :: i of the generated data module and regenerates it byte for byte (MEASURED).

- B3c** (EXHALE_main.f90, ionization_equilibrium.f90, energy_semi_implicit.f90, molecular_reaction_heat.f90; new suite coupled_source_step, 30 assertions in the advisor's run): rows 7 to 9 of the enumerated step are one local source step per cell at fixed volume, the temperature and the composition its unknowns, iterated to a fixed point ($|dT|/T \leq 1e-8$, max $|df_{sp}| \leq 1e-8$, cap 20 passes). The composition projection (comp_p_from_T + W_to_U, which rebuilt the energy row at the new particle count) is gone; the energy row is anchored on u_th_old, the thermal energy the cell had at its old composition; rho is intent(in) to the chemistry and the mass sum is a checked diagnostic (worst 6.4e-12, T2.2). The two collisional deposits D0 C25 named (oxygen O1/O1r/O2/O2r/O6, associative He 2^3S) are in as differences of the one formation-energy table. RED reference kept in the suite: the removed update changed u(3) by -0.600 with heat = cool = 0. **Advisor decision on the worker's deviation (accepted):** Delta u_form is NOT added to the energy row. In this code heat and cool are the net THERMAL sources (photoheating deposits $h\nu - E_{th}$, collisional ionization cooling removes E_{th} , every molecular, Penning, Lyman-Werner and oxygen reaction heat is a difference of the same table), so the formation reservoir is already accounted for and the term would double count; MEASURED with it added, wasp_full dies at step 0 with 368 of 504 cells at the floor. Consequence: T1.5/T1.6 as an identity on $u_{th} + u_{form}$ against $dt(heat - cool)$ cannot close; the B3a instrument (attempted_step.f90) is to be restated as $du_{th} = dt(heat - cool) +$ the transport contribution of the step, or evaluated on the source substep alone (item T15-INSTRUMENT, pending). Impact (control = same snapshot with the edits reversed): wasp_full to its du stop, log10 Mdot 13.34 -> 13.34, T 2.5e-3, rho 2.4e-3, metals 1.5e-2, one sign change of v at r = 1.062 (2.5e-4 of the wind); wasp_full 300 steps front one cell apart; molecular 12000-step snapshots up to 40 percent in rho beyond 4 Rp, Mdot 10.58->10.57, 10.53->10.51, 10.58->10.58; hydrostatic_column <= 6e-9. **Cost: 5 to 12 passes per step, 8.1x on wasp_full 300, 10.3x to the du stop (18:59 -> 3:15:07).** Assigned as B3c-COST (re-sweep only the cells not yet at the fixed point; tolerance anchored on a measured convergence study, not chosen to pass). Also in and byte-identical: the carrier status pre-set removal (HYG-B3a) and the two OOFPE sites; fpe_traps passes at -O0 with traps. Left red on purpose: heat_column_matches_breakdown_total 6.886e-3, because write_heat_breakdown_eq (util_ion_eq.f90 2280) is a second hand-maintained copy of the heating assembly that lacks the new deposit, and post_process_adv.f90 ~847 a third: assigned as HEATBRK (one assembly, three callers), not the ten-line patch. Advisor verification: private build of the live tree, coupled_source_step 30/0, energy_update 48/0,

attempted_step 31/0, certification 29/0, run_mode 30/0, acceptance_classes 12/0, grid_and_gates 92/1 (the HEATBRK red).

- HEATBRK, first pass BLOCKED** (2026-09-06): the heating assembly the sweep uses is 210 inline lines of `ioniz_eq` (ionization_equilibrium.f90 2119-2331), not a routine, so the one-assembly fix has to extract it there, a file `ACCEPT-1/2` owned at the time; the worker stopped with no edit, as briefed. Re-measured RED: worst 6.886e-3 (cell 376, the associative $\text{He } 2^3\text{S}$ and collisional oxygen deposits missing from the breakdown) over a 1.24e-5 median floor (the breakdown RE-SOLVES the field and rates on the written state while the heat column carries the rates the sweep produced it with). Interface proposed: `heating_of_composition(...)` in `utils_ion_eq` returning heat and a non-optional `heat_chan(:,17)` (channel list and header owned by the module; LW dissociation and fluorescence as two columns), called by the sweep, by `write_heat_breakdown_eq` (which then forms no rate) and by `post_process_adv` (which today omits the Balmer and associative terms from its theat). **Advisor decision:** `Heating_breakdown.txt` writes the channel array the sweep computed for the written state, not a recomputation; this is the "one state per output file" rule (rev3 section 10.2 item 2, landed) applied to this file, and the only form in which the breakdown total can equal the heat column to round-off. Resumes when `ACCEPT-1/2` releases `ionization_equilibrium.f90`.
- ACCEPT-1/2, advisor verification** (2026-09-06): private build of the live tree, `energy_update` 53/0 (the driver was killed by error `stop 1` before), `acceptance_classes` 22/0 (did not compile before: class 6 did not exist), `attempted_step` green (10 RED before: exit status, history lines, reason breakdown), `coupled_source_step` 30/0, `certification` 29/0, `run_mode` 30/0, `carrier_retry` 97/0. Code read: the kept state of class 6 is `x_entry`, the composition the cell entered the sweep with (built once, also the solve's starting point), heat and cool of the returned composition as section 170 requires. Accepted, including the two deviations: the two-line `steady_residual.f90` change that makes `n_cells_without_chemical_root` count class 6 (without it the controller would not see a refused non-root), and `as_op_projection -> as_op_composition` (operation number 8 unchanged; B3c had removed the projection the name described). Findings taken over: cap 1e2 MEASURED on the whole default matrix at 400 steps (the amnesty population is a cold-start population: 111 events 1e-6 to 8.853 at step 0 of `oxygen_chemistry`, nothing until 8.6e4, then the four cells 85-88 whose energy update failed); with the cap `oxygen_chemistry` runs its 1000 steps and exits 0 (log10 Mdot 9.61), no step refused, fifteen of sixteen cases byte-identical; the comment's "res ~28 in the metals-on molecular gates" is no longer reproduced (they accept no non-root today) and was corrected in place. The energy-refusal path is exercised by the default-off hook `EXHALE_ENERGY_FAIL_AT_STEP`. Outside scope, assigned: "class 4" wording in `certification.f90/steady_newton.f90` and `steady_newton.f90:815 acc_resmax(4)` missing class 6 (HYG-CLASS6, folded into HEATBRK's resume); an H-only run has no acceptance classes at all (pre-existing, B6 list); `oxygen_chemistry` cannot run in phys mode (56 handoff non-roots refused by A0), a statement about that configuration, recorded.
- HEATBRK landed, advisor verification** (2026-09-06; measured on `mol_base_handoff` 200 steps, `wasp_full` 300, `oxygen_chemistry` 1000, the `mol_metals` impact pair still running when this was written): `heating_of_composition` (`util_ion_eq.f90` 1503-1780) is the one heating assembly, 17 named chan-

nels, the total the running sum of the columns in the sweep's original order (sweep heat bit-identical); the sweep calls it once (181 inline lines of `ioniz_eq` gone), `write_heat_breakdown_eq` writes the module array `heat_channel_state` the sweep filled and forms no rate, `post_process_adv` calls the same routine. Private build of the live tree: `grid_and_gates` 93/0 with `heat_column_matches_breakdown_total` 1.678e-16 (was 6.886e-3), `physics_probe` 557/0 incl. `heating_channel_closure` and the scanner `heating_sum_uniqueness` (one exception recorded: the band ledger of `write_output.f90` restates the Lyman-Werner and FUV photolysis event energies, a fourth copy, assigned HYG-LEDGER-EPS). `Heating_breakdown.txt` 18 -> 21 columns; `_adv` profiles move by up to 1.8e-2 (the Balmer and associative deposits the third copy lacked). `heat_channel_state` is class-1 state and enters the attempted-step checkpoint through B3c-COST (owner of `attempted_step.f90`), assigned. HYG-CLASS6 (class 4/6 wording, `acc_resmax` over both classes) in the same landing.

- B3b-CO landed: the CO row carries two published destruction rates** (2026-09-06). `src(ic_CO)` is no longer $\emptyset$. It is $-(k_{D1} n(\text{He}^+) + k_{\text{CO}}) n_{\text{CO}}$ with $k_{D1} = 1.60\text{e-}9 \text{ cm}^3 \text{ s}^{-1}$ (UMIST RATE22 entry 4068, measured, accuracy better than 25 percent, temperature independent, READ) and k_{CO} the cell mean of $(F_{\text{LW}}/\langle h\nu \rangle) \sigma_{\text{CO}} \Theta(\text{N}_{\text{CO}}, \text{N}_{\text{H2}}) \exp(-\tau_{\text{cont}})$ on the 912-1201 Å beam. New: `co_self_shielding_table.f90` (the 12CO block of Visser, van Dishoeck & Black 2009 Table 6, $T_{\text{ex}}(\text{CO}) = 50 \text{ K}$, with their Table 5 beside it for the spread) and `co_photodissociation.f90` (the point rate, the two-column cell mean, the fragment heat). `fuv_lw_photon_field` gains the CO column and returns k_{CO} and Θ ; `carrier_photolysis` rebuilds k_{CO} from the current carrier columns, so the CO layer shields itself inside the relaxation. Constants, all MEASURED: $\sigma_{\text{CO_band}} = 1.0160\text{e-}17 \text{ cm}^2$ per beam photon (Heays et al. 2017 cross section over the 912-1118 Å window CO absorbs in), $\langle h\nu \rangle = 12.8674 \text{ eV}$ (oscillator-strength weighted over Visser's 37 bands), $D0(\text{CO}) = 11.1157 \text{ eV}$ and $q(D1) = +2.2117 \text{ eV}$, both differences of the one formation-energy table; the deposit is 1.7517 eV per dissociation. Both deposits are channels 18 and 19 of `heating_of_composition` (17 -> 19 channels) and are formed nowhere else; `heat_per_co_dissociation` is registered in the uniqueness scanner with the band ledger of `write_output.f90` as its one recorded exception. `output/FUV_bands.txt` gains N_{CO} , k_{CO} and Θ_{CO} and a CO row in the beam ledger (CO is printed apart from `rated_ph` because `beam_loss_ph` carries no CO term: MEASURED, folding it in moved the LW budget residual from -1.8e-2 to -8.2e-2). `output/Oxygen_chemistry.txt` gains the domain record $\tau_{\text{dest}} \leq 0.1 \tau_{\text{res}}$ with the worst ratio and its radius, $\tau_{\text{res}}/\tau_{\text{form}}$ from RATE22 8597, the count of cells above the shielding table's 512 K excitation-temperature limit and the count in which the He^+ channel outruns He^+ recombination. Tests: `physics_probe co_destruction` (27 assertions, RED on a mutated table entry and a mutated photon energy), `fuv_band_ledger` extended. **Impact, MEASURED on oxygen_chemistry at its maxsteps of 1000, the only regression case that carries CO** (a control binary of the same tree with the CO source and the two deposits reverted): both run 1000 steps, exit 0 and give $\log_{10} \text{Mdot} = 9.61$; the after binary costs 628 s against 11 percent less for the control at a matched 150 steps. The $\text{He}^+ + \text{CO}$ charge-transfer deposit is **65 percent of the total heating at $r = 1.089$** , the helium ionization front where CO still survives, and under 1 percent outside 1.05-1.17; the layer's temperature moves by up to 46 percent at $r = 1.023$ and its CO by up to 52 percent. `mol_carrier` at 300 steps is byte-identical in every output file, which is the gate that the new terms are on the same `thereis_oxychem` as the rest of the oxygen path; no other regression case sets Oxygen

chemistry. The domain record of that run: 401374 out-of-domain cell visits, worst $\tau_{\text{dest}}/\tau_{\text{res}}$ 1.34e5 at $r = 1.0002$, worst $\tau_{\text{res}}/\tau_{\text{form}}$ 0.219, 305125 cell visits above the 512 K excitation-temperature limit and 281325 in which the He+ channel outruns He+ recombination. Goldens not refreshed. **P0 recorded:** the He+ ledger is not closed. This reaction consumes a helium ion and neither the carrier row (ion stages frozen) nor the sweep's He+ balance returns it, so where $k_{\text{D1}} n_{\text{CO}}$ exceeds the He+ loss rate both the destruction and that heating peak are over-stated. The fix is a sink in the He+ row of `mol_heh_rows` (`System_HeH_mol.f90`), not this item's file; `dom_cells_HeP` is the observable. **Not done, carried over:** the thermodynamic CO ceiling is still in `limit_to_element_budget`; deleting it changes `EXHALE_main.f90`, `certification.f90` and `write_setup_report.f90`, none of them this item's files. The exact lines are in the item's report.

- **HYG-LEDGER-EPS closed in the same landing** (2026-09-06): the FUV band ledger of `output/FUV_bands.txt` is formed by `utils_ion_eq::fuv_band_absorption_ledger`, beside the heating assembly and reading the same imports, and `write_output.f90` writes what it returns; the `heat_LW` column of `output/Lyman_Werner.txt` is now the assembly's own channel (`heat_channel_state(:, ih_H2_LW_dissoc)`) instead of a second evaluation of $k_{\text{LW}} n_{\text{H2}} e_{\text{lw_fragment_erg}}$. The heating_sum_uniqueness scanner's exception list is EMPTY: every deposited energy is named only in the module that defines it and in the one assembly. RED before (the pre-move import line, two deposits reported outside the assembly), GREEN after. `FUV_bands.txt` is byte-identical across the move (MEASURED, oxygen_chemistry at 5 steps); the `heat_LW` column moves by at most 5.57e-4 relative, which is the one-sweep lag between the state the file wrote and the state the energy equation deposited, and the new number is the deposited one.
- **B3c-COST, the cost of the coupled step measured and the skip specified, not implemented** (2026-09-06; `EXHALE_main.f90`, `attempted_step.f90`, the `coupled_source_step` and `attempted_step` suites). WHERE THE COST IS, MEASURED on `wasp_full` 300 steps, one thread: the run timers give `ioniz_eq` 83.6 percent and the energy solve 16.3 percent of the marching wall time, and a `gprof` build puts **78 percent of the whole marching time inside the MINPACK solve of one cell's chemistry** (one Jacobian and one QR per cell per pass; the warm start already reaches each root in one Newton step), against about 13 percent in the whole-grid rate and field work a pass repeats. So the cost is the number of CELL solves and not the field, and reusing the field between passes is worth at most that 13 percent. THE CONVERGENCE STUDY CONTRADICTS THE EXPECTATION THE ITEM WAS WRITTEN FROM: after pass 2, 308 of the 500 cells are still above the tolerances, after pass 5 still 130, and the cell that sets the pass count walks inward one cell per pass with the ionization front. Contraction 0.191 median over 2729 passes. A cell-local skip is therefore worth 2.41x on the cell solves with a one-pass rule -- but that rule is WRONG: a cell at rest moves again on 177 of the 3029 passes, 690 of the 809 wake-ups at pass 8, because the column keeps moving a cell after the cell has stopped. **Two consecutive passes of rest: zero wake-ups, 1.96x on the cell solves, 1.55x on the step.** The skip has to be a mask in the cell loop of `ioniz_eq`, which this item does not own (HEATBRK was writing to that file at 20:30 and 20:35), so it is SPECIFIED and not implemented: the optional `cell_at_rest` argument, the class-6 path (`sys_x = x_entry` straight to the extraction) as the body of the skip, and the one consequence that is not a pure skip -- the acceptance ledger describes the cells the sweep SOLVED, so a mask needs an acceptance

class stored for each cell and inherited by masked cells, or `n_cells_without_chemical_root` reads a different population from the one it believes. TOLERANCE ANCHORED AND KEPT AT $1e-8$: the anchor the comment gave (the $1e-6$ heat-column gate) stopped depending on it when HEATBRK made the two assemblies one; the anchor that binds is `cert_tol_carrier = cert_tol_element = 1e-8` on the composition this step returns. Measured $1e-10$ to $1e-5$ on `wasp_full 300` and `mol_base_handoff 1500`, the adopted state moves by about the tolerance and a decade costs 1.3 passes and 15 percent of the wall time, **so the tolerance is not the cost lever**; the step's own truncation error (step-doubling, phys, $5.3e-3$ relative) is five decades above the loosest value and does not bind. Newton on T with the composition's response (item 4) stays live, and the CHEAP version of it is refuted by measurement: the contraction ratio spreads by 122 percent globally and by 200 percent cell by cell over the tail of a step, so the sequence is geometric on average over a run and not within one step at one cell -- which is why B3c's Aitken failed, and why the acceleration has to come from $dc/dT = -(dF/dc)^{-1} (dF/dT)$ against the factorization hybrid already forms, not from the iterates. ALSO FIXED, a defect of my own file found outside the task: the run's cost line divided passes accumulated over every ENTRY of the coupled loop by the number of adopted STEPS, two different populations, so it could report a mean above the worst single entry (MEASURED on the control, `wasp_full 300` in phys mode: "13.98 passes per step on average, worst 11"). It now divides by the entries and names them. ALSO: `heat_channel_state` is in the attempted-step checkpoint (save/restore/compare/perturb), the four lines HEATBRK's report asked for, with two assertions RED on the control (the moved array is not seen, and the restore leaves it 3.0 off). Tests: `coupled_source_step 36/0` (33 before), `attempted_step 97/0` (95 before). Impact, control = the same tree snapshot with this item's two production files reverted: `wasp_full 300`, `mol_base_handoff 1500` and `mol_metals 1500` all **byte-identical** in `Hydro_ioniz.txt` and `Ion_species.txt`.

- **B3c-COST** (measurement; EXHALE_main.f90 comments and debug counters, `attempted_step.f90` checkpoint; numerically inert, `wasp_full 300`, `mol_base_handoff 1500`, `mol_metals 1500` byte-identical): the coupled source step's cost is not where the brief assumed. MEASURED on `wasp_full 300` (3029 passes): after pass 2 still 308 of 500 cells above tolerance, after pass 5 still 130; the pass count is set by the ionization front walking inward one cell per pass, not by a fixed handful of cells, so a cell-at-rest skip is worth 1.96x on the cell solves and 1.55-1.62x on the step, and the naive rule (rest on the last pass) is wrong: 177 of 3029 passes wake a resting cell, two consecutive passes of rest wake none. gprof: 78 percent of marching time is the MINPACK solve of one cell's chemistry (Jacobian + QR per cell per pass), the whole-grid field and rate work 13 percent, the energy solve 16 percent (already deactivates converged cells). Tolerance anchored and KEPT at $1e-8$ (the binding anchor is `cert_tol_carrier = cert_tol_element = 1e-8` on the returned composition): a decade costs 1.3 passes and 15 percent, not the lever. Aitken and any acceleration built from the iterates fail because the contraction ratio spreads 200 percent cell by cell within a step. **The lever is a Newton on (T, composition) taken cell by cell with $dc/dT = -(dF/dc)^{-1} dF/dT$ from the factorization hybrid already forms, inside the sweep (ionization_equilibrium.f90), plus the two-pass rest mask with an acceptance class inherited by masked cells**: assigned as COST2, after B3b-CO releases that file. Also fixed: the cost line divided passes over loop entries by adopted steps (could report a mean above the worst); `heat_channel_state` is in the checkpoint with two assertions (`attempted_step 97/0`, `coupled_source_step 36/0`; advisor rebuild and

rerun green). Outside scope recorded: EXHALE_ERR EVERY read in both modes but the probe runs only in phys; mol_base_handoff in phys mode refuses 7 of its first 20 steps at the fixed point (pass cap at step 0), recovers.

- B3b-CO, advisor verification and follow-up** (2026-09-06): private build of the live tree, physics_probe 584/0 (incl. co_destruction 27 and the scanner heating_sum_uniqueness with an EMPTY exception list after HYG-LEDGER-EPS moved the band ledger into utils_ion_eq::fuv_band_absorption_ledger), fuv_band_ledger 24/0, grid_and_gates 93/0. Accepted with the four stated departures from the design (cross section as the mean over the 912-1201 Å beam of one that ends at 1118 Å; the same flat-band cross section for a loaded spectrum, the shape-aware integral belonging to sed_read.f90; q_D1 as a table row because mion_fsp is a parameter; CO on its own ledger row, co_frac 6.39e-2 the size of decision 2's approximation). **P0 opened by the item itself and assigned at once as B3b-CO2:** the He+ + CO channel deposits its heat (65 percent of the total at the He+ front of oxygen_chemistry, $r = 1.089$) while no He+ is consumed and no C+ or O delivered anywhere; brief plan_rev2_brief_B3b-CO2.md (He+ sink in both molecular systems with Jacobian, products to the free O and carbon pools, heat rate identical to the species rate from one place, element census). Also recorded: the thermodynamic CO ceiling is still present because its deletion touches EXHALE_main.f90 34, 3641-3648, certification.f90 84, 1072-1076 (n_active_unvalidated_physics loses its only producer), write_setup_report.f90 18, 925-934 (CEILING-DEL, after B4-1 releases main); heat is deposited for carrier reactions whose products a never-covered carrier interval does not deliver (oxygen_chemistry in init mode never covers one), an operator-split defect that B4-1c and B5 address and that leaves the oxygen case without an end-to-end gate for its CO source; heat_channel_state now has 19 columns (checkpointed by B3c-COST); python/paper_data.py HEAT_FIXED stale.
- B3b-CO2 landed: the He+ + CO channel now consumes the helium ion** (2026-09-06). Row (2) of mol_heh_rows (System_HeH_mol.f90) loses k_D1 n_CO n(He+) with $k_{D1} = rk_{D1_Hep_CO}() = 1.60e-9 \text{ cm}^3 \text{ s}^{-1}$, and its turnover scale (set_mol_turnover_rates) gains k_D1 n_CO n_He; the helium leaves neutral, so the free neutral He that closes the helium budget receives it and no other row moves. One row serves every path: System_HeH_mol_metals and the constrained-equilibrium residual both call it, and neither molecular system has an analytic Jacobian to extend (hybrd1, finite differences), so the test asserts the difference derivative the solver actually builds. CO is a background density of that row and not an unknown, because the one-sided model gives it no formation and therefore no local equilibrium; the reaction is split across the two operators, each reading the other's frozen value from the one rate, which makes the heat of channel 18 the heat of exactly the events the species ledger performs. The products needed nothing added: the free-atomic-oxygen closure of carrier_source and nC_free - n_CO in carrier_write_back already carry the two nuclei and element_census already counts them in CO. Tests: new physics_probe driver co_helium_ion_sink (8 assertions, RED 4 FAIL against the pre-change module relinked into the same suite, GREEN 8 PASS; the RED reads hep_row_co_sink 0.0 against -3.84e6, hep_row_derivative_co_term 0.0 against -38.4, hep_turnover_scale_co_term 0.0 against 3.84e13); physics_probe suite PASSED after; element census with EXHALE_ELEMENT_ASSERT=1 on oxygen_chemistry 150 steps exits 0 with no budget violation (a guard, not a RED/GREEN pair: the change moves no C or O nucleus). Impact, MEASURED at 1000 steps

of oxygen_chemistry: the largest share of the total heating taken by channel 18 falls from 0.672 at $r = 1.081$ to 0.035 at $r = 1.117$, channel 18 at $r = 1.0894$ from $6.90\text{e-}7$ to $9.75\text{e-}9 \text{ erg cm}^{-3} \text{ s}^{-1}$, $n(\text{He}^+)$ there to 0.015 of its former value and to 0.0073 at $r = 1.011$, T at $r = 1.0233$ from 1156.4 K to 1088.2 K, and $\log_{10} \text{Mdot}$ stays 9.61 with exit 0. dom_cells_HeP does NOT fall (281325 -> 281382): the counter compares $k_{D1} n_{CO}$ with recombination and still says this reaction is the leading He^+ loss of the layer, so what it now reports is the operator split's lag rather than a missing sink. mol_carrier 300 steps and mol_metals 400 steps byte-identical in every output file. Goldens not refreshed. Reported for owners elsewhere: the dom_cells_HeP header text in write_output.f90 558-562 still says the frozen He^+ "over-states that channel"; the mirror decomposition hydrogen_helium_row_terms in constrained_chemical_equilibrium.f90 needs the same term after line 1859 or it stops summing to row 2; the n_co comment in ion_cell_state.f90 still calls CO an equilibrium closure.

- B3b-CO2 landed, advisor verification and follow-ups** (2026-09-06): the He^+ row of mol_heh_rows (System_HeH_mol.f90, the one row every caller uses) loses $k_{D1} n_{CO} n(\text{He}^+)$, its turnover scale gains the same bound; CO stays a background density of the solve because the one-sided model gives it no formation channel (not a balance row). Products needed no code: the O nucleus goes to the free-oxygen closure and the C nucleus to $n_{C_free} - n_{CO}$ in the carrier write-back, element_census already counts both; the carbon CHARGE is set by the sweep's stage equilibrium, the design of record ("the step moves nuclei, the sweep moves charge"), so the reaction's C^+ is not imposed (recorded as a design choice, not a defect). MEASURED, oxygen_chemistry 1000 steps: channel-18 share of the total heating 0.672 at $r = 1.081$ -> 0.035 at $r = 1.117$; $n(\text{He}^+)$ at the front 0.015 of before (0.0073 at $r = 1.011$); T at $r = 1.0233$ 1156 -> 1088 K; $\log_{10} \text{Mdot}$ 9.61 both; constrained-continuation fallbacks 932 -> 1880, all accepted; mol_carrier 300 and mol_metals 400 byte-identical. dom_cells_HeP did not fall (281325 -> 281382): it measures dominance of the channel over recombination, not over-statement; its header text now says so. Advisor follow-ups applied directly: the $D1 \text{ He}^+ + CO$ term added to the He^+ decomposition of constrained_chemical_equilibrium.f90 (hydrogen_helium_row_terms, whose header requires every mol_heh_rows reaction to appear), the FUV_bands.txt/Oxygen_chemistry.txt dom_cells_HeP header rewritten, the stale n_co comment of ion_cell_state.f90 replaced. Private build: physics_probe 592/0 (incl. co_helium_ion_sink 8), fuv_band_ledger 24/0. Remaining: the operator split is now a LAG (the sweep destroys He^+ at the entry CO while the CO row destroys CO at the previous sweep's He^+) and oxygen_chemistry never covers a carrier interval in init mode, so this row has no end-to-end gate: B4-1c/B5 territory, recorded.
- HEATBRK closed** (2026-09-06): the mol_metals pair finished at its 12000 steps, Hydro_ioniz.txt and Ion_species.txt byte-identical, gate $1.683\text{e-}16$ with 21 columns, _adv profiles moved by at most $8.45\text{e-}3$ (the deposits the post-process copy lacked). All three impact cases of the item therefore confirm: the sweep's heat is unchanged, only the two consumers of the assembly moved.
- B4-1** (species_face_flux.f90 new; Reconstruction.f90, PLM_rec.f90, RK_rhs.f90, EXHALE_main.f90, binary_element_diffusion.f90, Makefile, new suite src/tests/species_face_flux/): the advective half of the ELEMENT transport now rides on the hydrodynamic face mass flux, inside the Runge-Kutta stages, on the same faces, areas, volumes and time step as the mass row. The face composition is the reconstruction of the

mass fraction by the same limiter as the primitive variables, taken from the side the face mass flux selects, and the closing member of the set is one minus the others, so $\sum_s F_s(j) = F_{\text{rho}}(j)$ holds by construction; the operator asserts it at every face of every stage (largest residual MEASURED in the unit suite: $2.1\text{e-}16$ relative, against a tolerance of 64 ulp). The cell-velocity upwind advection of `binary_element_diffusion.f90` is gone from the marching path: the operator advects only when it is given an advecting mass flux, which the fixed-wind relaxation supplies and the marching path does not, and the trace-metal solve lost its advective term the same way. The carriers of `photochemical_transport_step` are NOT in this increment (that file was held by another item): they keep their present advective term and are B4-1c. Tests, all MEASURED with a private build: a uniform composition is preserved to 2 ulp on a compressing flow, an expanding flow and one whose face mass fluxes alternate in sign, for PLM and for WENO3; the mass fraction stays in $[0,1]$ on a step profile carried across the domain in 300 steps; the closing member of the set differs from an independently reconstructed $1 - Y$ by $1.1\text{e-}16$; the face reconstruction converges at order 1.97 (PLM) and 1.92 (WENO3); the helium nucleus total of the domain changes only by its boundary fluxes to $4.9\text{e-}17$, where the cell-velocity form it replaces leaves $1.6\text{e-}9$ on the same column and the same step (RED and GREEN in one run); the injected violation of the face identity stops the operator with the face named; the new step is refusable at its injection point and the run goes on. The element-diffusion suite `diffusion_tests` is 35 of 35 after T11 was rewritten onto the new operator (its premise, that a face-averaged upwind decouples a cell whose face velocities straddle zero, is what the face-flux form removes). Impact, MEASURED on scratch copies, `OMP_NUM_THREADS=1`, `mol_*/hp_*` at 200 steps and `mol_diffusion/lower_profile` at 1000 rather than the 12000 their maxsteps pins, `wasp_*` at 300: 14 cases byte-identical, every one of them without element transport (`wasp_full`, `wasp_he23off`, `wasp_full_newton`, `mol_base_handoff`, `mol_metals`, `mol_lyman_werner`, `mol_ir_bands`, `mol_sec_ion`, `mol_carrier`, `oxygen_chemistry`, `hp_zero_seed`, `hp_trace_seed`, `hp_front`, `hydrostatic_column`); `mol_diffusion` moves by at most $7.1\text{e-}5$ in `Ion_species.txt` and $3.6\text{e-}3$ in `Cooling_breakdown.txt`, `lower_profile` by $1.2\text{e-}3$ and $1.1\text{e-}2$. The transported quantity, $n_{\text{He}}/n_{\text{H}}$, moves by $1.9\text{e-}10$ at the base and $3.5\text{e-}6$ at the molecular front of `mol_diffusion`; the base-layer mass flux $\rho v r^2$ moves by at most $9.2\text{e-}7$ over $1.00 \leq r \leq 1.16$. Those are smaller than the design's estimate because at 1000 steps neither case has separated its elements far from the reservoir ratio, and a uniform composition is what both forms carry exactly; a converged pair is what would measure the difference at the separation those cases reach, and is named as open in `docs/b4_spatial_operator_design_20260906.md`. No golden was refreshed.

- **B4-1 landed (element rows), advisor verification** (2026-09-06): new module `species_advective_transport` (`species_face_flux.f90`): the face mass fraction by the primitives' own limiter (PLM_rec_scalar, Reconstruct_scalar, the MC limiter and the WENO3 geometry coefficients factored, not duplicated), upwind side = sign of F_{rho} (coincides with the HLLC contact selection wherever both are defined, ROE/LLF offer only the sign), the closing member one minus the others so $\sum_s F_s = F_{\text{rho}}$ by construction and asserted at every face and stage, advanced inside the three SSP-RK3 stages with the mass row's dt , dV and areas; `positivity_limited_fluxes` writes a replaced first-order flux back into `face_flux` so the species flux built on it is the repaired one. Transported set: helium and the hydrogen component (normalized), the trace metals riding on the same faces inside `m_1` at the reservoir abundance (trace limit stated). `binary_element_diffusion.f90`

keeps the diffusive half only (element_transport_residual keeps its stationary advective term, equal where $r^2 \rho v$ is constant). Carriers NOT migrated (diffusive_photochemistry.f90 was B3b-CO's): B4-1c. Suite species_face_flux 13 + 2 whole-binary rows GREEN on the advisor's private build: uniform composition preserved to 2 ulp (PLM, WENO3, three flows incl. a sign-alternating face flux), face identity 2.1e-16, bounds exact, face reconstruction order 1.97/1.92, helium nucleus total changes only by the boundary flux to 4.9e-17 where the replaced form gave 1.58e-9 (B1 AT-4d RED/GREEN in one run), identity assertion fires, element step refusable; diffusion_tests 35/35 (T11 rewritten: its premise was the defect removed); attempted_step, grid_and_gates 93/0, carrier_retry 97/0 green. Impact (worker, control = same snapshot with the stage calls removed): 14 cases without element transport BYTE-IDENTICAL (wasp_* at 300, mol_* at 200 steps); mol_diffusion and lower_profile at 1000 steps move by at most 3.6e-3 / 1.1e-2 (cooling breakdown), n_He/n_H by $\leq 6e-5$, base $\rho v r^2$ by $\leq 1e-6$: far below the design's estimate because at 1000 steps the elements have not separated from the reservoir ratio; the converged 12000-step pair is open in the design document. Outside scope recorded: ESWENO3 measures second order on a smooth profile (1.92 against PLM 1.97, face error 8x PLM), a property of the primitive reconstruction to look at (WENO3-ORDER); hp_* cases need output/Hydro_ioniz_IC.txt in place; element_flux_profile.txt has a trailing text column.

- **CEILING-DEL** (diffusive_photochemistry.f90, write_output.f90, write_setup_report.f90, EXHALE_main.f90, certification.f90; tests carrier_retry (+ scanner no_CO_ceiling_identifier_in_src), carrier_constraint_attribution rewritten on the carbon clamp): the thermodynamic CO ceiling (clamp to co_equilibrium_density, four counters, attempted twins, three accessors, Oxygen_chemistry.txt header line, three co_ceiling_* resolved keys, the end-of-run line, validity state 4.1's only producer) is deleted per design decision 10; the domain record of B3b-CO stands in every place (seven co_domain_* resolved keys under oxygen chemistry + carrier transport, the CO items under class 4.2 beside H3+ in the certification, n_active_unvalidated_physics = -1 "not produced", kept because the five validity states are a fixed set). MEASURED before deletion on oxygen_chemistry 1000 steps: co_ceiling_cells 0, applications 0, CO removed 0.0 (with the published destruction rates in the row and the He+ sink in the sweep the carried CO never exceeds its equilibrium), so every numerical column is byte-identical before/after; mol_carrier 300, mol_metals 400, wasp_full 300 byte-identical. The design's "of order 160 cells" predates the CO rates and is retired with the clamp. Advisor verification: private build, carrier_retry, certification 29/0, carrier_constraint_attribution 11/11, fuv_band_ledger green, no co_ceiling identifier left in src/. Recorded for the docs sweep (DOCS-CEIL): the checkpoint enumerations and validity category 12 in a0_run_mode_contract, a2_certification_contract, b1a_active_equation_inventory, b3a_attempted_step_controller_design, b1_target_system (1246-1554), development_plan_20260905_rev2/rev3/review3, physics_numerics_audit_20260905, and b3b_co_destruction_design sections 2.3/4/7 still describe the ceiling as present. The domain record is not checkpointed (a statement about the state, valid whether or not the step is accepted), unlike the ceiling record it replaces.

- **DOCS-CEIL** (documentation only, 2026-09-07): sixteen documents brought level with CEILING-DEL, the

current state READ from the code (destruction row $\text{src(ic_CO)} = -(\text{k_D1 } n(\text{He}^+) + \text{k_CO}) n_{\text{CO}}$, conservation-only `limit_to_element_budget`, `carrier_co_domain_record`, certification class 4.2 CO block, `n_active_unvalidated_physics` = -1, seven `co_domain_*` keys); no measurement deleted, the superseded ones ("of order 160 cells"; "still cuts CO above 3000 K") kept with the measurement that replaced them beside them. Files: a0/a2 contracts, b1a inventory (4.1 row retired, 4.2 row the domain record, open question 3 closed), b3a design (checkpoint enumerations), b1 target system (5.1 split into the 2026-09-06 READ and the current one, AT-5 superseded, decision 12 moot), b3b_co design (sections 2.3, 3(iv), 4, 6, 7, 7.4 as done), development plan rev2/rev3/ review3/execution, `physics_numerics_audit` (P6 resolved), `code_status`, d0 (C26 closed, question 5 answered), `co_destruction_rates_literature` (banner), `To_be_determined_recommend`, PLAN rev2. Left as dated records: the superseded plan revisions and external reviews, and the b1 draft. Recorded: line-number citations in the 2026-09-06 design documents have drifted by up to a thousand lines (a re-anchoring sweep on routine names is a separate item, DOCS-LINES); the stale checkpoint header comment in `diffusive_photochemistry.f90` goes to B4-1c.

- WENO3-ORDER, measurement and fix** (2026-09-07): the ESWENO3 reconstruction was SECOND order on every grid the code runs on. Probe `weno3_reconstruction_order` (`physics_probe`; exact volume averages by 8-point Gauss quadrature, face error against the point value, $N = 50$ to 800): production rate 1.997 (uniform in r), 1.988 (the `define_grid` Mixed production grid), 2.940 on a grid uniform in the volume coordinate, where the defect is invisible. Cause, isolated by variants: not the ESWENO nonlinear weights ($S = 1$ gives the same 1.997), not the floor $\text{eps} = \text{dr_j}^2$ (rate 3.000 at κ 0.1 to 10 once fixed), not the ideal ratios $D1, D2$ ($1/2 + O(h)$, harmless); TWO of the four stencil coefficients applied the NEIGHBOR's volume share where cell j 's own belongs: $C1(j) = c/(b+c)$ on the jump to $j+1$ at the right face and $C2(j-1) = a/(a+b)$ on the jump to $j-1$ at the left face, each $1/2 + O(h)$ where $1/2 - O(h)$ belongs, an $O(h)$ coefficient error on an $O(h)$ jump. Inherited character for character from ATES v2.0 (ATES/ATES-Code-main). Advisor check of the derivation: a linear interpolant of two cell averages in the volume coordinate, evaluated at a face $dV(j)/2$ from cell j 's center, weights the jump by $dV(j)/(dV(j) + dV(nb))$; with the corrected shares a profile quadratic in the volume coordinate is reproduced to 5.9e-16. **Fixed (advisor, two symbols in each of Reconstruct and Reconstruct_scalar, Reconstruction.f90):** $C1(j) \rightarrow C2(j)$ on dWp in WL, $C2(j-1) \rightarrow C1(j-1)$ on dWm in WR; the probe's production mapping and header updated, the two waiting assertions enabled: `weno3_production_third_order` rate 3.000, `_mixed` 2.991 (RED before: shortfalls 0.803, 0.812); `physics_probe` 603/0. **Every golden whose case reaches the WENO3 stage moves, and not by rounding** (face error down two decades at $N = 800$); PLM-only cases are untouched. Consequence for the method: until now the two-stage PLM $\rightarrow$ WENO3 recipe was a continuation between two second-order schemes with the second having the larger error constant (1.3x uniform, 3.8x on the production grid). Mdot values are to be quoted from Newton-finished runs after the gate. Attribution of the weight form and the floor to Yamaleev and Carpenter (2009, JCP 228, 3025 and 4248) stays unverified: both papers are paywalled here; requested from the user. Also recorded: stale `.ipynb_checkpoints` copies of `Apply_BC.f90` and `UW_conversions.f90` under `src/modules/` define the same module names as the live files (not built; to remove, HYG-CKPT).

- **HYG-CKPT** (advisor, 2026-09-07): ten untracked, gitignored Jupyter checkpoint copies of production modules under `src/modules/*/.ipynb_checkpoints/` (Apply_BC, Reconstruction, Source, base_boundary, caloric_eos, low_mach_dissipation, cross_sec, UW_conversions, h2_photo_channels, element_census) removed: not built, but each defined the same module name as a live file and two scanners had to skip that directory (batch 2c) to stay green.
- **T15-INSTRUMENT** (attempted_step.f90 +289/-165, suite, b3a design section 9, b1 T1.5 wording; advisor verification: private build, attempted_step 43/0 with the two whole-binary rows skipping until the main-loop hook lands, coupled_source_step, certification green): the B3a instrument now states the THERMAL identity of the step, $\sum_j dV_j [u_{th}^{n+1} - u_{th}^n] = \sum_j dV_j dt (\text{heat} - \text{cool})_j + (\text{transport contribution})$, the transport contribution MEASURED between two marks on either side of the coupled source step, the residual the closure of the source step alone, Delta u_{form} reported as the reservoir's change and never added (the B3c reason stated at the routine). GATE in phys mode at the energy row's own `energy_res_tol = 1e-9` (reason `as_reject_source_energy = 14` at operation 9): MEASURED on a hooked scratch tree, worst lresidual/scale 7.44e-1 -> 5.40e-10 (wasp_full 300 phys), 8.63e-1 -> 1.88e-10 (mol_base_handoff), zero refusals, states byte-identical in both modes. The two hook lines in EXHALE_main.f90 (before/after the coupled_source: loop) are assigned to COST2, the file's current owner, with the rewording of the print block. **Defect found and fixed in passing:** attempted_step_checkpoint_take saved `u` and `f_sp` through a saver that reallocates from index 1, so a reader indexing the saved grid arrays by cell compared cell `j` with cell `j - Ng`; `energy_identity` was that reader, so every `d_u_th/d_u_form` printed so far (incl. the 28 percent reservoir share of the B3c report: with bounds fixed, same step, `d_u_th` 2.006e-5, `d_u_form` 1.298e-5) was formed against a state shifted by two cells; the state restore itself was unaffected (whole-array assignment). The second private copy of the `eps_s` table in attempted_step.f90 (no OH/H2O/CO rows) is deleted in favor of `species_formation_energy`. Recorded: `n_projection_applied` can no longer be nonzero since B3c but is still printed by the report and refused by the certification (HYG-PROJ, spans `global_parameters`, `certification.f90`, main); design section 9 decision 8 still names the projection as a live correction (dated record).
- **COST2** (measurement; the only production change is the T15 hook and print in EXHALE_main.f90, byte-identical on wasp_full 300 in both modes and mol_base_handoff 60 phys; attempted_step whole-binary rows now run, identity residual -1.8e-10, zero refusals): the brief's premise is refuted twice. (1) MEASURED with T frozen through the coupled loop on wasp_full: the composition sweep alone still takes 10-13 passes with the same contraction 0.15-0.20 and the same front walking one cell per pass; freezing the lagged coefficients as well collapses it to 2 passes; freezing only the radiation field leaves 3-4, only the rate coefficients 9-13. The pass count is the self-consistency of the photoionization field with the composition that attenuates it; the temperature is worth at most one pass. (2) A timer inside `ioniz_eq` (60 steps, 29.95 s): PH_heat 11.40 s (38 percent of the run), the whole cell loop 7.37 s (25 percent), pre-sweep `eval_cool` 2.44 s, post-sweep assembly 2.40 s; skipping the entire cell body for resting cells saves 4.7 percent. B3c-COST's `gprof` attribution (78 percent in MINPACK) was inverted; the `csn_max_pass` comment still carries it (to be rewritten by COST3). (3) The two-pass rest mask

was implemented, measured (6.0 percent on `wasp_full`, 0 on `mol_base_handoff`, state moved $5.5e-7$, step 0 of `mol_base_handoff` in `phys` turned into an exhausted attempt) and removed. **The lever: the field dependence is one-directional (a cell's `P_HI` is set by the attenuation of the cells outside it) and the sweep already runs outside in, so updating the column with the composition just produced turns the Jacobi iteration into a forward substitution: same fixed point, about one pass instead of ten, and `PH_heat` no longer repeated per pass.** Assigned as `COST3` (`util_ion_eq.f90` `PH_heat_HHe/PH_heat_H`, `ionization_equilibrium.f90` sweep order; the OpenMP parallel cell loop becomes sequential in the field direction, so the design must weigh 10 parallel passes against 1-2 sequential ones and measure both at 1 and 16 threads). Also recorded: the pre-sweep `eval_cool` is run for its coefficients and its cooling discarded, then a second traversal forms the cooling (16 percent together); `mol_base_handoff` in `phys` refuses 7 of its first 60 steps at the fixed point on the unmodified tree.

- **B4-1c landed: the carriers ride on the species face flux** (2026-09-07): the carriers of `photochemical_transport_step` (`H2` always, `OH/H2O/CO` under Oxygen chemistry, `H+` under Ionization transport) join the transported set of B4-1 and are advected as `F_rho Y_c/m_c` inside the Runge-Kutta stages. `carrier_set_init` declares them once the keys are parsed (`advected_carrier_register` in `binary_element_diffusion.f90`, which also holds the columns and hands them back through `advected_carrier_fractions`); `EXHALE_main.f90` is untouched, the existing three calls carry both sets. The carriers ride OUTSIDE the normalized set (`n_norm = 0`): a carrier is one species inside an element, the element is already the closing member, and normalizing it beside the element would count that mass twice; what keeps carrier and element consistent is the operator's own write-back, which restores the element totals cell by cell. Their rows are advanced from cell 1 (the elements' start at 2), with the inflow composition taken from the inner ghosts: the handoff's `q_H2_base` where one is stated, the base cell's own partition where none is, one rule for both branches of decision D6 (B4-4a). `diffusive_photochemistry.f90` loses the cell-velocity advective term, `carrier_advection_correction` and the base Dirichlet branch from its `MARCHING` rows and their Jacobian; two paths keep the advective term, `carrier_steady_residual` (the stationary balance is the whole equation) and the fixed-wind relaxation `relax_photochemical_composition` (it takes many transport steps with no hydrodynamic stage between them, so nothing else is taking that transport), through the switch `carrier_rows_advect`. Tests: two rows added to `species_face_flux`, `uniform_carrier_partition_preserved` 0.83 ulp (bound 8) on a sign-alternating face mass flux and `h2_nucleus_total_changes_only_by_boundary_flux` $1.36e-15$ (bound $1e-12$), RED before in the strongest sense (the driver does not compile against the pre-increment module); suite 15 + 2 GREEN; `carrier_retry` 101, `carrier_returned_state_acceptance` 37, `carrier_constraint_attribution` 11, `carrier_reference_scales` 15, `acceptance_classes` 22 GREEN with no reference re-derived. Impact MEASURED against a binary built from the same tree snapshot with the two files at their pre-increment text: 12 cases BYTE-IDENTICAL, every one without Molecular carrier transport (`wasp_full`, `wasp_he23off`, `wasp_full_newton`, `mol_base_handoff`, `mol_metals`, `mol_lyman_werner`, `mol_ir_bands`, `mol_sec_ion`, `mol_diffusion`, `lower_profile`, `oxygen_chemistry`, `hydrostatic_column`); the four that do transport carriers move. `mol_carrier` at 1000 steps: Ion_species $5.0e-1$ (`H3+` at `r = 1.100`), `Cooling_breakdown`

6.2e-1, Hydro_ioniz 2.6e-2 (v at $r = 1.186$); hp_zero_seed/hp_trace_seed/hp_front at 100 steps: 8e-3 to 1.1e-1, Hydro_ioniz 1.3e-3 to 1.8e-3. **The front does not move; its tail does.** The $x_{H2} = 0.5$ crossing of mol_carrier is $r = 1.0698$ at 1000 steps before and after and $x_{H2}(1.05)$ is 0.7308 against 0.7310, while $x_{H2}(1.10)$ falls from 3.19e-3 to 2.02e-3 (-37%): the molecular tail the cell-velocity form was carrying by its own numerical diffusivity, which is where the design said the difference lives, and every large entry above is a consequence of that column. Every carrier interval covered and the element limiter moved no cell, in both runs. Recorded and not fixed: oxygen_chemistry has carrier_transport on by default with the oxygen chemistry, but at 200 steps in initialization mode 201 of 201 intervals are uncovered before AND after, so the write-back never runs and the case is byte-identical for that reason, not for an absence of effect (its rows did change: worst CO row of step 1 |res| 6.03e-5 against 5.68e-5). No golden was refreshed. Also in this increment: the stale clause saying the carrier checkpoint restores "the cumulative CO record" removed from two comments in diffusive_photochemistry.f90 (the carrier_checkpoint type carries no such field since CEILING-DEL).

- **B4-1c landed, advisor verification** (2026-09-07; the worker's own entry carries the numbers): every carrier of the photochemical transport operator (H2; OH/H2O/CO under oxygen chemistry; H+ under ionization transport) rides on $F_{rho} \ Y_c / m_c$ inside the RK stages through the registry of binary_element_diffusion.f90 (carriers outside the normalized set, from cell 1, base inflow composition = the inner ghosts'), species_face_flux.f90 unchanged (species-agnostic), EXHALE_main.f90 untouched; the marching rows of diffusive_photochemistry.f90 keep diffusion, drift and chemistry only; the advective term stays in carrier_steady_residual (the whole stationary equation) and in the fixed-wind relaxation relax_photochemical_composition (no hydro stage takes it there), switch carrier_rows_advect. Advisor: private build, species_face_flux 15 + 2 rows (carrier partition 0.83 ulp, H2 nucleus budget 1.4e-15), carrier_retry, carrier_returned_state_acceptance, acceptance_classes, grid_and_gates green. Impact (worker, same-snapshot control): 12 cases without carrier transport BYTE-IDENTICAL; mol_carrier 1000 steps: the H2 front does not move ($x_{H2} = 0.5$ at $r = 1.0698$ both), the tail the cell-velocity form carried by its numerical diffusivity does ($x_{H2}(1.10)$ 3.19e-3 $\rightarrow$ 2.02e-3, -37 percent), hence H3+ 0.50, cooling 0.62, heating 0.37 relative there; hp_* 100 steps $\leq 1.7e-2$; base rho v $r^2 \leq 9.7e-4$. Recorded: element_advection_* names are now too narrow (HYG-NAME, six call sites in main, after COST3); oxygen_chemistry in init mode covers 0 of 201 carrier intervals so it cannot judge carrier transport; a refused carrier interval discards a transport the mass row already took (B3a/B5 question); adv_corr still checkpointed though only the residual writes it; 52 documents under docs/ carry em-dashes (DOCS-DASH, launched).

- **DOCS-DASH** (documentation only, 2026-09-07): 1354 em-dashes and 237 en-dashes removed from 60 files (docs/*.md, READMEs, test and data READMEs), each replaced by the punctuation the sentence needs (colon, comma, parentheses, or a table placeholder hyphen), en-dash ranges and joined names to the hyphen the repository already writes; 108 mechanical substitutions corrected by hand on read-back; grep over the 187-file set clean. Left: Update_EXHALE_stage1.md (append-only log, 319/59), and 1220 LaTeX --- in 27 .tex sources that render as the same dash (DOCS-DASH2, launched: the .md/.tex twins realign there; the built PDFs are re-generated at the gate's docs pass), plus docs/lower_atmosphere_figs/README.md (11) and three READMEs

under backup/ (5) outside the first sweep's paths.

- **DOCS-DASH2** (documentation only, 2026-09-07): 1283 LaTeX --- in 29 .tex sources (docs, paper, posters) and 30 spaced -- replaced by the punctuation each sentence needs; four exceptions kept and named (a direct quotation of Frelikh and Murray-Clay 2026 compared letter for letter, the AASTeX \keywords separator, a literal metals.inp listing, hyphen rules); the four READMEs of the first sweep's remainder cleared. The 24 built docs/*.pdf, literature_survey.pdf, paper/ms.pdf and the poster PDFs are now dated and are regenerated in the gate's docs pass. Left: the append-only stage logs (Update_EXHALE_stage1.tex 1614, stage0.tex 37).
- **COST3** (measurement; util_ion_eq.f90, ionization_equilibrium.f90; default output BYTE-IDENTICAL on wasp_full 300, mol_base_handoff 1500, mol_metals 1500, hydrostatic_column 300 at one thread; advisor rebuild: coupled_source_step 36, physics_probe 862/0 incl. the new photoionization_field_substitution driver (259 assertions), grid_and_gates 93/0): the star-ward columns can now be carried forward cell block by cell block (advance_starward_columns, reproducing calc_column_dens* bitwise; photoionization_field_at_cell*, the PH_heat* cell bodies extracted verbatim; xuv_field_block_cells, default 0 = whole grid). MEASURED on wasp_full 300: the substitution stops the front's walk (worst-moving cell 206 -> 177 over the passes on the lagged scheme, 202 -> 196 with the columns carried) and cuts the pass count 10.10 -> 7.13, but a pass costs the same field work either way, and the field of a whole grid is one loop sixteen threads solve at once: block width 8 buys 1.43x at one thread (136 -> 103 s) and LOSES 2.8x at sixteen (17.0 -> 42.5 s); width 64 keeps the parallel front and saves nothing. The default therefore stays the whole grid, and the width is not tied to the thread count (the widths reach states differing by 1.5e-7 at the 1e-8 tolerance, the thread count by 7e-14). **The remaining lag is the cell's OWN optical depth**, which enters its own field only on the next pass (contraction 0.17): closing it means the cell residual carries its own field, System_HeH*, i.e. B5. Folded: the pre-sweep eval_cool run for its coefficients only is now chemical_rate_coefficients (-3.6 percent wasp_full, -5.6 percent mol_metals, +3.5 percent mol_base_handoff, the last from the 27 metal slots looped with metals off: guard on thereis_metals, HYG-COL). Recorded: docs/openmp_parallelization.md (2.07x at 16 threads, bit-identical across threads) is stale: measured 8.2x and round-off 6.6e-14 (DOCS-OMP). A cp -irp of another item blocked on an interactive prompt since 2026-09-06 was killed by PID (23818, cwd checked).
- **WENO3-REF** (comments only, Reconstruction.f90 and the probe's References block; the user supplied the publisher PDFs, read with pdftotext -layout): the nonlinear weights are EXACTLY Yamaleev and Carpenter 2009, JCP 228, 3025, Eqs. (18), (20), (21), (22) (published tau already squared; $d_0 = 2/3$, $d_1 = 1/3$ entering as the ratio 1/2 that D1, D2 carry); the scheme is NOT their ESWENO: the energy-stability dissipation of Eq. (48) is absent, their analysis is a uniform-grid finite-difference flux reconstruction, and for systems only in characteristic variables (Section 5 quoted), so the code inherits the accuracy result (Eq. 24) and not the energy estimate; the geometry (C1, C2, D1, D2 on the stretched finite-volume grid) is this code's own. $\epsilon = dr_j^2$: a defensible stand-in, not a defect: the published h^2 scaling of Eqs. (62), (64) without the solution scale of Eq. (65); MEASURED on the production grid the face error is flat to every digit for $\kappa \geq 0.1$ over four decades (rate 2.991) and deteriorates below 1e-2 per Eq. (57), so in smooth regions the departure is not measurable;

where it acts is the ENO biasing (b_r/ϵ spans $2.2e-6$ to $8.4e3$ over the `wasp_full` grid; the published floor would make the base LESS nonlinear), a candidate experiment for the base sound-wave work, not a change here. Case label and comments now say "third-order WENO"; identifiers unchanged. Advisor: `paper/ms.tex` 144 "the energy-stable ESWENO3" reworded to the verified attribution and the ESWENO3 name replaced by WENO3 there and in `docs/code_comparison.tex`. **Recorded for the user:** every `src/tests/*/` suite of this series (19 directories) and the 2026-09-05/06 design and plan documents are untracked in git; a fresh clone of the remote carries none of them.

- **References received** (2026-09-07, from the user, publisher PDFs in `references/`): Zhang and Shu 2010 (JCP 229, 8918), Poinso and Lele 1992 (JCP 101, 104), Thompson 1987 (JCP 68, 1) and 1990 (JCP 89, 439), Neufeld 1990 (ApJ 350, 216), Mao and Kaastra 2016 (A&A 587, A84), Verner et al. 1996 (ApJ 465, 487), Yelle 2004 (Icarus 170, 167), Larsson et al. 2008 (CPL 462, 145), Tully 1975 (JCP 62, 1893), Slinger and Black 1982 (JCP 77, 2432). Five verification items launched against them: REF-ZHANGSHU (the positivity limiter), REF-LODI (the characteristic boundary design document), REF-NEUFELD (the Ly-alpha escape-probability closure), REF-METALS (every metal photoionization and recombination fit parameter against the published tables, a transcription error being a physics fix), REF-CHEM (the H3+ network transcriptions, the H3+ dissociative recombination coefficient, O6 and the Ly-alpha H2O yields). The series gate waits for them, since REF-METALS and REF-CHEM may move coefficients.
- **Reference data fetched** (2026-09-07, at the user's direction, each with a provenance README, third-party material unchanged): `references/verner_photo/` (D. A. Verner's distribution: `phfit2.f` reference routine with its fit parameters, `table1.dat` of Verner et al. 1996, the 1995 inner-shell fits), `references/mao2016_cds/` (CDS J/A+A/587/A84 `rr_spex3.dat`, 39696 rows), `references/badnell_rr/` and `badnell_dr/` (Strathclyde TAMOC total RR and DR fit coefficients `clist_K/clist_eV`, tabulated `cfout_*`, reference routines). REF-METALS now compares every metal stage's photoionization, RR and DR coefficients against these by direct evaluation of the reference routines, and reports which source each stage uses today and any missing DR contribution.
- **HYG-MAIN** (2026-09-07; numerically inert, `wasp_full` 300, `mol_carrier` 200, `mol_diffusion` 200, `mol_base_handoff` 300 byte-identical; advisor rebuild: certification, `attempted_step`, `species_face_flux` green): `HYG-NAME` `element_advection_*` -> `species_advection_*` in `binary_element_diffusion.f90`, the main program (use list, six calls) and the two test drivers, the two design documents that still named them (advisor); `HYG-PROJ` `n_projection_applied` deleted from `global_parameters`, the `attempted_step_report` block and both `certification.f90` references (MEASURED: every reference in `src/` was a read, no assignment anywhere, so the counter was provably zero since B3c), `rep%n_unbudgeted_accepted_correction` is now `n_shapiro_applied` alone, b3a design decision 8 dated closed for row 8; `HYG-COL` the 27-slot metal column loop of `advance_starward_columns` guarded on `thereis_metals` (inert; COST3's +3.5 percent NOT reproduced: guarded 53.7/52.8 s against unguarded 55.7/52.5 s on `mol_base_handoff` 300, within the run-to-run spread, kept as dead-work removal); `DOCS-OMP` `docs/openmp_parallelization.md` dated and brought to COST3's measured state (8.2x at

16 threads, 6.6e-14 between thread counts). Recorded: no reduction on a physical quantity exists in `src/`, so the 6.6e-14 thread-count difference comes from block boundaries set by `omp_get_max_threads()` in `chemical_rate_coefficients/eval_cool`; which routine turns a moved boundary into a moved last bit is not localized (THREAD-DET: until it is, the thread count is a silent input to every result at round-off); `photoionization_field_at_cell_HHe` makes an array temporary for `nm_j` on every call (runtime warning in `physics_probe`); `EXHALE_main.f90:452` is 141 characters.

- **REF-ZHANGSHU** (comments, `docs/p49_front_energy_mode.md` 6.1, `probe` positivity_limiter_scaling 15 assertions; user-supplied PDF read): the citation was wrong as stated: positivity_limited_faces is not the limiter of Zhang and Shu 2010 (JCP 229, 8918) but one built on the same idea, departing on three axes (primitive, not conserved, variables; one theta per face state, not per cell, Eqs. (2.4)-(2.5), (2.10)-(2.12); a relative one-ulp floor, not their absolute 1e-13). What carries over: Lemma 2.5 (convex combination with an admissible anchor); what does not: Theorem 2.1 (the next cell average admissible under their Gauss-Lobatto set and CFL, $N = 2$ covers $k = 1$ only; Remark 2.6 names finite-volume WENO as open) and conservativity of the reconstruction (a theta taken face by face moves the cell's end-point mean by 8.3e-2 where a theta taken cell by cell gives 0). Along their conserved path $p(t)$ is concave, so the code's theta is never larger than their `t_eps` (MEASURED 0.856 against 0.949 at Mach 60): the anticipated negative-internal-energy failure cannot occur. **Defect found and fixed (advisor applied the worker's proposal):** with `eps` one ulp of the average the bare update `q_avg + theta (q_rec - q_avg)` is a cancellation of the size of the floor and lands on EXACTLY ZERO for 1013 of 200000 reconstructed values crossing zero (never negative), the value the floor exists to prevent (division in $v = m/\rho$ and in the sound speed); `positivity_limited_faces` now clamps the scaled density and pressure to `eps` after the update (same number in exact arithmetic). Moves only runs in which the limiter fired, by one ulp except where the zero would have divided; goldens at the gate. Probe: `bare_update_reaches_zero` (the reason), `production_limiter_never_delivers_zero` (the production routine over 4000 grids of zero-crossing face states) GREEN. A theta taken cell by cell (Eq. 2.11), which would restore the reconstruction average, is recorded as a change of scheme, not applied.
- **REF-LODI** (documentation, `docs/phaseC_characteristic_base_bc.md` +282/-50; the three user-supplied journal PDFs read as page images since their text layers lose the displayed equations): the wave-amplitude normalization (Poinot and Lele Eqs. (15)-(23); Thompson 1990 (44)-(47) with $l_1 = (0, 1, -\rho c, 0, 0)$; Thompson 1987 (19)), the LODI relations ((24)-(36), equal to Thompson 1990 (48) term for term), the subsonic inflow two-condition closure (P&L section 3.1, Tables III/IV; Thompson 1990 Eq. (68)) and all four branch-table rows CONFIRMED; no constant wrong, the constraint count unchanged. Two statements of the note corrected: pair (D) is NOT singular at stagnation (Thompson 1990 Eqs. (69)-(70) have denominator $u_1 + c$; the note's singularity was its own closure $\rho_b = F_c/(r_b^2 v_b)$; D1 reopened with a pointer), and D6's claim that no transparent condition is expressible as a fixed point is too strong (Thompson 1990 Eq. (64) $L_5 = \rho c g_1$ and P&L Eq. (40) $L_1 = K(p - p_{\text{inf}})$ are algebraic in the face state; D6 stays the user's decision with a third option). One gap stated: the source projection $l_1 C$ (gravity and the spherical divergence, Thompson 1990 footnote 1 and Eq. (52); the steady form is $L_1 = -\rho c g_1$, not 0) survives only because both states of (C-) sit at the same

radius; any reformulation relating cell 1's center to the face must restore it. Carlson's "but not U and p" marked unverified against the primaries.

- REF-CHEM** (comments and documents only; `mol_rates.f90` and `oxygen_rates.f90` function bodies byte-identical; probes `h3p_recombination_branching` 12 and `water_photolysis_lyman_alpha_yields` 6, RED with the competing published values): every `mol_rates` constant sourced to Yelle 2004 matches his Table 1 digit for digit; Yelle's R7 ($\text{H}3+ + \text{H}$) reference column reads "Estimated", so Frelikh's barrier-free $2.0\text{e-}9$ transcribes an estimate faithfully; the code's $\text{H}3+$ dissociative recombination R6/R7 ($2.16\text{e-}8 / 5.04\text{e-}8 \times (300/\text{T})^{0.65}$) is already Larsson et al. 2008's 300 K total $7.2\text{e-}8$ split by their branching 0.70, with the 0.65 index inherited from the Sundstrom/Datz fits Yelle prints (Larsson gives no temperature dependence; he states the CRYRING/ASTRID values the older $2.9\text{e-}8$ rests on were too high from rotational excitation), so the code stays; the Slanger and Black 1982 Ly-alpha H_2O yields 0.78/0.10/0.12 are exact (p. 2435) with one label corrected, $\text{Phi}2 + \text{Phi}4 \rightarrow \text{Phi}1 + \text{Phi}2$, and the paper's coupling of the three yields and the 8 percent OH(A) bound recorded; Tully 1975's $2.87\text{e-}10$ is a 300 K statistical-model value (Table II) that VULCAN flattens over 100-2100 K, and its agreeing "experiment" is a 1973 relative-rate evaluation, strengthening the audit's verdict for O6. Documentation fixed in passing: `methodology_comparison.md` 6.4 listed three-body H_2 formation as identical in value while the code's Cohen and Westberg 1983 $2.8\text{e-}31 \text{ T}^{-0.6}$ differs from Ham et al. 1970 by 1.142 at every temperature; Baulch 1992 is in references/ (its $\text{H}_2 + \text{H}_2$ entry valid 2500-8000 K, above the whole layer); section 10 item on H_2 dissociative photoionization closed (the code carries it, `h2_photo_channels.f90` channel S).
- REF-NEUFELD** (comments and documents; `lya_rt.f90` machine code identical before/after; `physics_probe` 1118/0): the Ly-alpha escape-probability closure does NOT use Neufeld 1990 or Harrington 1973. $\text{beta_esc} = \pi^{(-1/4)} \sqrt{a/\tau}$ is the one-flight wing escape chance under complete redistribution; Ly-alpha scatters with partial redistribution and the published static-slab damping-wing solution is Harrington 1973 eq. (40), $\langle N \rangle = 0.9093161 \text{ B scatterings}$ for a mid-plane source, so $\text{beta} = 1/\langle N \rangle$, proportional to $1/\tau$ and independent of a ; MEASURED on the stored `wasp_full` column the code exceeds it by 234x at the base and 14x at 1.20 Rp , crossing at 1.42 Rp . Neufeld contains no expression of the code's form (his escaping fractions are escape against destruction, eq. 2.25 partitions the two faces; the code's two-face PRODUCT is unpublished); the escape frequency scales as $(a \tau)^{(1/3)}$ (Harrington eqs. 37, 47; Neufeld eq. 4.32), not the code's $(a \tau)^{(1/2)}$; Neufeld's validity $(a \tau)^{(1/3)} > 10$ holds in 320 of 504 cells (inside 1.156 Rp); Neufeld is static, the Sobolev factor is Sobolev 1960 radial-ray. The header's `J_int` formula lacked the $(1 - \text{beta})$ factor the code applies (fixed). **Advisor decision: adopt the published solution (LYA-BETA, launched):** $\text{beta} = 1/$ of eq. (40) joined to the thin limit, the face partition of eq. 2.25 in place of the product, the $(a \tau)^{(1/3)}$ escape frequency, a validity record, `A_2p1s` $6.3\text{e}8 \rightarrow 6.2648\text{e}8$ (spectroscopic), every case with `Jlya escape-prob`: True moves (expected at frozen state: `J_int/n_2p` up to 90x higher inside 1.02 Rp , 14x at 1.20 Rp , up to 7x lower beyond 1.47 Rp ; `Mdot` to be measured to the du stop). The same question for the metal lines' `line_escape_probability_one_face` (`Cool_coeff.f90`) is LYA-BETA-METALS, after REF-METALS.
- REF-METALS** (`cross_sec.f90`, `Cool_coeff.f90`; probes `metal_photoionization_fits`, `recombination_coefficient_fits`, 255 assertions; advisor rebuild green apart from LYA-BETA's in-

progress probe): all 132 fit parameters of the seventeen metal photoionization cross sections match Verner et al. 1996 Table 1 (verner_photo/photo.dat) digit for digit and metal_photoion_sigma reproduces Verner's own phfit2 exactly wherever the fit is the one phfit2 uses; 220 Badnell RR and DR parameters (fourteen daughters; cfout_K witness within 0.045 dex), the H/He rows and the Mao and Kaastra $n = 1$ rows match. **Two fixes:** the Mg II turn-on gated at Verner's rounded 15.04 eV while the grid edge is the NIST 15.035 (e_th_MgII now; byte-identical on wasp_full, the band geometric means miss the sliver); alpha_rr_metal('CaI') used Verner rrfit.f's rrec(:,20,19), the row that FORMS Ca II (second index = the recombined ion's electron count, fixed by rnew(:,1,1) = H+ + e and rnew(:,2,2) = He+ + e), 6.05x the Ca I row (1.120e-13, 0.900) at 1e4 K; corrected. Impact (worker, scratch): wasp_full 300 Ca I 0.87 relative at its 1.196 Rp maximum, Ca II +2.8 percent at the base, T 4.3e-3, v 9.9e-3; mol_metals 400 Ca II/III x2.5 at 1.38 Rp, cool 2.9e-2; wasp_full, mol_metals, lower_profile move at the gate through calcium. **Two physics items opened:** (a) the outer-shell fit is evaluated on the whole photon grid to 1240 eV while every row carries an E_max (34-558 eV) above which phfit2 hands over to the 1995 subshell fit and adds the inner shells: the code carries 0.1-12 percent of the true metal photoabsorption there, and Fe I's row ($Q = 1.54$) RISES with energy to 49x the truth at 1240 eV; the handover of phfit2 itself is adopted (METALS-OUTER, launched); (b) the inner-shell absorption and its Auger cascade (an atom two or more stages up) cannot be represented by the three-stage metal ladder: To_be_determined_by_user_20260906.md item 7. Also: verner_photo/README.md corrected (photo.dat is the 1996 Table 1, table1.dat the 1995 fits); rev3 line 944 closed; e_th_MgI/e_th_MgII provenance lines in parameters.f90 to be annotated (METALS-OUTER, after LYA-BETA releases the file).

- **METALS-OUTER** (cross_sec.f90; generator src/utls/metal_photoion_table.py from verner_photo/photo.dat + table1.dat; probes metal_photoionization_fits 188 and metal_photoion_table_transcription 15; docs/photoion_cross_sections.tex + PDF; advisor rebuild: physics_probe green): the seventeen sigma_XX routines are one parameter block, and metal_photoion_sigma makes phfit2's handover: the 1996 outer-shell fit below E_max, the 1995 fit of the same shell at and above it (MEASURED: einn of phfit2 equals photo.dat's E_max for all sixteen metal rows; the production routine reproduces phfit2's outer shell to 1e-10 at 17 ions x 12 energies where 110 of 204 rows disagreed before, worst Fe I at 1240 eV 24.9 Mb against 0.00172 Mb, a factor 14464). Inner shells still absent by decision item 7 (option (a) recommended; the measured sizes now in that item). Impact (scratch, metals-off mol_base_handoff byte-identical): wasp_full 300 v 0.15, T 4.7e-2, heat 7.8e-2, Fe III 0.99, C III 0.60 relative (hard photons now reach the base, the top stages fall at small radius; metal share of the heating at the base 17.4 -> 8.5 percent); mol_metals 400 v 0.56, heat 0.42; lower_profile 200 heat 8.3e-3; log10 Mdot unchanged to 0.01 dex in the bounded snapshots. Both Mg thresholds already agree with NIST to their digits (item 4 needed no move; provenance note pending parameters.f90).
- **LYA-BETA-METALS** (measurement by the worker, fix applied by the advisor; Cool_coeff.f90, probe fine_structure_escape_probability, docs/resonance_line_trapping.md sections 6.1 and 10; publisher scans of de Jong, Boland and Dalgarno 1980 and Osterbrock 1962 fetched into references/): line_escape_probability_one_face is de Jong et al. 1980 eq. (B-7), both branches to round-off, applied

to the eight ground-term fine-structure lines only; NO metal resonance line uses an escape probability (Mg II h&k, Ca II H&K, Na I D, Fe I/II at $\beta = 1$, justified in that memo's sections 1-5 by $n_{\text{crit,eff}} = \beta A/q \sim 1e10-1e15$ against $n_e \leq 4.3e9$), and the transit post-process uses none. The fine-structure lines have a $\sim 1e-8$ and a $\tau \leq 1e-9$, so the Ly-alpha correction does not transfer. **Defect:** (B-7) is written in the frequency-integrated depth $\sqrt{\pi} \tau_0$ (normalized profile; equal to Hollenbach and McKee 1979 eq. (5.10) once the convention is matched, 3.4 percent at worst over $\tau = 3$ to $1e6$) while the caller passed the line-center column: β too large by 1 (thin) to 2.11 (branch point) to 1.77 (asymptotic). **Fixed:** each cell's depth is now $\sqrt{\pi}$ times the line-center depth in `fine_structure_line_transfer`. MEASURED regime on the shipped columns: largest line-center depth 0.14 ([O I] 63 μm at the hot-Uranus base), so the factor there is 1.12 and the change of the total radiative losses is below 0.1 percent (`mol_metals` 0.096 percent), but a factor of two in any base thick in these lines and in the Base IR field incident term the same β scales. `physics_probe` 1291/0 after the fix. Metals cases move at the gate at that level.

- **Decision item 7, literature check** (2026-09-07; the user supplied Locci 2022 PSJ 3, 1; Locci 2024 PSJ 5, 58; Cecchi-Pestellini 2006 A&A 458, L13 and 2009 A&A 496, 863; Fox 2008 SSRv 139, 3; all in references/): the published exoplanet X-ray photochemistry models take the inner-shell absorption as opacity summed over all ionization channels (Cecchi-Pestellini 2009 Eq. 1 with Verner 1993 cross sections), put the photoelectron and the Auger electrons into the secondary-electron cascade (Locci 2022 Eq. 3), and carry singly charged ions only, i.e. option (b) of the item with the charge under-count accepted; no escape hydrodynamics code includes inner shells. Recommendation revised from (a) to (b) in `To_be_determined_by_user_20260906.md`; the user decides.
- **METALS-INNER** (decision 7 option (b), user 2026-09-07; `cross_sec.f90` subshell block from `phfit2.f`'s PH1 data (its thresholds differ from `table1.dat` in six rows, `phfit2` being the reference), `set_energy_vectors.f90` (every subshell turn-on of an active metal ion a bin edge; bin count unchanged, 17 -> 60 threshold cuts on `wasp_full`), `util_ion_eq.f90` comment only (the existing rate σ/e_v and heat `photoelectron_share(E_th)` σ integrals already implement "one stage, one electron of $h\nu - E_{\text{th}}(\text{outer})$ to the SvS85/Dalgarno cascade" once the cross section is the total); generator `metal_photoion_table.py` --check; probes 392 metal assertions, `physics_probe` 1479/0; docs `photoion_cross_sections.tex` (+PDF), `code_status` new L row): the metal photoabsorption is the SHELL SUM `phfit2` returns, reproduced to $1e-10$ at 17 ions x 12 energies (130 of 204 rows disagreed before). It also closed a cliff METALS-OUTER's handover had left at E_{max} where the 1996 fit stood for a valence complex the 1995 outer row does not cover (Fe I x0.012 and Fe II x0.0066 steps, now 1.23 and 1.13; genuine K edges C/N/O x19-21), and put E_{max} , a step of the cross section, on a bin edge (it was not). Approximation stated at the code site: ion advanced one stage, one electron of $h\nu - E_{\text{th}}(\text{outer})$ to the cascade (MEASURED with the degradation routine: 0.95-1.09x the heat and 1.10-1.37x the secondary H I ionizations of the true two-electron event; the un-spent I_2 is 24-35 eV of a 300-550 eV photon), fluorescence neglected (with $e_{\text{top}} = 1240$ eV only C, N, O, Na have K holes; the high-Z ions contribute L shells only). Impact (worker, same-snapshot control; metals-off `mol_base_handoff` byte-identical): `wasp_full` 300 the wind 1-3 percent hotter and faster, base heating +26 percent, T at most +9.5 percent, Fe II-

I/Mg III/N III at the base x74-160 from vanishing values, **Mg II falls 84 percent at $r = 1.178$** (the h&k transit line); mol_metals base heating +26 percent, T 1.4e-3; lower_profile heating +21 percent at the base, T 2e-5; metals' share of the 100-1240 eV absorption 2.2 -> 3.5 percent (wasp_full), the band 0.38 percent of the whole; log10 Mdot unchanged to 0.01 dex in the snapshots. Krause 1979 (JPCRD 8, 307) K/L fluorescence yields could not be obtained (NIST 503, AIP 403): requested from the user; no yield is quoted anywhere. Advisor verification: physics_probe 1479/0, grid_and_gates 93/0; spectrum_type segfaulted in loaded_sed_threshold_edges (its threshold array was sized $n_{\text{mion}} + 16 = 43$ while the active list now reaches 126), fixed by sizing it with the module's $n_{\text{thr_max}}$; suite green after. ($n_{\text{thr_max}}$ was a local parameter of set_energy_vectors; hoisted to module scope, public, so every caller of ionization_thresholds_active sizes its array by the one bound; spectrum_type 154/0, physics_probe 1479/0, grid_and_gates 93/0 after.)

- **METALS-INNER addendum, Krause 1979 yields** (comments and documents only; user-supplied Krause_1979_JPCRD_8_307.pdf, read from page images where the OCR is noisy, alignment checked on $\omega_K(\text{Fe}) = 0.340$): Table 3 ω_K for C/N/O/Na 2.8e-3/5.2e-3/8.3e-3/0.023, Mg/Si/S/K/Ca/Fe 0.030/0.050/0.078/0.140/0.163/0.340; Table 5 effective L1/L2 yields (Coster-Kronig included per the text, $\nu_3 = \omega_3$) for Mg to Fe all below 6.5e-3; Table 2 uncertainties (ω_K 40-10 percent at Z 5-10, footnote (a) on molecules and solids). With $e_{\text{top}} = 1240$ eV only C, N, O and Na I can hold a K hole, so the neglected radiative branch is at most 2.3 percent of a K-hole event (under 1 percent for the three coolants) and under 0.7 percent of any L-shell event; raising e_{top} past 1.3 keV (Mg, Si K) or 4 keV (Ca, Fe K) is the condition under which the approximation must be revisited, stated at the code site, in photoion_cross_sections.tex (new tab:kedge), code_status and decision item 7.
- **LYA-BETA landed, advisor verification** (lya_rt.f90, hydrogen_n2_rates.f90 ($A_{2p1s} 6.3e8 \rightarrow 6.2649e8$, the nonrelativistic dipole value times the reduced-mass factor, = NIST ASD's Wiese and Fuhr 2009 value), parameters.f90 comments; probe lya_escape_probability 24 assertions; advisor rebuild physics_probe 1479/0, certification 29/0): the escape probability is the published static-slab solution, $\langle N \rangle = \text{Neufeld 1990 eq. (3.27) at zero destruction} = \text{Harrington 1973 eq. (40)} (4 \sqrt{6})/\pi^2 u_2 B = 0.909316 B$ for a mid-plane source (the 0.9093161 of the REF-NEUFELD entry was a transcription slip), $\beta = 1/(1 + \langle N \rangle)$ (exact at both limits: a generated photon is emitted $1 + \langle N \rangle$ times, and the budget $n_{2p} A \beta = P$ wants the escape chance per emission), the two faces sharing one escaping population by Neufeld eq. (2.25) instead of the former product, $\Phi_i(x_i)$ summed in closed form; the Voigt parameter has left the escape probability. a still enters only lya_wing_penetration_coefficient (where an incident stellar photon crosses in one flight, a different question, judged at the code site). Defect found in passing: $\beta_{\text{tot}} = 1 - (1 - \beta_{\text{esc}})(1 - \beta_{\text{sob}})$ lost digits at the base (4.8e-9 relative), now $\beta_{\text{esc}} + \beta_{\text{sob}} (1 - \beta_{\text{esc}})$ (closure 2.2e-16). RED before 4.79e2 / GREEN 2.19e-16 on the end-to-end identity. **Impact, wasp_full to its du stop (2 h 59 m): log10 Mdot 13.34 -> 13.34 (+1.6 percent, 0.0067 dex); Jint/n_2p x84 at the base, x73 at 1.05, x37 at 1.10, x10 at 1.20, x0.75 at 1.50 R_p; T within 2 percent; the old min(1, .) cap had zeroed the internal field from 1.15 R_p outward. H(n=2) chord column x31.5 at b = 1.00, x2.8 at 1.20, x1.005 at 1.45: every H-alpha/H-beta transit number in this repository from a Jlya escape-prob: True run is superseded** (He 10830 not

affected). Validity $(a \tau)^{1/3} > 10$ holds in 349 of 504 cells, out to 1.193 R_p, exactly where the change is an order of magnitude. Advisor follow-ups applied: the second A_{2p1s} in `exhale_transit_lib.py` and `docs/transmission_spectrum.tex` set to 6.2649e8 (one value with the wind model), the stale $E = \min(\text{boost}, 1/\beta)$ comment in `parameters.f90` replaced by the implemented $1 + (\text{boost}-1)(1-\beta) T_{\text{star}}$. Pending: the B6-style print of the Ly-alpha domain record in `certification.f90` (the record and accessor exist; LYA-DOM, with the gate's follow-ups).

- **LYA-DOM** (advisor; certification.f90): the Ly-alpha damping-wing domain record (lya_wing_domain_record: cell visits below the $(a \tau)^{1/3}$ limit, visits seen, the smallest value, the limit) joins the out-of-domain closure count and is printed beside the H3+ and CO records, informational like them. certification, run_mode, attempted_step green. **Every worker item of the series has landed; the series gate begins (single build, make test, the 16-case matrix, movement table, one golden refresh, fcheck, docs pass).**
 - **Series gate, 2026-09-07** (PLAN rev2 steps A to C and the items above, one series): single build of the shared tree (make -q up to date, EXHALE.x md5 ba3d34e6eae); make test green apart from the two steady suites, which then passed on the wasp_full_newton run log (EXHALE_STEADY_RUNLOGS); the 16 default cases run against that binary (the harness holds a single-instance lock, so wasp_full ran through run_check.sh check and the other fifteen through a runner replicating its steps for each case, all OMP_NUM_THREADS=1); every case moves against the previous goldens, as every item of the series said it would. Movement of the physical columns over the cells where the golden value is at least $1e-3$ of the column maximum (max relative; T median; the change of $\log_{10} \dot{M}$ from the mean of $r^2 \rho v$ over 1.20-1.45 Rp): | case | rho max | v max | p max | T max | T median | dlog10 Mdot | log10 Mdot new | |---|---|---|---|---|---|---| | wasp_full | 2.97e-01 | 1.28e+00 | 1.65e-01 | 9.18e-02 | 3.8e-02 | +0.0095 | 13.35 | | wasp_he23off | 2.95e-01 | 1.30e+00 | 1.62e-01 | 9.18e-02 | 3.9e-02 | +0.0084 | 13.35 | | wasp_full_newton | 6.96e-01 | 3.51e-01 | 4.30e-01 | 2.35e-01 | 8.7e-02 | +0.0414 | 13.30 | | mol_base_handoff | 1.60e-01 | 2.50e-01 | 1.52e-01 | 2.33e-01 | 2.2e-02 | -0.0440 | 10.57 | | mol_metals | 1.51e-01 | 2.25e-01 | 1.53e-01 | 2.07e-01 | 2.3e-02 | -0.0476 | 10.58 | | mol_lyman_werner | 2.12e-01 | 4.87e-01 | 1.99e-01 | 2.55e-01 | 2.6e-02 | -0.0011 | 10.54 | | mol_diffusion | 1.59e-01 | 2.50e-01 | 1.52e-01 | 2.33e-01 | 2.2e-02 | -0.0440 | 10.57 | | mol_ir_bands | 1.51e-01 | 2.26e-01 | 1.53e-01 | 2.07e-01 | 2.3e-02 | -0.0475 | 10.58 | | mol_sec_ion | 2.16e-01 | 4.03e-01 | 1.96e-01 | 2.53e-01 | 3.0e-02 | +0.0149 | 10.51 | | mol_carrier | 1.54e-01 | 6.37e-01 | 1.44e-01 | 2.24e-01 | 1.9e-02 | +0.0106 | 10.56 | | lower_profile | 9.40e-03 | 3.22e+01 | 7.50e-03 | 9.77e-01 | 8.8e-03 | -0.0335 | 7.29 | | hydrostatic_column | 2.09e-12 | 2.81e-03 | 3.54e-12 | 1.40e-03 | 2.5e-14 | -0.0000 | 8.27 | | oxygen_chemistry | 8.62e-01 | 1.29e+02 | 9.76e-01 | 7.61e-01 | 3.7e-01 | +nan | 9.61 | | hp_zero_seed | 2.25e-05 | 1.81e-04 | 1.61e-04 | 3.20e-03 | 4.4e-05 | +0.0000 | 10.56 | | hp_trace_seed | 2.24e-05 | 1.81e-04 | 1.61e-04 | 3.20e-03 | 4.4e-05 | +0.0000 | 10.56 | | hp_front | 2.07e-05 | 5.77e-04 | 3.66e-04 | 1.00e-02 | 6.7e-05 | +0.0000 | 10.56 |
- The maxima sit at the fronts, which move by a cell; the T medians (2-9 percent on the wind cases) and the $\dot{M}$ changes (within ± 0.05 dex everywhere it is defined; oxygen_chemistry has a reversed flow in the window) are the state of the physics after B3c, B4-1/1c, LYA-BETA, the WENO3 coefficient fix, the metals (REF-METALS, METALS-OUTER/INNER), the LW band, the CO channels, ACCEPT-1/2 and the

rest. `wasp_he23off` exits 2 (its stationary claim refused by four inventory entries, as `A2tol2` measured at the default tolerance; outputs written in full); `wasp_full_newton` exits 0, certified, `log10 Mdot` 13.30. PDFs of the 25 dated documents, the paper and both posters rebuilt. **Golden refresh pending the user's hand:** the archive copy `golden` -> `golden_pre_series_20260907` and `run_check.sh golden` were refused by the session's permission classifier; the commands are in the report to the user. The final confirmation (`make check` against the refreshed goldens, serial) follows the refresh. **Golden refresh done by the user (2026-09-07 13:19):** `golden/` archived as `golden_pre_series_20260907/` (17 cases) and re-snapshotted from the case outputs of this gate (checked: `golden` equals output on `wasp_full`, `mol_metals`, `oxygen_chemistry`, `lower_profile`). The serial `make check` against the refreshed goldens runs next as the bitwise confirmation, then `run_fcheck.sh`.

- **User manual, environment-variable table** (user report, 2026-09-07): the fixed table float of the environment hooks (67 rows) overflowed page 11 into the footer; it is now a longtable with the caption at its head, breaking across pages 10-11 with a "continued" foot. PDF rebuilt. Pages 20-21 (user report): the keyword table's long file paths were unbreakable `\texttt` tokens and ran past the right margin; 18 such paths are now `\path{}` (url package, breaks at / and _), one inside a caption kept as `\texttt` (a moving argument). Rebuilt with `latexmk`, so the longtable column widths and cross-references are resolved (a single `pdfLatex` pass leaves both unresolved, which is what produced the "Table ??" and the first overflow the user saw).
- **Update log TeX twin** (user request, 2026-09-07): `docs/Update_EXHALE_stage2.tex` and its PDF now exist, GENERATED from this Markdown by `src/utils/update_log_to_tex.py` (pandoc body, the memo-class preamble of the stage logs); regenerate after appending here. 56 pages at this entry.
- **Decision item 4 closed** (user, 2026-09-07): candidate C, `wasp121b_composite_huang2023.txt` (solar XUV shape below 1700 Å rescaled to Huang et al. 2023's `F_XUV` and `Ly-alpha`, the MUSCLES WASP-17 photosphere above), is the WASP-121 b spectrum of record, as both `input.inp` files already state; A and B remain sensitivity inputs. No decision item of `To_be_determined_by_user_20260906.md` is open. **make check against the refreshed goldens: REGRESSION PASS, all 16 cases byte-identical (64 files), 13:21-16:25 KST 2026-09-07** (the harness runs the cases eight at a time, one thread each). `run_fcheck.sh` follows. **run_fcheck.sh: CLEAN** (1500 bounded steps of the HD 209458 b folder case under `-fcheck=bounds,do,mem,log10 Mdot 9.56`) after one harness fix: the scratch copy of the planet input named its spectrum by the folder's relative path, which the script now rewrites to the repository's absolute path for the `Spectrum` file and `Lower atmosphere` profile keys (the first run failed on "Cannot open file `../inputdata/sed/...`", not on a runtime check). Production EXHALE.x restored, `make -q` up to date, md5 `ba3d34e6eae`. **The series is closed:** goldens refreshed once, `make check` byte-identical, `fcheck` clean, docs and PDFs rebuilt. What follows is in `docs/session_handoff_20260907.md`.
- **HYG-PY** (advisor, 2026-09-07; `python/paper_data.py`, `python/make_struct_figures.py`): `HEAT_FIXED` (the headerless-file fallback) and `HEAT_CH_LABEL/HEAT_CH_ORDER` (the heating-budget panel) now carry all 19 deposit channels of `heat_channel_name` with the descriptive brackets stripped the way `_clean_name` strips them (`heat_He23S_assoc`, the two Lyman-Werner halves, `heat_mol_chem`, `heat_FUV_photolysis`, `heat_oxygen_collisional`, the two CO channels); the panel loop already skips a channel a run's file lacks. Both scripts parse; the loader reads the file header, so the fallback list was never what a real file was read by.

- **THREAD-DET** (advisor, 2026-09-07; `util_ion_eq.f90`): the thread count is no longer an input to any result. Mechanism, MEASURED: `Cool_coeff.o` imports glibc libmvec's two-lane `_ZGVbN2v_exp`, `_ZGVbN2v_log`, `_ZGVbN2vv_pow` (gfortran `-O3` vectorizes the rate-coefficient loops), and the vector variants agree with the scalar libm to one ulp, not bitwise; inside a block of cells the loop takes lane pairs from `j_lo` and an odd remainder by the scalar call, so where a block STARTS decides which cell meets which variant. `chemical_rate_coefficients` and `eval_cool` cut the grid into `2*omp_get_max_threads()` blocks, so the block boundaries, and with them the last bit of every rate, moved with the thread count (the $6.6\text{e-}14$ of COST3/HYG-MAIN). Fixed: both sweeps use blocks of the fixed even length `xuv_rate_block = 32` (16 blocks on the 504-cell grids); the threads decide only who takes which block. MEASURED on `wasp_full 300`: old binary at 1 thread, new at 1 thread and new at 16 threads are byte-identical in `Hydro_ioniz`, `Ion_species`, `Heating_breakdown`, `Cooling_breakdown` (the old single-thread partition happened to start both its blocks at even offsets, so the refreshed goldens are untouched). Shared EXHALE.x rebuilt from the tree (make `-q` up to date).
- **DOCS-LINES** (documentation only, 2026-09-07): every `file.f90:NNN` citation in the nine design and status documents (d0 922 tokens on 309 lines, b1a 127, b1 64, b3b 24, a2 7, a0 5, b3a 5, b4 2) re-anchored to file + routine, declaration or labeled comment block, read against the live tree; the step tables of d0 6.1 and b1 7.1 index by the `as_op_*` operation tag of `attempted_step.f90`; the convention is stated in each document. Citations of removed code (the composition projection, the two-iteration energy update, the CO ceiling, the cell-velocity carrier advection, the element advective term, DB96 eq. (39), the H3+ clamp, the second H2 partition function) kept as dated records naming the removing item. Stale values corrected in passing: `e_lw_photon_erg` $1.88021\text{e-}11$ erg (11.7354 eV) over 912-1201 Å, `binary_element_diffusion.f90` 3037 lines, a2 section 1's chemical count now passed by the marching caller. All 190 names introduced grepped present in `src/`. Recorded: pre-existing markdown table pipe misalignments (unescaped `|` in math) in a2, b4, `code_status`, d0; two markdown line citations of `grid_and_gates/README.md` left.
- **B5, carrier rows** (`steady_newton.f90`, `diffusive_photochemistry.f90`, `input_read.f90`; new suite `steady_species_rows` 41 + 8 rows; advisor rebuild: it and certification, `run_mode`, `carrier_retry`, `coupled_source_step`, `species_face_flux` green): the Newton unknown vector carries one row and one unknown per solved carrier (H2, OH, H2O, CO, H+) through a row registry in place of the H2-only `nvar = 4` special case; band geometry derived, `kl = ku = 3 nvar - 1`, `ncolor = 6 nvar - 1` (reproduces the 8/17 and 11/23 the code carried); the completion flag reads the species rows of the system it solved; EXHALE_SPECIES_JAC_TEST asserts `Jr` inside the solve (the run-level test builds its Jacobian before the registry exists). Ionization transport + Solver: Newton supported by derivation when the proton is a Newton unknown (`x_hp_fix` handed to the sweep), refused with the reason otherwise; MEASURED `nvar = 5`, Jacobian action $7.2\text{e-}7$ (all rows), $4.6\text{e-}5$ (species rows), 0 unresolved colors. Two pre-existing defects fixed: `frozen_residual` left the species slots of an `intent(out)` array unwritten (directional check relative error 1.0 from stale memory) and `n_headroom_iterate` was never assigned (the trial-admissibility threshold stood at 0; with the four oxygen balances as rows all 41 colors came back unresolved). `wasp_full_newton` stays certified (13.30, byte-identical); H2-only control byte-

identical; marching path byte-identical (wasp_full, mol_base_handoff 300). **Findings:** no regression case enters the steady solver, so every golden is a marching snapshot and cert_tol_carrier/cert_tol_element still have no anchor (contract section 11 records two rows from solves that did not meet their target, 7.46e-3 and 5.82e-3, not anchors). **Open, and a physical-consistency defect:** the stationary carrier row still uses the cell-velocity advective form while the marching rows ride on $F_{\rho} Y_s^{\text{face}} / m_s$ since B4-1c (two operators, two fixed points): B5b. Also open: the element rows (element_transport_residual must return a residual per element, not the worst at each cell), the cell's own field in the cell residual (interface written: FIELD-SELF), reload consistency, three EXHALE_main.f90 sites (the registry name, EXHALE_JAC_TEST coverage, the PTC refusal text).

- **B5b** (species_face_flux.f90, steady_residual.f90, steady_newton.f90, certification.f90, binary_element_diffusion.f90, diffusive_photochemistry.f90, three sites of EXHALE_main.f90; suite steady_species_rows 67 rows; advisor rebuild: it and species_face_flux, certification, carrier_retry, coupled_source_step, run_mode, acceptance_classes green): ONE discretization of the material advective term exists in the code: the RK stages, the stationary carrier rows, the fixed-wind carrier relaxation and the stationary element rows all form it as the divergence of the same face species fluxes (species_face_fraction, species_face_flux, new species_flux_divergence) on the mass row's faces, areas and volumes; the cell-velocity upwind form, its deferred van Leer correction (carrier_advection_correction, adv_corr, carrier_slope), the base Dirichlet record it needed and the advective coefficient ntv are gone from diffusive_photochemistry.f90 (RED 76.7 relative against a stage-1 update on a sign-changing face flux, GREEN 7.5e-14). element_transport_residual returns one residual per element (the certification prints one entry per element: lower_profile C 0.977, O 0.921, N 0.968, seven elements "not applicable"); rows srow_element_he and one srow_element_trace per metal element registered before the carriers, project_element_mass_fractions the write-back. mol_diffusion reloaded with Coupled carrier solve: nvar = 4, one element row, Jacobian action 4.9e-3 / 1.3e-3, 0 unresolved colors, the solve refused at the He/H row (0.9987) with no descent direction: the element system is expressible and its Jacobian right, not yet solved from that start (the element unknown moves the whole column's opacity, which the banded model does not carry; the "all rows" bound of the suite widened 1e-4 -> 1e-2 with the measurement at the site). Reload consistency asserted on the binary. EXHALE_main.f90: set_transportated_species_rows (not carrier-specific), the two EXHALE_JAC_TEST inserts, the PTC text; at that site two pre-existing defects fixed: the species slots of Yvec were read uninitialized (2.4e2 against 1.85) and the probe direction was unscaled (3.3e1 at nvar = 4); now 3.2e-6 / 4.5e-6. **By construction the stationary species rows now inherit the mass row's imbalance** ($\text{div}(F_{\rho} Y) = Y \text{ div } F_{\rho} + F_{\rho} \text{ grad } Y$; the old form was the second term alone), so on marching snapshots whose mass rows read 1.1-1.9 of their terms the rows read O(1) where they read 4e-4 to 0.4 before. Marching path byte-identical on wasp_full, mol_base_handoff, mol_carrier, mol_diffusion, lower_profile at 300 steps; wasp_full_newton reloaded certifies (13.30, 2.6e-9 of the stored state). Anchors still UNCONFIRMED (contract 11.3). Recorded: element_diffusion_step / relax_element_composition still advect on the smoothed steady mass flux, so the elemental Picard alternation and the elemental row are two operators (B5c); carrier_rows_advect kept as the on/off switch of one operator; Coupled carrier solve now also governs

the element rows (key not renamed); ionization_equilibrium.f90 cites carrier_base_state, which no longer exists (to FIELD-SELF). THREAD-DET confirmed by the official check: make check on the rebuilt binary, REGRESSION PASS, every case byte-identical (16:58-20:04 KST 2026-09-07; the binary of that check also carried whatever of B5 had landed by 16:58, and B5's marching path was measured byte-identical anyway). A further make check follows FIELD-SELF and B5c.

- B5c** (binary_element_diffusion.f90; suite steady_species_rows 60/0; advisor rebuild: it and species_face_flux, certification, carrier_retry green): the elemental relaxation and the stationary elemental row are one operator. element_diffusion_step takes the FACE MASS FLUX and forms its advective term as the divergence of the face element fluxes through the three shared routines (element_advective_divergence, one place); relax_element_composition reads that flux from the mass row of the state (face_mass_flux_of_state) and no longer smooths a steady $\dot{m}/(4\pi r^2)$; element_transport_residual takes both halves of the helium row from composition_residual and the trace rows from the new trace_composition_residual, the routines the step itself solves, so the fixed point is the row's zero by construction; solve_trace_element_in_hydrogen is a deferred-correction loop that takes one pass, term for term the old direct solve, when the advection is off (the marching path). MEASURED on a synthetic column relaxed to its own convergence: the row reads 8.65e-13 (He alone), 8.45e-13 / 5.38e-13 (He + ten trace elements) of its terms, the advection carrying 0.993 of them; the replaced form leaves 6.4e-3 on the same state. The smoothed-flux rationale does not survive: it repaired a form (the element equation minus X times the mass equation) that is gone, and its consequence is now a measured property of the equation: on a face mass flux that does not conserve mass the conservative balance has no bounded steady composition (X driven onto 0 and 1). Marching path byte-identical (mol_diffusion, lower_profile 300 steps); the direct-steady reload of mol_diffusion byte-identical, and its outer pass never reaches the relaxation (the solve ends info = 2, B5b 6.3), so the run-level RED/GREEN does not exist. **Found: diffusion_tests.x has not linked since B5b** (binary_element_diffusion now uses steady_residual_mod, whose chain the test's DIFT_SRC does not carry): a functions/ module depending on time_step/steady_residual is a layering inversion; the flux goes by argument (DIFT-LINK, launched), and make test's diffusion_tests row is red until it lands. Also recorded: the relaxation's convergence measure is helium-only (DIFT-LINK folds the trace drift in); the vanished-element write-back branch leaves a trace row at 1e-4 to 2e-2 where an element is driven to exactly zero (deliberate guard, measured, documented at the test site); element_transport_residual's dummy v is unread.
- FIELD-SELF** (ionization_equilibrium.f90 self-field loop, default one pass; util_ion_eq.f90 and System_HeH* comments; probe photoionization_field_self_consistency 12; advisor rebuild: coupled_source_step 36/0, physics_probe 1491/0, certification, acceptance_classes green): the cell's own optical depth CAN now be closed inside the cell solve (one pass leaves a 4.06e-2 self-consistency defect in the ionized fraction of a cell one optical depth thick, the loop closes it to 2.8e-15 in 13 passes, contraction 0.080; the field routines already formed the B4-6 cell mean from the cell's own densities, so no field code changed), **and doing so buys nothing: COST3's attribution is refuted**. MEASURED on wasp_full 300: coupled passes 10.09 -> 10.01 for 1.65x the wall time (2 self-field passes) and 1.87x (8); mol_base_handoff

1500: 7.19 -> 7.24 for 2.0x; the decisive binary with BOTH levers (carried columns at block 8 AND four self-field passes): 7.20 passes, 190 s, against COST3's 7.13 / 103 s for the columns alone. (**Superseded by COST4 below**: with both stellar lags closed at the settings that close them the count is 8.18, not seven, and the two couplings named here are worth 0.08 of it; the count is the slowest rung of a ladder of modes.) Same fixed point: state movement $\leq 1.4e-7$, at the $1e-8$ tolerance's level. Default output byte-identical on wasp_full 300, mol_base_handoff 1500, mol_metals 1500, hydrostatic_column; wasp_full_newton uncapped CERTIFIED, both files byte-identical to its golden (that binary carried B5b too). Kept in the tree at its inert default on COST3's precedent (the switch is the record of a measured negative result). Comment fixes in passing: System_HeH_metals.f90 said the sweep is serial (untrue since the parallelization; its arrays are threadprivate); the carrier_base_state citation of ionization_equilibrium.f90 rewritten to carrier_base_composition_imposed. Recorded: he_ground_singlet_density's clamp counter needs a lock when the self-field loop runs (a slot per cell would be free, composition.f90); f_vib_quench/e_vib_bound stay at the entry composition inside the loop; the COST3 attribution in that report, its log entry above and the csm_max_pass comment of EXHALE_main.f90 are superseded by this measurement (comment to be corrected when DIFT-LINK releases main).

- **DIFT-LINK** (binary_element_diffusion.f90 and its three call sites in EXHALE_main.f90, steady_newton.f90, certification.f90; suite steady_species_rows 62 rows; advisor rebuild: it and species_face_flux, certification, coupled_source_step green): the element module no longer uses steady_residual_mod; the face mass flux is a required argument of relax_element_composition and element_transport_residual, obtained by each caller with face_mass_flux_of_state. The layering inversion is gone and **make diffusion_tests links again with the Makefile and DIFT_SRC untouched: 35 passed, 0 failed** (MEASURED by the advisor on the live tree; it failed at the link since B5b). The relaxation's convergence measure now runs over every element it relaxed, each against its own reservoir (element_composition_distance): RED 5.99 / GREEN 0.0 on a column where Fe moves 1.78 while helium moves 0.25, so the helium-only measure returned a number 7x too small to bound the element that moved. Byte-identical on mol_diffusion, lower_profile 300 steps and the mol_diffusion direct-steady reload. Dead code removed (helium_mass_fraction, an unread element_mass_fractions call). **Advisor fixes on its findings**: the limiter of species_face_fraction divided by $Y_{av} - Y_f(j)$, which vanishes when the donor cell's average is itself negative (a trace species handed back at a small negative density; MEASURED as a SIGFPE under -ffpe-trap=zero): the anchor is now the average clamped into [0,1], which makes both denominators strictly positive and is what the comment already claimed; mol_diffusion 300 steps byte-identical, since at production flags the infinity was clamped to the same 0. The csm_max_pass comment of EXHALE_main.f90 and docs/steady_solver_design.md section 15 corrected to the FIELD-SELF and B5c measurements. Recorded: diffusion_tests T9 reads $1.098e-12$ against its $1e-12$ bound at -O1 -fcheck (an optimization-level rounding sensitivity of relative_settling_mass, 35/0 at -O3); the outer loop's under-relaxation damps helium only while the drift it tests can now be set by a trace element.

- **Batch gate after B5/B5b/B5c/FIELD-SELF/DIFT-LINK/THREAD-DET** (2026-09-07/08): shared tree re-

built (EXHALE.x md5 eba2d7f2260f, make -q up to date); make test green, the two steady suites passing on the wasp_full_newton run log and the diffusion_tests row restored; make check **REGRESSION PASS, all 16 cases byte-identical** (20:54 to 00:05 KST). Every item of this batch was designed to leave the marching path alone and the check confirms it: the goldens refreshed at the 2026-09-07 series gate still stand, and no golden was refreshed here.

- **HYG-T9** (advisor, 2026-09-08; src/tests/diffusion_tests.f90): T9's ambipolar-limit bound sat exactly on the arithmetic's noise floor, so the suite passed at -O3 and failed at -O1 with checks (DIFT-LINK's finding). MEASURED at both levels: the neutral limit is exact (no electron term), the H+ plasma reads 8.207e-13 (-O3) and 7.740e-13 (-O1), the He++ plasma 7.083e-13 and 1.098e-12. Mechanism, written at the site: the limits are exact algebra in the masses, so the deviation is arithmetic, and the ambipolar term of the plasma limits comes from a centered difference of $\ln \rho$ over dr whose logarithm is $O(b_0/r)$, so each ulp reaches dm_{eff} amplified by $1/dr$. The bound is now one decade above the largest measured ($1e-11$), the anchoring rule the certification tolerances use, and the test still catches a wrong ambipolar limit by ten decades (a wrong limit misses its target by 0.1 to 1) with the neutral limit still asserted exact. 35/0 at -O3 and at -O1 -fcheck.
- **B5d** (steady_newton.f90 only; suite steady_species_rows 77 rows; advisor rebuild: it and certification, species_face_flux, coupled_source_step, carrier_retry green): the item's own brief expected the element rows to fail because an element unknown moves the whole column's opacity, too non-local for the banded model. **MEASURED and REFUTED**: the scaled Jacobian column of an element unknown carries at most 6.3 percent of its 2-norm outside the band, the same order as a hydrodynamic column (mass 0.0018, momentum and energy 0.0000), so the diagonal correction, the rank-one column-integral update and the identity block the brief offered would each have repaired a defect that is not there; none was written. What does break the model, located by the same measurements and fixed: (1) the base element row was structurally EMPTY, cell 1 being the Dirichlet reservoir of the element operator, so the banded model was singular by construction (its LU pivot grew to 8.413e19 against a largest band entry of 5.229e3; now 1.433e6) and it now carries the boundary condition the operator states; (2) the element ROW SCALE collapses on the abandoned iterate (helium rows spanning 1.277e-29 to 2.360, so an ordinary $dF/dY = 46.7$ became a scaled band entry of 4.890e22 and $||grad\ merit|| = 1.104e44$), and the row scale is now floored at the flow-time rate of the quantity the row transports, the hydrodynamic convention with no chosen constant. Result: the same case reaches $||R|| = 1.723$ against 1.974 at outer iteration 18 instead of 8 with $||grad\ merit|| = 4.813e3$, 40 decades lower and a number the model's own entries can produce. **The element solve still does not converge and the item does not claim it does**; cert_tol_element and cert_tol_carrier are unmoved and UNCONFIRMED. New diagnostics: banded_model_conditioning (empty rows and columns stated once per registry, pivots and scales under EXHALE_BAND_REPORT), jacobian_column_reach_beyond_the_band (EXHALE_JAC_COLUMN_REACH). mol_diffusion 300 steps byte-identical; wasp_full_newton byte-identical, info = 0, CERTIFIED, 13.30, same 6 outer iterations. **Named as what is left: the globalization** (the non-monotone search accepts steps that raise $||R||$ from 1.7 to 25; the trust-region route gives up at radius 3.8e-16 after 12 ray-test refusals while the merit is still falling): item B5e. Also recorded: carrier_row_term_scale

has the same collapsing structure and was not measured (no carrier row in that case); the trace-metal case of the floor is covered by inspection only.

- **COST4** (measurement; NOTHING in the repository changed by the item, its four candidate files byte-identical to the snapshot it took; every experiment a private binary in a scratch tree whose no-switch build is byte-identical to the control): **the coupled loop's pass count is a ladder of nested modes and equals the SLOWEST of them, not their sum.** MEASURED on `wasp_full` 300, one thread, removing one composition coupling at a time: the production loop 10.09 passes; the stellar field's dependence on the composition removed, 7.16; the temperature removed as well (the energy solve replaced by its own converged answer), 4.94; the He recombination rates too, **2.00** (which reproduces, with a different probe, the 2 passes COST2 measured with every composition-dependent input frozen). Both hypotheses of the brief are refuted: the two diffuse couplings are worth at most 0.08 of the 10.09 (Balmer off 10.09, He rec off 10.01, both off 10.01, He rec frozen after pass 1 10.01, and on `mol_base_handoff` all three read the control's 7.19 to the last digit), and **the Balmer continuum is not a lag at all**: `excited_H_update` fills its arrays once BEFORE the loop, so they are constants of it, and refreshing it every pass costs 1.5 percent of wall for +0.01 passes. The stellar lag closed IN FULL for the first time (a one-cell block and eight self-field passes): 8.18 passes for 218 s against 10.09 for 139 s, the next rung, with freezing the field entirely giving 7.16. The temperature is not the residual either: the composition sweep alone, at the temperature the step converged to, takes 10.94 passes (6.32 against 7.19 on `mol_base_handoff`). **Decision: change nothing** (no loop-internal Balmer refresh, no self-consistent He recombination coupling, no default moved); to go below seven passes every rung must come out at once. Advisor: the three statements this refutes are corrected in place, the `esm_max_pass` comment of `EXHALE_main.f90`, the FIELD-SELF entry above (marked superseded) and open item 1 of the 2026-09-08 handoff. **The one remainder, now open item 1**: on `mol_base_handoff` the ladder is flat (freezing the field RAISES the count 7.19 -> 7.41) and everything the item could reach accounts for 0.6 of its 7.19 passes; the named candidate, READ and not measured, is `n_tot`, the third-body density of the three-body molecular reactions, built by `calc_ntot` from the ENTRY composition, and with it the H2 vibrational quench fraction and bound energy. Also recorded: `xuv_field_block_cells` and `xuv_self_field_passes` are compile-time parameters, so measuring the two together needs a rebuild; `heating_of_composition` calls `he_rec_coupling` a second time whole-grid every pass for its heating alone (both calls together bounded at 1.7 percent of wall); on `mol_base_handoff` the He recombination coupling is what keeps step 0 inside the pass cap.
- **COST5** (measurement; NOTHING in the repository changed by the item, its four candidate files byte-identical to its own snapshot; probe binaries byte-identical to the control at their default): the molecular remainder COST4 left is named, quantified, and the ladder reaches its bottom rung. **The count is arithmetic, and the law is measured and then tested**: $\text{passes} = 2 + \log_{10}(\text{the step's own excursion} / \text{the stopping tolerance}) / (\text{decades per pass})$. On `mol_base_handoff` the composition contracts by 0.171 a pass and the temperature by 0.072 (0.77 decades a pass) from an excursion of $7.0\text{e-}5$ to a tolerance of $1\text{e-}8$: 7.00 predicted, 6.93 MEASURED; on `wasp_full` 0.222 and 0.199 from an excursion seven times larger, 9.2 predicted, 10.09 measured (so the molecular cases are the cheap ones, not the pathological ones). Tested, not fitted: the law says the count depends

on the stopping tolerance through one logarithm at 1.30 passes per decade, and moving both tolerances together gives 8.06 / 6.96 / 5.68 / 4.04 at $1e-9$ / $1e-8$ / $1e-7$ / $1e-6$, i.e. 1.34 over three decades; and it is not the cell solve, six decades of hybrid tolerance leaving the count at 6.97 for up to 1.99x the wall. Of the four candidates only the named one is an iterate: `n_tot`, the third body of R12/R13/R15 (the ONLY composition dependence of the molecular network outside the field: of seventeen hoisted coefficients exactly two see it, both linearly). Holding it: `mol_base_handoff` 7.19 -> 6.03 at 1500 steps, `mol_metals` 6.99 -> 6.01; it limits the composition mode's RATE (0.171 -> 0.073) but the count falls only 6.93 -> 6.11 because the temperature mode then binds. The other three are entry state through the operator-split field (vibrational pair 6.96, electron density 6.94, Lyman-Werner 6.96, against 6.96), and the metal element totals are invariant across a sweep by construction (6.99 against 6.99, exact). With every input AND the temperature held the sweep needs **2.01 passes**, reproducing COST2's and COST4's atomic bottom rung on the molecular path. **Decision: nothing closed.** The mode that sets the count is the temperature/composition alternation (`EXHALE_main.f90`, `energy_semi_implicit.f90`, not that item's files); closing the `n_tot` lag exactly needs two sites in `System_HeH_mol.f90` (also not its), and the in-file alternative costs a second cell solve per pass, which the cell-tolerance scan bounds at more than the 0.80 passes it returns. The production treatment is not wrong: at the fixed point the entry and returned compositions are one state, so the third body the network is solved with is the third body of the accepted state, to `esm_comp_tol`. Advisor: the law is now in the `esm_max_pass` comment; the misleading indentation of `composition.f90`'s `calc_ntot` branches fixed. Recorded: the loop's stopping test compares an ABSOLUTE mass-fraction change against `esm_comp_tol`, so it is a relative $1e-8$ on H2 and no constraint on a $1e-12$ species. **The one lever that keeps both the fixed point and the tolerance is sequence extrapolation** on the (T, c) pair after the third pass (the decay is geometric with a ratio stable from pass 2 and close across cases): ESTIMATED at three passes of seven on the molecular cases and four of ten on `wasp_full`, item COST6.

- **HYG-4** (advisor, direct; the four hygiene items of the 2026-09-08 handoff): (i) `he_ground_singlet_density`'s run-wide clamp counter is incremented with `!$omp atomic update` inside the function, so the named critical section around its one call inside a parallel region (`ionization_equilibrium.f90`, self-consistent field loop) is gone and the three comments that justified serial column formation by the counter are reworded (the columns are still formed before the blocks are solved, for the ordering, not for the counter); (ii) `element_transport_residual` no longer takes the cell velocity it never read (four call sites: `certification.f90`, `steady_newton.f90`, two test rows), the balance being the divergence of the FACE mass flux handed in; (iii) the outer Picard loop of `steady_wind_with_element_diffusion` halved the element under-relaxation omega whenever `comp_drift` failed to fall, but `comp_drift` is the worse of the element drift and the carrier drift while omega damps the element relaxation alone (the carrier pass is bounded by `carrier_trust`), so a carrier drift that failed to fall slowed the element alternation for nothing; the damping test now reads the element relaxation's own drift (`elem_drift`) and the exit test keeps the worst of the two. No regression case runs both He_diffusion and the carrier transport, so the matrix cannot move. (iv) `f_vib_quench/e_vib_bound` staying at the entry composition inside the self-field loop is CLOSED BY MEASUREMENT (COST5): they are entry state through the operator-split field, and holding them at the first-pass value is worth 0.00 passes (6.96 against 6.96). Suites: `steady_species_rows` 74/0, `certification` 29/0, `diffusion_tests` 35/0.

- B5e** (steady solve globalization; steady_newton.f90, the steady_species_rows suite, design doc section 17, contract section 11.5): **the first stationary solve with a species row that meets its own target.** MEASURED on mol_diffusion reloaded with Coupled carrier solve: True (one element row, nvar = 4): the scaled trust region takes $||R||$ from 1.986 to **4.345e-09** against a target of 1e-08 in 66 outer iterations (54 accepted), where the pseudo-transient line search aborts at iteration 18 on "no descent direction exists for the banded model" at 1.702. Two defects were named by instrumenting the eleven exits of trust_region_step and repaired where the measurement pointed: (i) the radius collapse was the CAUCHY LEG ASCENDING the model, because its gradient is the transpose of the BANDED A and has no sign (MEASURED $r0 \cdot A g = -2.1e13, -2.2e19$ at the collapsing iterates, pred negative and exactly linear in the radius); where $r0 \cdot A g \leq 0$ the leg is now dropped and the dogleg reduces to the Krylov leg cut to the ball, which always predicts a reduction (GMRES gives $r0 \cdot A sN < -||A sN||^2/2$); and pred is formed as $-(r0 \cdot As) - ||As||^2/2$ instead of the cancelling difference of two squared norms, whose sign on a short step was rounding; (ii) the best-iterate ledger ranked on $||R||$ (three hydrodynamic rows) while the state is judged on those AND every species row, so it could hand back the iterate with the worse element balance; it now ranks on distance_from_certification, the largest of the judged row measures over their tolerances, formed once inside eval_residual under an optional argument; the three-unknown branch keeps its comparison character for character. **Refused by measurement:** bounding the STEP by the judged rows (implemented in all three step controls) stalled the element solve at $||R|| = 1.92$ on 13 refusals; and the banded-gradient Cauchy length $(r0 \cdot A g)/||A g||^2$, more defensible on paper, left the same solve crawling at 1.8e-4 after 81 iterations against 4.3e-9 in 66 with the original length, so the original stays, recorded at the site with both numbers. **Default moved:** use_tr = (nspec_row > 0), the trust region is the step control of the coupled route (EXHALE_TRUST_REGION=0 restores the line search for measurement); no regression case sets Coupled carrier solve: True, so no golden can move. **Anchor:** the elemental transport He/H row of the returned state reads **4.633e-11 against cert_tol_element = 1e-8** (215x inside), the first measurement under that tolerance; it is NOT moved, because contract sections 9 and 10 want a ladder (the same case at Resid tol 1e-9 and 1e-10, not run) to tell a target-limited row from one at its floor. The state is not certified: its cell-1 hydrodynamic mass row reads 6.2e-11 against 3e-12, a base-cell matter and not an element-row one. **The carrier-row solve is not rescued:** on mol_carrier the trust region stagnates at 1.873 (line search 1.855) and 18 of its 49 iterations end on "no Krylov direction could be sampled", so cert_tol_carrier stays UNCONFIRMED, and the measurement says it is not a step-length problem. Worker MEASURED wasp_full_newton byte-identical (it 6, $||R|| = 8.131e-09$, info 0, CERTIFIED, log10 Mdot 13.30) and the mol_diffusion trust-region run identical to the pre-item text under EXHALE_TRUST_REGION=1 (every landed change inert on a path that already worked). Recorded, not fixed: EXHALE_SPECIES_JAC_TEST=1 perturbs the run it measures (iterations 1-5 agree to every printed digit, iteration 6 differs, the two runs then diverge; some module state survives a probe eval_residual; the same root writes n_eq_sweeps_model and the frozen background); and the ray test can refuse a step that reduced the merit far MORE than the model promised (ratio = 7187), which Algorithm 4.1 of Nocedal and Wright never does, the obvious amendment (gate it on ratio <= eta_accept) left unmade for want of a control measurement. The worker was stopped by a rate limit after writing its report; advisor verified the tree: build clean; suites steady_species_rows 74/0 (six new driver

rows), certification 29/0, species_face_flux 16/0, carrier_retry 101/0, coupled_source_step 30/0; one "per-iteration" phrase in run.sh reworded; wasp_full_newton re-run by the advisor against its golden with the B5e binary: all four files byte-identical, info 0, CERTIFIED, log10 Mdot 13.30.

- **Batch gate 2 (2026-09-08, 05:04 to 08:08):** make check on the tree as of B5e + HYG-4 (before COST6 landed; the harness built at 05:04, COST6's first edit is 05:17) **PASS, all 65 files of the matrix byte-identical** to the goldens refreshed by the user on 2026-09-07. make test in the same script failed 7 suites on their build-staleness precheck only (the shared build/ had not been remade after HYG-4 when the suites ran, the harness remade it afterwards); the suites themselves were run on the advisor's private build of the same tree: steady_species_rows 74/0, certification 29/0, species_face_flux 16/0, carrier_retry 101/0, coupled_source_step 30/0 then 39/0 with COST6, diffusion_tests 35/0, physics_probe 1491/0, and energy_update, grid_and_gates, spectrum_type 53/0, 93/0, 154/0. Every suite of make test is therefore green on this tree. No golden touched.
- **COST6** (extrapolation of the coupled loop's geometric decay; EXHALE_main.f90 +532 lines, coupled_source_step +3 whole-binary rows; **default OFF**, EXHALE_CSM_EXTRAP=1 turns it on): implemented, guarded, measured. The object is the GLOBAL pair (T, f_sp), which is exactly the loop's iterate (the energy solve rewrites u(3,:) and W(3,:) in full from T and the entry anchor each pass, so an undo of the pair is exact, and the guard-forced-refusal row reproduces the off run byte for byte); one Rayleigh-quotient ratio per block over all cells and species, the projection onto the dominant mode COST5 measured stable, and one damped ray, so no cell can move in a direction the global mode lacks. That is how it differs from COST2's Aitken taken cell by cell (12.00 against 11.05 passes): a ratio taken cell by cell spreads 200 percent across the grid and cannot be tested for validity; the global one is tested ($||d_k - \theta d_{k-1}|| \leq 0.5 ||d_k||$) before use. Admissibility is by construction (fractions in [0,1], He 2³S inside its He I column, T inside the energy solve's own bracket, all linear in the step, so the largest admissible multiple is taken); MEASURED, the worst violation is identically zero on every case at 300 and 1500 steps, 1 and 16 threads, and the step was never damped below 0.95. The stopping test and its tolerances are untouched: a candidate is kept only if the pass taken FROM it meets them. Three things the measurement taught: the mode ALTERNATES (cosine of consecutive increments -0.85 to -1.00 on mol_base_handoff, +0.71 to +0.98 on wasp_full; the first implementation demanded $\theta > 0$ and refused 1188 of 1188 candidates); the one-mode model does not reach back into early passes (the ratio drifts 0.77, 0.47, 0.31, 0.22, 0.17, 0.14, 0.11 over one wasp_full step), hence a REACH guard, take the limit only where the modelled remaining error is already inside the tolerance ball (reach scan 6.61 / 6.15 / 6.36 / 7.81 / 7.74 / 7.75 at 1/2/3/10/30/100, control 6.96); and undoing a mediocre candidate costs more than it saves (7.29 strict against 6.94 lenient, control 6.99 on mol_metals). MEASURED, sweeps control -> on: mol_base_handoff 1500 10780 -> 10102 (**-6.3 percent**, matched pair one binary 285.4 -> 268.0 s), mol_carrier 300 -2.3, mol_metals 400 -0.75, wasp_full 300 -0.63, hydrostatic_column 0; state movement 8.3e-8 to 5.0e-7 relative, log10 Mdot unchanged. **The brief's estimate of three passes of seven is NOT reachable, and the reason is the item's real finding:** the stopping test measures the INCREMENT, and the error is $| \theta / (1 - \theta) |$ times it, one to two decades smaller, so the loop is one to two decades stricter than its tolerances

say and spends 1.0 to 1.6 passes on that; the extrapolation's ceiling is the last factor $1/\text{ltheta}$. Redefining the test on the estimated error is a change of what the loop returns and is put to the user as decision item 8 of docs/To_be_determined_by_user_20260906.md (recommendation (b)). Tests: coupled_source_step RED 2 rows against the control, GREEN 39/0 (extrapolation_lowers_the_pass_count 6.09 against 6.96 at 300 steps; refused_extrapolation_costs_nothing identical at 6.80; mean_passes_within_bound re-measured 6.80, unchanged); physics_probe 1491/0. Worker-introduced defect fixed inside the item: a ledger write with six integer descriptors for seven items. Recorded, not fixed: write_setup_report.f90 does not echo the switch; n_csm_x_kept can only count candidates the stopping test did not meet and reads 0 everywhere (the number that matters, met on the very next pass, is printed beside it); the csm_max_pass comment block is now some 60 lines of measured tables and belongs in a docs/ memo. Advisor: the measured tables and the item history that had grown to some 230 comment lines around the coupled-step constants are moved to docs/coupled_source_loop_cost.md (sections 1-7, decision item 8 linked); the code site keeps each constant's conclusion and a section pointer. The setup report does not echo environment switches (none of them, by design), so the switch is reported by the run counter line alone. Advisor verification: coupled_source_step 39/0 and physics_probe 1491/0 on the advisor build; default-off byte-identity against the GOLDENS at matrix length (which checks B5e, HYG-4 and COST6-off together): mol_base_handoff 12000 steps and wasp_full to its own stop, both Hydro_ioniz.txt and Ion_species.txt IDENTICAL to the goldens (both runs end with status 2, "not certified", exactly as the golden runs do).

- **COST7** (decision item 8, option (b), user 2026-09-08; EXHALE_main.f90, coupled_source_step suite, docs/coupled_source_loop_cost.md sections 7-8): **the coupled source loop stops on the ESTIMATED REMAINING ERROR of the pair (T, composition), not on the last increment.** The estimate is $\text{ltheta}/(1 - \text{theta})$ times this pass's increment in each of the two norms the test uses, theta the global Rayleigh-quotient ratio, formed in ONE routine (coupled_pair_geometric_error_estimate) that the COST6 extrapolation now reads too (its reach guard is literally the stopping test at a loosened tolerance; shared objects renamed csm_x_* -> csm_geom_*, env hooks EXHALE_CSM_GEOM_*). Where the sequence gives no estimate (passes 1-2, a guard failing) the increment test stands, and that fallback (EXHALE_CSM_GEOM_REFUSE=1) is byte-identical to the pre-COST7 binary on all 10 written files of wasp_full 300 and all 7 of mol_base_handoff 1500. **The item's finding:** the estimate is an L2 quotient against two max-norm tolerances, and MEASURED with a probe that runs the loop on to $1\text{e-}12$ and rolls back, the unfactored estimate accepts states OUTSIDE csm_comp_tol on 100 of 299 steps of mol_base_handoff (1.8x the tolerance at worst); a factor 2 leaves 1 of 299; **a measured safety factor csm_err_safety = 3 leaves none**, the accepted state's distance to its own fixed point reading $7.4\text{e-}9$ and below (composition) and $2.1\text{e-}9$ and below (temperature) on every one of the 2496 accepted states probed. The price is that almost none of the passes COST6 estimated survive: wasp_full 300 10.09 -> 10.01 passes (sweeps -0.9 percent), mol_base_handoff 1500 7.19 -> 6.96 (-3.1), mol_metals 400 6.99 -> 6.93 (-0.9), **mol_carrier 300 6.01 -> 5.01 (-16.6)**, hydrostatic_column 1.01 -> 1.01 (+1 sweep of 302: on its single three-pass step the estimate EXCEEDS the increment, so the new test is the stricter one there; it is not uniformly looser). log10 Mdot unchanged on every case; state movement $4.3\text{e-}10$ to $1.1\text{e-}7$ relative, at the tolerance level. What the change buys is the MEANING of the two tolerances, the decision's own first reason, not passes. The COST6 extrapolation un-

der the new test is worth nothing (6.79 against 6.82 passes) and stays default off. Tests: coupled_source_step RED 41/3 against the control, GREEN 44/0; mean_passes_within_bound re-measured 6.23 (was 6.80), both at the site; extrapolation_lowers_the_pass_count rewritten as extrapolation_does_not_cost_passes; the fallback-is-one-path row replaces the brief's byte-identity row (a suite cannot hold a golden of the previous binary; the identity is measured in the report). physics_probe 1491/0. Recorded, not fixed: the composition increment of mol_base_handoff has a floor near $5e-9$ (127 of 1500 probes cannot reach $1e-12$ in 59 passes), so csm_comp_tol = $1e-8$ is within a factor of two of what that case can reach at all; COST6's n_csm_x_kept still reads 0. **Every golden of the matrix moves at the 1e-8 level:** gate 3 below.

- **Gate 3 (2026-09-08, 09:29 to 12:00)**, tree after COST7 (with COST6 default off, HYG-4, B5e): shared build remade, **make test all suites green**; make check verdict FAIL against the $1e-3$ relative tolerance, as expected for a change of the stopping test, and every movement examined. log10 Mdot unchanged on all 14 cases that print one (13.35, 13.35, 13.30, 10.57, 10.58, 10.54, 10.57, 10.58, 10.51, 10.56, 7.29, 8.27, 9.61, 7.98). Movement table (largest relative change of any column, new against golden): wasp_full 6.6e-5 (v at the base, cell 136), wasp_he23off 3.8e-4 (v at cell 146), wasp_full_newton 1.1e-10, mol_base_handoff 2.8e-6, mol_metals 2.4e-6, mol_lyman_werner 4.1e-7, mol_diffusion 2.8e-6 (adv file 2.2e-3 at the outer boundary, cell 504), mol_ir_bands 2.4e-6, mol_sec_ion 4.5e-3 (v = 0.091 cm/s at cell 177, a near-zero base velocity, adv 1.5e-3), mol_carrier 5.2e-7, lower_profile 9.3e-6, hp_zero_seed / hp_trace_seed / hp_front identical, roundtrip within. Two cases move by order one and both are pre-existing base artefacts, not COST7's: **oxygen_chemistry** (1000-step snapshot) reads 42x in n(H II) at cell 22 because the GOLDEN carries an odd-even sawtooth there (cells 21, 24, 25, 27 at $7e3-9e3$ against $2e5$ in their neighbors) and the new run is smooth ($1.6e5$ to $3.9e5$ across cells 18-26; rho, p, T within $1e-6$); **hydrostatic_column** reads 0.999 in the adv file's T at cell 350 because the adv post-process gives $T_{adv} = 0.78$ K inside 1.40 Rp in this static column (the original T is 1124 K throughout) and the edge of that region moved by one cell; the non-adv files move by $4.6e-8$ (Ion_species by 0.16 at cell 2 on values of $1e-182$, denormal noise). The adv defect is recorded as an open item. **Goldens to be refreshed by the user** (archive first), then make check re-run.

- **ADV-STATIC** (post_process_adv.f90, new suite adv_static_limit, docs/postprocess_advection_validity.md): **the brief's premise is refuted and the 0.78 K of hydrostatic_column's _adv file is named.** MEASURED: the flow where it sits is Mach 0.23 to 0.82 (the case is a 300-step relaxation snapshot, not a static column: rho v r² spans seven decades); the IONIZATION correction is held at equilibrium in 503 of 503 cells (Ion_species_adv.txt equals Ion_species.txt to the bit) and its Damkohler condition already gives the continuous $v \rightarrow 0$ limit the brief asked for, so nothing there was touched. The number is the exact solution of the equation the _adv ENERGY solve poses: the steady internal-energy equation with the enthalpy flux of the mass-flux divergence DROPPED, $\rho v \frac{de}{dr} - p v \frac{d \ln(\rho)}{dr} = \text{heat} - \text{cool}$, which equals the exact $\dots + h \text{div}(\rho v) = \text{heat} - \text{cool}$ only where $\rho v r^2$ is stationary; with the radiation off it reduces to the discrete adiabat $T_j/T_{j-1} = \rho_{j-1}/[\rho_{j-1} - (\gamma-1)(\rho_{j-1} - \rho_{j-2})]$, and integrating that on the run's density reproduces the reported temperature to $6e-11$ over the 347 cells before the first fallback (1138 K $\rightarrow$ 0.78 K at 1.396 R_p; the

dropped term exceeds the kept ones in 480 of 502 cells). With heat/cool negligible the equation is homogeneous in v , so no $v \rightarrow 0$ condition can change its answer (99.96 percent departure at $|v| \leq 1.4e-3$ c_s). Landed: a validity condition of the energy solve's own, the thermal Damkohler number $Da = t_{\text{cross}} |heat - cool| / u_{\text{th}}$ (new pure thermal_damkohler_number; above one the cell keeps the run's own T), the counterpart of ionization condition (ii) that the bare $v \leq 0$ sign test lacked; the stationary-mass-flux assumption stated at the site with its exact dropped term and MEASURED at run time (largest ratio, where, how many cells exceed one; nothing acts on it yet); the first eval_cool writes a named tcool_in instead of a discard slot; the stale "is not evaluated" sentence of the validity memo rewritten with both tables. Impact: byte-identical everywhere measured (hydrostatic_column full case all four files; wasp_full, mol_base_handoff, lower_profile post-processed from the same golden state by both binaries, all four files; max thermal Da over tested cells 0.951 / 0.534 / 0.055 / $3e-19$, 0 cells above one anywhere). Suite RED/GREEN (rows 1-7 cannot link before; row D measured=silent before), advisor 10/0. Also MEASURED: no matrix case has a stationary mass flux by the cell-to-cell measure ($\rho v r^2$ over outflowing cells spans 1.27x its median in wasp_full, 2.2x in mol_base_handoff, 83x in lower_profile); the du stop is taken on a restricted window and does not bound this. **Follow-up decided by the advisor under the physical-correctness rule: restore the dropped term (item ADV-ENERGY)**, since an energy equation that is exact only for a stationary mass flux is wrong on every snapshot and at the $1e-3$ level on every converged wind; MEASURED worth on the static column 569 K at 1.40 R_p and 121 K at 2.96 R_p in place of 0.78 K and 6.0 K. The alternative, refusing the column where the assumption fails, would rewrite 162 of 502 _adv temperatures of wasp_full (median 5 percent) and is decision 14 (T8.1) of the target-system document, left to the user. Advisor fixed the stale provenance paragraph of the case README.

- **ADV-ENERGY** (T_equation.f90, ion_cell_state.f90, post_process_adv.f90, adv_static_limit +7 rows, docs/postprocess_advection_validity.md): the advection post-process's energy equation is now the EXACT steady internal-energy equation. The enthalpy flux of the mass-flux divergence, $h \operatorname{div}(\rho v)$ with $h = e + p/\rho$, enters T_equation through a new zero-default field of the cell record, teq_cell%div_rho_v, as a separate additive term in both branches (caloric $\mu_m \operatorname{div} \rho v (E(x_{H2}, T) + T)$, monatomic $\gamma \mu_m \operatorname{div} \rho v T$; μ_m, μ_p , coeff untouched; zero field an exact zero, tested to the bit). $\operatorname{div}(\rho v)$ is the mass row's own operator $(A_p F_p - A_m F_m)/dV$ over the control volume between the two points the upwind difference uses. MEASURED corrections to the briefs: T_equation has no caller outside post_process_adv (no marching solve to protect; the bitwise property holds anyway); and ADV-STATIC's 569 K prediction had put back $p \operatorname{div}(v)$ alone, not $e \operatorname{div}(\rho v)$; the whole term gives **1.7e7 K at 1.40 R_p and 1.6e9 K at 2.96 R_p** on the static-column snapshot, the closed form $w \sim \rho^{(\gamma-1)} F^{(-\gamma)}$ with $F = \rho v r^2$ falling 8 percent from cell to cell. Tests RED (nine "not a member" compile errors before) / GREEN (closed-form solution $T \sim r^{(-\gamma)}$ on a diverging column to $6.9e-4$ at 400 cells, first-order convergence $5.59e-3 / 2.79e-3 / 1.39e-3 / 6.95e-4$, the adiabat departing by 2.17); advisor adv_static_limit 17/0, physics_probe 1491/0. **Impact, MEASURED, same golden state post-processed by both binaries:** marching files bitwise in every case; T_adv movement wasp_full_newton 1.4 percent (the only Newton-converged state, He 2³S 1.9 percent), wasp_full (du stop) 16 percent at 1.088 R_p (He 2³S 16 percent), mol_base_handoff snapshot 301

percent, lower_profile snapshot 456 percent (He 2³S 238x), hydrostatic_column 4.2e9 (reaching 2.6e10 K). Restored term against the kept ones: wasp_full 67x at 1.033 R_p in 161 of 502 cells, lower_profile 25x in 299 cells, wasp_full_newton 24x in 2 cells. **The _adv temperature of a state that violates continuity is now unbounded above where it was bounded below, and the transit tools read those columns:** on such a state NO steady post-process has a meaning, and what to output there is decision 14 of the target-system document, now put to the user as decision item 9 with a recommendation. Golden: every _adv file moves by design; refresh deferred by the user to this item's gate.

- **B5g** (steady_newton.f90 +467/-35, steady_species_rows suite, design doc sections 18, 18.5): **a residual evaluation is now a function of its arguments.** Three class-(b) caches survived a probe eval_residual and were read back by the solve: the elemental rows and scales (erow_he/escale_he/erow_tr/escale_tr), the carrier measures (carrier_relnorm_last...), and the carrier arrays of diffusive_photochemistry (col_scale_car, row_scale_car, row_terms, headroom_car, the frozen backgrounds); the Newton ROW SCALE, the acceptance gate, the best-iterate ledger and the choice of the state handed back read them. Fixed on the reader side: eval_residual gained state_is_discarded and a probe holds/puts back that set (reusing the save_/restore_carrier_module_state pair certification.f90 already uses); the loop-top row scale is taken AFTER the iterate's own evaluation; the gates read a new carrier_relnorm_state. The two self-tests also tag their sweeps as candidate sweeps (under the iterate tag they were writing the run's acceptance ledger and ieq_nonroot_streak, whose limit stops the run). **PROVEN:** EXHALE_SPECIES_JAC_TEST=1 and the untested run are the same run, every (JFNK) it line identical over 17 iterations, Ion_species.txt byte-identical (RED: 82 differing lines, first at iteration 6). wasp_full_newton byte-identical (info 0, 6 iterations, CERTIFIED, 13.30). **What the fix uncovered is bigger than the item:** the mol_diffusion element solve that B5e brought to 4.3e-9 in 66 iterations now STAGNATES at 1.957 after 17; the runs agree for five iterations and part on a 3 percent difference of Drow between two evaluations of the SAME state, because the residual's seed dependence (resid_sc_probe) is **0.96 in units of Resid tol** against $||R|| = 1.986$, and tightening the sweep tolerance (1e-12, 1e-14 with pass caps 200, 500) never satisfies the increment test from either seed: **the eliminated composition residual of that state has no fixed point it reaches**, and B5e's convergence was a property of one sequence of evaluations. The change is kept (a row scale must be the row's own terms of the same evaluation). Part 2, the carrier rows: all 18 "no Krylov direction" exits and all 514 refused samples were ONE thing, the H2 carrier of cell 500 at -7.4e-19 (a species unknown on its lower bound), not the element budget (the counter's label named one screen of several; corrected, census added); jv_product now takes the backward difference $[F(Y) - F(Y - \text{eps } v)]/\text{eps}$ only after every forward halving is refused: 49/0/0 accepted/rejected/ refused where 31/18/18, GMRES truncations 18 -> 0, radius 8e-3 -> 2.05; $||R||$ unchanged at 1.873 because the dogleg cuts steps by 2⁻²⁵..2⁻⁴⁹ against the same bound from the trial side, a bound-constrained unknown space (decision, not repair). cert_tol_carrier UNCONFIRMED. Suites (advisor build): steady_species_rows 74/0 (88/0 with the self-test pair logs), certification 29/0, coupled_source_step 44/0, carrier_retry 100/0, species_face_flux green; advisor re-ran wasp_full_newton in full with the B5g binary: Hydro_ioniz.txt and Ion_species.txt IDENTICAL to the gate-3 output, CERTIFIED, log10 Mdot 13.30.

- B5f** (docs/a2_certification_contract_20260906.md section 11.6; no constant moved): the cert_tol_element ladder on mol_diffusion at Resid tol 1e-8 / 1e-9 / 1e-10 gives ONE state: the 1e-8 solve meets its target in 66 iterations (element row 4.633e-11, reproducing B5e digit for digit, so COST7 leaves the stationary solve untouched); the 1e-9 and 1e-10 solves abort after 88 iterations with the radius at 9.3e-23 (9 of 31 refusals on "the ray difference is below its cancellation floor") and hand back the SAME state, bitwise. The row is floor-limited at 4.63e-11 (knee at $||R||$ about 1e-7, 1.5 percent movement over the last factor 26); proposed value by the contract's rule 5e-10, NOT applied (two rungs are not solves that met their target; a decade above one configuration's floor is not a decade above the row's floor, which the operator's header puts between 1e-12 and 1e-5 across configurations; no second configuration exists). **Read together with B5g:** B5f ran on the pre-B5g text, and B5g shows the state it anchored on is reached by one sequence of evaluations only (with the probe caches removed the same solve stagnates at 1.957), so this rung stands as a measurement of that state's row, not as evidence that the solve converges. B5f also MEASURED that the coupled route's convergence depends on eq_sweep_reltol (1e-10 loses the 1e-9 solve at iteration 17), the same fact from the other side. Recorded: the JFNK of a reloaded 1e-8 state starts from $||R|| = 1.226$ because the marching loop takes two CFL steps first (the O(dt) kick, eight decades, seen for the first time from a stationary state); the certification prints no cell-resolved record of the species rows, which blocks any floor attribution.
- ADV-REFUSE** (decision item 9, option (a), user 2026-09-08; post_process_adv.f90 +227/-43, write_output.f90 +94/-10, examples/exhale_io.py, EXHALE_transit.py, README.md, README_HOWTO.md, user manual section 4 (PDF rebuilt, 76 pages), docs/postprocess_advection_validity.md, adv_static_limit +13 rows): **the advection post-process refuses the correction in every cell whose own energy-equation term ratio says the flow is not stationary; that cell keeps the run's temperature and the ionization equilibrium at it; a new integer column adv_status at the end of Hydro_ioniz_adv.txt records the reason row by row** (0 corrected; 1 thermal Damkohler; 2 not stationary; 3 ionization conditions (i)-(iii); 4 cell solve failed; five named module parameters in write_output, the # columns line extended, a # adv_status_counts header line counted from the array written). The criterion is enthalpy_flux_term_ratio (new pure function beside thermal_damkohler_number), the restored term over the LARGER kept term, refused on a STRICT > 1 so a stationary flux with round-off is never refused; formed ONCE on the state handed in over every cell, so the refused set cannot move with the iterate and the converged product does not depend on the iteration history; applied to temperature AND composition (a non-stationary cell has no steady advective ionization correction either). write_output gains an optional adv_status dummy (the four 'eq' calls untouched, Hydro_ioniz.txt keeps seven columns); exhale_io.load_hydro now takes names from the header (seven historical keys kept, Run.adv_status exposed, None for an equilibrium file); EXHALE_transit.py prints how many rows inside $1 \leq b \leq \text{Rib}$ are refused, by reason; EXHALE_plots.py and the four other header parsers adapt unchanged. MEASURED (same golden state, "Do only PP"; marching files bitwise everywhere): hydrostatic_column's corrected temperature falls from 2.6e10 K to **1182.32 K, exactly the run's own maximum, no row above it** (435 before); status counts 0/1/2/3/4: wasp_full_newton 501/0/2/1/0, mol_base_handoff 470/0/33/1/0, wasp_full 205/0/297/2/0 (160 of the 297 beyond conditions (i)-(iii), matching ADV-STATIC's route-2 estimate of 162; the larger total is the reference point, the state handed in

over every cell including the inflow cells), lower_profile 182/0/321/1/0, hydrostatic_column 0/0/481/23/0. Every refused row carries the run's T to the BIT and the equilibrium density of every species exactly (one $1.8\text{e-}16$ reassociation in lower_profile's He I, pre-existing). Movement against the pre-ADV-ENERGY goldens, T_adv: wasp_full_newton 3.3 percent, wasp_full 8.0, mol_base_handoff 64, lower_profile 1.8x, hydrostatic_column 1525x (0.78 -> 1182 K); species: He III 5.4 percent, He 2^3S 29 percent, He 2^3S 110x (at the base of mol_base_handoff), H II 11x (lower_profile base), He III 16 percent. **Transit effect on wasp_full**: 295 of 500 rows inside the impact-parameter range are refused (printed), and the metrics move at $1\text{e-}4$ relative or less (He 10830 red 4.102 percent both, blue 2.843 -> 2.844, FWHM 0.899 Å both; largest line movement O I 1302 +0.55 percent relative), because the refused rows sit in the base under the extended wind. Tests RED (seven "not found in module" errors; measured=7 columns on the old binary) / GREEN; advisor adv_static_limit 30/0, physics_probe 1491/0. Worker also fixed a dangling \ref{sec:retiredbase} in the manual (the four retired base keys named instead). Recorded: two documents describe the _adv validity conditions (manual, HOWTO) and had both drifted; four # columns parsers exist in python; dV_cv division now appears twice unguarded (a zero is a broken grid).

- B5h** (ionization_equilibrium.f90 +68/-2 diagnostics, steady_newton.f90 +211/-2 probe only, steady_selfconsistent_residual assertion 4; NO behavior changed, wasp_full_newton and the element run byte-identical to a control built from the same source with the additions removed): **why one state has two residuals**. MEASURED on mol_diffusion's element configuration with a lockstep two-seed trace of the composition elimination (EXHALE_RESID_SC_TRACE), a nine-seed basin ladder (EXHALE_RESID_SC_BASIN) and a cell acceptance report (EXHALE_IEQ_REPORT_CELL): the amnesty is never used (classes 4 and 6 zero at every pass, every cell class 1 with reaction residual $1\text{e-}25$ against $1\text{e-}6$), the self-field lag is not it (xuv_self_field_passes 1 -> 20, identical to every digit), the He recombination coupling is not it (off, identical to four digits), the stopping test is not it (a fixed count of one pass per call, chained, the difference frozen from pass 2 to 60). **Two things are**. (1) Two separated ROOTS of the network at the same incident field: from a seed displaced five cells the sweep reaches a composition 8.9x apart in x(H2) at the H2 front (cell 214) and 19 percent apart in x(H I) at the ionization front (cell 192), bitwise identical from cell 216 outward, both exact roots (class 1, $1\text{e-}25$). (2) **A marginal direction: the local-equilibrium elimination does not determine the shielded layer's H2 CONTENT**. Retention of a seed perturbation of the H2 partition is 0.77 at the base rising to 0.84 at $r = 1.064$ and falling to 0 outside cell 216, the SAME curve to three digits at perturbations of $1\text{e-}6$ and $1\text{e-}2$, so the sweep's map has eigenvalue 1 in that direction and the seed dependence is a derivative, linear over four decades with gain $3.2\text{e}3$ (the base cell's energy row amplifies the ghost composition by $\sim 4\text{e}3$), reaching Resid_tol at a seed change of $3\text{e-}12$. READ from the network: the fast $\text{H2} \rightarrow \text{H2}^+ \rightarrow \text{H3}^+ \rightarrow \text{H2}$ cycle returns every nucleus, so the H2 row as scaled by its turnover pins the PARTITION and leaves the total to $k15 \ n_{\text{HI}}^2$ against $k12 \ n_{\text{tot}} \ n_{\text{H2}}$, a detailed-balance pair that in the shielded layer is far below the turnover; and pinning it would pin the fully molecular limit (0.9997) where the state carries 0.376: the content is upstream and transport data, exactly why it is imposed at the ghosts. **The repair the code already owns, MEASURED**: with Molecular carrier transport: True ($n(\text{H2})$ a Newton unknown) the same ladder falls from $3.155\text{e-}3$ to **$9.736\text{e-}10$** at delta $1\text{e-}6$ and x(H2) of the front cell is bit-identical from all nine seeds; the residual is a function of its arguments below Resid_tol. What remains at

large seeds (0.98) is the ionization front's two roots, a physics decision (which root a stationary residual may land on, or a statement that the front is unresolved on this grid), reported. The element solve with H2 carried then meets B5g's carrier obstruction ($|R|$ flat at 1.986 over 19 iterations, carrier row 0.91, run stopped by PID at 4 min an iteration). Also MEASURED: eq_sweep_reltol below $\sim 1e-10$ is unreachable by construction, the increment measure $\max|df|/\max(|f|, 1e-12)$ reporting round-off of He I and H2+ fractions $\sim 1e-20$ in the ionized wind (plateau 6e-11 for sixty passes), which is what B5f's and B5g's "never met at $1e-12$, $1e-14$ " was. Tests: assertion 4 residual_is_one_number_for_basin RED 3.155e-3 with H2 eliminated, GREEN 9.736e-10 with H2 carried (the tolerance read from the run's own log); steady_species_rows 86/0 with the self-test pair; B5g's part 1 reproduced as a by-product. A defect of the item's own fixed inside it (a cell-report default of -1 was the lower ghost). **Decision put to the user as item 10:** refuse Coupled carrier solve: True on a molecular configuration without carrier transport, or turn the transport on for it (input_read.f90, not the item's).

- **B5i** (decision item 10, option (a), user 2026-09-08; input_read.f90 +51/-2, grid_and_gates new row coupled_carrier_h2 (+5 assertions), user manual (a Coupled carrier solve entry, which the manual had NOT had; the Molecular carrier transport entry corrected from "inert without oxygen chemistry" to "needs molecular chemistry"), README.md, README_HOWTO.md (new block; the oxygen paragraph's dangling "It" corrected), docs/steady_solver_design.md section 19): **Coupled carrier solve: True with Molecular chemistry: True and Molecular carrier transport: False is refused at startup**, after every key is resolved (so the oxygen default of the transport key is seen: MEASURED, Oxygen chemistry: True plus the coupled key is accepted, the same with the transport stated False is refused), with a message naming what was asked, why there is no root (B5h's numbers) and the remedy key; error stop 1, exit status 1 asserted (2 is the certification's refusal of a complete run). Scope exactly carrier_in_newton .and. thereis_mol .and. .not. carrier_transport, not gated on the Newton key (the direct steady route EXHALE_PTC=1 runs a coupled solve too, and it is the route B5b to B5h used); the atomic coupled solve with an element row is accepted (asserted). No shipped configuration is refused (MEASURED by grep: no *.inp in the tree sets the key; 62 of 81 B5-series scratch inputs would be, as the decision says). Two molecular cases byte-identical before/after at 200 steps. The parse-site comment that claimed "checked below" for a check that did not exist is corrected. A refused run leaves a zero-byte EXHALE_setup.out and no EXHALE_resolved.out, like every other input_read refusal. Tests RED (the eliminated-H2 coupled run completed and printed a Mdot) / GREEN; advisor grid_and_gates 98/0, physics_probe 1491/0. Advisor corrected docs/input_schema.md K15e ("needs K15d" now qualified to the molecular configuration). Recorded: EXHALE_main.f90 opens EXHALE_setup.out (truncating) before input_read, so any refusal erases the previous run's setup report (fix: open it after the input is accepted); the other "checked below" comments of input_read.f90 deserve a sweep against the checks that exist.
- **HYG-SETUP** (advisor, direct; EXHALE_main.f90): the setup report EXHALE_setup.out is opened AFTER input_read returns instead of before it, so a configuration input_read refuses no longer truncates the report a previous accepted run left in the same directory (B5i's finding). input_read writes nothing to that unit. grid_and_gates green on the advisor build; the marching path is untouched.

- HYG-CHECKED** (input_read.f90 comments plus one new report block, docs/input_schema.md, README_HOWTO.md, user manual (PDF 77 pages), grid_and_gates new row carrier_transport_inert +4 assertions): every requirement statement of input_read.f90 and every "Needs K.." / "refused with" clause of docs/input_schema.md checked against the code that enforces it. 40 enforced as stated. ONE not enforced (Molecular carrier transport: True with the molecular network off) is now a startup WARNING, not a refusal, by the code's own convention for a present-and-inert key (every consumer is guarded by thereis_mol, MEASURED inert; eight shipped inputs all carry the network). THREE enforced differently from the statement, one with the sense INVERTED: the schema and the HOWTO said Ionization transport is refused with Solver: Newton and with Coupled carrier solve, where the code REQUIRES the second whenever the first is set (K15e does carry a proton row; what is refused outright is the EXHALE_PTC=1 route); "Solver: PTC refuses it" named a solver value that does not exist; the base-grid resolution check lives in write_setup_report (warns below 10 cells per scale height). SIX stale statements corrected: known_keys also feeds the duplicate-key REFUSAL (MEASURED, all 99 entries used, no hole); the sub-Lyman absorbers are not mutually exclusive; Kzz_base/He_Kzz are NOT inert with He_diffusion off (the carrier transport reads kzz_cell), corrected in four places; Molecular reaction heat default True is guarded by with_molecules (a refusal would refuse every atomic run); the coronal cutoff window 0.08-0.13 superseded by coronal_cutoff_width.md 7.2; four K15b cross-references that pointed at the Stellar LW flux row corrected to K15. MEASURED byte-identical on wasp_full 100 and mol_base_handoff 200 (17 of 17 files). Advisor grid_and_gates 102/0, physics_probe 1491/0. Recorded: all 63 "F" line citations of the schema are stale by 300-1200 lines (marked stale at the convention's definition; recommend deleting the tokens); Coupled carrier solve: True is silently inert in an atomic run without He_diffusion (a report belongs where the row registry is); the user manual has no Ionization transport entry.
- Gate 4 (2026-09-08, 15:39 to 18:45)**, tree after COST7, COST6 (off), B5e, B5g, B5h, HYG-4, ADV-STATIC, ADV-ENERGY, ADV-REFUSE (B5i, HYG-SETUP and HYG-CHECKED landed during the matrix; they touch startup checks and a file-open order only, and were verified on the advisor build with the full grid_and_gates suite, 102/0): shared build remade, **make test all suites green**; make check FAIL against 1e-3 as expected. **The marching files are unchanged from gate 3 to the digit**: Hydro_ioniz.txt and Ion_species.txt carry exactly COST7's movement on every case (identical max-rel values in both logs, the three hp_* cases still identical), so B5g, B5h, the three ADV items, B5i and the hygiene items moved nothing on the marching path, and log10 Mdot is unchanged on all 14 cases. The _adv files move by design: Hydro_ioniz_adv.txt now has eight columns (adv_status, the harness reports SHAPE 8 vs 7), and Ion_species_adv.txt moves by 0.05 (wasp_full_newton, He I at the outer edge) to order one (the molecular cases and the hp_* cases, whose refused base cells now carry the equilibrium densities; the relative measure is element-wise, so a species the old correction had moved off its equilibrium value reads 1). **Goldens to be refreshed by the user, once, for COST7 + ADV-ENERGY + ADV-REFUSE together** (archive first), then make check re-run by the advisor.
- Golden refresh (2026-09-08 19:05, run by the advisor on the user's instruction, the user being away from the machine)**: the previous set archived to backup/regression/golden_pre_cost7_20260908/ (verified iden-

tical to golden/ before the refresh), then `run_check.sh` golden snapshotted the gate-4 outputs of all 16 cases. The refreshed goldens carry COST7 (the stopping test on the estimated error), ADV-ENERGY and ADV-REFUSE (decisions 8b, 9a). Gate 5, a make check on the current tree (B5i, HYG-SETUP, HYG-CHECKED on top, all startup-only), follows.

- **HYG-DOCS** (documents only: `docs/input_schema.md`, user manual (PDF 82 pages), `README_HOWTO.md`): the 91 stale "F" line tokens (77 lines) and the 37 GUI<n> numbers (MEASURED stale by a uniform 13 lines) are gone; a key is located by its label and the header says so; nine Notes cells that held only a citation got real notes READ from the parse sites. The manual gained the missing Ionization transport entry (in the sense HYG-CHECKED corrected) and the four other keys the schema documented and the manual did not (Reconstruction continuation, Atomic rate set, Caloric EOS, Photoelectron heating); MEASURED afterwards, all 97 schema keys have a manual entry. Three stale manual entries corrected (Resid norm is retired; Resid tol's norm is the row scaling of sections 143/145, not $|R|/\max|u|$; Flux spread tol default $2e-5$, not $5e-3$); the validity paragraph's "one of three" made four. The `_adv` validity conditions live in ONE place (manual section 4, `\label{sec:advvalidity}`), the HOWTO points there; the HOWTO names `exhale_io.py` as the loader every new script should use (four # columns parsers exist). Advisor: the same three stale statements in `parameters.f90`'s comments corrected (default $1e-5$; the row scaling; the retired base valve), and the four schema rows whose code spans carried a | (a strict renderer split them) rewritten; table widths verified.
- **B5j** (`steady_newton.f90` +1061/-41 only; `steady_residual.f90` and `diffusive_photochemistry.f90` bit-wise unchanged; `steady_species_rows` +14 rows, `steady_selfconsistent_residual` assertion 5): **the carrier row's lower bound, MEASURED as two mechanisms in sequence.** The outermost carrier row is linear in its own unknown: at the hand-off its root is $+2.1e-11$ with the iterate 4 percent above (positivity-preserving, the Newton right to reduce it); twenty iterations later the row's constant term has changed sign and the root sits at $-5.5e-13$, below the bound, so the row has no admissible root and the unknown is pinned; and at the cell that binds the solve (215) the root is $+2.75e-6$ while the Newton has driven the unknown to exactly zero, eleven decades below it, the row reading -1.00000 of its own scale. A species unknown at zero whose row wants it positive cannot be sampled at all (a two-sided difference leaves the box on both sides). Landed: the carrier mass fraction clamped onto $[0,1]$ in `write_species_rows_into_composition` (the rule the element fractions had); the trial projected onto the faces of the species box with the predicted reduction recomputed for the step taken; Bertsekas (1982) epsilon-active set held for the step and applied to every sampled direction; the probe step cut to the box with the side chosen by the room (subsuming B5g's backward sample); the Cauchy point as the step when the Krylov leg cannot be sampled; a defect of the item's own found and fixed (the accepted trial re-evaluated as $Y + D$ s rather than as the vector evaluated, whose round trip does not close for a projected component: a carrier at $-4.8e-35$). MEASURED on `mol_carrier`, three configurations, none converging: baseline 49 iterations $||R||$ 1.929, 527 refused samples, cut-backs to 2^{-49} ; (a) bound-aware region 63 it, 1.415, 0 refused, 2^{-4} ; (b) carrier as `ln n` (EXHALE_CARRIER_LOG_UNKNOWN=1 (the spelling at B5j; since B5k the log unknown is the default and $=0$ restores the density)) 65 it, **0.9972**, merit 17.6 $\rightarrow$ 0.924, smallest adopted unknown $2.2e-11$. (b) is better by every measure and is NOT the default (it changes the unknown space of a route a shipped key selects): decision

item 11. **Part 2:** the elimination's stopping test is now the change of the three quantities the residual reads the composition FOR (particle count, H2 caloric share, the radiative pair), EXHALE_EQ_MEASURE=species restoring the old one: at eq_sweep_reltol 1e-12 wasp_full_newton reaches it in 33/49 passes where the species measure never does (plateau 2e-11); the element case's remaining plateau (1.85e-11) is the radiative channel at cells 441-459 wandering, that case's own non-convergence (B5h), not the measure's floor. Seed dependence unchanged to four digits (9.736e-10). **wasp_full_newton moves 1.1e-9 relative**, isolated to the new measure by EXHALE_EQ_MEASURE=species reproducing the control byte for byte: the solve converges FURTHER ($||R||$ 8.1e-9 $\rightarrow$ 1.7e-9 in 8 iterations instead of 6, info 0, CERTIFIED, 13.30); below the 0.1 percent rule, the golden is not touched (the harness will read "within"). Suites (advisor build): steady_species_rows 93/0, steady_selfconsistent_residual 3/0, certification 29/0, coupled_source_step 44/0, physics_probe 1491/0; one "per-iteration" phrase in run.sh reworded. Recorded: a carrier row at an active bound is an error to the certification (row 1.00000 at cell 215) and a constrained optimum to the solver, so a bound-constrained stationary state has a multiplier there, not a zero row (decision item 12); 28 of 64 steps of configuration (a) are the Cauchy point alone (Bertsekas' two-metric projection not taken); load_IC refuses a molecular restart written at the input's own He/H by comparing printed digits for equality (7.9307E-02 against 0.0793). Advisor: wasp_full_newton re-run on the matrix path (marching then Newton) with the B5j binary against the refreshed golden: movement 5.9e-9 (Hydro_ioniz.txt) and 9.6e-9 (Ion_species.txt), 13 outer iterations to $||R|| = 4.3e-9$, CERTIFIED, log10 Mdot 13.30; below the 0.1 percent rule, golden untouched.

- **Gate 5 (2026-09-08, 19:03 to 22:02):** make check on the tree as of B5i, HYG-SETUP, HYG-CHECKED (built 19:03, before B5j landed) against the refreshed goldens: **REGRESSION PASS, all 64 files of the 16 cases byte-identical.** The refreshed goldens are therefore the current tree's own outputs, and the three startup-only items moved nothing. B5j's 1.1e-9 movement of wasp_full_newton (the Newton converging further under the new increment measure) is not in this gate's binary and will read "within" at the next one; below the 0.1 percent rule it refreshes nothing.
- **B5k** (decision item 11, option (a), user 2026-09-08; steady_newton.f90 +346/-85, steady_species_rows +137/-6 (114/0), design doc section 20, schema K15e, manual entry, PDF 82 pages): **the molecular carrier densities of the coupled steady solve are carried as $\ln n$ by default** (EXHALE_CARRIER_LOG_UNKNOWN=0 restores the density unknown; the element fractions untouched; no certification change, decision 12 (c)). The floor under $\ln n$ is the transport operator's own admissibility floor, 1e-20 of the carrier's element budget in that cell (carrier_headroom), the fraction carrier_residual already puts under its row scale; MEASURED 5.6e-28 to 4.6e-21 across mol_carrier's column (B5j's log unknown had $\log = -\text{huge}$, unsound: exp underflows to an exact 0). Stated at the site: the column factor n of the Jacobian (both the banded model and the Krylov action probe a RELATIVE change), the region asymmetric in the density ($n \rightarrow n \exp(+\Delta)$), and $\ln 0$. **A real finding at the arming point:** carrier_headroom does not exist when the solve reads its controls (allocated inside carrier_steady_residual, first reached by the opening eval_residual), so the first implementation silently fell back to the density unknown on every coupled run; the opening state is now packed as a density and rewritten as logarithms after that evaluation, sound because the residual is a function of the state and not

of its coordinates. Floor array made rank 2 (cell AND carrier row; two carriers of one cell have different elements). MEASURED, one binary, the two runs one variable apart: mol_carrier density 63 it ||R|| 1.415, 28 Cauchy-only steps, 10 unknowns on a bound, 716 adopted carriers at the floor; **ln n 66 it ||R|| 0.9957, merit 0.960, 0 Cauchy-only, 0 on a bound, 0 backward products, 0 at the floor, smallest adopted carrier 2.2e-11 = 2.5e15 times its floor**; mol_diffusion (H2 carried + element row) density unknown 0 accepted steps of 12, log unknown 50 accepted, ||R|| 1.986 -> 1.982. **Neither converges, and the next obstruction is named and is NOT the density unknown's**: the log unknown's steps are model-accurate (ratio 0.39-1.09) and SHORT (||s|| 0.09-0.17 against a radius of 2.5 sitting at its ceiling for 30 of 68 iterations), cut by 3-5 halvings before the trial is describable, and the refusing screen is the ELEMENT BUDGET (157 samples, against 0 for a negative carrier): the carrier CEILING, which species_unknowns_outside_their_bounds deliberately does not treat as a face of the box. Bertsekas' two-metric projection is not what the log unknown needs (it holds no bound). Gates: wasp_full_newton byte-identical twice (full case control vs new, and the reload identical to B5j's binary including every JFNK line); marching bitwise on mol_carrier 200 steps. New rows: default is ln_n, switch reaches density, the floor identical under both unknowns, carrier_is_above_its_own_floor (RED on the density unknown: 0, 716 at the floor). Advisor: steady_species_rows 93/0, steady_selfconsistent_residual 3/0, certification 29/0, physics_probe 1491/0; the two documents that named the old =1 spelling annotated. Recorded: gm_m = 40 is a literal at the call site in EXHALE_main.f90 and the log-unknown solve sits at it in 58 of 66 iterations (unmeasured whether raising it helps, the cycle reports iterations used, not the residual reached); carrier_headroom returns huge as an availability sentinel. **Next item B5l**: the element budget as a face of the unknown box.

- **B5l** (steady_newton.f90 +392/-82; diffusive_photochemistry.f90 +51/-6, the headroom accessor only; steady_species_rows suite +161 and +108; docs/steady_solver_design.md section 21; certification.f90, EXHALE_main.f90, steady_residual.f90, ionization_equilibrium.f90, input_read.f90 untouched): **the element budget is a face of the unknown box and the projection treats it exactly**. freeze_species_unknown_box is now the one place the admissible set of a species unknown is written down (one lower and one upper bound per unknown of the flat vector, formed once per outer iteration from the iterate and from the element budget the carrier operator froze at that iterate); the three routines that had each computed their own bounds (species_unknowns_outside_their_bounds, fix_active_species_bounds, largest_step_inside_the_species_box) read it. **A drift between the three was found and fixed on the way**: the projection had read the carrier ceiling from the trial's own density slot while the other two read it from the iterate, so the set a trial was written onto and the set the Krylov cycle was sampled under were not the same set. Faces of a carrier unknown: below, zero (carrier_log_floor in ln n); above, the cell density and the element budget (carrier_headroom, for H2 half the hydrogen the carriers may take), the tighter of the two. The budget is frozen for the step, not evaluated at the trial, which is the carrier operator's own choice (headroom_car assigned under weno_mode == 1, the one evaluation that IS the iterate; refreshing it per trial was tried there and is worse), so box and screen describe one state. The budget face is raised to the iterate where the iterate stands outside it (counted, n_budget_face_at_the_iterate; 0 on both coupled cases with the default step control); it is stored one floating-point step inside the budget (budget_face_inside = 1 - 8

eps, representability in $\ln n$, not a margin). An unknown whose box is a point (zero budget) is held without asking the gradient. EXHALE_SPECIES_BUDGET_FACE=0 restores the entry text's box (measurement switch, default on) [retired by N4b: the budget became a constraint row, EXHALE_ELEMENT_CONSTRAINT_ROWS=0 is the measurement switch now]. pgmres reports the relative residual it REACHED against the tolerance asked; EXHALE_GM_M overrides gm_m (the caller's 40 stays the default and the literal at EXHALE_main.f90 line 5343 is untouched). **MEASURED on mol_carrier (coupled, $\ln n$), face off vs on: 66 vs 52 outer iterations, both ended by the stall detector; ||R|| handed back 0.9957 vs 1.174; trials halved 3 or more times 38 of 68 vs 0 of 53; residual samples refused by the element budget 157 vs 0; trials written onto a face 0 vs 29; adopted carriers above their own budget 0 vs 0 (largest ratio 0.9886 / 0.9878); iterations ending "the model promises no reduction" 0 vs 19; worst Krylov relative residual reached 0.938 vs 0.569 where 0.1 was asked; log10 Mdot 10.63 both.** The halvings and refusals against the budget are gone exactly; ||R|| does not improve; the limit moved to a named place, the Krylov leg of the dogleg. **The Krylov budget is not the limit:** gm_m 40 / 80 / 160 give ||R|| 1.174 / 1.103 / 1.271, worst relative residual reached 0.569 / 0.630 / 0.614, ||s|| of the last accepted steps 0.04-0.24 in all three, log10 Mdot 10.63-10.64; the cycle spends every product and stops at the same 0.6 of its right-hand side, so it stagnates on this operator (preconditioner or operator, not subspace). On mol_diffusion (H2 carried) the face does the same to the step control (halved 6-11 times in 50 of 50 vs 0 of 41; budget refusals 388 vs 0) and the solve stalls where it always has (H2 front, cell 212, element row at 1.000 of its scale; ||R|| 1.982 vs 1.943; log10 Mdot 7.97 both), with the probe showing instead 419 backward-side products, 26 unknowns held on a bound, 12 Cauchy-only steps and 18 truncated cycles. A defect in the item's own first measurement was found and fixed (the adopted carrier was recorded against the PREVIOUS iterate's budget, which had reported 67 adopted carriers above budget, worst 1.058; the call now follows the iterate's own residual evaluation, arithmetic unchanged iteration for iteration). **Gates:** wasp_full_newton bitwise against the control binary (Ion_species.txt bitwise, Hydro_ioniz.txt identical but for the provenance line, every (JFNK) line identical, ||R|| = 4.299E-09, log10 Mdot 13.30, 13 iterations); marching bitwise on mol_carrier 200 steps; EXHALE_SPECIES_BUDGET_FACE=0 reproduces the entry-text binary exactly; steady_species_rows 118/0 with the new rows RED without the face and GREEN with it; physics_probe PASSED. **Advisor:** the five diffs match the tree against the worker's entry snapshots; forbidden-word scan clean; the current tree builds warning-free (build_lwv); the worker's wasp_full_newton output against the current golden moves 5.87e-9 / 9.58e-9, exactly the B5j movement already on record, so B5l itself moves nothing on the matrix path. No golden refreshed. **Recorded, not repaired (outside this item's files):** carrier_element_headroom states a box side per carrier but no joint bound for carriers sharing an element (the physical constraint $2 n(\text{H}_2) + n(\text{OH}) + 2 n(\text{H}_2\text{O}) + n(\text{H}^+) \leq n\text{H}_{\text{avail}}$ is a hyperplane the code does not state), and OH/H2O hold hydrogen nuclei their own headroom does not mention, so a state can pass every headroom and still leave the write-back's hydrogen remainder negative (the branch that zeroes H I, H II, H2+, H3+); neither reaches the {H2} carrier set both measured cases use. Also: the epsilon-active band (one finite-difference step) never engages on mol_carrier with the face (29 trials projected, 0 held; no switch, a wider band unmeasured); largest_step_inside_the_species_box returns zero for the whole direction when one component has no room (18 truncated cycles on mol_diffusion); note_adopted_species_unknowns is

not called for a converged state (pre-existing, cannot affect a stall). **Next obstruction on the carrier-row solve: the Krylov leg (banded preconditioner or operator), not bounds and not subspace.** Per the user's instruction of 2026-09-09 ("leave running things alone, stop code modification for now") no further item is started; B5m (the atomic element-row solve with no Krylov direction) stays planned.

- **Preparation of PLAN_20260909_rev1 (items N-1 and N17; advisor, documentation and file moves only, no source change; user's go-ahead 2026-09-09 with the instruction not to rerun regression checks after simple changes).** docs/PLAN_20260909.md was written from the review of the issues inventory (docs/ISSUES_20260909_review.md), reviewed (docs/PLAN_20260909_review.md) and revised to docs/PLAN_20260909_rev1.md, the plan of record (items N-1 to N17, stages A to E, decisions 13 to 19). Before writing rev 1 the advisor re-read the review's new source claims: the sampled integration-error estimate is printed and dropped and no production caller issues `as_reject_int_error` (EXHALE_main.f90 2806-2821); after a retry inside the full pass the half passes start from $0.5 \cdot dt_step$ and the clock advances by the saved `dt_step` (1706-1734, 2797-2799, 2820, 2836); the audit harness asserts the OLD failure (`if (ieee_is_finite(x(1))) error stop 4`); `use_tr = (nspec_row > 0)` (steady_newton.f90 5763) so `wasp_full_newton` never enters the trust region, which corrects a statement of the first plan. All confirmed. **N-1:** the B5m reproducer moved from the session scratchpad into the tree as `backup/regression/atomic_elem_newton/` (input, metals, setup and resolved reports, `run.log`, the two handed-back state files, `run.sh`, README; 6.8 MB, of which `run.log` 6.2 MB; not in DEFAULT_CASES); the common worker brief became `docs/worker_rules.md` with the session paths generalized and three rules added (one owner per source file per increment; affected-path validation only; named outcomes in reports); the review harness `docs/audit_20260905/run_issues_review_20260909.sh` was run once on the current tree, its output kept as `docs/audit_20260905/run_issues_review_20260909.out` (zero-radius step norm 0; Cauchy merit 0.5 to 40.5, line minimizer 0; identity GMRES 1; zero-operator GMRES non-finite with reported residual 0; enthalpy ratio 0.5 on a nonzero mass divergence; He/H pair 8.83e-5; hydrogen inventory 2 at the individual ceilings; outer-shell chord 11.83; "defects reproduced, not repaired"), the harness itself untouched. **N17:** ISSUES_20260909.md corrected per rev 1 section 3 (seed retention no longer stated as an eigenvalue of one; B5f marked a withdrawn, uncertified reference; 3.5 scoped to the present fixed-point map; the restart row now states the `heh_dev > 1e-6` branch and the 8.83e-5 discrepancy; `carrier_headroom_known` noted; `hydro_row_max` and `use_tr` bullets; the `_adv` statement in section 5 made conditional; new section 3.8 on the physical step never being refused for its integration error). `code_status_20260905.md`: the Huang et al. 2023 shell-opacity note added to the photoionization row; four pre-existing spelling/forbidden-phrase slips fixed on the way (per-cent to percent twice, two per-molecule phrases reworded). Verification: forbidden-word scans of the four documents; the reproducer's `run.sh` resolves its default binary path; nothing built or rerun.
- **N16a** (PLAN_20260909_rev1 Stage A, the physical step can be refused and the clock is right; EXHALE_main.f90, attempted_step.f90, `src/tests/attempted_step/{run.sh,attempted_step_tests.f90}`; user 2026-09-09 "N16a 병행"): **the sampled step-doubling comparison is one transaction over one macro-interval H:** the full pass covers H, the two half passes $0.5 H$ each and end at $t + H$ by construction (pass 3 no longer inherits whatever

dt held); a refusal inside any pass, a finite estimate above one, or a nonfinite estimate rejects the WHOLE macrostep (as_reject_int_error now issued by the production caller), restores as_entry, shortens H by the controller and recomputes both trajectories; exhaustion of the transaction's retry budget (n_macro_retry, cap 8) is a named failure that stops with status 2, names the pass and prints the checksum of the restored state (stop_on_exhausted_retry_budget, one report for both exits). The clock, the accepted count and the physical accumulations move only at the adoption boundary below the macrostep loop. **The order of the complete split update is MEASURED as $p = 1$:** at one fixed state the estimate scales as H^2 over five decades ($p + 1 = 1.964, 1.993, 1.9985, 1.9997, 1.9999, 2.0004, 2.0000$ for H from $2e-2$ s to $2.6e-7$ s; the coarsest $1e-1$ s is outside that regime and excluded), so attempted_step_error_estimate now returns the retained full step's error $\Delta/(1 - 2^{-p}) = 2 \Delta$ (err_order_p, err_full_step_factor, convention at the declaration) [superseded by N16d/N16e: $p = 1$ was the base-face leak; the conservative and electron/T rows are $p = 3$, factor $8/7$, the species rows $p = 1$] and the reduction is $0.9 \cdot e^{-(1/(p+1))}$. The brief's first procedure (one interval by 1, 2, 4, 8 prescribed steps, ratios of the output rows) was INCONCLUSIVE, p 0.4 to 4.0, because the third difference ($2.2e-8$ relative) sits at the coupled source step's fixed-point tolerance: the inner error is not below the temporal error there (review F3's condition, not yet enforced). Test hooks, default off: EXHALE_AS_ERR_INJECT[_COUNT], EXHALE_AS_REJECT_IN_PASS, EXHALE_AS_FIXED_DT (seconds; in initialization mode with local stepping it flattens the local steps, stated at the hook). EXHALE_STEP_CLOCK=1 now also prints an err-pass: line per pass and state= (checkpoint checksum) per accepted step. **Tests:** suite RED on the entry binary 117/16 (the sixteen new rows: passes cover one interval, forced error rejects, clock unchanged, shorter retry accepted, refusal aimed at pass 1/2/3 keeps the endpoint and names the pass, exhaustion in each pass and on the error is named and restores the accepted state by checksum), GREEN on the final binary 133/0; advisor reran the suite with the worker's final binary (md5 5ea530d3251a8b8ce77e72415ddef0ab): 133/0. **Runs:** mol_carrier marching 200 steps byte-identical apart from the provenance line (initialization mode never sets n_err_passes = 3; the guard lines quoted in the report); the suite's case in physical mode, 60 steps: identical through step 19, at step 20 $e = 1.215$ (raw 0.607, the number the old code printed and dropped) rejects, H 1.118 $\rightarrow$ 0.913 (e 1.77) $\rightarrow$ 0.618 s (e 0.89) accepted, t_{phys} 66.40 $\rightarrow$ 65.92 s; two rejections in 60 steps, all by the integration error. Not run: the matrix, wasp_full_newton, run_fcheck.sh (out of the affected path; the module compiles clean under the suite's -fcheck=all). No manual entry (the manual carries no run-mode or clock text; nothing in it became false). **Advisor:** file scope as briefed; six British per cent (two new, four pre-existing) changed to percent. **Reported, not fixed (outside the item):** (1) the accepted state of a retried macrostep depends on which pass was rejected, up to $9.1e-5$ relative in the base cell's velocity ($5e-6$ elsewhere), with the checkpoint round trip exact, so some state read by the coupled source step is neither saved nor rebuilt (seeds or caches of ionization_equilibrium, constrained_chemical_equilibrium, diffusive_photochemistry): a rejected attempt is not yet without consequence; (2) one interval of the order sequence (H $6.48e-5$ s from the 0.405 controller reduction) sits 2.3x off the H^2 law the other seven follow, unexplained, for N16b; (3) $\text{err_rtol} = 1e-3$, $\text{err_atol} = 1e-12$ are flat, unvalidated and cover no species (N16b); the gate now shortens physical-mode steps by about two at the sampled steps on this case.

- N0, N3, N1, N2 (PLAN_20260909_rev1 Stage A, one worker, four diffs; steady_newton.f90 +738/-143

over the four, new suite `src/tests/krylov_and_dogleg/`; no other file). **N0**: 14 rows calling the PRODUCTION `pgmres`, `dogleg_step` and the factored step-length and predicted-drop code (a test-operator hook in `pgmres` in the style of `carrier_headroom_set_for_test`); RED on the entry text 10/7, among them `zero_operator_step_is_finite = F`, `approximate_gradient_length_decreases_the_model = 40.5`. **N3**: seven named linear outcomes (`gm_tolerance_reached`, `gm_subspace_exhausted`, `gm_no_direction_sampled`, `gm_jacobian_action_unusable`, `gm_preconditioner_failed`, `gm_reduced_system_singular`, `gm_nonfinite`); a zero right-hand side returns before any factorization; hypot rotations; rank tested before the triangular division and the leading nonsingular block solved; LAPACK status and finiteness checked after each preconditioner apply and on the returned step; the TRUE scaled residual verified with one product at a breakdown; the cycle exits at a breakdown. Reorthogonalization not added: worst max $|V_i \cdot V_j|$ MEASURED $6.9e-13$ on the carrier fixture. **N1**: absent elements (reservoir abundance zero, the `melem_ab <= 0` test set `energy_vectors` and `load_IC` use) are no longer registered as rows; `initial_trust_region_radius` with named statuses, holding the active bounds before its own cycle; a positive floor `tr_delta_min = 1e-6 sqrt(neq)` in scaled coordinates (257x and 1157x below the first radius the two fixtures take); `tr_dmax` follows the radius; "no feasible descent" is `info = 3`. **N2**: the length along the direction used, $\tau = (r \cdot A g) / ||A g||^2$, the point named an approximate descent point, the paragraph defending the old length by B5e removed, a dropped Cauchy leg with a Krylov residual ≥ 1 exits as `tr_no_direction_reduces_model`. Suite after diff 4: 31/0, also under `-fcheck=bounds,do,mem`.

Root cause of B5m (R1's mechanism, its cause): the 1500 unknowns on a bound of the atomic element reload were the three elements `metals.inp` does not carry (Si, K, S x 500 cells), registered with an identity row and sitting exactly on 0, so the fraction-to-the-boundary rule zeroed ANY direction with a component there and the first cycle took 0 products; the radius became 0 and stayed so. MEASURED after (worker and advisor agree): unknowns on a bound 1500 $\rightarrow$ 0; initial radius 0 $\rightarrow$ $1.909e-2$ from a Krylov step that reached tolerance (3 products, $6.1e-2$ against 0.1); first step accepted, ratio 0.995; $||R||$ 1.888 constant $\rightarrow$ $8.3e-2$ at iteration 50, then an oscillation between $6.9e-2$ and $1.1e-1$ with the merit creeping $7.29e-3 \rightarrow 7.23e-3$ over 258 iterations (advisor run stopped by its 20-minute cap; 178 accepted, 80 refused; every cycle spends all 40 products). Not converged: the next obstruction is the model quality of the element rows near $||R||$ $1e-1$. **Carrier reload** (control reproduces B5l's run exactly): after, 54/1 accepted, $||R||$ handed back $1.174 \rightarrow 1.151$, worst achieved Krylov residual $0.569 \rightarrow 0.575$, radius range unchanged, $\log_{10} Mdot$ 10.63 both, outcomes now named (8 tolerance reached, 47 subspace exhausted). **The concurrent N16a edits are inert on every fixture** (paired controls bitwise). **wasp_full_newton** (once, final binary): not bitwise, largest relative movement $2.1e-10$ (`Ion_species.txt`, Ca I) and $8.8e-11$ (`Hydro_ioniz.txt`), `info = 0`, $\log_{10} Mdot$ 13.30; by code path the hypot rotation of N3 (`use_tr` is false there); below the 0.1% rule, identical, no refresh. `lower_profile` (the only default case with `He_metal_diffusion` and a partial reservoir) bitwise to its golden in both builds (it stops at its marching cap). Other suites (worker): `steady_species_rows` 106/0, certification 29/0, `steady_selfconsistent_residual` 3/0; `physics_probe` does not include `steady_newton` (`source_closure.py`). **Reported, not fixed**: `largest_step_inside_the_species_box` still annihilates a whole direction when any unheld unknown has no room on the side the direction points (an element

physically zero in one cell can trigger it; the outcome is now named and the radius survives, the direction is lost): an active-set rule decision, for N4b. Two dead locals removed from pgmres.

- **Stage A gate (advisor, 2026-09-09).** Tree after N16a, N0, N3, N1, N2. make OBJDIR=build_lwv EXE=EXHALE_lwv.x warning-free, md5 dd233d6a (identical to the worker's final binary, so the whole tree is what both workers measured). Suites on that build: krylov_and_dogleg 31/0, steady_species_rows 106/0, steady_selfconsistent_residual 3/0, certification 29/0, attempted_step 133/0 (with the N16a binary, md5 5ea530d3, same source). N7's first two iterations on the atomic reload logged above with named outcomes. Not run, by the affected-path rule: the matrix, physics_probe, run_fcheck.sh; the shared build/ and EXHALE.x remain behind the sources. No golden refreshed. Gate PASSED; Stage B (N4a, N5, N8a) and the N7 diagnostic list are next.
- **N13** (PLAN_20260909_rev1 Stage D, transit refusal census; EXHALE_transit.py, exhale_transit_lib.py, new Python suite src/tests/transit_census/; no Fortran, no output format change, so decision 17 is untouched): the census mask $(r \geq 1) \ \& \ (r \leq \text{Rib})$ was geometrically wrong (the ray integral uses every shell with $|r| \geq b$); one function chord_shell_indices now serves the census, resonance_depth and resonance_spectrum, and the census reports, line by line at eight impact parameters and at its maximum, the share of the line-center optical depth carried by rows whose correction was refused (refused_line_center_tau_share, the trapezoid rule rewritten as a sum over rows so that shares of disjoint subsets add to one), labeled a CONTRIBUTION, not an uncertainty. MEASURED: every tpm_*.txt of wasp_full (10) and mol_base_handoff (5) bitwise before/after; the defect is latent when Rib = $r[-1]$ (both cases as they stand) and real when the domain passes the stellar radius (wasp_full read at EXHALE_TRANSIT_RSTAR_RSUN=0.30: 419 -> 500 sampled rows, 81 above the cap); on wasp_full the rays near $b = 1$ take 99 percent of their Ly-alpha line-center depth from uncorrected rows. Suite RED on the pre-change copies (11 rows, the mask present and no shared selection), GREEN 11/0 on the tree; advisor rerun 11/0; +0.3 s on the whole script. Advisor: one per-line in the library header reworded. Reported, not fixed: the wavelength-resolved share and the file metadata wait for decision 17; np.loadtxt(..., max_rows=1) at line 896 warns on every run and duplicates the column count of line 325.
- **N12** (PLAN_20260909_rev1 Stage D, the upstream caloric energy at the upstream composition; T_equation.f90, post_process_adv.f90, ion_cell_state.f90 (two fields x_h2_up, e_up), src/tests/adv_static_limit/ (new driver upstream_caloric_energy.f90, rows U0-U4)): the upwind term of the _adv energy residual now receives the upstream cell's SPECIFIC internal energy $e_{\text{up}} = E(x_{\text{H2}}, \text{up}, T_{\text{up}})/\mu_{\text{up}}$ formed by the caller, and the caloric branch is taken when EITHER side carries H2 (an atomic cell below a molecular neighbor no longer loses the incoming rovibrational energy); the monatomic branch is verbatim when both sides are atomic. **A second, larger defect found and fixed in the same path:** x_h2_cell was assigned only inside the $k == 1$ stationarity loop (old 650) and read by the temperature loop for EVERY cell (old 1316), so the whole molecular layer was solved at the H2 fraction of the outermost ghost ($1.5\text{e-}7$ on mol_base_handoff, $2.3\text{e-}5$ on mol_carrier, against 0.84 at the base); advisor confirmed both lines in the pre-change source. The stationarity diagnostic's lower energy e_lo is likewise evaluated at the

upstream composition (the ratio's definition, threshold and operator unchanged). Energy convention unchanged: the caloric EOS carries the bound rovibrational ladder only, the 4.478 eV bond lives in the formation reservoir (molecular_reaction_heat.f90), no term is counted twice. **Tests** RED/GREEN (MEASURED): isothermal H2 gradient residual 1.508 -> 0; atomic cell below molecular upstream 1.006 -> 0; dissociation-layer march against the analytic discrete solution, largest departure 0.155 -> 2.4e-11 (RED marched 1000, 892, 782, 615, 560 K against the discrete 1000, 975, 863, 728, 596; GREEN both 1000, 974.76, 943.62, 880.65, 853.52); controls unchanged. Suite 36/0; advisor rerun on build_lwv (md5 8d027f8b, the tree with N12 and N13) 36/0 (a first advisor run reported one no_output FAIL from a relative EXHALE_EXE path, the advisor's mistake, not the suite's). **Impact** (post-process restart Do only PP: True of each case's own state; the before binary is the Stage A binary): wasp_full_adv files bitwise (atomic, both x_h2 zero, monatomic branch verbatim); mol_base_handoff and mol_carrier_adv move in 469 and 467 of 504 rows, T_adv by up to 94.2 percent (218 -> 424 K at r 1.0941) and 103.2 percent (169 -> 343 K), split about 35/45 and 41/46 percent between the stale-composition and the upstream-composition defects; both a warming of the molecular layer (5/2 kT per H2 instead of 3/2); p_adv, cool_adv follow, heat_adv at most 4.8 percent; adv_status changes in one cell of mol_carrier (row 26, 0 -> 2, from the e_lo composition in the ratio); run states (Hydro_ioniz.txt) bitwise in all three. **The molecular_adv goldens of the matrix will move; refresh at the Stage D gate, reported then, not now.** The writer's warning that molecular and oxygen columns are not advection-corrected stays; the product is still not a molecular stationary solution. Advisor: one per-thread phrase in ion_cell_state.f90 reworted; the worker corrected four British spellings in comments. Reported, not fixed: a stale scalar formed in a k == 1 block and read later is a class worth a sweep of the file; post_process_adv.f90 is named by both N11 and N12 (one owner at a time; N11 waits for decision 16).

- **N9** (PLAN_20260909_rev1 Stage D, the restart He/H discrepancy; load_IC.f90 +84/-12, new grid_and_gates row restart_round_trip (9 assertions); write_output.f90 untouched, no key, no tolerance change): **the reported failure was a message defect**: the refusal printed the He/H of the TOP physical cell while the branch tests the largest departure over all physical cells; on the failing state (B5j/scan500_stag, a coupled-carrier JFNK iterate of the hot-Uranus gate) the top cell agreed with the input to 17 digits and cell 199 was off by 3.96 percent, so the message read "7.9307E-02 against 7.9307E-02 ... cannot be rescaled". Review R5's 8.83e-5 is the distance between the INPUT values of two different runs, not a property of any state. Fixed: the comparison records the cell it attained and report_deviating_heh_cell prints cell, radius, file and input ratios at ES16.9, the departure and the threshold (heh_dev_tol = 1e-6, now named, value unchanged); RED/GREEN on a state with helium raised by half at cell 199. **The round trip is clean** (MEASURED on the 12000-step mol_base_handoff state with EXHALE_DUMP_IC=1): r, v, p bitwise; every species column and every inventory (H, He, O, C nuclei, charge, He/H, H2 ratio) within 4.2e-16; T 5.6e-16; the rho column 4.9e-12 because the loader never reads it and rebuilds mass from the species (perturbing the column by 1e-15 changes nothing, bitwise), so 4.9e-12 is the writer state's own mass-closure defect (4.1e-14 on a 40-step state); a second trip 5.0e-16, no drift; no clock advanced. **The producing path moved the ratio, not the loader** (MEASURED): He/H departs from the input by 1.0e-11 and 8.9e-12 after 12000 marching steps with carrier transport off and on, but by 3.9 to 7.9 percent on every coupled-carrier density-unknown JFNK state and 41 to 43 percent on the ln n

states (B5j, B5k): the carrier column is written after the element projection precisely so the Newton's carrier is not rescaled (write_species_rows_into_composition 1439-1469), the sweep reads the nuclei totals from the state (hydrogen_helium_nuclei_density), and with helium diffusion off no row re-imposes He/H. This is the element constraint N4a/N4b are about, seen from the restart side; reported, not touched. Also fixed on the way: parse_labels had no bound on 80-long label arrays (a file with more than 80 columns wrote past them); it now stops with a message (present schema at most 42). Atomic restart path MEASURED unchanged: wasp_full_newton's state reloaded before/after bitwise, no He/H line printed. Advisor: restart_round_trip 9/0 on build_lwv (conda gfortran; the worker's private build used /usr/bin gfortran 13, against the standing toolchain rule, so the advisor's run on the standard toolchain is the one on record). **Reported, not fixed:** the window integrals of the assembled residual are not Lipschitz in the state in the odd-even band above the base (one round trip at $4.9\text{e-}16$ moves the 1.03-1.10 mass window by 16 percent and flips cell residual signs at cells 183-187; one ulp on every species column reproduces it), so a round-trip residual budget must be stated on the row maxima (N5); restart_grid_guard.sh carries a stale header describing a fixed defect; nothing checks the writer's rho column against its species (decision 15's contract); EXHALE_RESIDUAL=1 runs one sweep unconditionally, so a true "no chemistry on load" evaluation has no entry point yet (N10).

- **N7 (PLAN_20260909_rev1 Stage C, the atomic element-row diagnostic after Stage A; steady_newton.f90 +477/-0, diagnostic prints under EXHALE_ELEM_DIAG=1 (iterations 1, 2, 60; =n moves the third), one extra Jacobian product per selected iteration with the refusal statistics held aside; no arithmetic of the step changed, MEASURED: every (JFNK) line identical hook off/on over 2027 lines, across four builds). **Verdict, MEASURED on the reload fixture:** the oscillation at $||R||$ $7\text{e-}2$ to $1.1\text{e-}1$ is NOT a model failure (predicted/actual ratio 0.9946, 1.0001, 1.0003 at iterations 1, 2, 60; the refused iteration's model is right to $3\text{e-}4$), NOT a bound effect (0 unknowns held, 0 samples refused in 293 iterations), and NOT limited by the linear solve although the solve fails its tolerance (40 of 40 products, achieved 0.6685 against 0.1, TRUE residual 0.6685 by one extra product, Arnoldi gap $2.3\text{e-}4$; the dogleg takes $2.1\text{e-}4$ of the Newton leg so that inaccuracy is second order). **Proximate cause: the ray test** (the finite-difference check of the model slope) refuses the step: at the refusing iteration model slope $-9.13\text{e-}9$ against ray slope $-1.48\text{e-}9$ (factor 6.2 against the allowed 4) with $|f2p - f2m| = 3.0\text{e-}12$ only 11x above the floor $\text{eq_sweep_reltol} * |f2| = 2.7\text{e-}13$ (at iteration 1 the same test has a factor 1371 of room and the slopes agree to 0.5 percent); 82 of the 91 refusals in 293 iterations are this test, 3 its sign form, 6 a real over-prediction. Limit cycle read off the radius: $9.78\text{e-}4$ accepted $\rightarrow 1.96\text{e-}3$ accepted $\rightarrow 3.91\text{e-}3$ refused by the ray test $\rightarrow$ cut to $9.78\text{e-}4$, period three; the merit falls $1\text{e-}4$ of itself per accepted step, so Resid tol $1\text{e-}8$ is seven decades away. **Underlying cause: the merit and the gate are different functionals.** At iteration 60 the eight element rows hold 0.437 of the squared merit (He 0.305) and NOTHING of $||R||$ (energy row holds 1.000 of $||R||$, 0.560 of the merit); their own certification measures sit at $1.2\text{e-}3$ to $7.4\text{e-}3$ against $\text{cert_tol_element} = 1\text{e-}8$; accepted steps swing $||R||$ by 40 percent ($7.8\text{e-}2$ to $1.09\text{e-}1$) while the merit stands at $7.291\text{e-}3$ to four digits. At iteration 20 the brief's option (c) is met outright: $||R||$ 1.400 set entirely by the mass row, the energy row 0.810 of the merit, seven of eight elements worst at cells 12 or 17. **The Cauchy leg is worth nothing here:** $||sU||$ $6.6\text{e-}9$ against a radius $3.9\text{e-}3$ ($g = B^T r_0$ from the band, $||A g||$ $1\text{e}4$ times r_0 . $A g$), predicted decrease $1.3\text{e-}9$ against the dogleg's $7.6\text{e-}9$; one product per iteration for $1.7\text{e-}6$ of the**

step. **The band is not the defect** (0 zero, nonfinite or round-off columns, dgbtrf info 0), though it conditions worse along the solve (dgbcon 7.3e-8 -> 9.4e-10, smallest pivot always a trace element of cell 500). **gm_m probe:** unlike the carrier-row solve, the larger subspace changes the achieved residual on this operator (first divergence at iteration 17: 40 of 40 at 0.1035 against 48 of 80 at 0.09999; deep in the stall 0.67 against 0.29) and the solve stalls WORSE (m = 80 plateau $\|R\|$ 0.27 to 0.34, merit 1.57e-2, against 7.7e-2 to 1.1e-1 and 7.3e-3 at m = 40); longer steps lift the ray probe off its floor and the refusal becomes an honest over-prediction, and the solve still goes nowhere. **Proposals, not implemented:** P1 measure the ray test's floor from two evaluations of one state instead of assuming $\text{eq_sweep_reltol} * \|f2\|$; P2 do not let the slope comparison overrule a reduction ratio within [0.998, 1.001] (every ray-size refusal inspected had one; the over-prediction guard stays live); P3 drop or rescale the banded-gradient leg where $\|sU\|$ is below a small fraction of the radius (one product saved); P4 a GMRES restart buys m = 80's reduction at m = 40's memory but does not fix the stall; **P5 the acceptance functional and the merit must be reconciled** (a merit that reads the element rows on their certification scales, or a separate control for them): a design decision, decision item 20 (section below). Also fixed in the same file: the `tr_exit_text/gm_outcome_text` buffers (48 and 64 characters) truncated two 67-character messages; all three are 80 now. **Reported, not fixed:** the worst `j`, `k` on the (JFNK) it line is the merit's largest scaled row while $\|R\|$ on the same line is a certification maximum, and the two need not be the same row (iteration 20: $\|R\|$ is the mass row's, worst is an energy cell); the brief's belief that Solver: Newton 100.0 caps the iterations is wrong (the third word is `newton_du_switch`; the JFNK cap is the literal 500 at `EXHALE_main.f90` 3509 and 3000 on the direct route, no key or hook; runs were capped by wall clock); `atomic_elem_newton/README.md` is right but incomplete (the refused trials have a correct model). Advisor: rebuild warning-free, `krylov_and_dogleg` 31/0 on the tree with N7 and N9.

- **N4a** (PLAN_20260909_rev1 Stage B, one elemental constraint map with three contexts; new `src/modules/lower_atmosphere/element_inventory.f90` (745 lines, one SRC line in the Makefile), `diffusive_photochemistry.f90` (its three element densities, hydrogen budget and five carrier box sides now derived from the map with unchanged arithmetic), `steady_species_rows` suite, `docs/element_inventory_contexts_20260909.md` corrected; `steady_newton.f90` untouched): **the map reports, nothing enforces it yet**, so the Stage B gate may not claim from this item that trial construction, chemistry and certification enforce one constraint (N4b). `nuclei_per_particle` is the one place a stoichiometric coefficient is read (`bsp_nH/nHe/nO/nC`, `mion_elem`); `element_inventory_of_cell` gives the full inventory grouped carried/locked/reserved/closure with the budgets and the shared-constraint breach as a MAGNITUDE (against the budget the sides spend, and against the budget the chemistry closes with, separately); `element_inventory_of_candidate` evaluates a trial's carriers against the frozen element densities; `EXHALE_INVENTORY_REPORT=1` prints it at the operator's entry. **Physical judgment:** the separate box sides are a wrong feasible set by stoichiometry (two carriers of one element spend one budget); a cell at both ceilings breaches by exactly 1.0 of the available hydrogen (MEASURED). **Seven corrections to the advisor's draft**, each checked in the source: `nO_free/nC_free` include the neutral stage while `cbg_nOion` excludes it (the F7 gap as a number: the oxygen box side spends the element total while `carrier_source` closes against the total less the ion stages, so the box admits states whose source rows clamp free O at zero; N4b's feasible set); `nH_avail`

is flooded; H I and He I are the closure, not rows; CO is fixed data of the local solve (its destruction row is the carrier operator's); **the shared sums ARE enforced on the marching path by limit_to_element_budget** (what has no shared statement is the coupled stationary solve, whose eval_residual compares a COUNT of cells); **max(rest, 0) in the write-back CREATES nuclei** (with rest < 0 the carriers keep what they hold and the closure cannot go negative, so the written element total exceeds the handed one by |rest|; only element_census_verify at EXHALE_ELEMENT_ASSERT 1 or 2 sees it, off by default); absent elements are no longer registered (N1). **Tests** 149/0 (advisor on build_lwv: 145/0, the four coupled-log rows unset), RED shown by EXHALE_INVENTORY_SHARED=0 (the three infeasibility rows fail with that setting); all nine molecular goldens classified: worst carrier breach exactly 0.0, tightest hydrogen occupancy 0.96 to 0.99 (oxygen_chemistry occupancy 1.0 is carbon's, 175 of 500 cells with all C in CO). **The N9 finding caught live:** inside a coupled carrier JFNK the He/H departure from the reservoir grows $1.4\text{e-}11 \rightarrow 4.9\text{e-}5$ (third iterate, the first with a Newton $n(\text{H}_2) \rightarrow 1.1\text{e-}1$ (thirteenth) while every carrier budget stays at breach 0.0, so the fix belongs at the write-back of each iterate (the element row or a projection there), not at the end of the solve; the two saved coupled states have carrier breach 0.0 and He/H departures $4.0\text{e-}2$ and $6.7\text{e-}1$. **Marching:** mol_carrier 200 steps byte-identical (two independent before/after pairs); mol_diffusion 12000 steps byte-identical over the 7511 steps both builds had covered at report time (step trace character for character, the step-7001 snapshot files byte-identical apart from provenance); the pair finished later in the session scratch and the advisor compared its end: Hydro_ioniz.txt and Ion_species.txt at 12000 steps IDENTICAL apart from the provenance line, log₁₀ Mdot 10.57 in both builds. Reported, not fixed: element_ratio_gate = $1\text{e-}9$ now exists in two modules because element_census_tolerance() is private. Rule breach reported by the worker: one pkill -f on its own shell pattern (it matched only its own shell; the other workers' runs were verified alive). Advisor: five British percent in the two files fixed; the advisor's earlier pgrep for the worker's binary name missed its renamed scratch copies, a false "finished" the worker corrected.

- **N7b** (N7 proposals P1 and P2, and the JFNK iteration hook; steady_newton.f90 +249/-40, krylov_and_dogleg +12 rows): **P2 opens the cycle; P1 is implemented and MEASURED inactive, and the measurement refutes N7's reading of the refusals.** The ray test's decision is extracted into ray_slope_verdict (public, tested), its floor is $\max(\text{eq_sweep_reftol} * |f_2|, \text{ray_floor_factor} * \text{measured_reproducibility})$ with the reproducibility measured once per outer iteration by two evaluations of the merit at one state (cost one residual evaluation per outer iteration, +1.5 percent on the carrier reload): MEASURED $1.8\text{e-}15$ at a merit of 14.95 and 0 to $4\text{e-}19$ at $2.7\text{e-}5$, four to seven decades BELOW the assumed part ($2.6\text{e-}13$), which therefore governs everywhere; the refused trials' $|f_{2p} - f_{2m}|$ ($1\text{e-}10$ to $3\text{e-}10$) stand two to three decades above the floor, so they are real differences and NOT the probe near its cancellation floor as N7 and ISSUES 3.1 (d) said (corrected there). What they are: at the refusing iterate the ratio is 0.999 while the model slope ($-5.5\text{e-}7$) and the ray slope ($+1.7\text{e-}7$) have OPPOSITE signs; the model slope is set by the banded-gradient leg, which is not in the step ($|s|$ $6.6\text{e-}9$ against a radius $3.9\text{e-}3$) but carries $|A_g|$ $3.1\text{e}3$ into r_0 . As, so the slope test is right that the slope is wrong and the leg is what is wrong (N7 P3). ratio_sound_band = $1\text{e-}2$: within it the ray verdict is formed and printed but not acted on; tr_ratio_below_acceptance untouched. EXHALE_JFNK_MAXIT read once per solve, resolved by jfnk_outer_iteration_cap, printed with its origin.

(base), $4.0\text{e-}3$ (front), 0.82 (wind); H_2^+ , H_3^+ , HeH^+ 3 to 12 decades faster than transport everywhere, so the transported- H_2 / eliminated-ion split stands and no promotion is proposed (conservation modes are removed by the parameterization itself: the neutral stage closes each element). **Rounding floor of the species rows in the judged measure is exactly epsilon** ($2.2\text{e-}16$) against tolerances $1\text{e-}8$, 7.65 decades apart, condition numbers 2 to $1\text{e}3$: compensated summation (the Al-Refaie et al. 2024 reading) is not a lever. Validation: krylov_and_dogleg 43/0, steady_species_rows 145/0, mol_carrier 200 steps bitwise, wasp_full_newton bitwise on all ten files (one new log line, N7b's cap). Advisor: rebuild warning-free, md5 913e6579 identical to the worker's final; krylov_and_dogleg 43/0; residual_determinism 5 PASS / 1 FAIL, the FAIL being the atomic closure-spread row by design (a first advisor run with the wrong variable name EXHALE_EXE tested the stale shared EXHALE.x, the advisor's mistake); two British percent in ionization_equilibrium.f90 fixed, the worker fixed five neighbor. The worker's session was cut once by the Opus limit (09:50 KST) and resumed; nothing was lost. **Reported, not fixed**: a discarded evaluation restores n_eq_sweeps_last, so a caller reads the previous state's pass count; report_accepted_cell_state writes inside the parallel region without a lock; carrier_diffusion_coefficient reads the marching operator's pct_Dco, which the stationary residual never fills (a pure stationary path would read zero; diffusive_photochemistry.f90); the global count shadows the intrinsic.

- **N16b** (PLAN_20260909_rev1 Stage E brought forward, species-aware integration error; attempted_step.f90 +546/-28, EXHALE_main.f90 macrostep code +125/-7, attempted_step suite +289): the step-doubling estimate reads four row classes on their own scales (conservative rows; transported species fractions, the f_sp columns carrier_state builds fc from: H^+ , H_2 , OH, H_2O , CO; element mass fractions under He_diffusion; electron fraction and T through the caloric EOS), the species scale being the species' own value with an absolute floor of element_ratio_gate ($1\text{e-}9$) of the largest density the species can reach in the cell, read from the N4a map's stoichiometry; the flat $1\text{e-}12$ no longer judges any species row (a carrier at $1\text{e-}20$ changed by 1 percent: $e = 20$ on its own scale, $2\text{e-}10$ on the flat floor). **The inner nonlinear error is collected cell by cell** (the coupled pair's safety-factored geometric factor on the cell's own increment, and the carrier substep's row residual) and the estimate carries the interval $e_{\text{lower}} \dots e_{\text{upper}}$ the three inner errors leave it in: reject when above one at the bottom, accept when below one at the top, UNRESOLVED in between (named outcome, counted in the end-of-run block); the brief's fraction rule for each class (1/10) is kept as the report, not as the verdict, because applied literally it called EVERY estimate of a healthy run unresolved (species difference $6\text{e-}17$ five decades below the pair's own error) and would have made F1's rejection inert. **p = 1 confirmed on every class** (atomic case, eight levels over five decades: species $p + 1$ 1.9987, 1.9799; electron/T 1.9943 to 1.9997; conservative 1.9809 to 2.0004; molecular case coupled ladder $H = 5, 2.5, 1.25$ s: species 1.949, 2.299, electron/T 1.912, 1.983), so the retained-step factor 2 holds for all rows. On the molecular case at $H \leq 0.4$ s the species class reports itself unresolved (inner over difference $2.7\text{e}5$): review F3's condition, MEASURED and detected. **The 60-step physical run is unchanged to 17 digits** (t_phys, checksum, both files): the conservative rows still decide it. mol_carrier marching bitwise. **N16a's 9.1e-5 state leak IDENTIFIED** (not fixed, files not owned): Apply_BC.f90 base_face_W and base_face_lower_W (lines 48, 50; written only by Apply_BC, read by Rec_BC in the first Reconstruct of a retaken pass BEFORE the pass's first Apply_BC, so the base face's left state is the

refused attempt's) and `caloric_eos.f90` `caloric_mixture_active`, `nk_per_mass`, `x_h2`, `molecular_cell` (125, 145-147; written only inside `get_species_densities`, which `rebuild_state_from_checkpoint` calls AFTER `U_to_W`, so the restored `p` and `T` use the refused attempt's caloric composition); the two reinforce each other at the base cell, which is the measured signature (v $9.1\text{e-}5$, T $6.8\text{e-}8$, cooling $1.7\text{e-}5$). Row `rejected_pass_leaves_no_state_behind` stays RED at $9.1162\text{e-}5$, labeled with the files and lines. N16a's 2.3x outlier reproduced and bounded (only at controller-formula levels, conservative and charge classes, always the base cell; 0.665 and 0.53 of the H^2 prediction), not explained; two candidates named (the unbounded inner error of the mass/momentum rows through the second half's coupling; the base-face leak). `err_rtol`, `err_atol` unchanged (the CFL step 20 would need $1.21\text{e-}3$; the estimate RISES at the first shortened interval, a second reason not to anchor on this case). Tests 155/1 (the one FAIL by design), RED on the entry text 150/6 (an estimate that was the solvers' own error rejected ten macrosteps there; now 0, named). Advisor: suite rerun with the worker's final binary (md5 7bd041dd) 155/1; four modelled in `EXHALE_main.f90` comments changed to modeled. **Reported, not fixed:** the mass and momentum rows carry no inner error; the carrier substep's number is a row residual, not a solution error (conditioning unmeasured); `NCO_col`, `k_co_diss`, `theta_co_shield` are outside the checkpoint by an unstated exception (read only after their write within a sweep).

- **N16d** (the checkpoint's restore is complete; `EXHALE_main.f90` `rebuild_state_from_checkpoint` 4 code lines, comments in `Apply_BC.f90`, `caloric_eos.f90`, `attempted_step.f90`, the suite row extended to both files): **closed by rebuilding, not saving.** The five leaking quantities (`base_face_W`, `base_face_lower_W`, `caloric_mixture_active`, `nk_per_mass`, `x_h2`, `molecular_cell`) are functions of (`u`, `f_sp`), so the rebuild now runs `rho = u(1,:)`, `get_species_densities`, `Apply_BC`, then `U_to_W` and `comp_T_from_p` (composition first because the pressure map and the boundary read it; boundary second so the face states are the restored state's). Every one of the six call sites follows a restore, so no restore-free path can move. **MEASURED:** `rejected_pass_leaves_no_state_behind` $9.1162\text{e-}5 \rightarrow 0.0$ bitwise on both files; suite 156/0 (advisor rerun with the worker's final binary md5 abdf17f7: 156/0). **The fix changes the measured order of the conservative rows:** the full-versus-two-half difference was 40x the real splitting difference (the leaked base face carried an $O(H)$ error into one face of one RK stage of an $H/2$ step, an $O(H^2)$ contribution) and now falls as H^4 ($p + 1$ 3.9966 to 4.0104 over the ladder), the local-error order of the three-stage Runge-Kutta; the electron/ T rows follow it; the transported species stay H^2 (unchanged to all printed digits). N16a's 2.3x outlier is EXPLAINED: the two controller-formula levels now sit on the H^4 law to 0.1 percent; the leaked base face carried the difference of two intervals whose ratio was not the ladder's. **Consequence not yet fixed (N16e, launched):** `err_order_p = 1` was calibrated on the leak; the conservative and electron/ T estimates carry a factor 2 where $8/7$ is right (inflated 1.75x) and the reduction exponent is $-1/2$ where $-1/4$ is; the species class is genuinely $p = 1$, so the order is a property of each class. `mol_carrier` marching 200 steps bitwise (0 refused attempts, no restore); with one injected refusal the initialization-mode state moves $2.1\text{e-}5$ (v at r 1.0185) and $1.2\text{e-}5$ ($He\ 2^3S$), the fix of the same defect with both channels live (atomic hydrostatic_column had the base-face channel alone). `wasp_full_newton` not run (no restore unless an attempt is refused; whether it has zero refusals was not measured). The reload route does not share the rebuild (checked). Advisor: two Per-cell phrases in `caloric_eos.f90` reworded. **Reported, not fixed:** the same composition-before-pressure ordering stands at

three other sites of EXHALE_main.f90 (1075-1078, 5162-5166, 5312-5326) and upmap_bc_roundtrip measures an identity its comment says is not one (N16e); the added Apply_BC adds one boundary evaluation per restore to the run-total diagnostic counters.

- **N20** (decision 20 a with N7's P3; steady_newton.f90 +314/-84, krylov_and_dogleg +13 rows, steady_species_rows +1 row; certification.f90 unchanged, cert_scale_floor was already public): **one functional, implemented exactly, with a measured consequence that is a decision, not hidden.** merit_row_scale_from_certification is the one denominator a species-row-route row is read through; cell_row_scales builds the hydrodynamic rows through residual_row_scale(k,j,u), the element rows through their certification scale, the carrier rows through the term sum the certification divides by; base cells of element rows keep the unknown's own scale (Dirichlet reservoir); the three-unknown route is untouched by construction ($D_r = D$ when $n_{\text{spec_row}} \leq 0$). MEASURED: on both reloads every element row's largest scaled merit row EQUALS its certification measure at the same cell and the merit's largest scaled row IS $||R||$'s row (before: factor 2.5 apart at iteration 1, 112 at iteration 60). **P3**: the banded-gradient leg is dropped from the step AND its image when shorter than $\text{cauchy_leg_min_fraction} = 1e-3$ of the radius (measured $1.4e-5$ to $8.7e-7$ on both fixtures, so everywhere), so the slope handed to ray_slope_verdict is the step's own; products saved: zero, because tau_c needs A g (a lazy leg would save one per iteration, P-N20-1, reported). A quotient verdict at $\text{ratio} \leq \text{eta_accept}$ is named tr_ratio_below_acceptance. tr_dmax capped at $1e3 \sqrt{\text{neq}}$ ($7.4e4$ on the atomic fixture; binds on neither fixture now, would have bound the $2.7e8$ of N7b). **The consequence**: the carrier reload improves 2.6x (best $||R||$ $1.151 \rightarrow 0.446$, both builds on the stagnation detector, +16 percent evaluations, dbgcon $5.4e-10 \rightarrow 1.9e-11$) while **the atomic element reload no longer solves**: the certification row scales span 11.4 decades ($3.7e-10$.. 92.7) against 8.3 before, dbgcon $7.3e-8 \rightarrow 8.9e-10$ at iteration 1, the Krylov point collapses to $||s||$ $2.5e-15$ inside a radius 19.6 at iteration 11 and the solve aborts on the no-descent watchdog at $||R||$ 1.777 (before: $7.38e-2$ at 150 iterations). A third scaling choice (element/carrier rows on certification scales, hydrodynamic rows on the state scales) does not collapse but is worse than the entry text on both fixtures (0.741, 1.026): the hydrodynamic reconciliation is what helps the carrier reload and kills the atomic one. The merit's row scaling IS the operator's ($r_0 = F/D_{\text{row}}$, $A = D_{\text{row}}^{-1} J D$), so the remaining freedom is the column scaling D and the preconditioner: N8a's, launched with the atomic collapse as its RED. Tests RED on the entry objects (the element-row scale row 4.0 against 1.0; the ratio-naming row), GREEN krylov_and_dogleg 56/0, steady_species_rows 145/0, certification 29/0; wasp_full_newton BITWISE on all ten files (two wall-clock lines). Advisor: rebuild warning-free, steady_newton.o md5 6f3aef50 identical to the worker's, the three suites 56/0, 145/0, 29/0. Two pre-existing defects fixed (a dead local; [diag 7] printed at cell 0 for the carrier). **Advisor decision**: (a) stays in the tree; the atomic reload's collapse is a conditioning problem of the scaled operator and is the first deliverable of N8a, not a reason to give the step control a functional other than the gate's; the state of the atomic reload is reported as WORSE than before N20 until N8a lands. Reported, not fixed: EXHALE_main.f90 1182 should pass u to cell_row_scales; docs/p54_base_layer_mass_flux.md 10.4 now has a second measurement behind it.

- **N16e** (the order of each row class is its own integrator's; attempted_step.f90 +106/-50, EXHALE_main.f90

+60/-21, attempted_step suite +75/-15): err_order_p(n_err_class) = [3, 1, 1, 3] (hydro, species, element, charge/T) with the measured ladders at the declaration (conservative 3.9966 to 4.0104; electron/T 3.9962 to 3.9867; species 1.9987, 1.9799; the element class is assigned the species' order by construction, said so); err_full_step_factor is a class's own (8/7 for the RK3-carried rows, 2 for the first-order split ones), the estimate records the deciding class (the one whose lower bound reached the gate) and attempted_step_reduced_dt uses that class's $-1/(p+1)$. MEASURED: the conservative and electron/T estimates fall by exactly 4/7 (inflation 1.7500004 removed), the 60-step physical run's trajectory does not move (N16d had already removed the step-20 rejection, so the brief's expected "1.21 rejected -> 0.69 accepted" belonged to N16b's entry text, not to this item's; stated); on the forced-error ladder the controller with exponent -1/4 asks for 0.846 and 0.226 of the interval where -1/2 asked 0.601 and hit the 0.2 floor, so the floor no longer does the sizing. Stale sites: EXHALE_RESIDUAL=1 and physical_handoff_check NOT reordered, the composition cannot have moved there (init and equilibrate_loaded_composition both end with get_species_densities), the invariant is now stated at both; update_map_begin_step's install branch reordered (composition before U_to_W; EXHALE_UPDATE_MAP on mol_carrier moves 0.0); upmap_bc_roundtrip renamed upmap_bc_interior, required exactly zero (Apply_BC writes only the ghosts), MEASURED 0. Tests 163/0 (156 before; the unit rows cannot be run RED against a scalar err_order_p, the RED is the entry suite's own contradiction, $p = 1$ asserted and $p + 1 = 4$ measured in one run; the whole-binary row 0 of 2 -> 2 of 2). mol_carrier marching bitwise. Advisor: suite rerun with the worker's final binary (md5 0ae8d8d9) 163/0. **Reported, not fixed:** update_map_begin_step's install branch owes the next Reconstruct an Apply_BC (same family as N16d, a diagnostic route); err_atol = $1e-12$ still floors the conservative rows on a code-unit density; the mass/momentum inner error is still unbounded (the deciding row's interval is a point); no fixture has the species class deciding a step, so the exponent switch is covered by unit rows only.

- **N8a** (PLAN_20260909_rev1 Stage B, scales, boundary equations, derivatives and the inner budget, with N20's atomic collapse as its RED; steady_newton.f90, krylov_and_dogleg +17 rows, steady_species_rows +2 rows; certification.f90 and steady_residual.f90 unchanged, no tolerance moved): **the collapse was one exactly empty Jacobian column, not the spread of the row scales.** At iteration 11 of the atomic reload the carbon unknown of cell 500 had been driven to zero, its column scale sat on the absolute floor $1e-20$, the probe step $\sqrt{\text{eps}} * 1e-20$ underflowed the residual difference, the column was exactly 0 and the banded preconditioner singular (dgbcon 3.98e-27, smallest pivot 3.7e-19); the $2.5e-15$ Krylov point is what a triangular solve with that factor returns. Iteration 1's $8.9e-10$ was only the 7.3-decade spread of the certification row scaling (no empty column). **Two repairs, the model untouched to the last bit** ($r0 = F/Drow$ and A unchanged, MEASURED by the scaled column norms): (a) element_unknown_column_scale = $\max(|x|, 1e-4 |reservoir|, 1e-20)$, the arithmetic at the function (the column is resolvable while the probe step stands far above $\text{eps} * X_{reservoir}$); (b) equilibrate_the_preconditioner: two-sided equilibration of the band that is FACTORIZED only, powers of two, banded_preconditioner_solve applying C dgbtrs(R v). Attribution (first solve, accepted/refused; empty columns at it 11; dgbcon at it 11): entry 14/16, 1, $4e-27$; floor alone 69/16, 0, $4.9e-11$; both 114/36, 0, $1.1e-9$. **Atomic reload (EXHALE_JFNK_MAXIT=150):** entry abort at $||R|| 1.777$ (no descent, 30 iterations); floor alone 1.189 (stagnation, 86); both **$3.800e-2$ at the 150 cap, still descending** ($2.5e-1$ to

3.8e-2 over the last thirteen iterations), dgbcon at iteration 1 8.9e-10 -> 1.65e-6, the brief's target (the pre-N20 7.4e-2) passed by 1.94x with one functional kept; refusals by name shift from ray-floor to ratio-below-eta (35) and one slope-size. **Carrier reload** 0.446 -> **0.1513** (2.95x), dgbcon 1.9e-11 -> 8.7e-6, +30 percent evaluations. The model's own column equilibration (EXHALE_COLUMN_EQUIL=1, default off) was built, MEASURED and NOT adopted: it moves the trust-region coordinates, the radius runs to the cap and trials ascend by 4e4 times the predicted decrease. **First cell rows declared once** (species_row_base_equation: element row = Dirichlet reservoir, READ from element_transport_residual; carrier row = control-volume balance, READ from carrier_residual's j == 1 branch; no equation changed). **Base derivatives**: the directional-derivative sweep at the base is a clean V with its minimum 4.8e-8 at sqrt(eps), the step build_banded_jac_full takes; whole-system rows 8.2e-8. **Closure map at the base** (measure_the_closure_map_at_the_base, four evaluations once per solve): the sweep exits after ONE pass at 9e-12, four decades inside the 1e-8 asked, so the reference is a pass count; quotients 2.1e-4 (atomic, attenuation) and 5.34 (carrier); outer_residual_closure_target (what the outer residual may keep, what the ray floor reads) is now separate from eq_sweep_reltol (what the sweep is asked = target / measured amplification; 1.87e-9 on the carrier reload). **B5h's ~4e3 base amplification is not reproduced on either fixture** and is not claimed. Manufactured states in krylov_and_dogleg (columns spanning eight decades, pivot ratio 5e-18 -> 0.54 after equilibration; exactness rows: the map and the unknown-space step unchanged to the last bit). Tests krylov_and_dogleg 73/0, steady_species_rows 147/0, certification 29/0; the new binary with both changes off reproduces the entry text to every digit; wasp_full_newton BITWISE (ten files; nspec_row = 0 short-circuits every new path, the hydrodynamic column scales untouched). **Advisor**: rebuild warning-free, steady_newton.o md5 c7621824 identical to the worker's, the three suites 73/0, 147/0, 29/0; the atomic reload rerun on the advisor binary reproduces the first solve to the digit (||R|| 3.800e-2 at 150, info = 1), and the run's SECOND stationary solve (the code re-enters the Newton from the handed-back state, another 150 iterations) reaches ||R|| **5.273e-4, flux spread 2.3e-6**, still not certified (log10 Mdot 10.10; exit 2), 32 min at 16 threads: the reload now descends four decades in 300 iterations where it stood still since B5m. Not converged. **Reported, not fixed**: n_eq_sweeps_model = 1 so the pass count, not eq_sweep_reltol, controls the closure on both fixtures; EXHALE_JFNK_MAXIT caps each solve, not the run; fix_active_species_bounds lost its D argument (the epsilon-active width now on state_column_scale); the [diag 4] true-residual row is not valid for long steps (its finite-difference product is out of range). The fixture README updated by the advisor.

- **Advisor measurement after N8a (atomic reload, EXHALE_JFNK_MAXIT=600, 16 threads, 18 min).** One solve with the cap at 600 stops on the stagnation detector at iteration 224 with ||R|| 2.671e-2 (info = 2; 170 accepted, 55 refused, all but three by "the reduction ratio is below eta"; log10 Mdot 10.06), WORSE than the same binary's 150 + 150 route, where the run's re-entry into a second solve from the handed-back state reached 5.273e-4. [The advisor's reading here was WRONG and N21 corrected it: the run does not re-enter the Newton from the handed-back state; info = 1 sends it back to MARCHING for 2000 steps (the log's Newton finish at step 2002), and the second solve starts from the marched state at ||R|| 6.06e-2. The two decades are gained by the marching, not by a fresh trust region; N21 measured every reset and none helps.] Not converged either way.

- **N4b** (decision 14 route (i), a step that moves along the element-budget surface; steady_newton.f90, element_inventory.f90 +94 (the derivatives of the map, element_constraint_derivatives), steady_species_rows +35 rows; diffusive_photochemistry.f90 untouched, the frozen budgets recovered exactly through the public carrier_headroom): **the acceptance row is met by ten decades**. On the carrier reload the He/H nucleus ratio against its reservoir stays at 1.39e-11 at every one of 68 iterates where the entry text drifted from 1.39e-11 to 2.016e-1 (20 percent of the ratio the run was given) over 79 iterates: the write-back of each iterate now returns the element totals it was handed (the N9 / N4a finding closed where those reports said it belonged). Best ||R|| 0.1513 -> **0.0899** at 14 percent fewer residual evaluations (5690 -> 4891). The shared constraint $2n(\text{H}_2) + n(\text{H}^+) + n(\text{OH})$

– $2n(\text{H}_2\text{O}) \leq n\text{H}_{\text{avail}}$ (and the O and C rows) replaces the five separate corners as the step limit (each corner is implied by the row, so nothing is lost and the direction along the surface is gained); a nonlinear feasibility screen by magnitude at every candidate; a restoration phase for an infeasible start (on the element-and-carrier reload 52 trials restored over 1165 cell-steps, worst violation 1.630 -> 2e-16; 6 candidates refused by magnitude; 167 write-back cells whose closure could not return an element total, counted). ****Stated, not claimed****: on both reload fixtures the shared hydrogen row is never within a probe step of being tight (occupancy 0.985 to 0.989, active band 1e-8), so the active-set PROJECTION onto the constraint hyperplane, the part route (i) is named for, fired on ZERO iterations of either run; its correctness is stated by the suite rows, and the measured improvement comes from the write-back, the budget leaving the coordinate box and the shared step limit. The atomic element reload (element rows, no carrier row) is identical line for line over its 150-iteration first solve (||R|| 3.800e-2 both). Tests steady_species_rows 182/0 (RED with EXHALE_ELEMENT_CONSTRAINT_ROWS=0, 20 rows fail), krylov_and_dogleg 73/0, certification 29/0; mol_carrier marching 200 steps bitwise; wasp_full_newton bitwise (||R|| 4.213e-9, info = 0, log10 Mdot 13.30). ****Advisor****: rebuild warning-free, steady_newton.umd542fcf223 and binary md59c9ba732 identical to the worker's; the three suites 73/0, 182/0, 29/0; wasp_full_newton's two matrix files compared by the advisor against the Stage A reference: IDENTICAL. Two earlier builds of the item were discarded and their runs rerun (report section 8). Certification unchanged (decision 12 c). ****Stage B gate****: N4a map, N4b constrained step and element-conserving write-back, N5's three contracts measured (contract 2 fails by design on two reloads), N8a scales and closure map: the gate's rows are in; what is NOT met is convergence of any species-row solve, which is Stage C's criterion, not Stage B's. N4b's user-visible switches: EXHALE_SPECIES_BUDGET_FACE is retired and replaced by EXHALE_ELEMENT_CONSTRAINT_ROWS(=0 restores the five corners), EXHALE_ELEMENT_WRITE_BACK added (the element-conserving write-back); no input key, output column or unknown layout changed. Also found by N4b: dc/d(nd) =

c/indexactly (the constraint is homogeneous of degree one in the density, so an active row cannot push the mass unknown); two earlier builds of the item were discarded on the atomic fixture's line-for-line check (a blocked probe component zeroed unconditionally made the sampled operator nonlinear, factor 33 regression; the probe sides compared before capping). The advisor annotated the retired switch in the B5l entry and indocs/steady_solver_design.md'.

- **N21** (what limits the species-row solves now; steady_newton.f90 +417, krylov_and_dogleg +31 rows): **no reset of the trust-region state helps, and the advisor's premise was wrong.** The run's second stationary solve does not start from the handed-back state: info = 1 returns the run to MARCHING for about 2000 steps and the second solve enters at $||R||$ 6.06e-2 (advisor's own log: Newton finish at step 2002, then step 4002, the third solve reaching 4.26e-4 on the stagnation detector), so the two decades the 150 + 150 route gains are gained by the marching between the solves. MEASURED on the atomic reload from the SAME 150-iteration prefix, EXHALE_JFNK_MAXIT=300: continue (control) 227 iterations, stagnation, $||R||$ 2.671e-2 (170/55, reproducing the 600-cap run); resetting the pseudo-transient shift, the acceptance memory or the closure map: bitwise identical logs (three exact no-ops: dtau already sits on its floor dtau0, the Grippo window is reported and never acted on on the coupled route, the closure map re-measures to amplification 1.000); resetting the radius and ceiling 2.851e-2; returning to the best iterate 2.921e-2; all five together (what a re-entry does) 3.161e-2; a second Krylov cycle of 40 2.927e-2; a restart every 10 iterations without improvement runs to the 300 cap at the same 2.671e-2. Carrier reload: control reproduces N4b (8.993e-2), periodic restart 1.153e-1, second Krylov cycle 1.728e-1, both worse. **The periodic restart trigger is delivered default OFF** (tr_restart_stall_default = 0) by the brief's own condition. **The linear solve is not the limiter:** after N8a the atomic cycle reaches a median 0.157 of its right-hand side over iterations 1 to 150 (N7: 0.57 to 0.69) but 0.74 to 0.82 over 151 to 225; the carrier cycle 0.963; doubling the products halves the linear residual and makes the outer solve WORSE on both fixtures; in the second atomic solve a cycle reaching 0.98 coexists with a two-decade fall of $||R||$. The radius reset is worse for a stated reason: the carried radius 9.85e-9 accepts steps at ratio 1.000 while a re-initialized 1.15e2 over-predicts by 39x then 3392x and costs four refused iterations. READ: the band is refactored every outer iteration (a re-entry changes the preconditioner only through the shift) and the Krylov cycle starts from zero every iteration. Tests RED (13 symbols absent from the entry module), GREEN krylov_and_dogleg 104/0, steady_species_rows 182/0; with everything disarmed the delivered binary reproduces the entry text's 20-iteration solve line for line; wasp_full_newton BITWISE (ten files). Advisor: rebuild warning-free, steady_newton.o md5 cdbbf535 and binary e140284a identical to the worker's, the two suites 104/0, 182/0; the advisor's own n8a_reload log confirms the marching between the solves. **The finding that names the next item (N22):** the predicted decrease stops scaling with the step below about 1e-7 (from iteration 140: the step falls 4x and pred stays 7.099e-4; every longer step of the resulting period-two cycle is an ascent of 3e-3 against a promise of 3e-4, ratios -4, -8, -13, refused, quartered, then accepted at ratio exactly 1.000); the production dogleg geometry is cleared by two new rows (quartered radius quarters the step and the predicted decrease to 2.5e-1 and 2.50009e-1), so what remains is trust_region_step's bookkeeping at those lengths or a precision floor in $r_0 \cdot A_s$ and $||A_s||^2$; the

separating measurement is one diagnostic line ($||A_{\text{ s}}||, r_0 \cdot A_{\text{ s}}$ beside $||s||$ at two radii of one iterate).
 README of the fixture updated by the advisor.

- N13b** (decision 17 option a, transit spectra carry their validity; EXHALE_transit.py, exhale_transit_lib.py, transit_census +8 rows; Python only): one reader read_adv_validity locates the status columns by name in the # columns line and distinguishes three cases: schema 2 (N11's two fields: a row is refused for the census when adv_T_status is not 0, the composition field reported beside it), schema 1 (today's single column), legacy (no schema: validity UNKNOWN, and both the console and the saved metadata say UNKNOWN rather than 0 refused). Every tpm_*.txt gains a comment metadata block (transit_schema 1, the input profile and its schema, its adv_input_certified / stationarity operator / model restrictions lines copied, its provenance lines, the tool's own identity by git hash and file mtime, the census for that line with the eight-point refused-share ladder and its maximum, the CONTRIBUTION-not-uncertainty sentence, the EXHALE_TRANSIT_* overrides in effect); the two hand-written products get the same block. MEASURED: numerical columns bitwise on all 10 wasp_full and 5 mol_base_handoff products; the schema-2 read tested on synthetic files and on a relabeled wasp_full profile (the relabeling is the worker's, not N11's mapping; the production file comes with N11); every consumer in the tree (python/make_transit_figures.py, the WASP-52b and LHS 1140 b notebooks, two LHS scripts) reads through np.loadtxt or skips #, none edited; the planet folders' tpm_*.txt untouched (mtime 2026-08-19). Suite RED 8 rows (no reader existed), GREEN 19/0; advisor rerun 19/0. Fixed on the way: the second np.loadtxt(..., max_rows=1) column count (NumPy warning on every run) replaced by one rule. Reported: the refused share is taken over the temperature field only (a companion share over the composition field would be the honest partner); still line center only; README_HOWTO.md should mention the metadata block (advisor added the sentence).
- N11** (decision 16 option a; post_process_adv.f90, the _adv writer of write_output.f90, examples/exhale_io.py, EXHALE_plots.py, adv_static_limit +driver M1-M5 and rows R8-R9): **stationarity of the input state is judged by the certification's own mass operator** (assemble_residual, residual_row_scale, the mass row $|R_1|/s_1$ against $\text{adv_mass_row_tol} = \text{cert_tol_mass}$), the enthalpy term ratio kept as a printed sensitivity diagnostic that refuses nothing; **two status fields** adv_T_status, adv_comp_status (0 corrected, 1 retained, 2 failed, 3 unsupported, 4 not_evaluated) in BOTH _adv files, columns 8 and 9, with write_adv_validity_header writing # adv_schema 2, the two legends, # adv_status_counts T <5> comp <5>, # adv_input_certified <T|F> <reason> (the certification's verdict on this very state), # adv_stationarity_operator face_flux_mass, # adv_product (a ONE-WAY correction on the input's density and velocity field; a converged scalar temperature solve does not make a row self-consistent), # adv_model_restrictions; adv_unsupported where the omitted molecular and oxygen species carry more of the cell's particle count than the carried ones (structural, no threshold); exhale_io exposes both fields and the header (load_adv_header, load_adv_status), a legacy file reads schema 1 with both fields ADV_UNKNOWN, never a translation of the old column; EXHALE_plots.py shades the refused rows. **MEASURED, the physics of the item:** the ratio screen and the mass row are not interchangeable and the ratio is the weaker: on the analytic outflow fixture the ratio is $1.1\text{e-}13$ everywhere while the mass row stands at $2.1\text{e-}7$ (RED of the

old criterion); on `wasp_full` the ratio refused 297 of 503 rows, the mass row refuses 503 of 503 (largest 1.04 at `r` 1.032); `mol_base_handoff` 33 vs 503 (0.245 at `r` 4.21); `wasp_full_newton` (the one certified state) 2 vs 3. So on the two relaxation snapshots every row is retained and the `_adv` product IS the run's own state (`T_adv` equal to `T` in 504 of 504 rows; the old product differed by up to 6.4 and 207 percent); on the certified state the change is confined to the base (`p`, `T` $7e-5$; species $7.7e-2$ at `r` 1.0002). Run states bitwise in all three; `rho`, `v` in the `_adv` files moved by exactly zero. Tests 48/0 (advisor rerun on `build_lwv`, md5 4320f5b7 identical to the worker's: 48/0). No golden refreshed; every `_adv` golden will move. **Advisor decision (N11b, launched)**: the criterion is the right OPERATOR but the wrong TOLERANCE for a conditional correction. `cert_tol_mass` = $3e-12$ is the tolerance of a certified stationary state; the advective correction is a first-order local correction whose own error is of the order of the mass row's measure (the fractional change of the face mass flux across the cell), so a du-stopped or snapshot state with a mass row of $1e-3$ is exactly what the tool was built for and is what review R6 calls the "approximate local use" with "a tolerance tied to the required error". N11b writes the mass row measure itself as a column of `Hydro_ioniz_adv.txt`, sets `adv_conditional_tol` = $1e-2$ (the correction is described as accurate to that fraction, stated in the header beside `adv_input_certified`), and keeps the certification verdict as the separate statement it is; the product is then a CONDITIONAL correction with its condition written in every row. Advisor: two per-field and two percent (one in a printed message) fixed. **Reported, not fixed**: `assemble_residual` leaves `R(:, 1-Ng)` undefined (`RK_rhs` assigns `2-Ng..N+Ng` only; no caller reads it today; `RK_rhs.f90/steady_residual.f90`); `README.md` 154, `README_HOWTO.md` 959 and the manual section 4 still describe the single `adv_status` column (N11b updates them with the final schema); a suite row for corrected needs a converged fixture.

- **N11b** (the `_adv` product is a CONDITIONAL correction with its condition in every row; `post_process_adv.f90`, the `_adv` writer of `write_output.f90`, `exhale_io.py`, `EXHALE_plots.py`, `EXHALE_transit.py`, `exhale_transit_lib.py`, `adv_static_limit` (53 rows), `transit_census` (21 rows), `README.md`, `README_HOWTO.md`, manual section 4): the row decision is taken against `adv_conditional_tol` = $1e-2$ (the fraction a corrected row is accurate to in the mass flux, the first-order argument at its declaration), the measure `m_j` = $|R_1|/s_1$ is written as column 10 `adv_mass_row` of `Hydro_ioniz_adv.txt` and named in # columns, the header carries # `adv_conditional_tol` and # `adv_mass_row`, and # `adv_input_certified` stays the certification's verdict with `cert_tol_mass`; schema stays 2 (readers are header-driven; the one place that counted columns from the end of the row was a test row, fixed to locate by name). MEASURED against the N11 binary: corrected rows (`T`) `wasp_full` 0 -> 215 of 504, `mol_base_handoff` 0 -> 341, `wasp_full_newton` 500 -> 502 (the two others are ghost rows below the energy loop's first cell); the column equals the certification's own printed measure on the same state to every digit (1.942392 at cell 432); run states and `rho`, `v` of the `_adv` files bitwise. `exhale_io` returns `adv_mass_row` and the tolerance; the transit metadata block copies the tolerance line and the largest measure; the census still refuses on `adv_T_status`. Tests 53/0 and 21/0 (advisor rerun on `build_lwv`, md5 3131cab0 identical to the worker's: 53/0, 21/0). **Every _adv golden moves, far on the two snapshot cases: refreshed at the Stage D gate**. For the record: the first-order argument bounds the dropped term; in some corrected cells the KEPT enthalpy-flux term of the mass divergence still dominates the energy equation (452 of 503 cells of `hydrostatic_column` against 390 retained), and whether one percent

of the mass flux bounds the temperature error of such a cell is not measured; a corrected row is not close to the run's own row and is not expected to be (the certified state moves T by $6.7\text{e-}2$ at m_j $3\text{e-}16$). Advisor: one percent in README.md fixed; docs/postprocess_advection_validity.md marked stale with a pointer.

- N10** (decision 15 option a, the restart contract; input_read.f90 +137, load_IC.f90 +551, write_output.f90 +13, EXHALE_main.f90 +166, grid_and_gates +1 row script of 20 assertions, docs/input_schema.md K43 and appendix D.2, manual entry; PDF not rebuilt): the key Restart intent: trajectory | relaxation | stationary [evaluate|equilibrate] with the three consistency refusals (default trajectory in phys, relaxation otherwise, so no existing input changes behavior); both state files carry the eight-line block (restart_schema 1, reservoir, species_columns, grid, constants, options with 20 switches, t_phys[s], source), built and parsed in load_IC.f90 (one definition), every number ES23.16; no block = legacy, loaded as before and marked provenance unknown in the log and in every file the run writes (the mark is inherited); a disagreeing reservoir (within heh_dev_tol), grid, constants or options is a refusal naming the field, species_columns and source are reported; a pair whose halves carry different blocks is refused. stationary_state_of_the_loaded_restart: WENO3 fixed, the N16d rebuild order, one equilibrium sweep (part of the residual's definition), the loaded composition's departure from that sweep's root printed, the residual and the certification held to the FILE's own certified=, then evaluate writes back and stops or (default) enters steady_wind_with_element_diffusion at once; no CFL step. **MEASURED:** the atomic element fixture AS LOADED stands at $|R|$ 1.8924 (the marched hand-off read 1.888: the two CFL steps); the certified wasp_full_newton state is NOT the fixed point of its own sweep (particle count moves $2.1\text{e-}11$, species $5.0\text{e-}10$, cooling $3.6\text{e-}10$: the cell solve's $xtol = \sqrt{\epsilon}$ and the file's digits), reported at every evaluation rather than equilibrated away; stationary evaluate takes 0 steps and returns the certification unchanged; stationary on that state enters the JFNK at $7.133\text{e-}9$ and exits $info = 0$ with no step. Tests 20 rows GREEN (RED on the entry binary: the key warned and ignored, one step marched, no block, no refusal); grid_and_gates 130/0 (advisor rerun 131/0 on build_lwv). mol_carrier 200 steps numeric-line bitwise (eight new # lines); run_check.sh line 297 excludes # lines, so the block cannot change a matrix verdict; wasp_full_newton cold numeric-line bitwise against a control binary of the same tree without N10 (its $5.8\text{e-}9$ / $9.4\text{e-}9$ movement against the golden is B5j's recorded movement, not N22's as the worker supposed). **Fixed on the way:** the header loop matched columns as a SUBSTRING (index(line, 'columns') > 0), so the _adv files' NOTE prose or the new species_columns field would have replaced the species labels silently; now the first token (comment_field_is); the same hazard in restart_round_trip.sh's Python readers fixed. Deviations from the design, stated: no EXHALE_STATIONARY_EVALUATE_ONLY env hook (the second word is the one answer); the provenance unknown mark is not yet in EXHALE_setup.out (write_setup_report.f90 not owned; one line); species_columns reported not refused (the by-label species map is a documented path); source is the git revision, not an executable md5. **Consequence for the user (decision item 21):** the exact options comparison forbids the option-ladder workflow (converge without an option, restart with it on); legacy pairs are unaffected, every new rung would be refused; the disciplined fix is a key naming the tokens allowed to differ. Also corrected: docs/input_schema.md appendix D listed the retired valve/fluxconst fields and the wrong base_bc values. Reported: Run mode is absent from input_schema.md; the EXHALE_PTC=1 route writes no setup report;

a re-evaluated state writes `sec_ion_step=0`.

- N22** (the predicted decrease stops scaling below $||s|| \cdot 1e-7$; `steady_newton.f90` +195/-10, `krylov_and_dogleg` +16 rows): **all three candidates of the brief refuted by measurement, and the cause named:** the trials that break the scaling are exactly the trials whose species unknown was written onto a face of its box (`nout = 1`), and that write is a move of the state no model of the step contains (11 of 11 such trials refused with a fixed ascent of $3e-3$ over five decades of step; 10 of 10 others accepted at ratio 1.000). When the radius is cut the SCALED step quarters but the step in the unknowns the residual reads, $||D \cdot s||$, moves 0.4 percent, so the model image and `pred` are unchanged to seven digits; $r0 \cdot A \cdot s = -||A \cdot s||^2$ to every digit (`pred = 0.5 \cdot (r0 \cdot A \cdot g)^2 / ||A \cdot g||^2, radius-free, the interior minimizer along the leg: not a defect); a fresh Jacobian action at the same step reproduces pred to every digit (no stale cache); the probe displacement is $\sqrt{\text{eps}}(1 + ||Y||) = 1.8e-7$ whatever the direction, nine decades above the residual's reproducibility (no floor; the reverse: the stalling step is $2.7e-3$ of the probe arc). The fix: hold_the_step_where_it_leaves_the_species_box zeroes the step at an unknown it would carry out of its box (the iterate is feasible, so the trial is, and the step is one the operator can be sampled along); probe_length_of_the_jacobian_action is the one definition of the probe arc; a diagnostic line for each trial under EXHALE_ELEM_DIAG. Atomic reload, cap 300, before -> after: 227 -> 113 iterations (both stagnation), accepted/refused 170/55 -> 91/19, negative ratios 49 -> 14, best $||R|| \cdot 2.353e-2$ -> 1.333e-2, handed back $2.671e-2$ -> 1.356e-2; the period-two cycle is gone (iterations 95-98: pred doubles with the step, all accepted at 0.998 to 1.001, steps $1e-2$ to $1e-1$ instead of $1e-9$); the closing refusals are ray/floor verdicts at POSITIVE ratios. Carrier reload BITWISE (the branch never fires); wasp_full_newton BITWISE (ten files; the probe-arc refactor is shared through jv_product and this settles it). Tests: the twelve-rung ladder rows are GREEN at entry (the geometry was never the cause), the new rows RED by four absent symbols, GREEN 120/0; steady_species_rows 182/0. Advisor: rebuild warning-free, steady_newton.o md5 d70a078b identical to the worker's delivered object, the two suites 120/0, 182/0. Not converged. Reported, the next item (N23): the approximate-gradient leg is 0.7 percent of the step and supplies ALL of its model image ($||sU|| / \Delta = 6.9e-3$ passes cauchy_leg_min_fraction = 1e-3 while $||A \cdot s|| = \tau_c \cdot ||A \cdot g||$ to four digits): the criterion must be the share of the IMAGE, not the length ratio (N7b's finding at a length the threshold admits).`
- N10b** (decision 21 option a; `load_IC.f90` +319/-20, `input_read.f90` +145/-1, `write_setup_report.f90` +62/-6, `grid_and_gates` +1 script of 11 rows, `docs/input_schema.md` K44 and D.2, manual entry): the key Restart option change: `<token>[, ...]`; the options field is compared TOKEN BY TOKEN from one table (`opt_name(20)`, `opt_value`, the emitted text bitwise unchanged), only the named tokens may differ, every other difference refuses the load naming the token with its from -> to; six layout tokens (`metals`, `mol`, `oxychem`, `carrier`, `carrier_newton`, `iontrans`) may never be named (they decide the rows the state carries: a cold start, not a restart); an unknown token, a grid/reservoir/constant word, the key with no token or without Load IC? True are input errors; the change is written into both halves of the new state as `# option_change from -> to at restart of <source>`, inherited (history of 32 lines); EXHALE_setup.out gains Restart provenance: UNKNOWN | UNKNOWN by descent | states its configuration and the permitted changes

(N10's deviation 2 closed; the "by descent" wording was added because the first wording was measured false on a ladder's second rung). **Correction of the advisor's brief:** LW is a FLUX (Stellar LW flux), not an option token, so the `heh_1_40k` -> `heh_1_lw_40k` rung (renamed 2026-09-16 from `armA_noLW` -> `armA_LW`) never needed this key; and every state written before the Jupiter-radius unification (those two included) is refused by the grid guard at $2.25e-2$ in the outermost cell, so the `heh_1_40k/heh_1_lw_40k` directories are configurations, not restart sources. The ladder step was measured on the `heh_1_40k` configuration with `base_ir`, `mol_ir` flipped: refused by token without the key, accepted with it, the provenance line in both halves. Tests RED 11/11 at entry (the key was an unrecognized line), GREEN 11/11; `grid_and_gates` 141/1 with the one FAIL attributed (below); `mol_carrier` 200 steps numeric-line bitwise (the rebuilt # options line bitwise). Also fixed: N10's test stage A read the LIVE `wasp_full_newton/output/`, which the matrix rewrites (a false failure of the legacy-mark row as soon as the pair carried a block); it now strips the block from its own copy. **Advisor:** rebuild warning-free, md5 `b10641e5` identical to the worker's; `grid_and_gates` 141/1, the FAIL being `stationary_evaluate_rebuilt_within_sweep_budget` reading the live `output/` of `wasp_full_newton`, which the running Stage D matrix had left as a mid-run relaxation snapshot ($3.2e-3$ against $1e-8$; the same pair evaluated by the entry and the N10b binary is bitwise, so not N10b's); the advisor pinned the certified state as `backup/regression/wasp_full_newton/IC/` (copied from the golden, `certified=T`, README there) and pointed the row at it. Reported: `restart_grid_guard.sh`'s stale header (third time); `sec_ion_step=0` on a re-evaluated state; whether `molbase`, `he_diff`, `he_metal_diff` belong with the layout tokens is a reviewer's question (the worker's reading: they add operators, not rows).

- **N23** (the approximate-gradient leg admitted by its share of the model image; `steady_newton.f90` +220/-5, `krylov_and_dogleg` +16 rows): two constants `cauchy_leg_image_share_max = 0.5`, `cauchy_leg_step_share_max = 0.1`, two pure routines (`cauchy_leg_shares_of_step_and_image`, `cauchy_leg_is_image_without_step`: image share above the first AND step share below the second, so a leg that IS the step keeps its image), and a branch in the dogleg's trial loop that DROPS such a leg from the step and its image and rebuilds the dogleg at no extra product (drop, not rescale: a rescale would either shorten A sU without sU, which is not a step of the operator, or move the step off the dogleg path); the model slope handed to `ray_slope_verdict` is the slope of the step taken in both branches. **MEASURED, the reading ([diag 9]):** at the reload's second solve quartering the radius moves the promise by 0.9959 with the leg admitted and by 0.2501 with it dropped (six iterates: 0.9959 to 0.8938 against 0.2501 to 0.2517), so the promise follows the radius again where it did not. **Honest impact: the new rule fires ZERO times on both fixtures and all three runs are BITWISE** (atomic 113 iterations, $1.356e-2$; carrier $8.993e-2$; `wasp_full_newton` 13.30); on the post-N22 trajectories the length test reaches every image-dominant leg first (117 length drops, 0 image drops); the shares printed at every trial say the image test is the right criterion nonetheless (0.9986 of the image at $3.4e-5$ of the step; eleven of seventeen read trials satisfy the predicate on their own), and the leg N22 saw at iteration 145 sat on a trajectory the N22 hold removed. Tests RED (four symbols absent), GREEN 136/0; `steady_species_rows` 182/0; -Wall warnings 6 before and after. **Advisor:** rebuild warning-free, the two suites 136/0 and 182/0 on `build_lvw` (the object md5 differs from the worker's delivered one by a post-measurement comment edit that moved line numbers, which the I/O statements carry; the worker's line-for-line reproduction of the solve covers

it). Not converged; the rule is not claimed to improve either fixture. Reported: `cauchy_leg_min_fraction` is now a proxy for a test the code can make directly (letting the image test alone decide is the next increment, not taken because no fixture exercises the difference); the trial diagnostic's evaluations enter the cost counters.

- **N24** (what refuses the remaining trials of the atomic element reload; `steady_newton.f90` +51/-2, `krylov_and_dogleg` +13 rows): **the obstruction is the linear model, and the row is named**. Atomic reload, cap 300: 110 trust-region iterations (the "113" of earlier reports counted three summary lines), 91 accepted / 19 refused. The 19 are two populations, neither a step-control defect: 14 ascents at LONG steps ($||s||$ 1.6 to 51, five with zero species unknowns held, so N22's mechanism is not what is left) and five closing refusals at positive ratios ($||s||$ $2.3e-4$ to $8.9e-7$ with the promise quartering exactly while the true decrease keeps a step-independent floor of $3e-11$ to $1.6e-10$). One accepted step (outer 59, ratio 0.948, $||R||$ $7.37e-2 \rightarrow 2.12e-2$) moves the binding row from the energy equation to **the sodium element row of the outermost cell (cell 500, r 4.06 Rp)** and the solve never leaves it: from outer 60 to 110 that row holds 0.435 of the squared merit and the whole judged distance ($5.3e-2$ of its own scale, $5.36e6$ at `cert_tol_element` $1e-8$) while $||R||$ stands at $1.333e-2$. At outer 105 everything else is sound (0 held, 0 refused samples, 0 empty columns, `dgbcon` $1.25e-6$, step ratio 0.9998); what fails is the linear model: 40 of 40 products, true relative residual 0.58 against 0.1, **the Arnoldi image 21 percent away from the operator** ($1.6e-6$ at iteration 1, $2.3e-4$ at N7's iteration 60), $||sN||$ $8.6e2$ against a radius $5.9e-2$, so the region uses $7e-5$ of an inaccurate direction; the stuck row is at once the smallest scaled column and the smallest scaled row of the band. The stagnation detector read the judged distance, which IS that row; $||R||$ was not falling ($1.337e-2 \rightarrow 1.333e-2$ over 32 iterations). **The image-only leg test** (`EXHALE_CAUCHY_LEG_BY_IMAGE=1`, default off) measured and NOT adopted: worse on both fixtures (atomic $2.30e-2$ against $1.36e-2$; carrier 0.324 against 0.090, to the cap), the two runs parting at outer 7, so N23's "the rules agree" was that trajectory's property. Deliverable 3 STOPPED as briefed (linear solve / model / one row's physics, not the step control). Tests RED (two symbols absent), GREEN 149/0; `steady_species_rows` 182/0; the delivered binary reproduces the entry solve line for line over 925 lines; `wasp_full_newton` BITWISE. Advisor: rebuild warning-free, 149/0, 182/0 on `build_lwv` (the advisor's binary of N23 had been built from an earlier revision of `steady_newton.f90`, as the worker showed; the suites and fixtures were the check). **Proposal (N25)**: (1) explain the 21 percent Arnoldi gap (`EXHALE_GM_ORTHO`; Jv reproducibility by two products); (2) ask the cycle for the direction ON THE BALL (Steihaug-Toint truncation) instead of solving to 0.1 a leg four decades beyond the radius; (3) the sodium row's own terms at cells 490 to 500: the element operator gives cell 500 a zero-gradient outer ghost, so nothing outward constrains it (a boundary question of the element operator, `binary_element_diffusion.f90`); (4) `cert_tol_element` $1e-8$ is what makes this row the whole judged distance (UNCONFIRMED, ISSUES 3.1 e; not moved). Reported: the (JFNK) it line prints `k=*` for a species row without naming the element (in this case the identity was the whole diagnosis); the stagnation messages quote $||R||$ where the detector read the judged distance.
- **Stage D gate and golden refresh (advisor, 2026-09-09 21:14 to 2026-09-10 00:27, make check on the shared build/; refresh 00:40)**. Tree after every item to N24 (N25 was running on a private build). Verdict against the goldens of 2026-09-08 19:05: on all 16 default cases the two marching files are **data identical** (15 cases

bitwise; wasp_ full_newton WITHIN at 5.78e-9 / 9.41e-9, B5j's recorded movement, which the 0.1 percent rule counts as identical) and all 32 _adv files FAIL by SHAPE (10 against 8 and 40/36/43 against 38/34/41 columns: the two status fields and the measure column of N11/N11b), so make check exits 2 as expected. The _adv movement is the physics of N12 (stale composition, upstream energy), N11 (mass-operator stationarity) and N11b (conditional tolerance): T_adv changes in 477 of 504 rows on mol_base_handoff (max 61.6 percent, 364 -> 588 K at r 1.093), 479 of 504 on mol_carrier (max 138 percent, 379 -> 901 K at r 1.098), 215 of 504 on wasp_full (max 2.3 percent, 5637 -> 5505 K at r 1.131). **Goldens refreshed:** the previous set archived as backup/regression/golden_pre_stageD_20260910/, run_check.sh golden snapshotted the outputs the gate had just produced (17 cases, including the three hp_* and oxygen_chemistry extras), and the advisor compared the 68 snapshot files against the case outputs with the comparator's own #-filter: 68 identical, 0 differ (this comparison, not a three-hour rerun, is the "check again" of the standing rule, since a snapshot is a copy; the user asked that regression checks not be repeated). The shared build/ and EXHALE.x are now current with the tree (md5 f6ccb498 at the gate's build; N23/N24 built after it touched steady_newton.f90 only, so the marching path of the matrix is the gate's). State files now carry the N10 metadata block and the _adv files the N11/N11b header; both are # lines the comparator ignores.

- **N25** (the Arnoldi gap, a direction on the ball, the sodium row at the outer boundary; steady_newton.f90, krylov_and_dogleg +25 rows, binary_element_diffusion.f90 a default-off print only, 33 lines, no arithmetic): **(1) the 21 percent Arnoldi gap is the NONLINEARITY of the finite-difference action across directions**, shown positively: every one of the 40 Arnoldi columns matches its own product to 1e-16 to 3e-16, two products of one direction agree exactly, the basis's orthogonality loss is 4.4e-13, yet the action of $V_1 + V_2$ differs from the sum of the two actions by 1.1e-4 and the assembled gap tracks the step length (8.5e-5 and 2.5e-4 per unit of $||sN||$ at outer 1 and 105; $||sN||$ 12 -> 865), the signature of a curvature term over the fixed probe arc $\sqrt{\epsilon}(1 + ||Y||) = 1.8e-7$. Reorthogonalization NOT adopted (loss 1e-13 to 3e-12 against the 1e-8 level; as a measured option EXHALE_GM_REORTH0=1 it is worse on both fixtures: 1.487 and 0.43). **(2) Steihaug-Toint truncation** is in as EXHALE_KRYLOV_ON_THE_BALL=1, default off: products per iteration 33 -> 7 (atomic), 40 -> 1.5 (carrier); atomic better on the judged distance (1.69e6 against 5.31e6, lowest $||R||$ 9.4e-3 against 1.33e-2) but worse on $||R||$ handed back (1.69e-2 against 1.36e-2); carrier worse (0.486 against 0.090); the brief's condition fails, default stays off. **(3) The sodium row of cell 500 is not a well-posed outflow condition, and the reason is a DEFECT of the stationary residual evaluation, not of the element operator:** the outer ghost of the element mixing ratio is FROZEN at the pre-solve composition (1.73111e-6, the metals.inp reservoir the handed-in state carried) throughout the solve while cell 500 falls to 1.0e-11, so at the stagnating iterate the ghost is 1.7e5 times the cell it "continues"; the marching path refreshes that ghost (the second solve, entered at step 2002, has ghost = cell 500 again); the accepted step of outer 59 did not move $fX(500)$ at all (it moved the outer face mass flux); the scaled Jacobian row is dominated by the momentum of cells 499/500 and the sodium of 499 while its own diagonal is 1.3e-4 of the largest entry (the row is nearly independent of its own unknown). The physically stated condition (zero gradient at the top, no diffusive flux) requires the ghost to follow the ITERATE; the repair is in the composition write-back / ghost policy of eval_residual, which is N26 (launched). **(4)** the (JFNK) it line names the element (worst row: element Na of cell 500) and

the stagnation messages quote the judged distance and `dj_best`. Tests RED (eight symbols absent), GREEN 174/0; `steady_species_rows` 182/0; default path unchanged (atomic 109 iteration lines identical to the digit; carrier identical); `wasp_full_newton` BITWISE. Advisor: rebuild warning-free, 174/0, 182/0 on `build_lwv`; three British neighbor in the element operator's comments fixed. Not converged; the on-the-ball run's best iterate is chosen by the judged distance while the control is `||R||` (the two rank iterates differently, which limits the comparisons).

- N26** (the upper ghost of a transported element or carrier follows the iterate; `steady_newton.f90` +135/-20, `steady_species_rows` +7 rows): in `write_species_rows_into_composition` (the ONE place a trial composition is assembled for the residual: `eval_residual` and the three best-iterate restores) the transported columns of the upper ghosts take the outermost cell's value by the rule the marching path uses (READ: $\text{Yetr}(N+1:N+Ng, q) = \text{Yetr}(N, q)$ in `species_advection_stage`; $\text{ftry}/\text{fc}(N+1:N+Ng, ic) = \dots(N, ic)$ in the carrier operator; `Apply_BC` never touches `f_sp`), the element totals of a ghost restored after the carrier copy, a ghost deficit clamped as the operator clamps its own; lower ghosts untouched (Dirichlet reservoir); the three-unknown route returns before reading anything. The mixing ratio follows the mass fraction exactly (equal He mass fraction at N and its ghost gives equal $n_{\text{He}}/n_{\text{H}}$, hence equal $n_{\text{X}}/n_{\text{H}}$; MEASURED $1.4\text{e-}16$). **MEASURED, atomic element reload (EXHALE_JFNK_MAXIT=300)**: `||R||` handed back $1.356\text{e-}2 \rightarrow 3.719\text{e-}4$ (36x), judged distance $5.36\text{e}6 \rightarrow 6.21\text{e}4$ (86x), flux spread $6.6\text{e-}4 \rightarrow 5.5\text{e-}7$, 91/19 - > 235/18 accepted/refused, the sodium row of cell 500 binds on **0 of 252** iterations (was 51 of 109) and no row of cell 500 is ever the worst; the solve now stagnates on the INTERIOR helium row of cell 246 (`r` 1.155), `info` = 2, uncertified. **Carrier reload**: binding cell (H2 of cell 205, interior) and judged distance ($4.807\text{e}7$) unchanged to four digits, `||R||` handed back $8.993\text{e-}2 \rightarrow 2.519\text{e-}1$ (worse by 2.8x, a path difference of a stagnating solve, reported not judged), flux spread down 12 percent. `mol_carrier` marching bitwise (no caller outside `solve_steady_jfnk`); `wasp_full_newton` BITWISE. Tests RED on the entry text with only the export added (4 rows: ghost mass fraction $6.7\text{e}4$ off, mixing ratio $1.0\text{e}5$ off, the outermost element row of a flat column $1.19\text{e-}2$ instead of 0, the carrier ghost 0.80 off), GREEN 189/0 (a non-vacuous row keeps the stale ghost and reads $1.19\text{e-}2$ in both builds); `krylov_and_dogleg` 174/0. Advisor: rebuild warning-free, `steady_newton.o` md5 4bd3138a and binary e56609ff identical to the worker's, 174/0 and 189/0 on `build_lwv`; one neighbors fixed. **Reported, the next small item (N26b)**: the relaxation solvers `solve_trace_element_in_hydrogen` and `element_diffusion_step` refill their working arrays' upper ghosts before every pass while `element_transport_residual`, documented as the same operator, reads the ghost from `f_sp` with no refill, so the operator's boundary depends on its caller (`binary_element_diffusion.f90`, two lines). Not claimed: convergence; an explanation of the collapse of the outermost cells to $1\text{e-}11$ of the reservoir.
- N26b** (the element operator refills its own upper ghosts; `binary_element_diffusion.f90` +26/-0, new suite `src/tests/element_operator/` with 5 rows): `element_transport_residual` now writes $\text{Xhe}(N+1:N+Ng) = \text{Xhe}(N)$, $\text{fX}(N+1:N+Ng) = \text{fX}(N)$ and $\text{wYtr}(N+1:N+Ng) = \text{wYtr}(N)$ (the third line the relaxation twin gets for free through `project_elements`) so the operator poses ONE outer boundary (zero gradient, no diffusive flux at faces 0 and N) whatever the caller left in the ghosts; the lower ghosts stay the Dirichlet reservoir; the header

states the boundary. Both outermost rows (N-1 and N, WENO3 reaches cell N+1 at face N-1) are covered. MEASURED: the relaxation's own fixed point handed to the residual with stale ghosts read $1.19\text{e-}2$ of its scale in the outermost row before and $1.5\text{e-}13$ after (the interior level is $1.3\text{e-}13$); the same column with stale and with zero-gradient ghosts now gives the two outermost rows bitwise. Suite RED 3 of 5 on the entry text, GREEN 5/0; certification 29/0; the atomic reload's first two JFNK iterations identical but for the merit's reproducibility going $2.8\text{e-}14 \rightarrow 0$ exactly (the residual no longer reads a ghost the sweep leaves at a different last bit), the outputs moving at $7.9\text{e-}12$ (T) and $1.6\text{e-}10$ (an `_adv` status/measure column) because N26's write-back hands over the ghost to $2\text{e-}16$ and the refill to the bit; `mol_diffusion` marching 200 steps BITWISE (the marching path reaches the relaxation twin, never the residual). Built and measured in an isolated source tree with `steady_newton.f90` pinned, because N27 was editing it. Advisor: rebuild warning-free, `element_operator` 5/0, certification 29/0 on `build_lwv`; one percent fixed. Reported: the write-back's ghost is the cell's to round-off, not exactly (any other consumer of those ghosts reads a $2\text{e-}16$ gradient).

- **N27** (the helium row of cell 246 and the H2 row of cell 205 read side by side; `steady_newton.f90` +191/-3 diagnostic only, `steady_species_rows` one row restated): **the helium row of cell 246 is held by the LINEAR SOLVE, not by its physics, its boundary, a bound, the closure or the tolerance.** At outer 245 the row is $+2.43\text{e-}22 = \text{diffusive (settling)} + 2.40\text{e-}22$ plus advective $+3.3\text{e-}24$, same sign (they add); the Newton removed the advective term (a factor 7200 since outer 1) and barely touched the diffusive one ($\times 1.28$); the measure $6.17\text{e-}4$ is the maximum of a flat plateau ($5.9\text{e-}4$ at 240 to $5.4\text{e-}4$ at 253), not a bad cell; 0 held, 0 refused samples in 246 iterations, band healthy (`dgbcon` $7.5\text{e-}7$), merit reproducibility exactly 0, the trust region inert (ratio 0.933, 0 cuts, `aN` 1.0) and the Krylov cycle spends 40 of 40 products on a direction whose true relative residual is **0.9917** against 0.1 with the Arnoldi image 14.7 percent off the operator; the step buys $4.7\text{e-}5$ of the merit per iteration. The helium Jacobian triple is a proper tridiagonal stencil with the diagonal largest (the unknown controls its row; the wind dominates the row's entries). The element rows hold 0.98 of the merit (He 0.63, Fe 0.36); $||R||$ is the energy row at $1.37\text{e-}4$; the stagnation detector is honest (merit and $||R||$ fall while the judged distance rises $6.12\text{e}4 \rightarrow 6.21\text{e}4$). **A number that does not fit N25's reading:** the Arnoldi gap is 0.147 at $||sN||$ $1.8\text{e-}2$, 8 per unit against N25's $2.5\text{e-}4$ per unit, so the gap is not a function of the assembled length alone. **The H2 row of cell 205** is the same shape on a coarser front: measure 0.480 (not a cancellation, half its own terms), `f(H2)` falling $1.4\text{e-}1 \rightarrow 1.1\text{e-}3 \rightarrow 4.9\text{e-}4$ over cells 195-207 (logarithmic slope -757), `f(HI)` flat (1.7 percent), so cell 205 is NOT a two-root HI front (consistent with N5); linear 40/40 at 0.97, gap $3.1\text{e-}2$. Deliverable 3 STOPPED (nothing in the step control to change at the binding iterate). Deliverable 4: `cert_tol_element` would have to be $6.2\text{e-}4$ (4.7 decades; far above the $1\text{e-}5$ the operator's header gives as its loosest floor); the closure's reproducibility ($3.1\text{e-}7$ in the row's measure per unit seed, N5) is not what holds it. Proposals: re-measure the Arnoldi gap at three iterates (it no longer tracks $||sN||$); the untried freedom is the COLUMN preconditioner (row scaling is fixed by decision 20 a); whether the detector should stop a solve whose merit still descends while the judged row does not is an acceptance decision; the carrier front is a grid statement. Tests 189/0, 174/0; advisor rerun 189/0, 174/0. **Incident, found by N27 and CONFIRMED by the advisor:** with N26b's ghost refill in `element_transport_residual` the atomic element reload parts from N26's run at outer 10 and stands at $||R||$ 1.199 at iteration 40 where N26's run was at $3.21\text{e-}2$ (advisor run, cap 40, current tree: 1.888, 1.891, 1.677, 1.484, 1.518, 1.582, 1.199 at iterations

1, 5, 10, 15, 20, 30, 40 against N26's 1.888, 1.891, 1.678, 1.955, 1.758, 0.912, 0.0321); N27's own diagnostic is inert (hook on = hook off). N26b's validation (the first two iterations, the certification suite, mol_diffusion marching) did not reach it, and the advisor's verification of N26b did not rerun the reload: the advisor's mistake. N26b's refill is therefore NOT accepted as it stands; N26c (launched) finds why an operator that poses its own boundary breaks a solve whose caller already hands it that boundary to 2e-16, or reverts the refill. N27 restated the row a_stale_upper_ghost_is_read_ by_the_outmost_row to its negation because N26b made the original false; N26c reconciles the two rows with whichever text stands.

- N26c** (why N26b's ghost refill broke the atomic solve; binary_element_diffusion.f90 refill REVERTED with the header restated, element_operator rows restated (6), N26's steady_species_rows row restored): **the refill changes the residual by one element's outermost row in its twelfth digit and nothing else, and the atomic element reload does not survive a last-bit change of the upper ghosts, whatever makes it.** MEASURED: the caller's ghosts already equal the cell to 1 to 4 units in the last place (Xhe 9e-16, fX 1.9e-15, wYtr 6e-16 over the whole solve); the two texts' outermost rows agree to 17 digits on every element but nitrogen's, which differs by 9.8e-13 relative (6.8e-17 in its own measure); no other ghost of the routine is read (face coefficients stop at N-1); the band's stencil already carries the dependence of rows N-1, N on cell N through the write-back. **A control that multiplies the same three ghost quantities by $1 + 2e-16$ and poses no boundary at all breaks the reload as badly** (||R|| at iteration 40: caller's ghosts $3.212e-2 = N26$ to the digit; refill 1.199; one-ulp control 1.125; the three part at outer 10-11 and from each other). Where it amplifies: the Krylov leg at its one-digit tolerance (relative residual reached $8.86e-2 / 8.89e-2 / 8.93e-2$ at outer 10, pred 0.4 percent apart, then 2.7x apart at outer 11, then different branches of the acceptance at outer 12: accepted 0.874 / REFUSED -1.98 / accepted 85); the solve's own merit reproducibility prints $1.4e-14$, a hundred times the ghost movement. **The refill is physically right and is reverted anyway:** no version of it can be bit-inert on the production caller, and the reload's acceptance is a trajectory no last-bit change preserves. Delivered text reproduces N26's atomic reload line for line over 252 records (||R|| 3.719e-4, judged 6.22e4, He of cell 246, 27543 samples, exit 2) and N26's carrier reload over 63 (0.2519, 4.806e7, H2 of cell 205); mol_diffusion marching bitwise; suites certification 29/0, element_operator 6/0 (restated to ASSERT the measured caller dependence, 1.194e-2, so the suite exits 0; the advisor accepts that choice), steady_species_rows 189/0, krylov_and_dogleg 174/0; advisor rerun of the four on build_lwv (md5 3201310a, the worker's delivered binary): 6/0, 29/0, 189/0, 174/0. **The finding that matters beyond this item:** the atomic element reload is NOT a reproducible quantity; its ||R|| at a fixed iteration count is set by which Arnoldi vector crosses a one-digit tolerance at the last bits of the residual, so "reproduces the previous run" rejects correct changes as readily as wrong ones, and what the reload can carry is named outcomes (binding rows, refusal counts, flux spread), not ||R|| at iteration N. Recorded in ISSUES 3.1 (d) and in the fixture's README. The asymmetry of the element operator (the relaxation twins write their ghosts, the residual takes the caller's) stays in the tree, documented and measured ($1.19e-2$ on a stale caller). Advisor: a cp -irp of the N4b worker hung on its interactive prompt since 2026-09-09 (PID 61163, cwd the repository root, copying from the session scratch) was killed by PID. CORRECTION within the hour: killing the cp let the worker's shell run its NEXT statement, which re-applied N4b's first (discarded) patch onto the delivered element_inventory.f90 (832 -> 926 lines, a duplicate routine, would not compile); the N4b worker,

woken by the stale task, rebuilt the file from its saved entry copy plus the delivered patch and it reproduces `diff_element_inventory.diff` line for line; the advisor confirmed 832 lines, one routine, a warning-free build (md5 3201310a, N26c's delivered binary) and `element_operator` 6/0, `steady_species_rows` 189/0. Lesson written into `docs/worker_rules.md`: `\cp -f / \mv -f` always, and a stale command is canceled as a task, not killed as a process.

- **Direction after N26c (user, 2026-09-10: "권고안대로 진행").** The advisor's recommendation adopted: (b) N8b, anchoring the species-row certification tolerances on the two candidate states by the review's five measurements (reproducibility floor, derivative convergence, a manufactured steady solution of the element operator, grid refinement of the candidate state, conserved-flux error), reported per row class and regime with a PROPOSAL for the user (decision item 22) and no value moved; and in parallel (c) N28, the code items of ISSUES 3.7/3.8 (run-wide JFNK cap, one `element_ratio_gate`, the update map's install branch owing an `Apply_BC`, the undefined innermost ghost column of `assemble_residual`, the PTC route's setup report, `sec_ion_step` on a re-evaluated state, the stale test header, `Run` mode in the input schema). (a), the column preconditioner, waits for N8b's numbers. The user manual PDF was rebuilt by the advisor (85 pages; it was two keys behind its source: `Restart` intent, `Restart` option change).
- **N8b** (anchoring the species-row certification tolerances; `certification.f90` +173 additive, no value moved; `steady_newton.f90` one diagnostic; `certification` suite 29 -> 42 rows; memo `docs/certification_tolerance_anchoring_20260910.md`): **1e-8 is refused by no floor of the arithmetic, the closure or the representation (five to eleven decades below it); what refuses it is the DISCRETIZATION and the ELEMENT FLUX CONSERVATION.** MEASURED on the two candidates: reproducibility exactly 0; closure-seed floor $5.3\text{e-}15$ (the state's own sweep seed $1.7\text{e-}10$); state-perturbation amplification 0.23 to 0.30 in the wind but $1.1\text{e}3$ to $2.0\text{e}3$ at $r < 1.01$ (floors $5\text{e-}17$ and $4.4\text{e-}13$); derivative: a clean V at the base ($4.5\text{e-}7$), a PLATEAU at $6.4\text{e-}5$ over four decades of probe length at the binding cell 245 (the part of the derivative outside the band) so the smallest row a Newton step resolves there is $3.8\text{e-}8$; manufactured relaxed column at 801/401/201 cells $5.4\text{e-}13$ / $2.47\text{e-}6$ / $7.42\text{e-}6$, order 1.59 (the flat column $2.7\text{e-}14$); grid refinement of the candidate at r 1.1515: the He/H row $5.97\text{e-}4$ ($N = 500$) -> $2.48\text{e-}4$ ($N = 1000$), remap control $1.6\text{e-}5$, so **about 78 percent of the $6.0\text{e-}4$ is discretization** (a C2 interpolant is needed; below r 1.03 the remap dominates); element flux spread 39.3 in the layer, $2.6\text{e-}2$ in the wind, $2.2\text{e-}3$ over the escape window against the $5.5\text{e-}7$ mass-flux spread the candidate is quoted with. Memo section 7 is the proposal (decision item 22, added by the advisor to `To_be_determined`): (a) $1\text{e-}5$ in the wind with the layer reported, (b) keep $1\text{e-}8$ and state the fixtures cannot certify, (c) two regimes gating; no proposal certifies either candidate. **Two defects found outside the item's files, reported:** (1) **the candidate's species do not carry the run's own density:** mass closure $6.68\text{e-}3$ at the top against $1.9\text{e-}9$ on the three-unknown control; reproduced in the suite on a column closed to $6\text{e-}16$ that reads $3.37\text{e-}3$ after `ONE relax_element_composition`, so `project_elements` does not return to hydrogen the mass it takes from helium (N29, launched); (2) the writer-to-loader round trip does not return the state: the atomic candidate reloads at `||R||` 0.486 instead of $3.7\text{e-}4$ because `load_IC` rebuilds `rho` from the species (a consequence of 1). Tests 42/0 (RED at the build level: the suite does not compile against the entry

objects); the default certification report byte-identical (anchoring under EXHALE_CERT_ANCHOR=1). Advisor: rebuild warning-free, certification 42/0; one neighborhood and two percent fixed.

- **N28** (the code items of ISSUES 3.7; EXHALE_main.f90, RK_rhs.f90, steady_residual.f90, element_census.f90, element_inventory.f90 and attempted_step.f90 (the duplicate constant), grid_and_gates +2 rows and headers, docs/input_schema.md K45): (1) EXHALE_JFNK_MAXIT caps the RUN: the three call sites of steady_wind_with_element_diffusion pass one named default jfnk_outer_iterations_default = 3000 (the marching hand-off's literal was 500, the one default that moved; wasp_full_newton converges in 13 and cannot reach either), and under the variable the run takes its FIRST stationary solve and refuses later entries by name (MEASURED: three solves of 2 iterations -> one); (2) element_ratio_gate has one owner, element_census.f90 (public, element_census_tolerance() public with its override), read by element_inventory.f90 and attempted_step.f90; (3) update_map_begin_step's install branch calls Apply_BC in the N16d order (the EXHALE_UPDATE_MAP diagnostic's second block on mol_carrier moves 5 percent in the mass and energy rows: the fix of a defect, a boundary of one state under an interior of another); (4) assemble_residual defines the innermost ghost column (dF, S, R zero at 1-Ng; two callers read it in whole-array expressions); -fcheck clean on the certification driver and on a 20-step mol_carrier (the run that reaches it); (5) the EXHALE_PTC=1 route writes its setup report (was 0 lines; new row RED/GREEN); (6) a Restart intent: stationary evaluate keeps the file's sec_ion_step (2402 -> 2402, was 0; new row RED/GREEN); (7) restart_grid_guard.sh's stale header rewritten and the script made to honor EXHALE_TEST_OUT; (8) Run mode documented as K45. Item 9 (n_eq_sweeps_last on a discarded evaluation) is steady_newton.f90's, reported. Validation: mol_carrier 200 steps and wasp_full_newton cold BITWISE on every output file (the one log difference: cap 500 -> 3000, from the caller); grid_and_gates 146/0; attempted_step 162/1 with the FAIL proven pre-existing (rebuilt from entry sources: same row); advisor rerun on build_lvw: grid_and_gates 146/0, attempted_step **163/0** (the N28b fix below had reached the tree by then). **Found, reported, N28b launched:** bg_cell (type(ion_rates)) is allocated without initialization and the type has no default values, so before the first sweep the checkpoint checksum reads 8.7e244 and the row a_perturbed_state_is_not_the_checkpoint could never fire (a perturbation of 1e2 lost in the round-off of 1e244). Advisor: three forbidden phrases fixed (per-element in a printed line, neighboring, per-region). One deviation stated by the worker: an extra private -fcheck build (the shared run_fcheck.sh runs make distclean).
- **N28b** (bg_cell allocated and never initialized; ion_cell_state.f90 the type only, ionization_equilibrium.f90 the allocation site, attempted_step suite +1 row and a deterministic dirty-heap setup): **the defect is real, wider than bg_cell, and includes a LOGICAL.** Three grid-sized objects of ionization_equilibrium came out of allocate holding the recycled block: bg_cell and ieq_rate_cell (both type(ion_rates), no default values) and the real array heat_chem (the one array left out of the routine's zero list). MEASURED with a probe that frees a 16 kB block before the allocation: T_K, P_HI 8.7e244 and **x_h2_fixed TRUE** (an imposed H2 partition the caller never asked for, read from garbage, standing in the array the carrier transport reads), on a fresh heap all zero (which is why N28 saw the dead

row fail and the advisor's rerun saw it pass: the defect appears only when a freed block is recycled). Fix: ion_rates carries its zero in every one of its 37 fields (34 reals, 3 logicals) as default initializers, stated as the invariant "a cell not yet swept has no gas, no rates and no imposed partition", which reaches bg_cell, ieq_rate_cell, bg_cell_adopted, bg_cell_best, the closure scratch and the four checkpoint copies at once; heat_chem = 0 at the allocation. Two comments that claimed a default initializer would not reach threadprivate copies were corrected (MEASURED with a four-thread program: it does, as OpenMP specifies). Tests: RED on the entry text with the new driver 68/2 (the_unswept_background_carries_no_rates -3.4e250; a_perturbed_state_is_not_the_checkpoint h1 == h0), GREEN 164/0 (the intermediate build located heat_chem at 8.7e244 by a field-by-field peek). mol_carrier 200 steps bitwise (every reader of the three objects is gated on bg_ready/ieq_rates_ready, set after the loops that write them; heat_chem has no physics reader, only the checkpoint); wasp_full_newton not run (atomic: the arrays never enter a physics path). Advisor: rebuild warning-free, attempted_step 164/0. Reported: three test fixtures enumerate all 37 fields of bg_cell by hand (now redundant) and element_census_tests.f90 set two fields and left the rest to the allocator (now covered by the type).

- **N29** (the element projection does not conserve mass; binary_element_diffusion.f90 +258/-113, element_operator suite 6 -> 14 rows): **the leak is the mass the write-back is told to hand back, not the write-back.** project_elements took the mixture mass as msum = m_1 n_H + m_He n_He with m_1 = mass_per_H_nucleus_without_He(), the RESERVOIR mass per hydrogen nucleus (the trace metals at melem_ab), while the cell carries the metals at whatever the trace transport left; the projection then wrote the difference into hydrogen at every call. MEASURED at one cell of the certification column (metal/H 0.024401 against the reservoir's 0.028227): sum m_i f_i 1.0000000 -> 1.0000636 after one step, helium's gain exceeding the hydrogen and metal loss by 6.4e-5: mass created. A second leak of the same kind: the trace-metal diffusion wrote metal densities with no counter-flux in the hydrogen it diffuses through (3.0e-3 on the suite column with the metals frozen). **Fix:** mixture_mass_split reads the mass the cell's species actually carry (bsp_mass and melem_A, no species weight restated), project_elements(f_sp, Xhe, msum, scale_metals) scales component 1 as a whole, every helium-free species included (CO too), the metals written first and the hydrogen group closing the mixture; one projection after the metal loop puts the metals' mass change back on hydrogen; settling_coefficient takes its component-1 mass from the same split. Metals off is bitwise by construction (m_1 = m_H = 1 exactly). **MEASURED:** certification column closure after the relaxation 3.37e-3 -> 2.3e-15; the handed-back atomic candidate 6.44e-3 -> 1.94e-9 (the file's own precision, the three-unknown control's 1.93e-9); the writer-to-loader round trip of that state now returns it (writer ||R|| 1.112 and reload 1.1123 on the SAME row, against 3.2e-2 vs 0.4385 on different rows before: memo section 6 closed); mol_diffusion and wasp_full_newton BITWISE (metals off / no element row); **lower_profile MOVES and should:** its stored golden is 3.6e-4 out of closure (O/H 4.61e-4 against the reservoir 4.90e-4), a 1500-step cold start goes 6.77e-3 -> 3.14e-9 in closure with T at 1.6 Rp moving 1.3 percent and H I 6.3 percent: **a golden that moves for a physics fix, refreshed at the next gate.** Atomic reload cap 40 by named outcomes: closure 6.44e-3 -> 1.94e-9, 32/8 -> 31/9, refusals 0 in both, binding row energy of cell 2 -> cell 4, flux spread 2.3e-3 -> 1.4e-2, same stop. Tests RED 4 rows on the entry text (3.44e-3,

4.78e-4, 3.44e-3, 3.82e-4), GREEN 14/0; certification 42/0, diffusion_tests 35/0, krylov_and_dogleg 174/0. **The worker's control discipline:** the advisor's EXHALE_lwv.x is rebuilt at every verification and is a build of the live tree, so the worker built its own control (the delivered objects with only the entry-text binary_element_diffusion.o) and discarded three comparisons that had used the advisor's binary as "before"; workers are to build their own controls (added to the rules below). **Advisor:** steady_species_rows_tests.f90 referenced element_ratio_gate through element_inventory, which since N28 imports it with use, only and does not re-export it, so the suite would not build (N28's validation did not include this suite; the advisor's did not either): one use element_census, only: element_ratio_gate line added by the advisor; rebuild warning-free (md5 a9627a14), element_operator 14/0, certification 42/0, steady_species_rows 189/0, krylov_and_dogleg 174/0; one percent fixed. Reported: the metal diffusion's counter-flux is closed on hydrogen but not on momentum or internal energy (a trace approximation, named); element_census_verify gates the element ratios and only reports the mass closure (the 6.4e-3 state passed every gate it has); whether it should gate is a decision.

- **N8c** (the five anchors re-measured on the states after N29; the memo docs/certification_tolerance_anchoring_20260910.md only, 344 -> 524 lines, no source file changed): **the reading is unchanged and is now taken on states worth measuring.** Post-N29 candidates (advisor-verified: no src file newer than the brief): atomic reload 275 iterations, 254/22, ||R|| 2.271e-4, binding row He/H of cell 245 (r 1.153), flux spread 9.8e-7, **mass closure 1.95e-9** (was 6.68e-3), 11 of 13 entries refuse; carrier reload byte-identical to N26's (no metals, no element row; the corrected projection is never reached), closure 1.93e-9. The anchors: two evaluations of one state 0 exactly; the state's own sweep seed 1.7e-10 -> 1.6e-14 (the loader no longer repairs a composition whose mass missed the density), closure floor 5e-19; state-perturbation amplification 1.2e3 to 2.0e3 at $r < 1.01$ and 0.4 to 0.7 elsewhere; derivative: a clean V at the base (7.6e-8), the PLATEAU at 6.6e-5 at the binding cell stays (smallest resolvable row 3.9e-8 there, 6.5e-9 in the wind; both taken at the first JFNK iterate of a relaxation restart, two CFL steps from the loaded state, stated); manufactured column order 1.589, closure 2.3e-15 (was 3.4e-3), extrapolated 1.52e-6 at the production spacing; grid refinement of the candidate 5.93e-4 ($N = 500$) -> 2.47e-4 ($N = 1000$), ratio 2.40, remap control 1.5e-5, Richardson limit 1.31e-4: **78 percent of the row is discretization**; element flux spread in the layer **39.3** -> **9.79** (He), 37.3 -> 9.09 (H), the wind 2.6e-2 -> 2.7e-2, the escape window 2.0e-3. Memo: sections 3 (mass closure) and 6 (the reload) marked CLOSED by N29 with the readings; section 5 rewritten; section 7's proposal re-stated with the new floors and **option (a) supported as written** (1e-5 gating at $r \geq \text{cert_regime_wind_r}$, the layer reported and not gating); two of its three prerequisites discharged, the layer's flux conservation (9.8) standing. No proposal certifies either candidate (wind element row 3.05e-4 and 7.2e-2 against a proposed 1e-5). certification 42/0. Reported: the iron wind row of the atomic candidate rose 18x between the two chaotic states (a size, not a trend); the derivative anchor is not measured at the loaded state itself (needs the probe under stationary evaluate, steady_newton.f90).
- **Decision 22 (user, 2026-09-10): option (a).** cert_tol_element and cert_tol_carrier become 1e-5, gating, for cells at $r \geq \text{cert_regime_wind_r}$ (1.20); the layer's element and carrier rows ($r < \text{cert_regime_layer_r}$,

1.10) are reported and do not gate until the layer's element flux conservation (9.8 after N29) is repaired; the hydrodynamic tolerances do not change; a wind-certified state says so in one line. Anchor: the operator's stated floor and a decade above the manufactured column's discretization at the production spacing (1.52e-6); no existing state is admitted by it (3.0e-4, 7.2e-2). Implemented as N30 (launched).

- Gate after N29 and the lower_profile refresh (advisor, make check 2026-09-10 07:58 to 11:03; refresh 11:10).** Tree after N28, N28b, N29 (N30 running on a private build). Against the 2026-09-10 00:40 goldens: **15 of 16 default cases data identical on all four files** (wasp_full_newton now identical too, its golden being the 00:40 snapshot); **lower_profile FAILS on all four, by design** (N29's mass fix: its golden was 3.6e-4 out of closure; v moves up to 4.8 percent at r 3.18 in 85 of 504 rows, rho 1.0 percent in 4 rows, T 0.5 percent in 2 rows; the _adv measure column 57 percent at one row; log10 Mdot 7.29). The failing case was identified by its own section of the log (the advisor's first parse attributed the summary line to hp_front, which PASSES). Refreshed: golden/lower_profile archived as golden_pre_n29_20260910/lower_profile, run_check.sh golden_lower_profile snapshotted the gate's output, the four files compared identical to the case output with the comparator's filter. No other golden touched.
- N30** (decision 22 a; certification.f90 +296/-54, steady_newton.f90 +50/-30, certification suite 42 -> 57 rows, steady_species_rows 189 -> 194, docs/input_schema.md the cert_reason row): cert_tol_element and cert_tol_carrier are gone as single numbers; cert_tol_element_wind = cert_tol_carrier_wind = 1e-5 with one accessor each (cert_tol_*_at(r), a step at cert_regime_wind_r = 1.20; cert_tol_reported_only = huge below), the anchoring written at the declaration with the memo cited; **one reduction** certification_species_row_gate decides a species row over the GATED cells (no gated cell = refused, never a satisfied row); the entries carry gated and reported maxima with their cells; the report prints gated at r >= 1.2 against 1e-5: <measure> at cell <j> and, for a certified state with species rows, the one line certified IN THE WIND ... the rows below it are reported and do not gate: worst <measure> at cell <j>; the state file carries cert_reason=certified_in_wind (the field already existed). **The band 1.10 <= r < 1.20 is reported and does not gate** (the decision was silent; the band holds the candidates' binding cell at r 1.153 where the discretization is 4.6e-4, so a 1e-5 gate would ask the residual to fall 1.5 decades below the error of its own discrete equation; no anchor for the band exists). distance_from_certification and certified_row_measures read the gated measures and the accessors, so **the trust region's ranking changes and both reloads move** (that is the decision's intended reach): atomic reload 275 -> 166 iterations at the same log10 Mdot 10.10, binding row He/H 2.896e-4 at cell 288 (a WIND cell) against 1e-5, NOT certified; carrier reload 63 -> 76 iterations, H2 7.284e-2 at cell 285 against 1e-5, NOT certified; the hydrodynamic tolerances, the H(n=2) level balance (4.8e-10 against 1e-6) and steady_gates_met's own carrier gate (carrier_resid_th, a volume ratio, not this tolerance) untouched. wasp_full_newton bitwise, still certified with no reason. Tests RED (the superseded rule refused a layer-1e-2 / wind-5e-6 row and a wind-5e-5 row alike, on a layer cell; four judged-distance rows moved onto the new divisor), GREEN 57/0, 194/0, krylov_and_dogleg 174/0. No key added (the radii are properties of the flow, constants of certification.f90). Advisor: rebuild warning-free, 57/0, 194/0, 174/0 on build_lwv; the stale comment in EXHALE_main.f90 (580) rewritten; write_output.f90's

`adv_input_certified` T branch now carries the reason (a wind-certified state keeps its qualification in the `_adv` header); superseded notes appended to `a2_certification_contract`, `steady_solver_design` and `element_inventory_contexts`; one neighborhood fixed. Reported: whether `steady_gates_met` should read the anchored tolerance is a separate decision on a shared interface.

- **DOC1 to DOC3** (the documentation brought to the stage-2 state, 2026-09-10, three concurrent documentation workers, no source touched, no binary run): **DOC1** `EXHALE_BC_and_IC.tex` (29 pages): the whole lower-boundary chapter described the retired component-wise ghost closure (Dirichlet density, one-way valve, isothermal ghost pressure, four keys that `input_read.f90` now refuses with error `stop`); rewritten as the characteristic (p, s) face condition of `base_boundary.f90` (branches by face Mach number, the smoothstep of width `base_face_mach_blend`, the Gauss-Legendre ghost averages, the three surviving controls), the upper boundary as `free_outflow_ghost`'s isothermal continuation, a new section on the transported columns' boundaries (N26, N26c, N8a, N29), the checkpoint rebuild order (N16d), and the old test matrix retitled as the retired closure's, not reproducible by today's binary. `composition_restart_and_base_handoff.tex` (18): thirteen base species in the nuclei count (OH, H₂O, CO added), every `load_IC.f90` line citation re-derived, a new section for the restart contract (intent, the eight-line metadata block field by field with refuse/report on each, provenance unknown, the grid guard, `Restart option change`), the round-trip and projection evidence of N9 and N29 labeled as read from this log. `transmission_spectrum.tex` (11): a new section for `read_adv_validity` (schema 2, the five statuses, `adv_conditional_tol`), the `chord_shell_indices` census with the CONTRIBUTION statement, and the `transit_schema 1` block. **DOC2** `steady_solver_memo.tex` (16): a new section 7 for the species-row solve as it stands (merit on certification scales, the trust region with named outcomes and the leg rules, the constraint rows, the empty-column repair and the equilibrated preconditioner, the three options measured and disarmed, the linear solve as the limiter with the additivity finding, the ulp-level chaos, the residual contract, the certification by regime), eleven hooks added to its table. `molecular_hydrogen_treatment.tex` (23): the `_adv` row corrected (the energy residual reads the upstream caloric energy since N12), the layer table RE-MEASURED on the current `mol_lyman_werner` output (144/184/476 cells, H₂ peak 5.15e12, 1 percent edge 4.48e9 against the stale 227/251, 4.63e13, 4.42e9), thirteen of sixteen cases pinned, a new section on the element budget, the H₃⁺ departure factors marked STALE in place, the `mol_ir_bands` peak re-measured (2.148e-8 at r 1.0059 against the stale 2.934e-7). `lower_atmosphere_coupling.tex` (28): a new section 7.4 on the element operator (N29's conservation, N26b/N26c's boundary, N8c's flux spread table as the open item); two pre-existing overfull boxes fixed. `EXHALE_user_manual.tex` (88): certification by regime in section 2.7, the `Run mode` key entry, eleven hooks, the coupling header example replaced by the line the code writes (`cert_reason`), `Cooling_breakdown.txt` columns 12 to 14 (H₂, H₂O, CO infrared) restored to the column list. **DOC3** `README.md` (sixteen-case matrix, 25 suites, the restart contract, the transit metadata, certification stated, nineteen examples), `README_HOWTO.md` (`Resid tol` default 1e-5 not 1e-3, `Flux spread tol` default 2e-5 on the face flux not 5e-3, a certification table, a restart section, an eleven-row hook table, the sixteen cases and 25 suites), the `NEW docs/code_status_20260910.md` (418 lines, replaces the 2026-09-05 file as the read-first document; section 3.3 states what is and is not certified and what limits the rest), the project `CLAUDE.md` pointer line, `TO_BE_DONE.md` (an index of what is open, three headings corrected, 35 em-dashes removed),

steady_solver_design.md section 22. All eleven PDFs rebuilt, `grep -c '^!' 0` on every log. Advisor: every default and constant quoted checked against the source (`flux_spread_th_default 2.0e-5`, `resid_max 1e-5`, `cert_tol*_wind 1e-5`, `cert_regime_wind_r 1.20`, `ratio_sound_band 1e-2`, `tr_delta_min_dimensionless 1e-6`, the `Cooling_breakdown` header); one constant name corrected in the memo; pre-existing British spellings fixed in the workers' files, in `ISSUES_20260909.md` and in the handoff; the 51 em-dashes of the workspace `CLAUDE.md` replaced. **Reported by the workers and CONFIRMED by the advisor** (ISSUES 3.7): the base reservoir is stated once at startup and the refreshes of `dp_bc` and `ntot_bc` after every sweep are dead writes (the boundary itself is right: the reservoir equals the user's base pressure by construction); `golden/mol_ir_bands/` carries no `Cooling_breakdown.txt`; `examples/README.md` covers part of nineteen folders under the wrong planet; the .md twins of two .tex documents have parted from them; the `max(rest, 0)` write-back clamp (N4b) restated. ISSUES section 5 rewritten to the state after N30; the handoff opens with a dated 2026-09-10 state block. The carrier reload is still a scratch recipe (`B5k/mkrun.sh carrier`), not a pinned fixture: to pin with N31.

- **N31** (why the finite-difference Jacobian action is not additive across directions, and two probe options; `steady_newton.f90 +563/-3`, of which the arithmetic is two probe rules and the rest is a default-off hook, `krylov_and_dogleg 174 -> 190` rows, new fixture `backup/regression/carrier_elem_newton`): **the action is not additive because the residual carries a NON-SMOOTHNESS FLOOR five to nine decades above the double-precision rounding of its own rows, and the difference quotient along the PRECONDITIONED Krylov directions divides that floor by an increment five decades smaller than a generic direction produces.** N25's reading (curvature over the fixed arc) is refuted at the iterates that matter and confirmed at one that does not. MEASURED on the atomic element reload with the new hook `EXHALE_JV_ADDITIVITY=1` ([diag 13], at outer 1, 20 and the last of a capped run, on the first two Arnoldi directions and on two directions off the basis, at a tenth, one and ten times the probe arc): the defect of the Arnoldi pair scales as the arc to the power **-1.026** at outer 1 and **-0.999** at outer 40 (rounding), the off-basis pair as **+1.125** and **+2.109** (curvature, and $3.2e-8$ and $2.1e-8$, negligible), and at outer 20 both are curvature at cells 1 and 2. The second difference of the residual on the defect-holding rows confirms it directly: ratios 0.774 and 1.676 at outer 1 where a smooth residual gives 4 and 4, and the floor stands **9.5e5** (outer 1) to **2.0e9** (outer 40) times epsilon times the residual on those same rows, so it is not the arithmetic of the assembly. READ, the only tolerance in the atomic path that can set a floor of that size: `ionization_equilibrium.f90` line 969 hands `hybrd1` an `xtol` of `sqrt(dpmpar(1)) = 1.49e-8` (lines 1432, 2110, 2114), a statement about the step and not the residual, so the composition is a piecewise map of the state; `eval_residual` fixes the number of composition passes but not `hybrd1`'s own iteration count. **A decision of the feasible set is ruled out:** on all six probes of the three iterates, components blocked 0, forward side, and the step taken equal to the nominal step. **The localization gives no unknown class to fix:** at outer 1 the squared defect is 0.31 mass, 0.64 energy, 0.03 momentum, 0.02 helium, under $1e-4$ in the eight trace elements, and nine tenths of it sits in 35 of the 500 cells between 1 and 75. The scale reading is a uniform shortfall: the largest unknown of the state is displaced by $2.2e-8$ of itself (about `sqrt(epsilon)`) and almost every other by $1e-10$ to $6e-9$, one to two decades short, with no unknown near a bound (nearest face $2.9e-5$ against a displacement of $4.5e-14$). The model that follows, `defect ~ floor / (eps || Jv ||)`, predicts both pairs at outer 1 within a factor three. **Two options,**

both default off. EXHALE_JV_COLUMN_SCALE=1 (probe_step_on_the_column_scales, $\epsilon_0 = \sqrt{\epsilon}$) ($1 + ||Dc^{-1} Y||$) / ($||Dc^{-1} v||$) is the brief's remedy and it is **WORSE**, for the reason the diagnosis gives: the Krylov direction's weight sits on unknowns of small column scale, so the rule **SHORTENS** the arc by 42 and the defect rises $8.430e-6 \rightarrow 1.466e-4$ at exponent -0.965 , the $1/\text{arc}$ law reproduced by a change of the arc made for another reason. EXHALE_JV_PROBE_ARC=<x> (jv_probe_arc_scale, default 1) is the lever the measurement points at, and it wins the linear question and loses the Newton by the same change: at $x = 100$ the defect of the Arnoldi pair falls $8.430e-6 \rightarrow 2.240e-6$, the exponent turns $+1.139$ and the second differences become 4.065 and 4.022, so the rounding regime is gone and the quotient is truncation-limited, while the atomic reload then **ABORTS** at outer 33 (info = 2, $||R||$ 1.504, flux spread $5.902e-2$, 12 steps refused by the model with the actual reduction 26.4 times the predicted, radius $2.028e-7$) where the control runs 144 iterations to $3.641e-4$ and $5.557e-6$ with no model refusal. At $x = 20$ it aborts at outer 43 at 1.524 with 21 model refusals and a radius of $2.076e-13$; the carrier reload runs its 40 iterations at both values and hands back 1.006 at $x = 100$ and $9.076e-1$ at $x = 20$, each with 2 model refusals, against the control's $9.572e-2$ with none. **At $x = 5$, which does NOT stall**, the binding regime is reached (outer 58, $||R||$ $3.087e-3$, element Fe of cell 146, against the control's $3.386e-3$ and element Fe of cell 147 at outer 60) and **THERE** the Krylov cycle still spends 40 of 40 products at a relative residual of 0.898 to 0.980 against the 0.1 asked, indistinguishable from the control's 0.935 to 0.959. **The substance of the item:** the additivity of the linear operator wants a **LONGER** arc (defect $\sim \text{floor} / \text{arc}$) and the trust region's model wants a **SHORTER** one (truncation $\sim \text{arc}$, and the ray test refuses a step whose model and true slope differ in size), the arc already sits at the Newton's balance point, and every direction was measured to lose, so **NO** probe rule fixes this. What can is lowering the residual's own floor, which is not double precision. Column-scale named outcomes at cap 40, option against control: atomic $||R||$ at 40 1.674 against $7.607e-2$, handed back 1.633 against $5.377e-2$, flux spread $4.985e-1$ against $3.555e-3$, 30/10 against 32/8; carrier $||R||$ at 40 $1.357e-1$ against $1.070e-1$, handed back $1.263e-1$ against $9.572e-2$, flux spread $7.718e-2$ against $4.635e-2$, 31/9 against 33/7, **SAME** binding row (carrier H2 of cell 205), no new gm_no_direction_sampled and no refusal of any class in either. The one thing the column scaling improves is the count of Krylov cycles reaching the tolerance asked (40 of 40 against 26 of 40), which is the **REDUCED** problem's residual on an operator that is not the Jacobian and is precisely why that count must not be read as progress. **NEITHER OPTION IS ADOPTED:** both fail the first clause of the acceptance rule outright. **Default path unchanged:** with every hook off the atomic reload's 40 iteration lines are identical to the control's to the digit, the carrier reload's binding row and $||R||$ at 40 are the control's, and wasp_full_newton is BITWISE but for the provenance timestamp; with EXHALE_JV_ADDITIVITY=1 on, the 40 iteration lines are still identical, so the hook is inert armed as well as disarmed. Tests: krylov_and_dogleg **RED** on the control objects (seven symbols absent), **GREEN** 190/0; steady_species_rows 192/0, certification 57/0, element_operator 14/0, residual_determinism 5/1 with the same row failing by design in the control build. **Fixture pinned (deliverable 0):** backup/regression/carrier_elem_newton now holds the carrier reload that existed only as a scratch recipe, with run.sh, a README and the control of the unmodified tree at cap 40 ($||R||$ 1.969, 1.984, 1.921, $6.800e-1$, $1.070e-1$ at 1, 5, 10, 20, 40; handed back $9.572e-2$; flux spread $4.635e-2$; info = 1; 33 accepted, 7 rejected; 40 of 40 products every cycle; binding row carrier H2 of cell 205; 2754 samples admitted, none refused). A capped run of that fixture never reaches a mass-loss rate. **Defect**

fixed on the way: the new probe rule was first written with a local integer `n`, which is the grid size `N` of `global_parameters` under Fortran's case-insensitive rules, so its own size guard fell through on every call; the new suite rows caught it at once and the local is now nunk. Not claimed: that the floor is `hybrd1`'s `xtol` -- that is `READ` from the source and not measured, and the experiment that closes it (tighten that `xtol`, re-measure the floor) belongs to `ionization_equilibrium.f90`, which this item does not own. The comment of the suite row `the_probe_arc_stands_above_the_residual_noise` was restated: the arc IS nine decades above the residual's reproducibility as a length of the whole vector, but the conclusion drawn from it about the quotient does not follow. Advisor: rebuild on `build_lwv` (one linker warning, pre-existing), `krylov_and_dogleg` 190/0, `steady_species_rows` 194/0, `residual_determinism` one row FAIL by design; atomic reload at cap 40 with every hook off, 8 threads: 1.888, 1.891, 1.819, 1.583, 1.861, 8.964e-1, 7.607e-2 at 1, 5, 10, 15, 20, 30, 40, handed back 5.377e-2, flux spread 3.555e-3, identical to the worker's control; the carrier fixture's README control checked against its own `run.log`; the fixture README is the reference for every later carrier-reload number (the 0.25 quoted after N26 was a scratch-recipe run, not this fixture). Accepted. **Next item N32:** the residual's floor (the inner solves of the composition and of the temperature stop on step or bracket tolerances, `newton_dense`, `hybrd1` `xtol` `sqrt(eps)`, `brent_root` 1e-10); measure whether the additivity floor follows them.

- **N32** (which inner tolerance sets the residual's non-smoothness floor; two default-off tolerance hooks and a widened [`diag` 13], `ionization_equilibrium.f90` +47/-2, `T_equation.f90` +42/-3, `steady_newton.f90` about +130 inside the additivity hook, `krylov_and_dogleg` 190 -> 203 rows): **NEITHER. The floor follows neither the composition solve's stopping tolerance nor the temperature bracket's, and it is not in the chemistry at all: it sits in the HYDRODYNAMIC rows of the residual.** New hooks `EXHALE_IEQ_TOL=<x>` (replaces `tol = sqrt(dpmpar(1))` for the run; new function `composition_solve_tolerance`, cached once by `ioniz_eq`, announced only when armed) and `EXHALE_TEQ_TOL=<x>` (the bracket of `brent_root`, now a module variable at its old 1e-10 through `temperature_bracket_tolerance`). MEASURED on the atomic element reload at outer 1 (cap 40, 8 threads, one entry state for every run): the additivity defect of the Arnoldi pair is 8.174e-6 at exponent -1.063 (default 1.49e-8), 7.452e-6 at -0.995 (1e-10), 9.348e-6 at -1.059 (1e-12) and 9.546e-6 at -1.064 (1e-14), and the second difference on the defect rows stays on its floor at 5.6e-12 to 1.6e-11 with no trend in the tolerance: **the law is FLAT**, over six decades. The measurement is exactly reproducible (three control runs, and 1 thread against 8, give 8.174e-6 and 1.362e-11 to every printed digit), so the +-17 percent between runs is a real movement of the state and not a fall. `EXHALE_TEQ_TOL=1e-13` is inert to the bit through all 40 iterations, which confirms as MEASURED what was READ, that `brent_root` is reached only from `post_process_adv` and never from `eval_residual`. **Why the composition cannot be the floor:** quadratic convergence pins its root at the SQUARE of the step tolerance. On the reported cell the accepted `x(H+)` is 2.037266074305163e-12 at the default and 2.037266074305300e-12 at 1e-14, agreeing to 13 significant digits (7e-14 relative), and the pinned fixture's own `run.log` records an accepted reaction residual of 2.91e-16 at maximum for the steady sweeps; the floor implies a jitter of about 1e-9 relative, five decades above that. The tightest tolerance costs 30 percent more wall time per residual evaluation (0.090 s against 0.066 s at 4 threads, 17 evaluations each) and buys nothing. **Where the floor IS**, from the widened hook: the energy SOURCE, which is where excited_hydrogen, `lya_rt`, the cooling table interpolation, the write-back clamps and `comp_T_from_p` all end up, is smooth on the

arc and five decades below the floor (second difference of heat - cool on the defect cells $3.039\text{e-}17$, $2.474\text{e-}16$, $1.046\text{e-}15$ at half the arc, the arc and twice it, ratios 8.1 and 4.2, textbook h squared, against the residual's own $1.362\text{e-}11$ on the same rows); by row class over the whole column the floor is mass $1.801\text{e-}11$, energy $3.090\text{e-}11$, momentum $5.946\text{e-}12$ against element He $4.269\text{e-}12$ and every trace element at $1\text{e-}16$ to $2\text{e-}13$; nine tenths of it sits in 29 of 500 cells between cell 1 and cell 58. The mass row carries no source at all, so no part of the chemistry can be its cause. The competing explanation, that the row is a small difference of large terms and its floor is their rounding, is REFUTED: the largest of heat and cool on those cells is $3.424\text{e-}3$ against $||F||$ $5.507\text{e-}3$, a ratio of 0.62, and epsilon times it is $7.6\text{e-}19$. **Deliverable 2 (the stationary-solve-only polish) was NOT BUILT:** the brief conditions it on deliverable 1 naming a tolerance, and none is named. **A correction to the brief and to N31:** on this fixture `solve_ieq` and `newton_dense` never run. The helium triplet is on (the fixture's `EXHALE_setup.out`) and `metals.inp` is present, so the sweep calls `hybrd1(ion_system_HeH_TR_metals, ...)` directly; MEASURED `nt_calls` 0 over the whole hook and no "ioniz-eq solver" line in the fixture's reference run.log. `hybrd1` returns `info = 1` at the default, at $1\text{e-}10$ and at $1\text{e-}14$ alike, so there is no `info = 2/3` and no fallback to count, and `newton_dense`'s hardcoded step test $1\text{e-}11 * \text{xscale}$ is not in this path. **Default path unchanged:** `wasp_full_newton` BITWISE in `Ion_species.txt` and identical but for the provenance timestamp in `Hydro_ioniz.txt`, every (JFNK) line identical; the atomic reload's 40 iteration lines identical to the control's with every hook off. Tests: `krylov_and_dogleg` RED on the control objects (both new functions absent), GREEN 203/0 with 13 rows added; `steady_species_rows` 194/0, certification 57/0, `element_operator` 14/0, `adv_static_limit` 53/0, `coupled_source_step` 30/0, `physics_probe` 1491/0, `residual_determinism` 5/1 with the same row failing by design as in the control. Noticed, not fixed: the fixture README does not say the helium triplet is on, which is what selects the direct `hybrd1` branch and is what sent this item's brief to the wrong routine; and `newton_dense` applies the caller's tolerance to `fnorm` while its step test uses a hardcoded $1\text{e-}11 * \text{xscale}$ no caller can set, which would be the real floor on a fixture that does reach it. **Next item:** the floor is in the flux assembly, not in `ionization_equilibrium.f90` -- the candidates left are the reconstruction where the limiter is not the frozen WENO3 weight, the Riemann wave-speed estimate, `Apply_BC`, and the transport coefficients of the diffusion operator, which are the one remaining way the composition reaches a hydrodynamic row. Advisor: rebuild on `build_lwv`; `krylov_and_dogleg` 203/0, `steady_species_rows` 194/0, `adv_static_limit` 53/0, `residual_determinism` the same single row FAIL by design; atomic reload at cap 40 with every hook off, 8 threads: the N31 ladder to the digit (1.888, 1.891, 1.819, 1.583, 1.861, $8.964\text{e-}1$, $7.607\text{e-}2$; handed back $5.377\text{e-}2$, spread $3.555\text{e-}3$), so the worker's 1.112 at iteration 40 is the thread-count sensitivity of N26c, not a change of the path. The atomic fixture README now states the triplet is on and which branch that selects; `newton_dense`'s hardcoded step test recorded in ISSUES 3.7. Accepted; deliverable 2 correctly not built. **Next item N33:** the jump itself, scanned along the Arnoldi direction across four arcs and split by term and operator (base boundary, reconstruction, Riemann flux, caloric inversion, composition read, transport sources).

- **N33** (where the residual's non-smoothness floor is: the term, the operator and the branch; a default-off multi-scale jump scan [`diag 14`], `steady_newton.f90` +710, `krylov_and_dogleg` 203 -> 220 rows): **THERE IS NO BRANCH. The floor is the ROUNDING OF THE FLUX ASSEMBLY, raised above the last bit of the row by two cancellations that compound.** The Roe flux of a face is built from the JUMPS of the two

reconstructed states, and in the nearly hydrostatic base layer those jumps are $3.2e5$ to $5.9e6$ times smaller than the states themselves, so the interface flux carries the last bit of $O(1)$ quantities and not of its own value; the row is then the flux difference divided by the cell volume, $r^2/dV = 5.12e3$, one over the $1.955e-4$ width of a base cell. The bound $\epsilon \times (\text{face state}) \times r^2/dV$ is what a correctly rounded assembly cannot go below, and MEASURED it IS the floor: $1.675e-12$ against $1.240e-11$ at cell 1, $2.520e-13$ against $1.644e-12$ at cell 84, $4.326e-15$ against $2.366e-14$ at cell 167, $4.241e-17$ against $4.409e-16$ at cell 251 -- the bound falling 4.6 decades with the measured floor following it, ratio 5.5 to 12 throughout, and the ten cells holding the largest second difference are cells 1 to 23, every one a base cell. Beyond cell about 250 the bound falls below the residual's own curvature and stops binding, which is what a floor does. New hook EXHALE_RESID_JUMP_SCAN=1 samples F at 65 points of $Y + t \cdot v$ on ten windows, each a quarter of the one before, from two probe arcs down to two arcs over 4^9 ; one window is not enough, because at the standard arc every interval already holds thousands of steps and none stands above its neighbors. **The measurement:** on the atomic element reload at outer 1 the tracked row is the energy row of cell 1, its median first difference falls by exactly four whenever the spacing does (the smooth part is a straight line), and its second difference does not fall at all -- $1.0e-11$ to $1.6e-11$ over 5.4 decades of spacing, $1.8e-5$ down to $6.9e-11$ -- which is a step of F and not a kink or curvature. Isolated steps appear once the smooth increment per interval falls below the floor: 28 of 64 intervals at $3.05e-5$ arcs and 61 of 64 at $7.63e-6$ arcs, at least $4.0e6$ stepping intervals per probe arc, the largest moving its row by $2.625e-12$, $2.749e-9$ of the row. **The term and the operator,** at the largest step (cell 9, mass row): S of the mass row is identically zero and the whole step is in dF ; every quantity upstream of the flux -- the ghosts, T , n_{tot} + n_e and the four reconstructed face states -- moves by one to eight units in the last place of an $O(1)$ number with no outlier, in exact multiples of $2.220e-16$; the first quantity that steps is the interface flux, by $4.803e-16$ where its neighbors move by $3.2e-17$, and $4.803e-16 \times 5116.3 = 2.457e-12$ against the row's measured $2.625e-12$. **No branch fires anywhere near it:** faces falling back from Roe to HLLE 0 and unchanged, face states the positivity limiter scaled 0 and unchanged, so the base boundary's Newton, `characteristic_branch_weight`, the Mach cap, the ghost quadrature, the stencil selection, the wave-speed branch and `comp_T_from_p` are all continuous here. The off-basis direction carries the SAME floor, $9.1e-12$ to $1.4e-11$ flat over 5.4 decades, and isolates no step at any window, because its smooth increment never falls below it: the floor is a property of the residual and not of the direction, and what the preconditioned direction does is shrink the numerator of the difference quotient by five decades. The carrier reload reproduces the law (floor $1.532e-11$, bound $1.382e-12$, ratio 11.09), so it is the discretization at the base and not one fixture. **N31 corrected:** the floor's size stands, but "five to nine decades above the rounding of its own rows" compared it against $\epsilon \cdot |F|$ on the row, which is the wrong reference for a row built as a flux difference over a cell width; against the right one it is a factor of ten. N32's localization is confirmed and made quantitative. **Deliverable 2 NOT BUILT:** the brief conditions the deliverable on a branch being named and there is none. The treatment the measurement does name, for the advisor, is a WELL-BALANCED flux difference at the base -- assemble the rows from the departure of the face states from the local hydrostatic isentrope, which `base_boundary.f90` already integrates, so the balance cancels analytically instead of in floating point -- in `Num_Fluxes.f90`, `RK_rhs.f90` and `Source.f90`, none of them this item's files; the cheaper partial lever is the base cell width, on which the floor depends as

one over it. **Default path unchanged:** the atomic reload at cap 40 gives all 338 (JFNK) lines identical to the control's through done info=1 ||R||= 1.112E+00; with the scan armed the outer-1 line is the control's to every digit; wasp_full_newton BITWISE in Ion_species.txt, Ion_species_adv.txt and the three breakdown files, differing only in the provenance timestamp in the two hydro files, log10 Mdot 13.30 on both. Tests: krylov_and_dogleg RED on the control objects (resid_jump_scan_on absent), GREEN 220/0 with 17 rows added; steady_species_rows 194/0, element_operator 14/0, certification 57/0, residual_determinism 5/1 with the same row failing by design and the same measured 3.354e+01 on the control binary. Noticed, not fixed: global_parameters declares integer :: count, which shadows the intrinsic COUNT in every scope that uses the module, so count(mask) written anywhere fails to compile -- the same class as the t0 and g2s incidents and worse in that the shadowed name is an intrinsic; renaming a public global was out of scope. Advisor: rebuild on build_lvw; krylov_and_dogleg 220/0, steady_species_rows 194/0; atomic reload at cap 40 with every hook off, 8 threads: the N31 ladder to the digit (7.607e-2 at 40, handed back 5.377e-2, spread 3.555e-3), no [diag 14] line; two British spellings in the new comments fixed. Accepted; deliverable 2 correctly not built (there is no branch to smooth). The finding is structural and reaches the marching discretization, so it is put to the user as DECISION 23 (well-balanced flux differencing recommended; quad accumulation of the flux difference as the control experiment; coarser base cells; accept the floor). The count global shadowing the intrinsic recorded in ISSUES 3.7 (HYG). **Next item N34** (runs without a decision, changes no default): the control experiment, quad accumulation of the face fluxes and their difference inside the stationary residual only, as a measured option.

- **N34** (the control experiment of decision 23: the hydrodynamic rows of the stationary residual in quadruple precision). EXHALE_RESID_QUAD=1 sends the STATIONARY evaluations of assemble_residual -- the tag ieq_sweep_state_kind puts on an iterate or a probe; the marching stages reach RK_rhs directly and are untouched -- through a quadruple-precision instantiation of the whole flux assembly, from the conserved state through the caloric EOS, the PLM limiter and the WENO3 weights, the Roe flux with its entropy fix and HLLE fallback, the geometric and gravitational sources and the division by the cell volume, rounded to double once, at the row. One source text (hydrodynamic_rows_body.inc, 1111 lines, kind-generic) instantiated twice by hydrodynamic_rows.f90; the composition, the heating and the cooling stay double and enter converted exactly. **The double instantiation is the production operator BIT FOR BIT**, asserted three ways: a suite row on a stated grid (largest difference exactly zero in dF, S and the four face states, tolerance 0), and two whole 40-iteration reloads under EXHALE_RESID_QUAD=2 -- the atomic (all 355 (JFNK) lines identical) and the carrier, which is the molecular one and so exercises the generic caloric EOS (all 364 identical). The quadruple instantiation reaches the analytic mass row of a uniform state to 1e-30 before its rounding, where the double one stands at its own. **Verdict: the floor is confirmed and is NOT removed.** On the atomic reload at outer 1 the tracked row's non-smoothness floor falls only 1.240e-11 to 5.917e-12, and cell by cell it lands on epsilon_DOUBLE x face state x r^2/dV -- ratio 0.93 to 1.4 over the base cells against 5.7 to 8.9 in the control -- and not on the quadruple bound, which is 1.224e-30, 4.8e18 below what is measured. So N33's amplifier (the face cancellation 3.2e5 times the r^2/dV 5.1e3) is confirmed a second way and its attribution completed: four fifths of the floor is the arithmetic INSIDE the assembly, the rest is the double representation of what the assembly is handed and hands back -- the ghosts Apply_BC writes, the sweep's temperature and particle count, and the double row itself -- which precision

inside one operator cannot reach. The detector isolates no step at any of the ten windows (control: 28 of 64 and 61 of 64 at the two narrowest), but the second difference is still flat over 5.4 decades and the row still changes sign 39 times in 64 intervals: a staircase with a smaller rise. **What the quadruple-precision accumulation does buy is the fidelity of the Krylov model:** the additivity defect falls $8.174\text{e-}6$ to $1.009\text{e-}6$ with the exponent in the arc still negative (-1.063 to -0.901 , so still rounding and not curvature), and at the binding iterate of the atomic reload at cap 40 the relative gap of the Arnoldi image falls $1.188\text{e-}1$ to $3.813\text{e-}3$, a factor 31, the returned relative residual then being the true one to four digits (0.2114 against 0.2114 ; the control returned 0.1122 for a true 0.1617). The element rows, which do not pass through the assembly, are unmoved to four digits ($C\ 1.367\text{e-}13$ to $1.366\text{e-}13$), which is the control inside the control. **The Newton does not converge and the Krylov cycle still stalls.** Atomic cap 40, quad accumulation on: 40 of 40 products at 0.21 against the requested 0.10 , $||R||\ 8.413\text{e-}1$ handed back against the control's 1.112 with trajectories parting at iteration 9 (the solve is chaotic at the ulp level, so that is not by itself a result). Atomic cap 250: both stop on the stagnation detector, control at iteration 167 at $3.047\text{e-}4$, quad at 136 at $4.096\text{e-}4$, both with 40 of 40 at 0.983 to 0.989 , both binding on the element Fe rows of cells 132 to 137, both NOT CERTIFIED with 11 entries of the inventory refusing. Carrier cap 40: unchanged in every named outcome -- binding row the H2 carrier row of cell 205 throughout, 40 of 40 at 0.9886 against 0.9729 , $||R||\ 9.336\text{e-}2$ against $9.427\text{e-}2$ -- as it must be, since that row is assembled by `carrier_steady_residual` and never sees the flux. Cost MEASURED: the flux assembly alone 46x at 8 threads and 74x at one (500 cells, $3.628\text{e-}5$ s to $1.681\text{e-}3$ s), the whole atomic reload at cap 40 1.11x (271 s to 302 s), because a stationary residual is mostly the ionization sweep. **Nothing adopted, no default changed:** `wasp_full_newton` bitwise but the provenance timestamp, `log10 Mdot` 13.30 on both; the atomic and carrier reloads identical line for line with the option off. Tests: `krylov_and_dogleg` RED on the control objects, GREEN 232/0 with 12 rows added; `steady_species_rows` 194/0, `element_operator` 14/0, `certification` 57/0, `energy_update` 53/0, `acceptance_classes` 22/0, `adv_static_limit` 53/0, `species_face_flux` 1/0, `spectrum_type` 154/0, `grid_and_gates` 146/0, `attempted_step` 70/0, `coupled_source_step` 30/0, `residual_determinism` 5/1 with the same row failing by design and the same $3.354\text{e+}01$ on the control binary. One file outside the brief's list: `caloric_eos.f90` gained three read-only accessors (the mixture ratios and the H2 rovibrational table nodes) with no arithmetic and no behavior change, because the generic caloric routine cannot evaluate the molecular EOS without them; the molecular whole-run bitwise comparison is the evidence that they changed nothing. Noticed, not fixed: `backup/regression/atomic_elem_newton/README.md` still names the helium row of cell 246 as the binding row at cap 250 from N26/N27, where the measurement on this tree is the element Fe rows of cells 132 to 137 at iteration 167 and $||R||\ 3.047\text{e-}4$. **For decision 23:** extra precision confirms the diagnosis and does not cure it; the amplifier has to go (well-balanced flux differencing, or a coarser base cell, which buys one factor of the width), and even with the hydrodynamic rows made model-faithful the two rows that actually bind -- the element rows of the atomic reload and the H2 carrier row of the carrier reload -- are not hydrodynamic rows and are untouched by any of it. Advisor: rebuild on `build_lwv` (the new `hydrodynamic_rows.f90` in SRC); `krylov_and_dogleg` 232/0, `steady_species_rows` 194/0, `energy_update` 53/0; atomic reload at cap 40 with every hook off, 8 threads: the N31 ladder to the digit ($7.607\text{e-}2$ at 40, handed back $5.377\text{e-}2$). The `caloric_eos.f90` accessors (read-only, outside the brief's list) accepted on the molecular reload's bitwise evi-

dence. Accepted; nothing adopted. Decision 23 restated with this result: option (b) retired as a remedy, (a) still the right discretization but not by itself the cure of the species-row solve; the atomic fixture README gained a 2026-09-10 state section (binding rows Fe of cells 132 to 137 at cap 250 on this tree). **Next item N35** (runs without a decision): the conditioning of the preconditioned operator on the element and carrier rows at the binding iterate, and what the banded preconditioner misses there.

- **N35** (what holds the Krylov cycle on the species rows once the rounding floor is removed: the conditioning of the preconditioned operator and what the banded preconditioner misses). Three default-off measurement hooks on the linear system of one outer iteration, adopting nothing: EXHALE_KRYLOV_SIZE_SCAN=1 ([diag 16]) reruns the SAME cycle at 40, 80, 160 and 320 products and the last two again with the basis orthogonalized twice, with EXHALE_GM_HISTORY=1 printing the reduced residual after every product and the true one every twenty; EXHALE_PRECOND_SPECTRUM=1 ([diag 15]) takes 200 Arnoldi products of $A_z M^{-1}$ from a deterministic start, diagonalizes the Hessenberg (dgeev) and reports the Ritz values of the whole operator and of its compressions onto the species and the hydrodynamic rows, with the localization of the three smallest Ritz vectors; EXHALE_BAND_DIFFERENCE=1 ([diag 17]) measures $(A - A_{\text{band}}) v$ on those Ritz vectors and on the first two Arnoldi directions, splits the columns they live on into the part inside the band, the band's disagreement there and the part outside, with the radiation live and frozen, and reads the binding row's band row by class as a fraction of its diagonal. **Verdict: the band misses nothing, and the operator itself is near-singular on the species rows.** Its action agrees with the full finite-difference one to $4.0e-5$ to $1.5e-3$ relative on every direction tested on both fixtures; the columns hold 99.4 to 100.0 percent of their norm inside the band (the largest outside part is 0.6 percent, element He of cell 308); and what remains is not on the rows that bind at all, but 0.999 in the momentum row of the LAST cell (carrier) and 0.83 to 0.89 momentum plus 0.11 to 0.17 energy in five to seven cells (atomic), the element and carrier rows taking 0.0000 to 0.003 of it. So the two remedies the brief conditioned on -- a rank-few radiation correction by the Woodbury identity, or a wider kl_jac -- would each repair something already right, and **nothing was built. What IS wrong, MEASURED at the binding iterate of both fixtures.** (1) The preconditioned operator is well conditioned on the hydrodynamic rows and near-singular on the species rows: the Ritz spectrum of the compression onto the hydrodynamic rows has a magnitude ratio 43.4 (carrier, carrier H2 of cell 205 binding) and 158 (atomic, energy of cell 52 binding), none and three of 200 below $1e-2$; onto the species rows, $6.56e5$ (carrier, smallest $1.493e-6$, nineteen below $1e-2$, one below $1e-4$) and $1.06e4$ (atomic, smallest $9.366e-5$). The smallest Ritz vector is 0.71 to 0.91 carrier rows on the front cells 199 to 205 and 458 to 466 (carrier) and 0.9955 element rows over 145 cells (atomic), and the carrier fixture's near-zero part is a complex pair, not a real null direction. The attribution is there from outer 1 ($2.17e5$ against 39.6). And the discretization statement behind it: the band row of carrier H2 of cell 205 has diagonal $1.2396e-1$ against off-diagonal magnitudes $4.1026e2$, **3.304e3 times the diagonal to the hydrodynamic unknowns of its own stencil and 5.65 times to the neighboring carriers**, so the binding row's own unknown is three decades below its coupling to another class and there is no diagonal for a preconditioner to stand on. (2) **A larger subspace makes the step worse, not better**, which is neither of the brief's two alternatives. The reduced least-squares residual falls monotonically with the size while the TRUE residual of the returned step stops falling at 60 to 80 products and rises: atomic with quad accumulation on, reduced $2.658e-1$, $1.781e-1$,

1.144e-1, 9.958e-2 at 40, 80, 160 and 182 products against TRUE 2.851e-1, 2.575e-1, 3.006e-1, 3.203e-1 -- the cycle announces its tolerance reached at 182 and hands back a step leaving 0.32, three times what was asked; carrier, reduced 9.677e-1, 8.001e-1, 4.639e-1, 1.434e-1 against TRUE 9.731e-1, 8.638e-1, 9.668e-1, **1.472e+00**. With quad accumulation off the true residuals are a factor 3 to 6 worse at every size (atomic 7.856e-1, 3.905e-1, 7.580e-1, 2.004e+00) and the shape is the same. Reorthogonalizing the basis changes nothing at any size, the measured loss of orthogonality being 0.000e+00, and the Arnoldi image of each column at the same iterates is exact to 3.30e-16 and 3.77e-16: the recursion is not the defect. What fails is additivity -- the action of the sum of the first two directions departs from the sum of their actions by 1.357e-3 (atomic) and 7.601e-5 (carrier), the assembled 40-product step's image by 1.068e-1 and 2.256e-1 -- because the matrix-free action is the secant of a nonlinear residual over a fixed arc and the step it is extrapolated to grows from 0.31 to 11.4 while the arc stays 5.2e-6. Tests: krylov_and_dogleg 251/0 with 19 rows added (RED on the control objects, which do not carry the routines), among them the Ritz values of a stated upper triangular 6 by 6 spanning six decades reproduced to 1.171e-12 with the recursion carried to the dimension of the space, the column split summing to the whole difference with an empty band and with an exact one, and the four hooks off by default; steady_species_rows 194/0, element_operator 14/0 (the three suites whose driver links steady_newton). Default path: the atomic reload at cap 40 with every hook off, new binary against my control, all 354 (JFNK) lines IDENTICAL through ||R|| 5.377E-02, flux spread 3.555E-03, the N31 ladder to the digit; wasp_full_newton bitwise but the provenance timestamp, log10 Mdot 13.30 on both. Cost MEASURED: the size ladder is 1082 products per diagnostic iteration (the carrier reload at cap 40 from about 3.5 to about 9 minutes at 8 threads), the spectrum 600 and the band difference about 15. Noticed, not fixed: the frozen-radiation column of element He of cell 410 carries 7.267e-4 of its 2.816e-1 norm outside the band where the WENO3 stencil says it should carry none, 0.26 percent and two decades below anything in this item's conclusions; the likely route is the base boundary state recomputed from the iterate, not confirmed. Also corrected in the reading: the preconditioner is build_banded_jac_full, a coloring of the FULL residual, not of the frozen one -- build_banded_jac on frozen_residual is called only by the standalone check in EXHALE_main.f90 -- so "inside the band" carries the coloring's contamination by the non-local response, and the frozen column is a separate control rather than the band's own approximation. **For the advisor:** the next question is not the preconditioner. The species rows' row scaling (the binding row's diagonal 3.3e3 below its hydrodynamic coupling is set by the certification row scale of decision 20 a divided by the column scale, and using one scale for the merit and another for the model is a decision); the cycle has an optimal length around 60 to 80 products on both fixtures and the step control asks for a tolerance the operator does not honor; and gm_resid_rel is verified against the operator only at a breakdown or a rank loss, where the measurement says it needs verifying whenever the cycle ran more than a few tens of products. Advisor: rebuild on build_lwv; krylov_and_dogleg 251/0, steady_species_rows 194/0; atomic reload at cap 40 with every hook off, 8 threads: the N31 ladder to the digit (7.607e-2 at 40, handed back 5.377e-2). Accepted; deliverable 4 correctly not built. The two levers named are the LINEAR model's row scaling (a freedom decision 20 a did not fix: the merit and gate keep the certification scales, the linear inner product may differ) and a Krylov cycle that returns the step its TRUE residual chooses (the optimum lies at 60 to 80 products; the reduced residual lies past it). **Next item N36:** both as measured options, default off.

- **N36** (the two levers N35 named: the row scaling of the LINEAR model, separate from the merit's, and a Krylov cycle that returns the step its TRUE residual chooses). Two default-off options on the linear solve, adopting nothing. EXHALE_MODEL_ROW_EQUIL=1 (unit_infinity_norm_row_scaling_of_the_band) forms the diagonal E that brings every row of the banded model to unit infinity norm as a power of two, applies it to the band that is factored and, inside pgmres, to the right-hand side and to every action of the operator, and maps the image and the relative residual BACK to the certification scales at every exit, so that the merit, the gate, the trust region's predicted decrease and the certification keep the scales decision 20 a fixed and only the linear model's inner product changes; ab itself is untouched, so the merit's gradient is where it was. EXHALE_GM_TRUE_RESIDUAL=1 makes the cycle measure its candidate step against the operator from 20 products on, every 10 and at every point where it would otherwise end, keep the best, and stop on two consecutive rises or on the tolerance IN THE TRUE NORM; what comes back is that step, its own image from the product that measured it, and the true residual as gm_resid_rel, under a new eighth outcome gm_true_residual_turned. **Verdict: both options work as specified, both fail the acceptance, both stay off.** (1) **The row scaling is a real freedom and taking it is worse.** The cycle then minimizes a functional the merit does not read, and the two disagree by a factor 1.2 to 673 MEASURED ([row equil], equilibrated against certification: atomic 9.956e-2 / 4.111e-1 at outer 2 and 9.937e-2 / 1.218e-1 at outer 40; carrier 2.353e-1 / 5.730e+1 and 1.286e-1 / 8.657e+1), so the cycle announces the 1e-1 it was asked for while handing back a step that leaves 86.6. It cannot touch what binds: the preconditioned operator $A_{eq} M_{eq}^{-1} = E (A B^{-1}) E^{-1}$ is SIMILAR to the unequilibrated one, and the measurement at the same iterate agrees (hydrodynamic Ritz ratio 39.62 against 39.60 carrier, 57.02 against 56.36 atomic; the count of species-row Ritz values below 1e-2 unchanged at 18 and 2), while the banded factorization is bit for bit the same matrix because equilibrate_the_preconditioner already equilibrates both sides; and a row scaling leaves every ratio WITHIN a row where it was, so the binding carrier row's diagonal 1.2396e-1 against 4.1026e2 of coupling, 3.304e3 times the diagonal to the hydrodynamic unknowns of its own stencil, is exactly N35's number still. What it costs: the carrier reload at cap 40 goes from ||R|| 9.572e-2, info=1, binding carrier H2 of cell 205 to 1.955, **info=2 with four entries of the inventory refusing**, and the true residual of the returned step at 40 products from 0.973 to 1.050e2. (2) **The true-residual cycle is the right instrument and it does not turn the solve.** At the binding iterate it caps the cycle where the step stops improving: on the ladder 40/80/160/320 the control's returned step gets WORSE with size (atomic TRUE 2.851e-1, 2.575e-1, 3.006e-1, 3.203e-1; carrier 9.731e-1, 8.638e-1, 9.668e-1, **1.472e+00**) while the true-residual cycle stops at 80 and at 120 products and returns 2.660e-1 and 8.244e-1, never worse than the control's best at any size and at fewer products. Two cycles of the atomic reload at cap 40 name the new outcome. But the best true residual any subspace reaches is 0.27 and 0.82, the Newton does not descend past its plateau, and ||R|| handed back at cap 40 is 1.060 against the control's 5.377e-2 (atomic; the trajectory parts at iteration 15, the first cycle to pass 20 products) and IDENTICAL on the carrier, where a subspace of 40 always makes the last candidate the best and the option changes only the number reported (true 9.870e-1 against reduced 9.839e-1). **Acceptance, by the named clauses:** the true residual below 0.1 within 80 products on both fixtures fails for both options (option 1 atomic 8.79e-2 but carrier 7.51e+1; option 2 atomic 2.66e-1, carrier 8.64e-1), and the refusal count at cap 40 does not fall by half (atomic 9 to 10 for both options; carrier 8 to 7 and 8 to 8), with no new refusal class

and no new gm_no_direction_sampled anywhere. Corrected mid-item and re-measured: the first true-residual cycle left the reduced tolerance in place as a stop and so discarded a step it never measured (carrier outer 1, the 80-product cycle ended at product 49 and handed back product 40's step at TRUE 1.259e-1 where product 49 reaches 9.724e-2); with the reduced residual no longer ending a cycle, the numbers above are the corrected cycle's. Tests: krylov_and_dogleg RED on the control objects (nine symbols absent), GREEN 264/0 with 13 rows added, among them the equilibrated solve returning the same step as the unequilibrated one to 9.90e-16 on a stated dense system solved over the whole space, the equilibrated cycle reporting the CERTIFICATION residual of a truncated solve to eight digits, and a stated operator whose action stops being one linear map after 40 products, on which the cycle names gm_true_residual_turned and stops before its subspace is spent; steady_species_rows 194/0, element_operator 14/0, residual_determinism 5/1 with the same row failing by design and the same measured 3.354e+01. **Default path:** the atomic reload at cap 40 with every hook off, all 354 (JFNK) lines IDENTICAL to my control through $||R||$ 5.377E-02, flux spread 3.555E-03; the carrier reload all 364 identical; wasp_full_newton bitwise but the provenance timestamp, log10 Mdot 13.30. With the options ON wasp_full_newton stays CERTIFIED: option 1 $||R||$ 3.043e-9 against 1.667e-9 with a largest relative movement 5.2e-10 in the output, option 2 bitwise in Ion_species.txt with 8 extra residual evaluations, its cycles never reaching the 20 products the needs. **For the advisor:** what the two options rule out is the linear algebra. The norm the cycle minimizes is free and wrong to change; the length of the cycle is free and already at its optimum; the operator's species block is near-singular at every scaling and the step it can produce leaves 0.27 to 0.82 of the residual whatever the subspace. The remaining lever is the one N35 named and N34 confirmed the arithmetic cannot reach: the DISCRETIZATION of the rows that bind. Advisor: rebuild on build_lwv; krylov_and_dogleg 264/0, steady_species_rows 194/0; atomic reload at cap 40 with every hook off, 8 threads: the N31 ladder to the digit (7.607e-2 at 40, handed back 5.377e-2). Accepted; both options stay off. **The linear algebra is exhausted as a lever** (N31 to N36: probe rule, inner tolerances, extended precision, band coverage, row scaling of the linear model, cycle length). STOPPED here for the user: decision 23, and the next question is the discretization of the species rows that bind at the front (the carrier row of cell 205 whose own diagonal is 3.3e3 below its hydrodynamic coupling). code_status_20260910.md section 5 task 1, ISSUES 3.1 and 5, the handoff and decision 23 restated accordingly.

- **N37** (decision 23 (a): a well-balanced flux difference for the near-hydrostatic layer, as the default-off option Well balanced:). With the key on the reconstruction, the Riemann jumps and the pressure force carry the DEPARTURE from the cell's own local hydrostatic equilibrium instead of the state: within cell j that equilibrium is the constant-density one through its own (ρ_j, p_j) , $p_{eq,j}(r) = p_j - \rho_j (\phi(r) - \phi(r_j))$, which assumes no thermal, entropy or compositional stratification and so preserves ours (Kaeppli and Mishra 2016, A&A 587, A94, sections 2.1.1 and 2.1.3; the momentum source as the face-pressure difference of the cell's own equilibrium is their 2014 paper, J. Comput. Phys. 259, 199, eq. 2.26). The pressure stencil acts on the neighbor's pressure measured against that equilibrium continued THROUGH the face (this cell's density up to the face, the neighbor's beyond it), which is what vanishes on the discrete equilibrium; the two cell pressures are differenced first, while that is exact, and the hydrostatic terms added to the difference. The momentum row is then assembled from the face pressure measured against the same equilibrium and source returns $S(2) = 0$: with A+

$P_{up} - A - P_{dn} - (A^+ - A^-) p_j = -\rho_j [A^+ (\phi_i(j) - \phi_c(j)) + A^- (\phi_c(j) - \phi_i(j-1))]$
 identically, the equilibrium's flux difference and its gravitational source cancel in the ALGEBRA and the cell's own $O(1)$ pressure never enters the row. Mirrored in the kind-generic copy, so EXHALE_RESID_QUAD=2 is the production operator bit for bit under both key values (suite row, largest difference exactly zero, tolerance 0).

Exact preservation, MEASURED: on the scheme's own discrete equilibrium (the isothermal solution of the face-matching condition, the spherical form of their eq. 42) the momentum row falls from $7.0e-5$ (PLM) and $2.5e-3$ (WENO3) of the local weight ρg at $N = 250$ to $4.1e-14$ and $3.9e-14$, and stays at $4.5e-14$ to $5.5e-14$ at $N = 500, 1000, 2000$ while the base scheme falls at its design order; the mass and energy rows reach $1e-16$ against $1e-8$. Twelve new assertions in `hydrostatic_residual`, plus eleven rows in `krylov_and_dogleg` (with no gravity the well-balanced option is the base scheme to $8e-16$ on a uniform state and to $1.4e-16$ on a Sod tube, for both reconstructions; and the same exactness on a second grid and geometry).

What it does NOT reach, and this was written into the design before the code: N33's non-smoothness floor does not move (energy row of cell 1, $1.009e-11$ with the option against $1.046e-11$ without; cells 84, 167, 251 the same to a factor 1.3; six to ten times $\epsilon \times \text{face state} \times r^2/dV$ either way), because a cell pressure is itself a rounded $O(1)$ double and every jump built from two of them steps by its last bit whatever the grouping. What does move is the additivity defect at ten times the probe arc, $2.410e-7$ with the option against $3.151e-5$ without.

The stationary solves are worse with the key on, and the reason is not the option: `store_row_terms` scales the momentum row by $\max(|dF(2)|, |S(2)|, |S_{mom}|)$, the option returns $S(2) = 0$, so the row IS its own scale and every normalized momentum residual is exactly 1 (MEASURED on the carrier reload as loaded: $k=2$ is $1.000000E+00$ with the option against 1.954 without, while the mass row is the same number to seven digits and the energy row to four). The merit then carries no momentum information: $||Fs||^2$ rises from 16.5 to 28.1 at the same state, and $28.1^2 - 16.5^2 = 517$ is the 500 momentum rows pinned at unity. The atomic reload at cap 40 reaches 1.415 against the control's 1.112 and the carrier reload 1.732 against $9.34e-2$; at cap 250 the option's solve aborts at iteration 61 on "no descent step in 12 consecutive iterations" with $||R||$ and the worst scaled row both exactly 1.000 and that row the MOMENTUM of cell 1, which is what a merit whose momentum rows are all pinned at one looks like from inside the line search.

`wasp_full_newton` with the key on says the same from the other side: it is NOT CERTIFIED on exactly one entry, the momentum row at $1.000E+00$, while its flux spread is $1.539e-15$ against the control's $6.233e-14$, its energy row $9.911e-12$ against $1.667e-9$, its mass row $1.228e-13$ against $1.453e-13$ and $\log_{10} Mdot$ 13.30 either way, the profiles moving by at most $1.2e-4$ (velocity).

The mechanical column drains by the same amount with the option as without (`hydrostatic_column`, $\max |v|/c_s$ 0.796 against 0.792 at 300 steps and 2.776 against 2.777 at 3000): that case's state is not the scheme's equilibrium and its drain is set by the outer boundary, as B4 recorded.

Default path: with the key absent or False every path is bit for bit my control (`wasp_full_newton` reload, the atomic reload's 196 (JFNK) lines, `hydrostatic_column` at 300 steps, `mol_carrier` and `mol_diffusion` at their maxsteps); the one line that moves in every output file is the # options field, which now carries the 21st token `wellbal`, nameable in `Restart option change`.

`EXHALE_resolved.out` and the parse dump each carry one new line. Key documented as K46.

For the advisor: the option is kept in the tree and stays off; before it can ever be default on, the momentum row's reference scale has to become the pressure force the option cancels analytically, $|\rho_j (A^+ (\phi_i(j) - \phi_c(j)) +$

$A - (\phi_c(j) - \phi_i(j-1))|/dv$, which is two multiplications from what RK_rhs already holds but sits in an acceptance interface this item does not own. Advisor: rebuild on build_lwv; krylov_and_dogleg 279/0, grid_and_gates 162/0 (four rows FAIL when the shell rows are pointed at the stale shared EXHALE.x: pass EXHALE_EXE with EXHALE_OBJDIR), steady_species_rows 194/0, certification 57/0, element_operator 14/0, residual_determinism the same single row FAIL by design; atomic reload at cap 40 with every hook off and the key absent, 8 threads: the N31 ladder to the digit $(7.607e-2)$ at 40, handed back $5.377e-2$). The regression comparison strips comment lines, so the new wellbal token in the # options header is invisible to make check. Accepted as a default-off option; the momentum-row reference scale in store_row_terms (the option's $S(2) = 0$ makes the row its own scale) is recorded in ISSUES 3.7 as the prerequisite of any default-on decision; the publisher PDFs of the two Kappeli and Mishra papers remain wanted (the institutional versions were read in full).

- **N38** (the discretized element and carrier rows at the front: why the row that binds is nearly independent of its own unknown). One default-off measurement hook, adopting nothing. EXHALE_FRONT_ROW=1 ([diag 18]) takes the species row that carries the largest scaled residual and its two neighbors and prints, at the selected outer iteration: every TERM of the row on its certification scale, with the advective term split per FACE by masking the face mass flux (the divergence of a face species flux is exactly zero where the mass flux it multiplies is, so the masked call leaves one face standing alone and the SAME operator is used, never a copy); the reconstructed face fraction read back out of that masked divergence beside the donor cell's own; the chemical source of a carrier from the sweep's own rates; the band's entries for the row, to its own unknown, to the same unknown in the neighbors and to every hydrodynamic unknown of the three cells, each as a number and as a multiple of the diagonal; those same entries SPLIT BY TERM, by one-sided differences of the row's own terms along the same column with the same step build_banded_jac_full uses; the cell Peclet number, the logarithmic gradient and the chemical against the advective time scale; and the column scale of every unknown of the binding cell beside the unknown itself. **Verdict: the binding row's diagonal is NOT small -- it is the size of the row -- and the 3.3e3 of N35 is a CHOICE OF COLUMN SCALE that the Krylov cycle cannot see.** (1) **Every hydrodynamic entry of the row is its ADVECTIVE term**, on all three fronts measured: carrier H2 of cell 205 (carrier reload, outer 40) momentum of cell 205 $-7.40478e+1$ of which the diffusive part is $2.0e-5$ and the chemical $-2.3e-5$; element He of cell 191 (atomic, outer 1) momentum of cell 190 $-1.86758e+4$ with a diffusive part $1.9e-6$; element Fe of cell 137 (atomic cap 250, outer 140, one of the rows N34 named) momentum of cell 136 $-1.25980e+4$ with $2.0e-6$. Summed over the reported stencil the hydrodynamic columns carry **3389, 4176 and 2424 diagonals** and the neighboring species unknowns 5.855, 1.052 and 1.005, which is N35's $3.304e3$ and 5.646 reproduced. Each species sub-block is meanwhile a clean diffusive tridiagonal ($15.385/-7.772/-7.612$ helium, $17.432/-8.861/-8.564$ iron): N27's reading of an interior element row holds at the front too, and it is the wind columns that bury it. (2) **The size of that coupling is one over the Mach number**, because the momentum column scale is $|\rho v| + \rho c_s$, the SOUND speed in a deeply subsonic layer: MEASURED scale over unknown **3.839e+2** (carrier front, momentum $6.83274e-5$ against scale $2.62332e-2$), **1.361e+5** (helium front) and **5.149e+4** (iron front), against 1.000 for the mass, the energy and every species unknown. The arithmetic closes to a factor 1.4 to 1.6: one face carries -0.5322 of the carrier row and $383.9 \times$

0.5322 = 204 against the measured 129; one face carries -0.1969 of the helium row and $1.361e5 \times 0.1969 = 2.68e4$ against the measured $1.87e4$. (3) **The three other candidates are measured and are not the cause.** The species column scale's reservoir floor is INACTIVE everywhere it was looked at (the carrier unknown is $\ln n$ with column scale exactly 1.0, the default; the helium and iron unknowns' scales are their own magnitudes to five digits), so raising or lowering `element_column_scale_floor` by a decade changes the diagonal by nothing. A stiff advective/chemical cancellation in the diagonal is a factor 1.8 and not decades (carrier diagonal $1.19228e-1 = \text{diffusive } +2.17367e-1$ and advective $-9.89614e-2$, with the chemical derivative $+9.44223e-4$, 0.8 percent, although the chemical time scale is 220 times the advective one), and there is no cancellation at all in the helium diagonal. The limiter is a live switch (the reconstructed face fractions stand at 0.587 and 0.657 of the donor at the carrier front, and frozen at first order the advective part of the diagonal changes SIGN, $-9.89614e-2$ to $+2.50409e-1$) but it moves the ratio by less than 2, because freezing it raises the momentum coupling by the same kind of factor. N5's two-root front was not re-run: cell 205 is not one. (4) **The test that settles it:** the existing default-off column equilibration (`EXHALE_COLUMN_EQUIL=1`) is exactly the remedy branch (i) names, and MEASURED at outer 1, where both runs stand at the same iterate, it brings the binding row's hydrodynamic coupling from 3389 to **0.993** (carrier) and 4176 to **2.926** (atomic) while the Krylov cycle returns the SAME step at the SAME residual to every printed digit (carrier 40 of 40, reduced $1.257e-1$, TRUE $1.259e-1$; atomic 8 of 40, $7.821e-2$, TRUE $7.824e-2$) and the species-row Ritz count below $1e-2$ is unchanged at 18 and 2. It cannot be otherwise: a column scaling E is applied to the operator and to the band built FROM it, and $(A - E)(band(A) - E)^{-1} = A - band(A)^{-1}$ IDENTICALLY. N36 showed a row scaling is a similarity; this shows a column scaling is an identity, so **both sides of the scaling are now exhausted**. What the column equilibration costs, since it also reshapes the trust-region ball and the probe steps: cap 40 ends at $||R|| = 1.822$ (carrier) and 1.592 (atomic) against $9.572e-2$ and $5.377e-2$. Acceptance by the named clauses: the Ritz ratio below $1e3$ FAILS ($2.07e5$, $1.45e3$), the cycle reaching 0.1 within 40 products FAILS on the carrier, named outcomes are lost on both; it stays off. **Deliverable 3 built nothing**, by the branch N35's deliverable 4 took: the remedy branch (i) names is either already the default ($\ln n$) or already measured to be invisible. Tests: `krylov_and_dogleg` RED on the control objects (`attributed_jacobian_entry` absent), GREEN **279/0** with 6 rows added (the hook off by default; a stated term set composing its own row and its two faces composing its advective term through `species_row_from_its_terms`, the expression the measurement itself uses; the attributed entry of a term linear in the unknown equal to its coefficient times the scaling, with a zero probe step and a row with no scale attributing nothing) -- 279 rather than 270 because N37 added nine rows to the same file, a brief overlap reported and not resolved; `steady_species_rows` 194/0, `element_operator` 14/0, `steady_selfconsistent_residual` 3/0 on the `wasp_full_newton` log, `residual_determinism` 5/1 with the same row failing by design and the same $3.354e+01$. Default path: the atomic reload at cap 40 with every hook off, all 354 (JFNK) lines IDENTICAL to my control through $||R|| = 5.377E-02$, `flux_spread` $3.555E-03$; the carrier reload all 364 identical at $9.572e-2$; `wasp_full_newton` done `info=0` $||R|| = 1.667E-09$ on both binaries with all 28 lines the same; and with the hook ON both reloads still end at the control's values, so the measurement is not an operation on the state. Cost MEASURED: one residual evaluation for each of twenty columns at each of three outer iterations, about four minutes on the carrier reload at cap 40 and 8 threads. Also corrected in the reading: N35's "there is

no diagonal for a preconditioner to stand on" does not follow from the band row -- the diagonal is the size of the row, the disparity beside it is a column scale, and the near-singularity of the species-row compression of A^{-1} is a separate fact that the band-row ratio is not evidence for. Advisor: verified on the same final build (the suites above); atomic reload identical. Accepted; deliverable 3 correctly not built. N35's sentence "no diagonal for a preconditioner to stand on" is withdrawn: the diagonal is the size of the row, the 3.3e3 is the momentum column scale (the sound speed) against a layer Mach number of 1e-3 to 1e-5, and column scalings are invisible to the cycle, so the near-singularity of the species-row compression is a fact of its own.

- Close of the species-row solver program** (2026-09-10 evening, user's instruction: "어떤 것을 해도, 끝이 없어 보이니, 엄청난 차이가 나오지 않는 한 이번 건만 마무리 하고 끝내기 바래", then "있는 그대로 정리하고, md tex 파일들 update 후 commit/push"): N37 and N38 were finished as briefed and no further item is started. State of the tree at the close, verified on the advisor's rebuilt build_lvw: krylov_and_dogleg 279/0 (174 at the morning commit), grid_and_gates 162/0, steady_species_rows 194/0, certification 57/0, element_operator 14/0, residual_determinism one row FAIL by design; the atomic reload at cap 40 with every hook off and every new key absent gives the N31 ladder to the digit (1.888, 1.891, 1.819, 1.583, 1.861, 8.964e-1, 7.607e-2; handed back 5.377e-2, flux spread 3.555e-3); the wasp_full_newton reload ends info = 0, ||R|| 1.667e-9, the number every worker's paired control gave (its output is not compared with the cold-start golden, which is a different path: sec_ion_step 2402 against 0). Everything N31 to N38 added is a default-off hook, option or key: EXHALE_JV_ADDITIVITY, EXHALE_JV_COLUMN_SCALE, EXHALE_JV_PROBE_ARC, EXHALE_IEQ_TOL, EXHALE_TEQ_TOL, EXHALE_RESID_JUMP_SCAN, EXHALE_RESID_QUAD, EXHALE_KRYLOV_SIZE_SCAN, EXHALE_GM_HISTORY, EXHALE_PRECOND_SPECTRUM, EXHALE_BAND_DIFFERENCE, EXHALE_MODEL_ROW_EQUIL, EXHALE_GM_TRUE_RESIDUAL, EXHALE_FRONT_ROW, and the input key Well balanced: (K46); the marching path and the goldens are untouched (the # options header gains the wellbal token, which the regression comparison strips with every comment line). New fixture backup/regression/carrier_elem_newton. Documents closed to this state: ISSUES_20260909.md 3.1, 3.7 and 5; code_status_20260910.md 3.3 and task 1; session_handoff_20260910.md (new, the handoff of record); To_be_determined_by_user_20260906.md decision 23 with its outcome; steady_solver_design.md 22.4; input_schema.md K46 and the manual's key table (PDF rebuilt, 90 pages); README.md (the key). The analysis the user asked for, of why the coupled solve fails and what a different approach would be (a segregated solve, the CETIMB way read from Koskinen et al. 2013 section 2.1.3 and 2022 Appendix B, a different treatment of the layer), is docs/solver_approach_analysis_20260910.md. Reported, not fixed (ISSUES 3.7): the count global shadowing the intrinsic; newton_dense's hardcoded step test; the dead dp_bc/ntot_bc refreshes; store_row_terms's momentum scale under the new key; fortdep.py blind to include; golden/mol_ir_bands/ without Cooling_breakdown.txt; examples/README.md; the two .md twins.

8. PLAN_20260911: the partitioned stationary route (2026-09-11)

Plan of record: docs/PLAN_20260911_partitioned_solver.md. User instruc-
tion of 2026-09-11: read docs/solver_approach_analysis_20260910.md, its review
docs/solver_approach_analysis_20260910_review.md and docs/solver_partition_experiment_20260911.md,
correct the errors they establish, run the solver experiments, and adopt the partitioned route in the production solver if
it measures better than the coupled species-row solve. This reopens the stage-2 solver program that the 2026-09-10
instruction had closed after N37 and N38. Items P1 to P9, then P10 to P12 on the same day's further instruction to fix
the open items the plan had left to the user, then P14 and P15 on the two findings of that afternoon's Codex adversarial
review and P16 to P18 on the user's two decisions of the same afternoon, then P20 and P21 in the evening and the
terminology sweep A1 to A4 on the user's instruction of that evening (P13, P19 and P22 are the record itself); the rules
every worker on this tree follows are docs/worker_rules.md.

The items

- **P1 (defect D1): a failed element composition solve is no longer handed back as an update** (binary_element_diffusion.f90 +311/-23, src/tests/element_operator/element_operator_tests.f90 +190/-13; every line count of this section is MEASURED from `git diff --numstat` on the tree at the close). `solve_mass_fraction` returns an explicit success result and no longer accepts the last halved trial of a line search that never descended; `element_diffusion_step` takes an optional status and, when asked, restores the composition it was handed unless the candidate is finite, inside $[0,1]$ to $1e-12$ before the range clip, and no farther from the density the step held fixed than the entry composition was, to $1e-10$; `relax_element_composition` steps on a copy, discards a refused step, retries at half the length within a bounded budget and reports three outcomes by name. MEASURED (P1) on the archived D1 reproduction (atomic_elem_newton, species mode, omega 1): a returned composition missing rho by $1.768571764708571E-01$, a nonfinite chemical refresh and exit status 1 become the entry composition restored at $1.046888214610824E-15$, an admissible refresh at $1.10297235E-15$ and exit status 0. The first inadmissible operation of that failure, which the experiment report left unlocated, is now named: the composition solve drives X out of $[0,1]$ and the range clip used to absorb it (`element_step_out_of_bounds`), and the mass discrepancy is downstream of the clip. Tests: `element_operator` 14/0 at entry, **24/0** after, **22/2** on a RED build with the four acceptance actions disabled (`an_unsolved_step_is_refused`, `an_unsolved_step_leaves_the_entry_composition` measured $1.31725E-02$); `diffusion_tests` 35/0, `steady_species_rows` 194/0, `certification` 57/0 unchanged. Impact (MEASURED, P1, control built from the entry text of the one file): `mol_diffusion` and `lower_profile`, 300 steps, one thread, `Ion_species.txt` byte-identical and `Hydro_ioniz.txt` identical but the provenance `run=` timestamp, on both cases; the marching caller passes no status, so none of the new tests runs there. One fixture of the same suite was corrected while there (`the_relaxation_fixed_point_under_a_callers_ghost` built a column whose species missed rho by $1.321E-01$): on the suite's own mass-closed column the worst interior element row of the relaxation fixed point moves from $1.62384E-02$ to $1.35062E-13$, so the fixed point

on a mass-closed column really is a zero of the stationary element row.

- **P2 (defect D2): the carrier movement bound now constrains the returned state** (diffusive_photochemistry.f90 +225/-81, src/tests/carrier_retry/carrier_retry.f90 +261/-1). The displacement is defined once (the largest change of any solved carrier column over the physical cells, divided by the largest H2 mixing ratio of the outer entry state, a quantity that cannot move while the pass runs), it is enforced before acceptance on every trial, a refused trial is undone and halved down to $2^{**}(-10)$ of the cell transport time, and the pass ends on one of five named outcomes. MEASURED (P2) on carrier_elem_newton through the archived driver: the entry text returns drift = $2.8126757844770485e-02$ at trust = 0.01, reproducing the 0.02812676 of docs/solver_partition_experiment_20260911.md section 7.2 exactly; the repaired routine returns $9.9956436739254070e-03$ at the same setting and $2.9845242387380568e-03$ at trust = 0.003, so the bound holds and it scales (a factor 3.33 smaller setting gives a factor 3.35 smaller movement). The halving ladder was measured on an instrumented copy: 0.4412 at the full cell crossing time, then 0.2523, 0.1445, 0.0779, 0.0404, 0.0205, 0.0103, $5.155e-3$, $2.581e-3$, $1.291e-3$ and $6.458e-4$ at the tenth halving, so the movement is proportional to the trial length and a bound of $1e-4$ keeps nothing on this column. Tests: carrier_retry **111 PASS / 10 FAIL** on a RED control carrying the entry logic behind the new interface, **122/0** after; carrier_returned_state_acceptance 36/0, attempted_step 70/0, carrier_constraint_attribution 10/0, carrier_reference_scales 14/0, species_masses 9/0, certification 57/0, adv_static_limit 53/0. Impact: mol_carrier, 300 steps, one thread, Ion_species.txt byte-identical and Hydro_ioniz.txt identical but the provenance timestamp, which is the expected result because the marching loop calls photochemical_transport_step directly and never this routine. The cost is stated: a bounded pass advances less, so the frozen H2 residual after the composition update is larger ($4.4708e-01$ at trust 0.01 against the entry text's $3.1897e-01$, from $4.9275e-01$ before), and the state handed on is correspondingly closer to the entry state.
- **P3 (defect D3): the stop test of the stationary solve is now the acceptance test of the state** (steady_newton.f90 +203/-29, src/tests/steady_completion_flag/run.sh +61/-15, src/tests/krylov_and_dogleg/krylov_and_dogleg_tests.f90 +127/-1). The loop top of solve_steady_jfnk and of solve_steady_ptc no longer stops for success on steady_gates_met alone: it stops only when the gate AND every row the certification judges the carried system by are within their own tolerances, measured on the iterate by the same evaluation and the same formula the acceptance at return uses. A gate met while a certified row refuses prints one line naming the row, its measure, its own tolerance and its cell, and the iteration continues within its budget. MEASURED (P3) on the D3 reproduction (carrier_elem_newton, first hydrodynamic solve): the entry text stopped at the loop top of iteration 14 with info = 0 and was refused at return on a mass row of $5.954e-12$ against $3e-12$; after the repair the same solve continues, 27 of its 40 iterations meet the gate with a row refusing, and it ends info = 1 on its pass budget with mass $5.945e-12$, momentum $5.991e-13$, energy $2.481e-09$. So the second of the two admitted outcomes: the mass row of this solve does not come below $3e-12$ in 27 further iterations (it stands at $5.9e-12$ to $7.9e-12$ throughout, consistent with the N33 rounding floor), and the flag is now honest about it. Cost of the added test, MEASURED with a temporary clock pair in a private build: 27 calls, mean 3.19 ms against a mean outer iteration of 0.73 s, 0.44 percent of one outer iteration, and only behind the gate, so an iteration below the gate

pays nothing. Tests: krylov_and_dogleg 279/0 on the control objects and **290/0** after (11 new rows asserting the predicate the stop test reads); steady_completion_flag RED on both real logs at entry (an info = 0 log with no statement that the stop test read the certified rows; a refusal at return with no stop statement) and GREEN after; certification 57/0, carrier_returned_state_acceptance 37/0, attempted_step 71/0 unchanged. Impact: the wasp_full_newton reload ends info = 0, ||R|| 1.667e-09, flux spread 6.233e-14, 8 outer iterations, log10 Mdot 13.30 on both binaries, the certification block line for line identical, every output byte-identical but the provenance line of the two Hydro_ioniz* files and one extra loop-top stop: line in the log; the carrier_elem_newton reload at cap 40 gives every (JFNK) line identical because the gate is never met on that reload. No golden can move: READ from the ten DEFAULT_CASES, none of their golden logs carries a (JFNK) start or (PTC) start line at all.

- P4: the momentum row's reference scale under Well balanced: True** (RK_rhs.f90 +90/-2, steady_residual.f90 +59/-2, src/tests/grid_and_gates/hydrostatic_residual.f90 +318/-11, that suite's README.md +44/-1). Under the well-balanced option store_row_terms scaled the momentum row by $\max(|dF(2)|, |S(2)|, |S_{\text{mom}}|)$ with $S(2) = 0$ by construction, so every scaled momentum residual was exactly 1 (N37, ISSUES 3.7). The reference is now $\max(|dF_2|, |S_2|, |S_{\text{visc}}|, \text{equilibrium_pressure_force}(j))$, the last term the magnitude of the pressure force of the cell's own hydrostatic equilibrium, which is the right-hand side of whichever cancellation identity the assembled row belongs to (PLM $|\rho_j| [A^+ (\phi_i(j) - \phi_c(j)) + A^- (\phi_c(j) - \phi_i(j-1))]/dV$, WENO3 $|\rho_j| (\phi_i(j) - \phi_i(j-1))/dr$; Kaeppli and Mishra 2016, A&A 587, A94, eq. 16 in spherical geometry). It has one definition, equilibrium_pressure_force_of_state, called by RK_rhs, by positivity_limited_fluxes and by assemble_residual (the last so that the kind-generic EXHALE_RESID_QUAD assembly, which returns the well-balanced row but not the term it is read against, does not scale the row by a stale weight). The numerator is untouched; with no gravity the term is zero and the scale is the row's own terms exactly (MEASURED 0 for all four scheme/solver pairs). MEASURED (P4): wasp_full_newton reloaded with the key ON is **NOT CERTIFIED** on the entry text on exactly one entry, the momentum row at **1.000E+00** at cell 1 (which reproduces the N37 record to the digit, spread 1.539E-15, energy 9.911E-12, mass 1.228E-13, Mdot 13.30), and **CERTIFIED** after, info = 0, ||R|| 9.719E-10, momentum row 9.719E-10 against 1.0E-08 at cell 500, log10 Mdot 13.30. Tests: grid_and_gates hydrostatic_residual **28 PASS / 12 FAIL** on the entry-text objects (the twelve failures at exactly 1.000 for all four pairs) and **40/0** after, of which 16 are new rows measuring the row divided by residual_row_scale(2,...), the one expression the certification reads; the whole suite 162/0 at entry (READ, N37) and **178 PASS / 0 FAIL** after; krylov_and_dogleg 279/0 and certification 57/0 unchanged. Default path (key absent): wasp_full_newton reload and mol_carrier at 300 steps byte-identical on both binaries apart from the provenance line, every (JFNK) line identical, key-off info = 0, ||R|| 1.667E-09, Mdot 13.30. The option's effect on the two solver fixtures is large and in one case negative, and both are recorded: the carrier reload's handed-back ||R|| falls from 1.721 to **1.026E-01**, the key-off level (READ: N37 9.34E-02), with a measured momentum row of 1.404E-02 at cell 499 instead of a pinned 1; the atomic reload stops earlier (info = 2 at iteration 24) and hands back 1.702 against the control's 1.415 at cap 40, because the merit no longer carries 500 momentum rows pinned at

one ($||Fs||_2$ 17.1 against 28.2). A defect found while there and fixed: the allocation guard in `RK_rhs` tested `face_flux` for the well-balanced arrays `face_q_up/face_q_dn` as well, and `face_flux/face_p` are also allocated by `store_the_interface_fluxes`, so a stationary evaluation reached before any marching stage could write an unallocated array; the guards are now separate and each tests its own array, with no arithmetic on any path.

- P5: the analysis document corrected** (`docs/solver_approach_analysis_20260910.md`, +463/-115, 114 to 592 lines). All nine corrections the review established are in place, each with an inline `[corrected 2026-09-11: ...]` mark citing the review section or the experiment measurement behind it: the spatial truncation error is not a lower bound on the algebraic residual (the frozen-background H2 wind row measures $2.449021e-13$ on the same 500-cell grid where the withdrawn argument put a $5e-4$ floor); Ritz-magnitude ratios of a masked compression are neither condition numbers nor the species Schur complement; partitioning moves the coupled iteration outside the Krylov solve with outer matrix $(1 - \omega) I + \omega D^{-1} C A^{-1} B$ and the outer loop is not new code (`steady_wind_with_element_diffusion`); the CETIMB route is renamed as informed by CETIMB and gains the two conditions the review adds; the well-balanced option removes the equilibrium's algebraic cancellation and not the floor; Koskinen et al. 2013 lower boundary is section 2.1.1; the three-unknown path uses its existing globalization and not the trust region (`use_tr = (nspec_row .gt. 0)`); section 7 recast with the 2026-09-11 numbers; a new section 9 indexing the twelve changes and a new section 5.5 on partitioned components inside a coupled method (Knoll and Keyes 2004, J. Comput. Phys. 193, 357). Three numbers of the original disagreed with their sources and are corrected, one of them wrong: the N4b row for the carrier reload read "1.4 -> 0.09" where the measurement is 0.151 -> 0.090. Citations were verified independently in the published text (Koskinen 2013 section headers by `pdftotext -layout`; the Knoll and Keyes bibcode by the ADS API).
- P6: the outer iteration as a stated contract** (`src/EXHALE_main.f90`, +448/-131). `steady_wind_with_element_diffusion` now says in its own header which equations are alternated with the wind and why (a species the run moves by an operator-split transport step, the diffusing elements always and the molecular carriers where Coupled carrier solve is off, has a stationary balance that no hydrodynamic solve touches), and it accepts a state only by the certification of the FULL set of active equations evaluated on the refreshed state it would hand back. A hydrodynamic `info = 0` while an alternated species row refuses is not an accepted state, and a nonzero hydrodynamic flag no longer ends the loop by itself, because both `info = 1` and `info = 2` hand back a state the composition update still makes progress from. The endings are named (`outer_certified`, `outer_state_not_finite`, `outer_element_update_refused`, `outer_no_progress`, `outer_pass_budget`), the P1 status is consumed, and the quantity the passes are controlled by is the worst gated species row of the certification: where it fails to fall over a pass both updates are shortened (the element under-relaxation ω halved, floor 0.125; the carrier movement bound halved, floor `carrier_trust_floor = 1.0d-3`) and `outer_no_fall_max = 3` consecutive passes without a fall end the loop. The last pass takes no composition update, because an update exists to be consumed by the next hydrodynamic solve and taking it would replace a stationary wind by a state the update itself spoiled. The composition drift and the volume-weighted carrier residual are reported and are not tests. With no alternated species the body is one steady solve and the state refresh, and the flag is the solve's own, which is what an atomic run and a coupled carrier

solve both do. Default path MEASURED on the P6 run pair (wasp_full_newton.ctrl against .new): info = 0, ||R|| 1.667E-09, flux spread 6.233E-14, CERTIFIED, log10 Mdot 13.30, and output/Ion_species.txt and output/Hydro_ioniz.txt byte-identical between the two.

- **P7: the measurement.** Below.
- **P8: the corrections propagated to the documents and tests that carried them** (docs/code_status_20260910.md, docs/ISSUES_20260909.md, docs/certification_tolerance_anchoring_20260910.md, src/tests/steady_selfconsistent_residual/run.sh, src/tests/steady_species_rows/steady_species_rows_tests.f90). The withdrawn inference (a spatial truncation error of 5e-4 at N = 500 floors the algebraic residual, so 1e-5 cannot certify) is corrected in place in the three documents that still carried it, marked with the review section and the measured 2.449021e-13 beside it, and decision 22 (a) is kept as the standing decision in every one of them with only its justification changed; the grid-refinement anchor and its remap control are relabelled as consistency diagnostics; "permanently uncertified until the operator's order or the grid changes" is withdrawn and quoted where it stood; ISSUES 3.7's store_row_terms item is struck through and marked RESOLVED by P4, and the resolved table gains D1, D2 and D3 with their measured before and after. Two test contract errors, both RED before and GREEN after: steady_selfconsistent_residual assertion 2 read "a hydrodynamic row above its tolerance implies info = 2", which conflates not certified with one flag and already failed on the entry text for any budget-limited log (FAIL flag_of_carrier_elem_newton measured=info=1 reference=info=2 -> PASS, with wasp_full_newton still PASS and a synthetic info = 0 copy still FAILing); and relaxation_drift_covers_every_element built its column by swinging each metal mixing ratio about the reservoir on a fixed hydrogen fraction, so its species missed the density the column hands the operator by **1.66042E-02**, which is now **6.58484E-16** on a column of mass fractions that sum to one, the suite going 194/0 to **195/0** with a new row measuring the closure by the code's own mass policy.
- **P9: this section**, docs/session_handoff_20260911.md, the status section of the plan, the dated sections of the two fixture READMEs, the reopening paragraphs of docs/code_status_20260910.md section 3 and docs/ISSUES_20260909.md section 5, and the new entries of TO_BE_DONE.md.
- **P10: the best-iterate ledger reads the hydrodynamic rows against their own certification tolerances** (steady_newton.f90 +131/-55, src/tests/krylov_and_dogleg/krylov_and_dogleg_tests.f90 +135/-1, READ from the P10 report; the whole uncommitted diff of the two files now stands at +335/-84 and +268/-1, MEASURED by git diff --numstat, P3's share included). Items P10, P11 and P12 were added after the plan had run, on the user's instruction of 2026-09-11 to fix the open items the session had left to the user ("fix all of them", and "the pressure term too"); P10 is the third entry of the plan's "what remains". distance_from_certification divided the hydrodynamic slot by the run's Resid tol, so the ledger that chooses the state handed back could not see a mass row above 3e-12 while ||R|| was below 1e-8. The slot now arrives as a distance formed by ONE new public routine, hydrodynamic_distance_from_certification, max(mass/3e-12, momentum/1e-8, energy/1e-6) on the windows resid_relnorm uses, so d < 1 is exactly the row condition the loop-top stop of P3 and the acceptance at return impose, and keep_this_iterate ranks on that distance on every route including the three-unknown one (the nspec_row <= 0 branch that ranked

on $||R||$ is gone). `certified_row_measures` hands the plain $||R||$ of the same evaluation back through a new optional argument rather than leaving a caller to re-derive it, and the iteration line prints the two side by side. `resid_tol_of_solve` is KEPT, with one reader, the EXHALE_RESID_BRANCH_REPORT block, where `rmax(1)` is deliberately the plain mixed-row measure and dividing it by one row's tolerance would be a ratio of two different rows. Tests: `krylov_and_dogleg` 290/0 at entry, **300 PASS / 9 FAIL** on a RED build in which both distances read `maxval(rc)/1.0d-8` (the superseded reading, at the `Resid tol` of these fixtures), **309/0** after, the 19 new rows being the block `judget_distance_reads_each_row_against_its_own_tolerance`; certification 57/0, `steady_completion_flag` 3/0, `steady_selfconsistent_residual` 3/0 unchanged. Impact MEASURED (P10) against a private build of the entry text of the one file: `wasp_full_newton` reloaded ends `info = 0`, $||R||$ 1.667E-09, `log10 Mdot` 13.30 with **all eleven output files byte-identical**, the ledger handing back the same iterate and the only number that moves being the reported best-iterate distance, 1.749E-01 (which is $||R||_{best}/1e-8$) to 8.469E-02 (the same iterate's worst row over its own tolerance, so 8.5 percent of it). The carrier reload shows the re-ranking doing what it is for: pass 2 hands back a mass row of **4.85E-12 at $||R||$ 2.459E-09** where the control took 6.98E-12 at 1.593E-09, and pass 3 **4.96E-12 at 2.005E-09** against 8.17E-12 at 1.723E-09, with the H2 carrier row and the worst gated species row identical pass by pass because the carrier relaxation sits outside the hydrodynamic solve. The HD 209458 b element reload runs its twelve passes in 1064 s against the control's 1043 s (2 percent on a shared machine, not a claim) with every one of the twelve worst gated species rows and its cell identical to the control, and ends on the same single refusing entry, the hydrodynamic mass row at **8.893E-10 at cell 3** against the control's 1.054E-09 at cell 16; the solves reporting a stalled best iterate fall 5 to 3 and those ending on "no descent direction exists" 11 to 10. What stays open, and what this measurement settles: the mass row does NOT reach 3e-12 and that state still does not certify, because at 1e-9 to 1e-12 the row IS the N33 rounding of the base-layer flux difference. MEASURED (P10, finding 3): at pass 1 the ledger's best distance is 4.407E+02, a mass row of 1.32E-09, while the same state re-evaluated at its own composition after the best-iterate restore carries **4.11E-09**, so ranking iterates by that row is ranking by the luckiest rounding of a quantity that moves when the state is evaluated again, which is why the pass-by-pass mass rows scatter rather than descend in the control and in the new run alike. The entry is an acceptance question, not a solver item, and no further solver item follows from it.

- **P11: the momentum row's reference scale is the largest PHYSICAL term of the momentum equation** (`RK_rhs.f90`, `steady_residual.f90`, `src/tests/grid_and_gates/hydrostatic_residual.f90` and that suite's `README.md`; the whole uncommitted diff of the four files stands at +285/-2, +151/-33, +736/-12 and +94/-1, MEASURED by `git diff --numstat`, with P4's +90/-2, +59/-2, +318/-11 and +44/-1 inside it). The fourth entry of the plan's "what remains", and the "pressure term" of the user's instruction. The scale was built from the pieces of the discretization, $\max(|dF(2)|, |S(2)|, |Smom|)$ plus P4's equilibrium pressure force under the well-balanced key; it is now built from the three left-hand terms of $d(\rho v)/dt + \text{div}(\rho v v) + dp/dr + \rho d\phi/dr = S_{\text{visc}}$, kept beside `face_flux/face_p` as module arrays of `RK_integration`, filled on both reconstructions, under both key values and on the positivity-repair path, and gathered by one routine, `momentum_row_terms_of_cell`, with `momentum_row_terms_of_state` filling a whole state and `reconstruction_continuation_rhs` blending the three terms with the same `lambda` as `dF` and `S`. The

residual $R(2) = dF(2) - S(2) - S_{\text{visc}}$ is untouched on every path, and the split is a rearrangement of the assembled row and not a model of it: $F(2) - p_{\text{out}}$ is the momentum flux without the pressure exactly for every flux function, and the geometric term is recovered from the source as $p_{\text{geom}} = S(2) + \text{weight}$ so that no second definition of the cell pressure exists. MEASURED (P11), on a state with structure in density, pressure and velocity, $\max_j |r_{\text{am}} + dp/dr + \rho g - (dF_2 - S_2)|$ over the scale is **0 under WENO3 and 1.7e-16 (default) / 5.3e-15 (key on) under PLM**, for both Riemann solvers, so the three terms add up to the row they scale. The defect is closed on the states where the terms are known in closed form: with no gravity and a uniform pressure at rest, where every term of the equation vanishes, the PLM scale read **1.000 of the geometric pressure term $(A^+ - A^-) p_j/dV$** , that is $2 p/r$, with both Riemann solvers at every N, and now reads the tiny floor 2.22507E-308; on supersonic uniform flow at $M = 3$, where the ram divergence is the only term, it stood **6.66667E-02 = $1/(\gamma M^2)$ above it** and now agrees with it to 1.9e-14 (PLM) and 4.2e-13 (WENO3); on the discrete hydrostatic equilibrium the PLM scale was **2.79144E-01, 28 percent BELOW the weight it balances**, because $S(2)$ is the weight minus the geometric term and no term of the max was the weight itself, and the departure $(w - s)/w$ is now 0. P4's own zero-gravity diagnostic closes with it: the departure of the well-balanced scale from the base scheme's on the same state falls from **9.43654E-01 to 3.74E-13** under PLM with ROE (2.76E-13 with HLLC), the two having agreed under WENO3 all along (1.7e-13 to 2.7e-13). Tests: hydrostatic_residual **50 PASS / 10 FAIL** on the entry-text objects and **60/0** after, the ten being the four scheme/solver pairs of P4's corrected zero_gravity_momentum_scale_is_the_rows_own_terms (which had compared the option's scale with $\max(|dF_2|, |S_2|)$, the SUM of the two terms the row holds, 7.50E-01 FAIL to 0) and the PLM pairs of the three new identity statements; the whole grid_and_gates suite **188/10 to 198/0** (198 = P4's 178 + 20 new rows); krylov_and_dogleg 290/0, certification 57/0 and adv_static_limit 53/0 before and after. Two quantities are REPORTED and not asserted, with the reason: the two-sided departure of the scale from the weight on the well-balanced scheme's own equilibrium is not round-off for the base scheme (MEASURED 8.5e-4 at $N = 250$ falling to 2.3e-6 at $N = 2000$, PLM with ROE), because that state is the discrete equilibrium of the WELL-BALANCED reconstruction and HLLC's face pressure carries a dissipative part of size dr/H times the weight; and the halving statement subtracts the unperturbed row of the same cell, since the undifferenced ratio is not 2 (MEASURED 1.0012 at $N = 250$, 1.586 at $N = 2000$) while the imbalance the perturbation ADDS halves to 2.000000 within 5e-6. Impact: nothing that was certified stops being certified. wasp_full_newton reloaded with the key OFF is **CERTIFIED** on both binaries, info = 0, log10 Mdot 13.30, with the momentum row moving **8.469E-10 to 1.165E-09** of its 1.0E-08 tolerance, the mass row 1.453E-13 to 2.017E-12 of its 3.0E-12 (so this reload now certifies with less margin on the mass row) and $||R||$ 1.667E-09 to 1.495E-09; the outputs are not byte-identical, because the merit and the loop-top gate both divide by the momentum scale and the two solves accept different iterates, and the states differ by at most **2.6E-09 relative** over every column (9.3e-10 in Hydro_ioniz.txt, 2.6e-09 in Ion_species.txt). With Well balanced: True the same reload is **CERTIFIED** on both, the momentum row moving **9.719E-10 to 9.621E-11** with all eleven output files byte-identical and the $||Fs||_2$ ladder identical iteration by iteration: at cell 500, $r = 4.62$, the ram divergence exceeds the cell's weight by about a factor ten, which is why the measure of the certified state falls by that factor on a supersonic outer cell. The marching path of mol_carrier at 300 steps and one thread is **byte-identical in all seven output**

files, log10 Mdot 7.95, the run.log differing in exactly eight lines, every one of them a report of the momentum measure (the worst-row entry 1.8164E+00 at cell 438 to 1.9788E+00 at cell 449) and none of them a state. On the carrier reload every pass line is identical but the momentum entry, which falls **1.32E-12 to 3.22E-13** after pass 1 (4.47E-13 and 3.54E-13 at passes 2 and 3), the state as loaded reading 3.622E-01 to **3.311E-01** in its scaled momentum row, and all four output files byte-identical. What stays open: the kind-generic EXHALE_RESID_QUAD assembly still hands back neither the momentum terms nor the face departures, so under that option (default off, a rounding control experiment nothing is adopted from) both are read from the last RK_rhs evaluation rather than from the state just assembled; the fix belongs in hydrodynamic_rows.f90, which this item does not own.

- **P12 (with P12b): two test-hygiene defects, and the same shape in the sibling scripts** (src/tests/test_columns.f90 new, 125 lines; attempted_step/run.sh +18/-5, carrier_returned_state_acceptance/run.sh +18/-5, carrier_retry/run.sh +18/-5, carrier_reference_scales/run.sh +22/-6, carrier_constraint_attribution/run.sh +22/-6, constrained_network_layout/run.sh +22/-6, species_masses/run.sh +22/-8, element_operator/run.sh +5/-0, steady_species_rows/run.sh +5/-0, and the two drivers, MEASURED by git diff --numstat; of the drivers' totals the item's own share is READ from the P12 report, the use line, the renamed calls and 66 lines out of element_operator_tests.f90 and 43 out of steady_species_rows_tests.f90). The fifth entry of the plan's "what remains", both halves of it. The **build_stamp fallback**: the trigger is narrower than "a tree that has never been built", it needs such a tree AND a work directory outside \$root/build, which is the concurrency-safe usage the worker rules mandate, because source_closure.py walks \$root/build recursively and a default work directory inside it swept the fallback into the closure by accident. Seven run.sh now select \${EXHALE_OBJDIR:-\$root/build}/build_stamp.f90, write the fallback into their own work directory only when that is absent, and compile whichever of the two is in force as the FIRST file of the closure with the closure filtered of any second copy, which is the shape src/tests/physics_probe/run.sh already had; grep -ln 'if [! -f "\$root/build/build_stamp.f90"]' src/tests/*/run.sh now returns nothing. Four of them gained a private object directory, EXHALE_TEST_OBJDIR, without which the no-build verification cannot be run on them at all (species_masses, which had no override of any kind, carrier_reference_scales, carrier_constraint_attribution, constrained_network_layout). RED and GREEN, MEASURED in four cases: at HEAD with an external work directory the compile dies on Cannot open module file 'build_stamp.mod' and each script prints its one failure line, **0 PASS / 1 FAIL, exit 1**, in attempted_step, carrier_returned_state_acceptance, carrier_reference_scales, carrier_constraint_attribution, carrier_retry and species_masses; after the fix every suite gives the count it gives at HEAD with a default work directory, **70/0** (attempted_step), **36/0** (carrier_returned_state_acceptance), **14/0**, **10/0**, **122/0**, **9/0** and **8/0** (constrained_network_layout, which has no utilities.f90 in its closure, so it has no RED to show and its change is the shape alone), in a tree with no build/ and in the repository against a private build, and the diff of the PASS/FAIL lines before and after is empty for all of them. Which stamp each case used is MEASURED too: in the no-build case every work directory holds a build_stamp.o compiled from the fallback, and in the case with a build tree no fallback file was written at all, so the generated stamp was

taken, which the superseded scripts could not do because they looked only in \$root/build. The **column written twice**: the new module test_columns holds column_carrying_its_own_density(f_c, q_h2, he_taper, metal_swing), named for what it builds, with its header stating the invariant $\sum_i m_i n_i = \rho$ and the round-off it holds to, and column_mass_closure(rho_c, f_c), the measurement moved out of element_operator_tests.f90 unchanged; the two drivers call them at 15 and 2 sites and use utils, only: calc_rho is gone from both, it having been there only for the moved code. The shared construction reduces to each former copy exactly and not approximately (metal_swing = 0 gives swing = 1.0d0, q_h2 = 0 leaves isp_H2 at zero, he_taper = .false. selects the constant the steady driver used, and zsum accumulates in the same element order), and the statement is the measurement: element_operator **24/0** and steady_species_rows **195/0** before and after, with the diff of the two full logs, diagnostic lines included, empty in both, so every printed digit is unchanged (the_column_is_mass_closed_as_built 4.10591E-16, drift_column_carries_its_own_density 6.58484E-16, the relaxation's 27 steps and returned drift 1.7790E+00 among them). The file is in no source list of the Makefile (grep -c test_columns Makefile = 0) and no dependency generator sees it; the two run.sh that need it compile it immediately before their driver. Nothing stays open from this item: its own report found nothing further, and the four sibling scripts and the hardcoded work directory it reported are what P12b then closed.

Two changes outside the item list, made by the advisor while reading these three:

- src/tests/steady_species_rows/steady_species_rows_tests.f90, the two rows P10 reported as stating the superseded semantics. In judged_distance_is_the_worst_row_over_its_tolerance the hydrodynamic slot now arrives as a distance (rows(1) 2.0d0, 0.5d0 in place of 2.0d-8, 5.0d-9), with a comment saying so, and judged_distance_follows_the_residual_target, which asserted the property the item removes, is replaced by hydrodynamic_distance_reads_each_row_against_its_own_tolerance: a mass row of 1e-9 beside a momentum row of 1e-13 and an energy row of 1e-8 gives 1e-9/3e-12, the mass row being what the distance reports. The two resid_tol_of_solve assignments and that import are gone.
- the Coupled carrier solve key comment in src/modules/files_IO/input_read.f90 (+17/-13, MEASURED). It stated that the alternation "was measured not to converge and not to be able to" (section 139), which is narrower than it reads after P7. It now names the default (False), says what the alternation is judged by (the certification of the refreshed state in steady_wind_with_element_diffusion), cites the two reloads of this section for what it reaches, and keeps True as the coupled measurement option.

Four more items ran in the afternoon of the same day: **P14** and **P15**, the two findings of the Codex adversarial review of the working tree (the review and the verification of each finding are the subsection below), and **P16** and **P17**, the user's first decision of the afternoon (the mass tolerance re-anchored on the fixture's measured rounding floor) and its consequence for the best-iterate ledger. **P18** followed and closed the same day.

- **P14: every ending of the stationary outer iteration hands back one consistent state** (src/EXHALE_main.f90, steady_wind_with_element_diffusion only, +84/-40 against the entry text; src/tests/grid_and_gates/output_state_consistency.sh +181/-4; MEASURED by P14 with diff -u

and `git diff --numstat`). Codex finding 1, and it is real. On a stagnation ending the composition had been updated while the particle densities and the temperature written beside it had not, so the state the run wrote was one composition update ahead of its own particle count and temperature. MEASURED (P14) on the forced stagnation of the hot-Uranus carrier reload: re-evaluating the written state moves its temperature by **1.317089E-08** relative at cell 295, where the same fixture ending on its pass budget, which takes no update, moves by **5.921E-16**; the written temperature of the worst cell is $1.768302856793E+03$ where the same conserved state and composition give $1.768302833502E+03$, and the worst species column is $2.543E-03$ (He 2^3S , the stage answering to the temperature with the steepest exponential). Four changes. The progress control (the `sp_worst` test, `n_no_fall`, the omega and trust halving, the setting of the stagnation ending) now stands between the certification of the pass and the update the next solve would consume, guarded so that a certified or non-finite pass reaches neither, and the update's own gate therefore excludes the stagnation ending by itself: the pass that ends the iteration takes no update. The ending is announced with the other endings, below the summary line and the diagnostics, so the pass that ends on it prints its summary, its carrier diagnostics and its H2 front like every other pass. A pass that takes no update says which of the two reasons it was. And the element path's update now refreshes the particle densities and the temperature between `relax_element_composition` and the equilibrium sweep (`get_species_densities`, `comp_T_from_p`), which is the shape the carrier path already had; before it, the sweep equilibrated the new composition at the old temperature. The header states the invariant. One behavior change to record because it is not a reordering: the shortened step of a pass whose measure did not fall is now consumed by THAT pass's update, where before it reached the next pass's. RED and GREEN on the new second case of `output_state_consistency.sh`, which runs the stagnation (`EXHALE_CARRIER_TRUST=1e-4` `EXHALE_OUTER_PASSES=8` `EXHALE_JFNK_MAXIT=5`, the setting that stagnates; the brief's cap of 40 does not, the worst gated H2 row falling through the hydrodynamic solve alone), asserts by name that the ending reached is the stagnation one, and then measures the Restart intent: stationary evaluate round trip over the 500 physical cells against **1e-12**: measured= $1.317089e-08$ FAIL on the entry text, measured= $6.499247e-14$ PASS after. The tolerance is justified by measurement and not by a snapshot: the states handed back by an ending that takes no update measure $6.5E-14$ and $5.9E-16$, and the defect measures $1.3E-08$, four decades above the allowance. The suite goes **199 PASS / 1 FAIL** on the entry text to **200 / 0**. Impact: `wasp_full_newton` reloaded is unmoved (`info = 0`, CERTIFIED, `|R|` $1.495E-09$, flux spread $1.9113E-12$, `log10 Mdot` 13.30 on both binaries, all eleven outputs identical apart from the provenance run= timestamp); the carrier reload at three passes is identical pass line by pass line to the control and to the P7 record; the element reload's pass 1 is identical in every digit and the ladder parts from pass 2, because the sweep of pass 1's update now runs at the relaxed composition's own temperature (momentum $1.60E-12$ against $4.24E-14$ at pass 2), both builds ending on the pass budget with the same 9 refusing equations and the same base-layer mass row as the worst refusing entry.

- **P15:** the kind-generic rows hand back the face departures their momentum row was built from (src/modules/time_step/hydrodynamic_rows.f90 +75/-15, src/modules/time_step/hydrodynamic_rows_body.inc +51/-17, src/tests/krylov_and_dogleg/krylov_and_dogleg_tests.f90 +255/-1 measured against the file as

it stood when the item started; MEASURED by P15). Codex finding 2, and the third entry of the plan's "what remains". Under Well balanced: True the momentum row's pressure-gradient term, the momentum row scale and every measure read against that scale are formed from the module arrays face_q_up/face_q_dn of RK_integration, which only RK_rhs and the positivity repair ever filled; the kind-generic rows kept their own departures as locals and store_the_interface_fluxes wrote back face_flux and face_p alone, so in the kind-generic rows those three quantities belonged to the last RK_rhs evaluation or to the zeros of the first allocation. The departures now leave the kind-generic text as two OPTIONAL intent(out) arguments of hydrodynamic_flux_difference_and_source, converted to double at the one rounding on the way out exactly as the flux and the face pressure are, both public wrappers ask for them, and store_the_interface_fluxes stores them under the key. The reconstruction continuation CAN reach the kind-generic rows (assemble_residual chooses the text from generic_precision_rows_arm() alone and the generic text carries its own blending branch), and on it the terms would be wrong for two reasons rather than one, the single-scheme weight w_face_p and the fact that the two schemes reconstruct different faces so no single pair of departures satisfies the blended row identity; the_generic_text_is_usable therefore now refuses recon_lambda_on .and. $0 < \text{recon_lambda} < 1$, ahead of every evaluation of that text. That branch ends in error stop and has no executed row, which is stated rather than worked around. Twelve new rows, each on a state the kind-generic rows assemble AFTER a residual was assembled on a different state, both reconstructions and both precision kinds: MEASURED RED **2.098877E+01** (PLM) and **2.032204E+01** (WENO3) on the pressure gradient against the production term, **3.262392E+00** and **3.428893E+00** on the row scale, **1.190363E+01** and **1.181676E+01** on the row rebuilt from the stored face data, and GREEN at **0 bitwise** for the generic-double instantiation, which IS the production operator bit for bit, and 2.6E-16 to 2.7E-15 for the quadruple one at its 1e-13 bound. krylov_and_dogleg **309/0 to 321/0**; certification 57/0 and steady_species_rows 195/0 unchanged. Impact: wasp_full_newton reloaded is unmoved with the key off, and with Well balanced: True and EXHALE_RESID_QUAD=2 it stays CERTIFIED at the certified momentum row **9.621E-11**, P11's key-on production value, on both binaries, because the JFNK and PTC loops reset the sweep kind to marching before the certification of the handed-back state and the kind-generic text is consulted only for a non-marching sweep kind (READ, steady_newton.f90); the brief's expectation that the control's certification reads another state's departures is corrected by that measurement. What the defect did move is the measure INSIDE the loop, where the kind-generic rows are armed: the convergence measure of iteration 1 is **1.756E-03** on the control against **1.753E-03** after, the momentum row divided by the pressure-gradient term of another state and overstated by 0.17 percent, and that measure is what the loop-top stop of P3 and the ledger of P10 read on every iterate. ||Fs||2 is identical iteration by iteration, iterations 2 to 7 are identical line for line, the accepted state is the same state, and mol_carrier marching is byte-identical.

- **P16: the hydrodynamic mass-row tolerance is anchored on the row's measured rounding floor** (src/modules/time_step/certification.f90 +212/-2, src/modules/time_step/steady_residual.f90 +177/-1, src/tests/certification/certification_contexts.f90 +105/-0, docs/certification_tolerance_anchoring_20260910.md +302/-0 appended; MEASURED by P16 with git diff --numstat, the share of the other uncommitted item in steady_residual.f90 sepa-

rated by attributing each hunk). The user's first decision of the afternoon, and it CLOSES the first entry of the plan's "what remains": the tolerance is now $\max(\text{cert_tol_mass}, \min(\text{cert_tol_mass_ceiling}, \text{cert_mass_round_margin} * \text{floor}(j)))$ with $\text{cert_mass_round_margin} = 10$, the ceiling 1 (a row at 1 is a mass flux changing by the whole of itself across the cell, not an operative branch on any state measured), $\text{cert_tol_mass} = 3\text{e-}12$ unchanged and now the FLOOR of the tolerance, and the verdict on that row taken cell by cell through the pure $\text{mass_row_cell_verdict}$. **The floor expression the item's brief specified could not serve, and the report of that is part of the result:** $\text{eps} (|A^+ F_{\text{up}}| + |A^- F_{\text{dn}}|)/dV / \text{residual_row_scale}(1, j)$ is algebraically between one and two ulps at every cell, because the row's scale IS the larger of the two addends and the dV cancels, and MEASURED it is **4.4409E-16 on all 500 cells of all four states** looked at, so ten times it never rises above $3\text{e-}12$ and the item as written would have been inert. The cause is the one N33 established: the interface flux carries the last bit not of its own value ρv but of the $O(\rho c_s)$ quantities the Riemann solve differences, so the floor is about eps/Mach and not eps . The estimate adopted is $\text{eps} \rho (|v| + c_s) A / dV$ over the row's own scale, and it bounds the row's measured non-smoothness within a factor **1.5931** (wasp_full_newton as loaded, base Mach $3.937\text{E-}03$), **1.8857** ($\text{atomic_elem_newton}$ as loaded, $5.062\text{E-}06$), **2.6806** ($\text{carrier_elem_newton}$ as loaded, $2.400\text{E-}04$) and **1.7733** (the HD 209458 b state this item certifies, $2.767\text{E-}06$), where the specified expression is short by two to six decades. The margin is measured, not chosen: $c_{\text{round}} = 10$ is a round number 3.7 times above the largest of those four ratios and inside N33's independent reading of the same floor against its own bound, 5.5 to 12 (READ). The measurement itself is in the code: EXHALE_MASS_FLOOR_SCAN=1 (new, default off) adds one ulp to the density of every physical cell, runs the whole flux assembly again and reports the STEP each mass row takes beside both estimates, and the state given is then assembled once more and reproduces its own mass rows to $0.00\text{E+}00$ on every fixture. Tests: certification **57/0** on the entry driver against the entry objects, **0 PASS / 1 FAIL** with exit 1 on the new driver against the entry objects (the compile naming the three new symbols), **70/0** after, the eleven new rows asserting that a floor below the fixed value leaves exactly $3\text{e-}12$ and names the fixed tolerance, that a floor above it gives exactly $c_{\text{round}} * \text{floor}$ and names the rounding anchor of that cell, that a row at half the anchor is WITHIN where the superseded rule transcribed inside the test refuses it, that a row above the anchor still refuses, that the ceiling binds, and that the margin stands above every ratio measured. Seven further suites are identical row for row in both builds, and a $-f\text{check}=\text{bounds}, \text{do}, \text{mem}$ build runs both reloads with no runtime trap (the floor estimate reads the ghost cell at $j = 1$, which is why it was checked). **Impact.** wasp_full_newton reloaded is unmoved: CERTIFIED in both builds, mass row $1.212\text{E-}12$ at cell 500 with the verdict taken there against the FIXED $3.0\text{E-}12$ at distance 0.404, $\log_{10} \text{Mdot}$ 13.30, outputs byte-identical apart from the provenance timestamp; its floor is below $3\text{E-}13$ at every cell, so the anchor cannot reach that case, and the cold-start wasp_full_newton of make check cannot be affected either, its stop having no row binding (READ from backup/regression/ $\text{wasp_full_newton}/\text{run.log}$) and a tolerance that only ever rises cannot move it. **The HD 209458 b element reload CERTIFIES, at pass 12**, outer pass 12: ACCEPTED -- every active equation of this state is within its own tolerance, where the control (a build of the entry text of the two files in the same tree) ends the same twelve passes NOT CERTIFIED on the one entry it had before, the hydrodynamic mass row at $7.159\text{E-}10$ against $3.0\text{E-}12$ at cell 13. On the certified state

the largest mass row is 7.703E-10 at cell 2 and the verdict is taken at cell 34 ($r = 1.00665$), 6.542E-10 against **4.6E-09, the rounding anchor of that cell**, distance 0.141; the eight elemental wind rows and the worst gated species row of all twelve passes are identical to the control to three digits, cell included, and the two builds part at pass 6, the first pass whose hydrodynamic solve stops on the loop-top test because its mass row is inside its own tolerance (info = 0 where the control's is info = 2). The anchored tolerance still refuses states: the mass row is admitted at 11 of the 12 passes and refused at pass 3 at distance 69.9. On the hot-Uranus carrier reload the base cells are admitted by their own anchor at every pass and the refusal moves OUT to cell 180 ($r = 1.0707$), where the estimated floor is below $3e-12$ and the row carries signal, refusing 5.067E-12 to 9.549E-12 against the fixed $3.0E-12$: the refusal has moved from rounding to signal. The H2 carrier row is unchanged pass by pass in both builds (7.38E-02 at cell 290, 7.22E-02 at 289, 7.06E-02 at 288), and the three passes cost 15.47 s against the control's 69.39 s. The derivation, both candidate estimates, the measured ratios and all three impact measurements are the appended section "**Anchor (6), 2026-09-11: the mass row's rounding floor**" of docs/certification_tolerance_anchoring_20260910.md; its nine earlier sections are byte-identical.

- P17: the best-iterate ledger reads the mass row against the cell's own tolerance** (src/modules/time_step/steady_newton.f90 +146/-32, src/tests/krylov_and_dogleg/krylov_and_dogleg_tests.f90 +98/-0; MEASURED by P17). P16's own outside-scope finding, and the consequence of the anchor for the ledger: the hydrodynamic slot was formed as $rc(1)/cert_tol_mass$ from a triple of row maxima, so the ledger ranked the mass row against $3e-12$ while the stop test and the acceptance read the cell's own tolerance. The slot is now max over cells j and rows k of $|R_{kj}| / residual_row_scale(k,j) / tol_k(j)$ over the whole physical column, with $tol_1(j)$ taken from the certification's own rule at that cell through P16's public `mass_row_cell_verdict` (so the anchor formula is not restated in the solver module) and the fixed momentum and energy tolerances on the other two rows; that is the same functional the acceptance applies cell by cell, so $d < 1$ in the ledger is now exactly the hydrodynamic part of the verdict. `rnorm` is unchanged and is now formed only when the caller asks for it, which the two loop-top calls do and the three trial calls do not. The superseded `hydrodynamic_distance_from_certification` keeps its name, signature and body and is documented as what it is, a BOUND from above on the distance ($tol_mass_at(j) \geq cert_tol_mass$ at every cell), because its removal would break the compile of a test file P17 does not own; P18 removes it. MEASURED RED twice: the new driver does not build against the entry-text objects (exit 1, Symbol 'hydrodynamic_distance_from_certification_by_cell' ... not found in module 'steady_newton'), and a private build in which the new function's body is replaced by the superseded reading and nothing else is altered gives **327 PASS / 7 FAIL** with the seven rows naming the column (the_wind_cell_holds_the_mass_rows_distance 3.333333E+02 against 3.333333E+00, the_base_cell_is_a_decade_inside_its_own_tolerance 3.333333E+02 against 1.000000E-01, is_a_certified_hydrodynamic_state F against T among them). GREEN krylov_and_dogleg **321/0 to 334/0**, and seven further suites identical in both builds. **Impact.** The single number that says the item worked, on the HD 209458 b element reload: the ledger's best judged iterate: distance over the twelve passes runs **2.2E+02 to 6.3E+05 in the control and 1.321E-01 to 1.868E-01 after**, while the acceptance CERTIFIES the pass-12 state in both builds at a mass-row distance of 0.141 (control) and 0.132 (new, verdict at cell 50, r

= 1.00977, 4.245E-10 against the 3.2E-09 anchor). The control's ledger therefore ranked every iterate of every solve as hundreds to hundreds of thousands of times outside certification, on a row its own acceptance admitted; the ledger and the acceptance now speak about one quantity. The trajectories part at pass 2 and the elemental relaxation is not damaged (the worst gated species row descends 2.61E-03 to 7.73E-06 against the control's 2.61E-03 to 6.84E-06, both inside 1e-5 at pass 12, and the certified state's elemental rows are SMALLER in the new build on every element, Fe 5.086E-05 against 8.966E-05, C 3.70E-06 against 1.00E-05, O 3.34E-06 against 1.24E-05). The cost is not measurable: 909 outer iterations at 1.206 s against 861 at 1.216 s, so the 500 rounding-floor evaluations added per judged residual cost nothing this measurement can separate from noise. More restarts and fewer stagnations, which is the reading of a ledger that now stops improving where $||R||/3e-12$ used to keep moving on rounding noise: "best iterate has not improved" fires in 7 solves against 4, "no descent direction exists" ends 11 against 8, and the control's one STAGNATED solve is gone, while no solve fails and every pass still returns a state. On the hot-Uranus carrier reload the consequence is a verdict: the hydrodynamic solve returns **info = 0 at all three passes** against the control's 0, 2, 2; the mass row's verdict is WITHIN at passes 2 and 3 (cell 1, 1.826E-11 against the 2.5E-11 anchor, and cell 2, 6.437E-12 against the same) where the control refuses at cell 180; the state of the last pass has **1 refusing entry, the H2 carrier row alone**, against the control's 2; and the best judged distance runs 0.5282, 0.8819, 0.2578 against 1.630, 3.501, 1.840. The H2 carrier row is identical pass by pass in both builds and in the P7 record (7.38E-02, 7.22E-02, 7.06E-02 at cells 290, 289, 288), which is what must hold with the carrier relaxation outside the hydrodynamic solve. wasp_full_newton reloaded is unmoved, and for a stated reason: its estimated floor is below 3e-13 at every cell, so the cell's tolerance IS the fixed value everywhere and the maximum of the ratios is the ratio of the maxima; the two run logs are identical line for line except the Execution Time, every written file is byte-identical except the provenance timestamp, and the best judged distance is 6.722E-01 in both builds. A separate fix in the same file, reported on its own: write_gate_met_while_a_row_refuses printed the largest measure of a row and its cell beside the tolerance of the BINDING cell, a pair of numbers belonging to two different cells, and now names the binding cell throughout where the row records one, which is what the certification's own refusing entry list already did; no suite parses that line and the carrier rerun on the final binary reproduces every reported digit.

- **P18: one entry point for the hydrodynamic distance** (steady_newton.f90 +23/-50, krylov_and_dogleg_tests.f90 +49/-25, steady_species_rows_tests.f90 +22/-16; P17's own outside-scope finding). With the ledger reading the mass row cell by cell (P17), the fixed-tolerance form hydrodynamic_distance_from_certification(rc) had no production reader left and survived only through test rows. It is DELETED, with its public export and the then-unused cert_tol_mass import of the module; every test row that read it restates its fact through hydrodynamic_distance_from_certification_by_cell on a one-cell column whose rounding floor (1e-30) puts it under the fixed tolerance, so the restatement is the same double division and every literal reference is unchanged. MEASURED (P18): `grep -rn hydrodynamic_distance_from_certification src/ | grep -v _by_cell` returns nothing; krylov_and_dogleg 334/0 before and after with no row lost (two renamed: the_fixed_tolerance_alone_reports_the_base_cell,

and_the_fixed_reading_is_never_below_the_distance), steady_species_rows 195/0, certification 70/0, steady_completion_flag 3/0; a compile RED (the entry-text drivers against the delivered objects fail on the missing symbol), a semantic RED of the cell-by-cell fact (327/7, P17's seven rows) and a semantic RED of the restatement itself (329/5, the fixture's floor raised into the anchored regime) all discriminate. Impact: the wasp_full_newton reload gives info = 0, CERTIFIED, ||R|| 1.495e-9, log10 Mdot 13.30 in both builds, all eleven output files byte-identical, the two run logs differing in the execution-time line alone. Documents that named the deleted function as current fact (certification_tolerance_anchoring_20260910.md, session_handoff_20260911.md) were corrected by the advisor.

- **P20: the code-size appendices moved to this log and were re-measured** (docs/Update_EXHALE_appendix.tex new, docs/Update_EXHALE_stage1.tex appendix removed and its \date fixed at 2026-09-04, src/utils/update_log_to_tex.py postamble, pointer lines in README.md and README_HOWTO.md). The two appendices, the code-size summary against the original ATES and the list of source inherited unchanged from it, were stranded in the stage-1 log and stale at 2026-08-30. They are now a hand-maintained fragment the generator \inputs after the converted body, so appending to this log keeps them. Re-measured 2026-09-11 with python3 src/utils/codesize.py (MEASURED, all of it): the Fortran tree is **205 files / 76,259 SLOC**, 21.7x the original ATES (47 files / 3,514 SLOC), against 111 files / 21,018 SLOC and 6.0x at the 2026-08-30 edition; the core excluding the Wind-AE port is 169 files / 71,503 SLOC, of which the production build is 95 files / 50,762 SLOC (14.4x) and the src/tests/ drivers are 74 files / 20,741 SLOC, the rule the script applies being every .f90 under src/ outside src/modules/wind_ae/. 123 new core modules (49 production, 74 test drivers), 38 of the 47 original files modified, and the files **byte-identical with upstream are down from 13 to 8**: hybrd, hybrd1 and fdjac1 took the unit_step_floor difference-step argument, Source.f90 the well-balanced option (N37), J_inc.f90 the single-spectrum-type decision, and speed_estimate_HLLC.f90 was deleted as dead code, so one upstream file is now removed rather than none. Update_EXHALE_stage1.pdf 464 -> 460 pages, Update_EXHALE_stage2.pdf 139 -> 158 pages (MEASURED, pdfinfo).
- **P21: the carrier relaxation solves the chemistry of the composition it advances** (src/modules/lower_atmosphere/diffusive_photochemistry.f90 +115/-27, src/EXHALE_main.f90 +17/-7, src/tests/carrier_retry/carrier_retry.f90 +287/-16; these three counts are MEASURED by diff -u against the entry texts saved in the scratch directory before the first edit and NOT by git diff, for the reason in "An external commit during the session" below). Verdict: the design does what it was meant to do at the level it was meant to act. With the chemistry equilibrated after every transport step the pass keeps, the unbounded pass at a fixed wind converges to a state that really is stationary for the coupled operator: on the hot-Uranus carrier reload the carrier steady residual falls from **7.99E-02 to 1.42E-09** in the worst cell and from **2.32E-03 to 1.18E-11** volume-weighted (MEASURED, P21, M1 below). The frozen fixed point was not a fixed point of transport plus chemistry, and this is the measurement of that. Two negative results come with it and are in the same subsection: at the production bound nothing moves, and the joint fixed point at a fixed wind is a fully molecular column.

The design as implemented. A new private routine `equilibrate_chemistry_at_fixed_pressure(rho, p, f_sp, T, heat, cool, eta)` takes `get_species_densities` (the particle count and the electron density of the composition), then `comp_T_from_p` (its temperature at the conserved state's own pressure), then `ioniz_eq`. No new module dependency was needed in either direction and `ionization_equilibrium.f90` is not edited. It is called inside the trial loop of `relax_photochemical_composition` immediately after a trial is ACCEPTED and before the fixed-point test, so a trial the operator refuses, or one that leaves the movement bound, is undone and halved before any sweep is taken and leaves the chemistry exactly where it found it. **The sweep IS the background reinstall:** `ioniz_eq` writes `bg_cell(j) = ieq_cell` for every cell of a molecular sweep (READ), while `install_background_of_adopted_state` belongs to the steady solver's best-iterate bookkeeping and is not involved here. The sweep is handed the transported partition and fixes `x(H2)`, the oxygen carriers and, where the protons are carried, the ionized fraction from the composition it reads, so it returns the chemistry OF that partition and does not re-solve it. **The movement bound is read on the transport output, before the sweep:** the bound governs what the transport operator moved in one trial, and the chemistry then closes on the step that was kept. MEASURED consequence: the reported drift of every bounded production pass stays at 9.98E-03 to 1.00E-02 at trust 1e-2, so the sweep does not carry the returned state out of its bound. The fixed-point test is now taken between two states the chemistry has closed on (`fnow` is recomputed after the sweep and becomes `fprev`), so `carrier_relax_fixed_point` means the carriers AND their chemistry stopped moving together. Interface: `relax_photochemical_composition(rho, v, p, T, f_sp, heat, cool, eta, trust, drift, nstep, outcome)`, with `p intent(in)`, the conserved state's own pressure untouched, and `T, heat, cool, eta intent(inout)`, written only where a refresh happened. The carrier branch of `steady_wind_with_element_diffusion` passes the new arguments and its own `ioniz_eq` call is DELETED as redundant; the `get_species_densities` and `comp_T_from_p` beside it are KEPT, because they are not a refresh but the filling of the density columns the routine keeps beside the composition, and they repeat the arithmetic of the pass's last internal call exactly.

Tests: `carrier_retry` 122/0 at entry, 124 PASS / 3 FAIL on a RED build carrying the delivered module with the one call `equilibrate_chemistry_at_fixed_pressure` line removed, 127/0 after. The new section (11) has five rows and states why the joint fixed point cannot be asserted on the synthetic 12-cell column at all: the shortest admissible trial still moves 6.46e-4 of the largest entry H2 mixing ratio, six decades above `relax_tol = 1e-10`. What it asserts instead is the consistency the joint fixed point rests on: the background third-body density read from the returned composition (GREEN 4.7244E-07 against the entry composition's 5.5008E-03; RED 5.4938E-03 against 1.2191E-16), the background temperature (GREEN 2.5675E-07 against 5.4837E-03; RED 5.5241E-03 against 0), and one further sweep moving the composition by 1.3637E-10 against `hybrdl`'s own `xtol = sqrt(eps) = 1.5E-08` (RED 2.0110E-04). Every other suite whose driver links `diffusive_photochemistry` is unchanged before and after: `carrier_returned_state_acceptance` 36/0, `attempted_step` 70/0, `steady_species_rows` 195/0, `certification` 70/0, `carrier_constraint_attribution` 10/0, `carrier_reference_scales` 14/0, `element_operator` 24/0, `species_masses` 9/0.

- **P22: the record of P21, of the terminology sweep and of the external commit** (this section with its two new subsections, the status rows of docs/PLAN_20260911_partitioned_solver.md, docs/session_handoff_20260911.md, the rewritten H2 front entry and the new outside-scope entry of TO_BE_DONE.md, and one sentence in docs/code_status_20260910.md section 3).
- **D1 to D3: docs/EXHALE_physics_and_algorithms.tex, the physical equations and the algorithms as the code applies them** (user instruction 2026-09-11, evening: a document at the depth of Caldiroli et al. 2021 sections 2 and 3, written from the source, not from the memos). Main file with natbib; eleven section files in docs/physics_overview/ (introduction and notation, hydrodynamics, radiation, ionization, thermal source, molecules and metals, numerical scheme, boundary and initial conditions, stationary state, output and post-processing, the table of physics keys with the reference list, 69 entries). D1 wrote the hydrodynamic, numerical, boundary and stationary sections, D2 the radiation, ionization, thermal, molecular and post-processing ones, D3 made them one document (55 inline citations converted, one symbol for one quantity with a notation table, about 20 cross references, repetition and run narrative removed). MEASURED: latexmk rc 0, 57 pages, no LaTeX or hyperref warning, every \texttt{} file name present in src/ (74 checked). Code facts established while writing, each stated in the document where it belongs: no escape-probability suppression acts on the Ly-alpha cooling channel ($\lambda_{\text{coex_HI}}$ in `Cool_coeff.f90` is the optically thin fit; the escape probability enters the metal fine-structure statistical equilibrium and the $\text{H}(n=2)$ populations only); `equation_T` (`T_equation.f90`) is called by `post_process_adv.f90` and test drivers, the marching loop's temperature being `solve_energy_semi_implicit`; `Numerical flux`: is a mandatory key with no default and every shipped input (70 of 70) says HLLC, ROE being refused with molecular chemistry unless `Caloric EOS: monatomic`; 2D approximate method: `alpha` applies neither the dayside dilution nor the mass-loss factor, which match the `Rate/...` strings only. Pointers added in README.md and the project CLAUDE.md.
- **The measurement of record after P14 to P17** (advisor, private snapshot build of the tree with P14 to P17, 8 threads, MEASURED): the HD 209458 b element reload, partitioned, 12 passes, ACCEPTED -- every active equation of this state is within its own tolerance at pass 12 (worst gated species row $7.73\text{e-}6$, He/H partition at cell 289; mass $9.26\text{e-}10$, momentum $2.44\text{e-}14$, energy $2.17\text{e-}8$); the hot-Uranus carrier reload, 3 passes, hydro info = 0 at every pass (mass $7.33\text{e-}12$ / $1.29\text{e-}11$ / $6.44\text{e-}12$ admitted by the cell's anchor), the one refusing entry the H2 carrier row ($7.06\text{e-}2$ at cell 288).

Three changes outside the item list, made by the advisor while reading P14 to P17:

- `assert_written_state_is_the_accepted_one` (`src/EXHALE_main.f90`) carries an ABSOLUTE floor of $1\text{e-}12$ beside its $1\text{e-}10$ relative test. P14 reported the false alarm: the spread is a difference of $\mathcal{O}(1)$ numbers over their mean, so two evaluations of one state agree only to the rounding of that difference, and on the partitioned carrier reload a state flat to fourteen digits read **3.0982E-14 as written against 2.7487E-14 accepted** while the relative test alone called that a change of u. $1\text{e-}12$ stands a decade above that rounding and four decades below the smallest spread a gate has ever accepted (READ, the comment at the test).
- the three rows of the `grid_and_gates` suite that wrote to a fixed directory under `$ROOT/build/tests/grid_and_gates` and ignored `EXHALE_TEST_OUT` now honor it

(momentum_row_from_fluxes_only.sh, base_level_single_statement.sh, sed_coverage_stop.sh; READ, each now takes `${EXHALE_TEST_OUT:-$ROOT/build/tests/grid_and_gates}`). Two concurrent runs of the suite, which is what a before-and-after measurement is, collided there. P14 fixed `output_state_consistency.sh`, the fourth, because its item owned it.

- the paragraph at `assemble_residual` (`src/modules/time_step/steady_residual.f90`) that stated what the kind-generic rows do not hand back is rewritten to what P15 made true: the kind-generic rows return the momentum row and store the interface fluxes, the face pressures and, under the well-balanced option, the face departures they were built from, but neither the three terms of the momentum equation nor the equilibrium pressure force the row is read against, so the one expression of each is evaluated there for the state just assembled; and a reconstruction continuation never reaches them, the generic text refusing it.

Two changes outside the item list, made by the advisor while reading P21:

- the H2 front: diagnostic of `src/EXHALE_main.f90` printed `r(max(j50,1))` and `r(max(j01,1))`, so a front that had left the domain was reported as the radius of the first cell. On P21's M1 run that read `x2=1e-2` at `r=1.0002` for a column in which no cell is below `1e-2` at all, beside a correct `x2=0.5` at `r=2.2367`. Each level is now printed as `> r(N)` when no cell of the column is below it.
- the noun of the 2026-09-11 terminology rule removed from the comment at `src/EXHALE_main.f90` line 2022 and from `diffusive_photochemistry.f90` line 248, and from the two fixture READMEs (about 26 lines), which item A4 left to whoever was editing those files.

The pseudo-time start of the stationary restart route

Found while running P6 and settled by the advisor. Restart intent: stationary entered the stationary solve at the CFL interval of the reloaded state. A state written by a run is already relaxed, and a continuation started at the CFL step from it does not reach the root: MEASURED, 8 threads, the hot-Uranus carrier reload's hydrodynamic solve is handed back at `||R|| 2.685E-01` after 40 iterations and the next four passes enter at 1.133, 1.018, 1.026 and 1.025 and hand back 2.976E-01 to 3.579E-01 (`<scratchpad>/P6/runs/carrier_p6/run.log`), while the HD 209458 b element reload stops on its stagnation detector at `||R|| 1.490` and the pass is then refused because the element relaxation finds no admissible advance (`<scratchpad>/P6/runs/atomic_partition.new/run.log`). With the start at 1.0, the value the marching hand-off uses, the same two solves reach `||R|| 2.507E-09` at iteration 13 and hand back **1.498E-09** (carrier) and hand back **1.167E-08** (atomic). The advisor therefore made 1.0 the default of the Restart intent: stationary route (`src/EXHALE_main.f90` line 5234), with `EXHALE_PTC_DTAU0` as the override for measurement; the direct steady route keeps its CFL start, which is the pseudo-transient continuation it was designed as. Verified with no environment variable set: the run prints stationary restart: `dtau0 = 1.00E+00` and reproduces the first pass to every printed digit (`<scratchpad>/P7/carrier_default_dtau0/run.log`). No regression case sets Restart intent: at all (MEASURED: the grep over `backup/regression/*/input.inp` returns nothing), so this default moves no golden; it moves any user run that sets the key.

P7: the measurement of record

One binary (EXHALE_P6.x, built from the merged tree), both fixtures, Restart intent: stationary, EXHALE_PTC_DTAU0=1.0, OMP_NUM_THREADS=8, 500 cells. The partitioned route is Coupled carrier solve: False with EXHALE_OUTER_PASSES setting the pass budget and EXHALE_JFNK_MAXIT the hydrodynamic cap; the coupled route is the same fixture with Coupled carrier solve: True. Wall times of the partitioned runs are the sum of the seconds the outer pass lines report, which agrees with the run directory's file timestamps to under a second; the coupled carrier control prints no total, so its time is the timestamp span of its run directory. Every number MEASURED from the log named.

route	fixture	passes (hydro cap)	wall time	hydro mass / momentum / energy	worst gated species row	certified?
state as loaded	carrier, hot Uranus	--	--	2.284e-1 / 1.954 / 1.871	H2 1.166e-1 at cell 492	no, 4 entries
partitioned	carrier	8 (40)	173 s	8.75e-12 / 1.66e-12 / 2.41e-9	H2 6.217e-2 at cell 282	no, 2 entries
partitioned	carrier	40 (40)	831 s	8.121e-12 / 3.377e-12 / 2.730e-9	H2 2.234e-2 at cell 443	no, 2 entries
coupled	carrier	one solve (250), stopped by the stagnation detector at iteration 105	321 s	1.231e-2 / 1.009e-1 / 5.801e-2	H2 7.815e-2 at cell 283	no, 4 entries
state as loaded	atomic, HD 209458 b	--	--	1.509 / 3.348e-1 / 1.892	He/H 5.840e-1 at cell 291	no, 11 entries
partitioned	atomic	5 (80)	528 s	1.05e-9 / 4.40e-13 / 1.13e-8	O 4.64e-4 at cell 268	no, 9 entries
partitioned	atomic	12 (80)	1043 s	1.054e-9 / 1.665e-13 / 1.501e-8	He/H 5.468e-6 at cell 288	no, 1 entry
coupled	atomic	3 (250), ended in the fourth	2392 s	1.16e-6 / 2.89e-6 / 2.83e-6	He/H 8.29e-5 at cell 295	no, run not completed

Sources: <scratchpad>/P7/carrier_elem_newton_part1/run.log, carrier_part40/run.log,

carrier_coupled250/run.log, atomic_elem_newton_part1/run.log, atomic_part12/run.log, atomic_coupled250/run.log. The atomic coupled control alternates the element relaxation too, because He_diffusion is on, and its three completed passes cost 1380.59 s, 482.04 s and 528.98 s; it was ended in the fourth. The (EXHALE_main) outer pass N: line of every pass carries the hydrodynamic flag, the worst gated species wind row with its cell, the three row maxima, omega, the carrier trust and the seconds, and the closing (certification) block names what refuses. Those <scratchpad>/ paths are the session scratch root and are TEMPORARY: they will be gone, and everything quoted from them is recorded here, in the two fixture READMEs and in docs/session_handoff_20260911.md. The runs reproduce from the pinned fixtures by the recipes in those READMEs.

What the ladders show. On the atomic fixture the worst gated species row falls $2.61\text{e-}3$, $3.81\text{e-}3$, $1.77\text{e-}3$, $9.45\text{e-}4$, $4.64\text{e-}4$, $2.36\text{e-}4$, $1.20\text{e-}4$, $5.97\text{e-}5$, $3.07\text{e-}5$, $1.50\text{e-}5$, $7.94\text{e-}6$, $5.47\text{e-}6$ over the twelve passes (the one rise, pass 1 to pass 2, is the alternation's own feedback across a hydrodynamic solve and is why a single rise does not end the loop), the refusing entries fall 9 to 1, and in the last pass every elemental wind row is within $1\text{e-}5$: He/H $5.468\text{e-}6$ at cell 288, and the metals $1.582\text{e-}6$ (Mg) to $3.720\text{e-}6$ (O). On the carrier fixture the H2 wind row falls $7.38\text{e-}2$ to $2.23\text{e-}2$ over the 40 passes with the binding cell moving 290 to 443, which is the H2 front advancing about one cell in a pass and not a residual settling.

The decision. The plan's adoption rule is that the route is adopted only if it certifies where the coupled solve does not, or reaches a strictly lower joint residual in no more wall time. Neither fixture certifies, so the first branch is not met on either. The second branch is met on both: on the atomic fixture the partitioned route stands strictly below the coupled control on every judged row (mass $1.054\text{e-}9$ against $1.16\text{e-}6$, momentum $1.665\text{e-}13$ against $2.89\text{e-}6$, energy $1.501\text{e-}8$ against $2.83\text{e-}6$, worst gated species row $5.468\text{e-}6$ against $8.29\text{e-}5$) in 1043 s against 2392 s for three passes; on the carrier fixture the partitioned route at 8 passes stands strictly below the coupled control on every judged row ($8.75\text{e-}12$ against $1.231\text{e-}2$, $1.66\text{e-}12$ against $1.009\text{e-}1$, $2.41\text{e-}9$ against $5.801\text{e-}2$, H2 $6.217\text{e-}2$ against $7.815\text{e-}2$) in 173 s against 321 s, and the 40-pass continuation goes on to $2.23\text{e-}2$ for 831 s. The partitioned route is therefore what the production outer loop runs. Coupled carrier solve already defaults to False (carrier_in_newton = .false., src/modules/init/parameters.f90 line 419), so no default changes; True remains the coupled measurement option, and what this session added on the partitioned side is the joint acceptance contract of P6 and the three defect repairs P1 to P3 that made a pass mean something.

What did not certify, and why. Entry 1 was CLOSED the same afternoon by the user's first decision (P16, anchor (6)); entries 2 and 3 are stated as they stand:

1. **CLOSED 2026-09-11 by P16.** The atomic element reload refused on the hydrodynamic mass row alone, $1.054\text{E-}09$ against $3.0\text{E-}12$ at cell 16 ($r = 1.0031$), on the three-unknown solve of this fixture. Every one of its twelve hydrodynamic solves ends with $||R||$ between $8.980\text{E-}09$ and $1.501\text{E-}08$ against the run's Resid tol of $1\text{e-}8$, and the last one ends on no descent direction exists for the banded model at this state at $||R||$ $1.501\text{E-}08$. So the alternation has done its work (the element rows are inside their tolerance) and what is left is the base-layer mass row of the hydrodynamic solve, which is the N33 rounding floor territory and not a species-coupling question. **P10 settled which of the two it is.** With the best-iterate ledger reading that row against its own $3\text{e-}12$ (P10), the same twelve passes end at $8.893\text{E-}10$ at cell 3, the same order, and the row is

MEASURED to carry no signal at that level: the ledger's best distance of pass 1 is $4.407\text{E}+02$, a mass row of $1.32\text{E}-09$, while the same state re-evaluated at its own composition after the best-iterate restore carries $4.11\text{E}-09$. At $1\text{e}-9$ to $1\text{e}-12$ the row is the rounding of the base-layer flux difference, so ranking iterates by it is ranking by rounding noise and no ledger can bring it down. The entry was therefore an ACCEPTANCE question, the $3\text{e}-12$ tolerance against the base-layer rounding floor of this fixture, and not a solver item; the user decided it the same afternoon, and with the mass row judged at $\max(3\text{e}-12, 10 * \text{floor}(j))$ the same twelve-pass recipe **CERTIFIES at pass 12** (MEASURED by P16 and again by P17, which certifies at pass 12 in both of its builds). The verdict is taken at a wind cell, not at the base: cell 34, $r = 1.00665$, $6.542\text{E}-10$ against the $4.6\text{E}-09$ anchor of that cell, distance 0.141 (P16), and cell 50, $r = 1.00977$, $4.245\text{E}-10$ against $3.2\text{E}-09$, distance 0.132 (P17). The base-layer floor of this fixture is what the anchor is measured on: see the decisions subsection below and docs/certification_tolerance_anchoring_20260910.md, "Anchor (6)".

2. The carrier reload refuses on the H2 wind row, and the user's second decision of the afternoon ran the continuation that says what that row IS. MEASURED over 110 passes (the table and the recipe are in the decisions subsection below): the H2 front does NOT stop, $x_2 = 0.5$ moving 1.0726 to 1.1230 R_p over 108 passes at about $5\text{E}-04$ R_p in a pass with no sign of slowing, so the continuation is unfinished after 110 passes and $0.05 R_p$. And the worst gated row stops being the front after pass 40: it stands at **$2.2\text{E}-02$, nearly uniform in r** , and its cell walks outward through the wind, 443 to 500, reaching the outer boundary at pass 108 while the value does not fall. That floor is a property of the operator and not of the front: the bounded carrier pass (trust $1\text{e}-2$ on the H2 mixing-ratio maximum, met by one transport step) advances the front but leaves the wind cells, whose H2 mixing ratio is orders of magnitude below the maximum, at one implicit step in a pass, so their own H2 balance never relaxes, and the drift measure, absolute in the grid maximum, does not see them. The consequence for the method, recorded and not adopted: a bounded front advance and a relaxation of the wind cells' H2 balance are two different operations, and the second needs its own step (a cell-relative measure, or a wind-window relaxation after the bounded pass) before the H2 wind row can be judged at $1\text{e}-5$. Whether the front stops at all inside $2 R_p$ is a physical question and stays open (the P23 comparison against Koskinen et al. 2022 Model A).
3. The hot-Uranus mass row sits on the same floor throughout the carrier run, $4.48\text{E}-12$ to $1.57\text{E}-11$ against $3.0\text{E}-12$ over the 40 passes, with the momentum row at $4.14\text{E}-13$ to $5.13\text{E}-12$ and the energy row at $1.75\text{E}-09$ to $4.56\text{E}-09$. The mass row of that fixture is the rounding of the base-layer flux difference recorded in N33, which is the reading P16 then made operative: under anchor (6) the base cells of this fixture are ADMITTED by their own anchor at every pass and the refusal moves out to cell 180 ($r = 1.0707$), where the floor is below $3\text{e}-12$ and the row carries signal.

The Codex adversarial review (2026-09-11)

Run on the working tree after P1 to P13, that is on the merged text of the morning's items and not on the committed tree. Two findings, both verified in the source by the advisor before any item was written, and both closed the same afternoon. The findings are quoted as they were written.

1. **[high]** "정체 종료 시 조성과 출력용 상태가 서로 달라집니다 (src/EXHALE_main.f90:6503-6514)": on a stag-

nation exit the composition had been updated while the particle densities and the temperature were not refreshed on the element path, so the certified state and the written state differed; the recommendation was to move the progress control before the composition update. Verified in the source, reproduced as a measured inconsistency of $1.317089\text{E-}08$ in the temperature of the written state, and **closed by P14** with exactly that ordering: the progress control now stands between the certification of the pass and the update, so a pass that ends the iteration takes no update, and the element path's update refreshes the particle densities and the temperature before its equilibrium sweep.

2. **[medium]** "고정밀 경로가 다른 상태의 압력 정보를 인증에 사용합니다 (src/modules/time_step/steady_residual.f90:256-258)": with EXHALE_RESID_QUAD the generic rows stored only face_flux and face_p, so momentum_row_terms_of_state read the previous RK_rhs call's face_q_up/face_q_dn (zeros on the first call); the recommendation was to return the terms from the generic computation and to refuse the continuation combination. Verified in the source, and **closed by P15**, both halves: the kind-generic rows now hand their own face departures back along the route the fluxes take, and the continuation combination is refused, because the kind-generic rows CAN be reached on a continuation (assemble_residual chooses the text from generic_precision_rows_arm() alone and the generic text carries its own blending branch) and no single pair of departures satisfies the blended row identity.

Neither finding moves a certified state of the default path: P14's wasp_full_newton reload and P15's, with the key off and with Well balanced: True and EXHALE_RESID_QUAD=2, are CERTIFIED with every output byte-identical apart from the provenance timestamp. What each moves is stated in its own bullet above.

User decisions of 2026-09-11 (afternoon)

Two decisions were taken after the open items of the plan were put to the user.

Decision 1: the mass tolerance is re-anchored above the fixture's measured rounding floor (item P16, and the ledger consequence P17). The tolerance of the continuity row is now

$$\text{tol}(j) = \max(\text{cert_tol_mass}, \min(\text{cert_tol_mass_ceiling}, \text{c_round} * \text{floor}(j)))$$

with cert_tol_mass = $3\text{e-}12$ unchanged as the FLOOR, the ceiling 1, and c_round = cert_mass_round_margin = 10. The floor estimate is $\epsilon \rho (|v| + c_s) A / dV$ over the row's own scale, which is one ulp in the wind and about ϵ/Mach in a layer whose flux is a tiny residue of the momentum the cell carries at its own fastest signal speed. The brief's first expression for the floor, $\epsilon (|A+ F_{\text{up}}| + |A- F_{\text{dn}}|)/dV / \text{residual_row_scale}(1, j)$, is inert, and that is a measured result and not a judgement: the row's scale IS the larger of the two addends and the dV cancels, so the expression is algebraically between one and two ulps at every cell, MEASURED at **$4.4409\text{E-}16$ on all 500 cells of all four states** examined, and ten times it never rises above $3\text{e-}12$. The adopted estimate bounds the step each mass row actually takes when one ulp is added to the density of every physical cell (EXHALE_MASS_FLOOR_SCAN=1, new and default off) within a factor

state	base Mach	step over the adopted estimate	step over the brief's expression
wasp_full_newton as loaded	3.937E-03	1.5931	1.109E+02
atomic_elem_newton as loaded	5.062E-06	1.8857	2.4365E+05
carrier_elem_newton as loaded	2.400E-04	2.6806	5.0915E+03
the HD 209458 b state P16 certifies	2.767E-06	1.7733	1.4201E+06

All MEASURED by P16, largest ratio over the 500 cells of each state. $c_{\text{round}} = 10$ is a round number 3.7 times above the largest of those ratios and inside N33's independent reading of the same floor against its own bound, 5.5 to 12 (READ). The outcome: the **HD 209458 b element reload CERTIFIED at pass 12**, where the control built from the entry text of the two files ends the same twelve passes NOT CERTIFIED on the hydrodynamic mass row (7.159E-10 against 3.0E-12 at cell 13), and **wasp_full_newton unmoved** in both builds (CERTIFIED, mass row 1.212E-12 at cell 500 judged against the FIXED 3.0E-12 at distance 0.404, $\log_{10} \text{Mdot}$ 13.30, outputs byte-identical apart from the provenance timestamp), its floor being below 3E-13 at every cell. The anchor still refuses states: the HD 209458 b mass row is refused at pass 3 at distance 69.9, and on the hot-Uranus carrier reload the refusal moves from the base cells, admitted by their own anchor, out to cell 180 where the row carries signal. The derivation and all three impact measurements are the appended section "Anchor (6), 2026-09-11: the mass row's rounding floor" of docs/certification_tolerance_anchoring_20260910.md.

Decision 2: the long continuation of the hot-Uranus carrier reload. The pass budget was set to 400 and the run went to pass **110**, where the progress control ended it ("the worst gated species row has not fallen in 3 consecutive passes"), the carrier trust having been halved 1.0E-02 to 5.0E-03 to 2.5E-03 at passes 109 and 110. Recipe: backup/regression/carrier_elem_newton, Load IC? True, Coupled carrier solve: False, Restart intent: stationary, EXHALE_OUTER_PASSES=400 EXHALE_JFNK_MAXIT=40 OMP_NUM_THREADS=8. Binary: the advisor's snapshot build of the merged tree after P1 to P12 and BEFORE P14 to P17, so every hydrodynamic solve of this run returns $\text{info} = 1$ on the base-layer mass row (5E-12 to 1E-11 against 3E-12) and the run predates anchor (6). All numbers MEASURED from its run.log.

pass	worst gated H2 row	cell	H2 front $x_2 = 0.5$ [R_p]	$x_2 = 1e-2$ [R_p]
1	7.38E-02	290	1.0726	1.0966
10	5.86E-02	281	1.0755	1.1017
20	4.07E-02	274	1.0796	1.1070
30	2.62E-02	267	1.0827	1.1140
40	2.23E-02	443	1.0871	1.1199
50	2.20E-02	473	1.0917	1.1262

pass	worst gated H2 row	cell	H2 front $x_2 = 0.5$ [R_p]	$x_2 = 1e-2$ [R_p]
70	2.19E-02	482	1.1017	1.1413
90	2.18E-02	490	1.1126	1.1563
100	2.17E-02	494	1.1184	1.1643
108	2.17E-02	500	1.1230	1.1727
110	2.17E-02	500	1.1230	1.1727

Two regimes. Passes 1 to 30: the worst gated row is at the front's outer edge, its cell moving 290 to 267 (r about 1.28 to 1.25), and it falls $7.4E-02$ to $2.6E-02$ as the front relaxes. From pass 40 the worst row is no longer at the front: it sits at $2.2E-02$ and its cell walks outward through the wind, 443 to 500, reaching the outer boundary at pass 108 while the value does not fall. The front itself keeps moving outward at about $5E-04$ R_p in a pass, $x_2 = 0.5$ from 1.0726 to 1.1230 over 108 passes, with no sign of stopping.

So three statements, the first two measured and the third a reading of them. (a) The front has NOT reached a stationary position in 110 passes: 0.05 R_p traveled and the continuation unfinished. (b) The remaining wind residual of $2.2E-02$ is nearly uniform in r and is NOT the front. (c) The reading: the bounded carrier pass, trust $1E-02$ on the H2 mixing-ratio maximum and met by one transport step, advances the front but leaves the wind cells, whose H2 mixing ratio is orders of magnitude below the maximum, at one implicit step in a pass, so their own H2 balance never relaxes, and the drift measure, absolute in the grid maximum, does not see them. The progress control ended the run because after pass 40 the worst gated row measures that wind floor and not the front.

Consequence for the method, recorded and NOT adopted: a bounded front advance and a relaxation of the wind cells' H2 balance are two different operations, and the second needs its own step, a cell-relative measure or a wind-window relaxation after the bounded pass, before the H2 wind row can be judged at $1e-5$. The physical question whether the front stops at all inside 2 R_p is separate and stays open (the P23 comparison against Koskinen et al. 2022 Model A).

P21: the fixed point at a fixed wind, and the production loop

All of this is MEASURED by P21 on backup/regression/carrier_elem_newton (Coupled carrier solve: False, Restart intent: stationary, EXHALE_JFNK_MAXIT=40, 8 threads, scratch copies), against a control binary built from the entry text of the three files.

M1, the fixed point at fixed hydro. EXHALE_CARRIER_TRUST=1e6, that is the movement bound disabled, and EXHALE_OUTER_PASSES=2 rather than 1, because the outer loop takes no composition update on its last pass (P6) and at 1 the relaxation never runs. Both binaries end the pass on carrier_relax_fixed_point.

quantity	control (frozen chemistry)	new (refreshed)
transport steps kept	38	57
composition drift	9.79E-01	9.95E-01
carrier steady residual after the pass, volume-weighted	2.32E-03	1.18E-11
worst cell	7.99E-02 at cell 266, $r = 1.211$	1.42E-09 at cell 2, $r = 1.000$

quantity	control (frozen chemistry)	new (refreshed)
H2 front, $x_2 = 0.5$	1.1327 R_p	2.2367 R_p
$x_2 = 1e-2$	1.2056 R_p	nowhere in the domain
largest x_2 on the grid	0.9774	0.9848
cells clamped onto the element budget	none	none
pass 1 wall time	18.64 s	16.04 s
pass 2 wall time (hydrodynamics only)	16.66 s	37.06 s

The wind H2 row before the pass is 7.38E-02 on both, at cell 290, reproducing the experiment's 7.380744e-2, and the control's 7.99E-02 reproduces its 7.989781e-2 to the digit. The experiment's three-number sequence has only two members in the production loop, the diagnostic being taken after the caller's refresh in the control, so the frozen intermediate (2.449021e-13) is not a state the production loop ever holds; in the new build there is no intermediate at all, which is the point. Wall time of the new pass is LOWER despite 50 percent more steps and one sweep a step; the plausible reading, offered as a reading, is that a transport step taken on the background of the composition it advances converges in fewer inner iterations than one taken far from a frozen background.

M2, the production loop: at the production bound nothing moves. Same fixture and recipe with the movement bound at its default 1e-2, EXHALE_OUTER_PASSES=12 and a 60-pass continuation, against the control's first three passes and against the 110-pass ladder of the same day's decision 2 (both READ from the P7 run logs).

pass	worst gated H2 row, cell	recorded ladder	$x_2=0.5 / x_2=1e-2$	recorded
1	7.38E-02, 290	7.38E-02, 290	1.0726 / 1.0966	1.0726 / 1.0966
10	5.86E-02, 281	5.86E-02, 281	1.0755 / 1.1017	1.0755 / 1.1017
20	4.07E-02, 274	4.07E-02, 274	1.0796 / 1.1070	1.0796 / 1.1070
30	2.62E-02, 267	2.62E-02, 267	1.0827 / 1.1140	1.0827 / 1.1140
40	2.23E-02, 443	2.23E-02, 443	1.0871 / 1.1199	1.0871 / 1.1199
50	2.20E-02, 473	2.20E-02, 473	1.0917 / 1.1262	1.0917 / 1.1262
60	2.20E-02, 477	(not in the ladder)	1.0953 / 1.1327	

Twelve and sixty passes reproduce the recorded ladder to every printed digit: the worst gated H2 row, its cell and both front positions are identical at passes 1, 10, 20, 30, 40 and 50. Every pass ends on the movement bound with 3 to 6 kept steps and a drift of 9.98E-03 to 1.00E-02, and every hydrodynamic solve returns $info = 0$. So **the 2.2E-02 wind floor does not go away and the frozen chemistry was not its cause**; the bound's measure, absolute in the grid maximum, remains the standing explanation. Cost is not a reason to hesitate: a production pass costs 3.07 to 3.72 s against the control's 3.10 to 3.54 s, and one equilibrium sweep on this 500-cell grid costs below 0.029 s at 8 threads (MEASURED from 300 marching steps of `mol_carrier` in 8.66 s including initialization), so 3 to 6 of them cannot be resolved against that scatter.

M3 and M4, the default paths. `mol_carrier` marching, EXHALE_MAXSTEPS=300, one thread: `Hydro_ioniz.txt`, `Ion_species.txt` and both `_adv` files identical except the two `git=` provenance lines, and `Cooling_breakdown.txt`,

Heating_breakdown.txt and IC_dump.txt byte-identical, which is the expected result because the marching loop calls photochemical_transport_step directly and never this routine. wasp_full_newton reloaded, one thread: the two run logs are byte-identical, both info = 0 and CERTIFIED at $|R| = 1.920E-09$, ten of the twelve written files byte-identical and the other two differing in the same provenance line alone. No molecules, so the branch is never entered.

Does the front stop at fixed hydro, and where? No, and the way it fails to stop is new information. At a wind held fixed and with the chemistry solved on the advancing composition, the pass runs to a genuine fixed point of the coupled operator ($1.42E-09$) whose H2 fills the whole domain: $x_2 = 2n(H_2)/n_H$ is 0.98 at 1.1 R_p, 0.96 at 1.2, 0.84 at 1.5, 0.60 at 2.0 and 0.33 at 3.0 R_p, with no cell below $1e-2$ anywhere. The largest x_2 is 0.9848, so this is NOT the element ceiling and no cell is clamped in either run, and P2's frozen artifact (x_2 driven to the element ceiling in 123 cells) does not appear. A plausible reading, offered as a reading: the H2 column and its self-shielding grow together, so re-solving the photodissociating field on the advancing composition removes the sink that held the front, and nothing but the wind can stop it, which it cannot while it is held fixed. What this says is that **the front's position is set by the hydrodynamic response and not by the carrier operator alone**, consistent with the bounded advance being the only usable form of the update; the physical question, whether the front stops inside 2 R_p in the coupled system, stays exactly where P23 has it.

Does the wind floor go away? No, as M2 records. The $2.2E-02$ row that appears at pass 40 and walks outward through the wind is reproduced pass for pass and cell for cell with the chemistry refreshed. The frozen background was not its cause; the bound's measure, absolute in the grid maximum, still is.

The terminology sweep (A1 to A4, 2026-09-11)

The user asked whether "arm" is a word physics uses for what this tree was using it for, and instructed that terms belonging to software engineering, clinical trials or management not be used as the names of physical or numerical concepts at all. The rule is now in `~/claude/CLAUDE.md` under "금지 표현", in two bullets. The first: avoid terms that physics, chemistry, mathematics and astronomy do not use, in documents, comments, log messages and identifiers alike, and write instead what the thing is, the standard term of those fields or plain English, naming what is compared ("control build", "measured build"), which physics option is meant ("the well-balanced option"), or which quantity ("the mass row", "the H2 front"); where such a term already stands, fix it where it is visible. The second: the NOUN "arm" is forbidden, for one side of a comparison, for a selectable code path and for one branch of a fixture alike, and code identifiers (ARM_*, *_arm) are held to the same rule; the VERB "arm / armed / arming / disarm", meaning to enable a device, is plain English and stays.

Four workers swept the tree. Every count is MEASURED by the worker that owns the files, by `grep -w -i -E "arms?"` against the git HEAD text, and no number, file name or measured statement was changed anywhere in the four diffs.

- **A1, the 51 top-level docs/*.md outside the two update logs: 396 to 17.** All 17 that remain are allowed: 14 are the verb, 3 are the literal directory glob arm*. The replacements are by sense: a side of a comparison became "the two runs" or "both builds"; an optional code path became its key or its physics ("the well-balanced option", "the

kind-generic residual assembly"); a fixture's solve became "the element reload" or "the carrier reload", which is the word those same documents already use; a rung of a ladder of runs became "rung"; one chemical network of a comparison became "network". One code span changed, the placeholder <arm> of a run-directory path in `vulcan_photochem_comparison.md`, which named nothing on disk and is now <network>, the names that do exist there being `vulcan/`, `ncho/`, `zahnle/`, `zahnle_s/` and `eqfit/` (READ from the tree).

- **A2, docs/Update_EXHALE_stage2.md: 176 to 1** (the one is the verb, "when both stellar keys arm H(n=2)"), and `Update_EXHALE_appendix.tex` 1 to 0. The largest class was "in both arms" of a two-binary comparison, now "in both builds"; then the optional paths ("the well-balanced option", "quad accumulation", "the kind-generic rows"), the fixtures ("the atomic element reload", "the carrier rows"), and the unknown space of B5j/B5k ("the density unknown", "the log unknown"). `Update_EXHALE_stage2.tex` was regenerated and the PDF rebuilt at **158 pages**, the same count as before the sweep (MEASURED, `pdfinfo`). Thirteen British spellings were corrected in the same pass, twelve in the Markdown and one in the appendix (modelling, centre/centred, labelled, cancelled, behaviour, travelled, NEIGHBOUR), as the American-spelling rule for documents asks.
- **A3, the hand-maintained LaTeX and the stage-1 log:** `Update_EXHALE_stage1.md` 211 to 4 and `Update_EXHALE_stage1.tex` 196 to 7 (verb uses and four `heh_1_x2matched` (renamed 2026-09-16 from `arm_heh1_x2matched`) case-name literals), and nine other `docs/*.tex` files, of which `lit_wasp121.tex` was deliberately not edited: its "Fe I (blue arm), VLT/UVES" is the spectrograph's own blue arm and its "spiral arms" are the outflow's, both standard astronomy and not the forbidden sense. Nine PDFs were rebuilt with every page count unchanged (MEASURED, `pdfinfo`: `Update_EXHALE_stage1` 460, `EXHALE_user_manual` 90, `lhs1140b_exhale_vs_pwinds` 76, and six smaller memos).
- **A4, the source, the tests, the scripts and the top-level documents: 35 files, 328 to 12**, the 12 all the verb plus two references to the `arm*` case directories by name. Identifiers renamed: `generic_precision_rows_arm` to `generic_precision_rows_selected`; `ARM_OFF`, `ARM_QUADRUPLE`, `ARM_GENERIC_DOUBLE` to `ROWS_PRODUCTION`, `ROWS_QUADRUPLE`, `ROWS_GENERIC_DOUBLE`; `arm_carrier_log_unknown` to `form_carrier_unknown_space` (named for what it forms, since it runs in both coordinates and always forms `carrier_log_floor`); `arm_departure_of_the_rows` to `key_departure_of_the_rows`; the local `arm_name` to `kind_name`; the local `e_arm` to `e_wb`; and eleven printed test-row names. Environment variables and input keys were NOT touched (`EXHALE_RESID_QUAD`, `Well balanced:`, `EXHALE_ELEMENT_CONSTRAINT_ROWS`). Every old row name and changed message was grepped over the whole working copy and matched only the file that defines it, so nothing parses them. MEASURED: the eight suites that link the changed modules keep their counts exactly (`krylov_and_dogleg` 334/0, `certification` 70/0, `grid_and_gates` 200/0, `steady_species_rows` 195/0, `residual_determinism` 5/1 by design, `steady_completion_flag` 3/0, `steady_selfconsistent_residual` 3/0, `element_operator` 24/0); and the `wasp_full_newton` reload is byte-identical between a build of the delivered text and a control build of the entry text, both plain (`info = 0`, `||R|| = 1.495E-09`, flux spread `1.911E-12`) and under `EXHALE_RESID_QUAD=2` with `Well balanced: True` (`info = 0`, `||R|| = 7.364E-10`, flux spread `1.591E-13`), so the renamed selector still selects. That the selection is not inert was measured beside it: `EXHALE_RESID_QUAD=1` gives `||R|| = 7.363E-10` at a flux spread of `1.588E-13`, different in the last digits from mode 2, while mode 2 against no selection at all is byte-identical, which is the design assertion for the

generic-double instantiation.

Three further occurrences were removed by the advisor at the close, outside the four items: the noun in the comment at `src/EXHALE_main.f90` line 2022 and in `diffusive_photochemistry.f90` line 248, and about 26 lines of the two fixture READMEs (`backup/regression/atomic_elem_newton/README.md`, `carrier_elem_newton/README.md`), which A4 left to the item then editing them.

What stays by design, everywhere: the verb; the `arm*` case directories under `backup/regression/` and the two glosses that now point at them as file names; and the astronomy senses in `lit_wasp121.tex`.

An external commit during the session

Commit **a7c18a6**, "**remove build_***" was made by the user at 21:39:24 KST on 2026-09-11 from another session, while item P21 was mid-edit (READ, `git log` and `git show --stat a7c18a6`: 149 files changed, 39,652 insertions, 201 deletions). It removed the `build_dift/` and `build_fcheck/` dependency stamps; it added the 141 files of `docs/audit_20260905/partition_experiment_20260911/runs/`, 38 MB of copied inputs, `.state` and `.faces` dumps and logs (MEASURED, `du -sh`); and it carried P21's in-progress edits of `src/EXHALE_main.f90`, `diffusive_photochemistry.f90` and `carrier_retry.f90` as they stood at that minute. Two consequences are recorded because they are visible in the measurements above. P21's line counts are `diff -u` against the entry texts saved in its scratch directory before the first edit, `git diff` no longer being able to produce that item's diff. And the provenance `git=` line of every output written by a build made after the commit reads `a7c18a620f1b` where the control binary's reads `0608382f639c`, which is the only difference in P21's M3 and M4 comparisons. The "What the repository carries" paragraph of `docs/audit_20260905/partition_experiment_20260911/README.md`, which said `runs/` was kept in the working copy only, now records that it is committed since `a7c18a6`.

Gates

Advisor verification of the merged tree (READ from the closing brief): snapshot build with the default toolchain; `element_operator` 24/0, `carrier_retry` 122/0, `krylov_and_dogleg` 290/0, `certification` 57/0, `steady_species_rows` 194/0 and 195/0 after P8; `grid_and_gates` 158 PASS with 3 FAIL, all three missing files of the snapshot copy (`inputdata`, `examples`) rather than assertions (P4 measured the same suite 178/0 in the live tree); the `wasp_full_newton` reload `info = 0, ||R|| 1.667e-9`, CERTIFIED, `log10 Mdot` 13.30, unchanged. make check (advisor, 2026-09-11 09:40 KST, the shared build rebuilt from this tree): REGRESSION PASS, every DEFAULT_CASES output and `wasp_full_newton` byte-identical to the goldens; no golden moved, because no default case enters the stationary solve or the fixed-wind relaxations.

After P10 to P12 (advisor, merged tree, private snapshot build, 8 threads, MEASURED 2026-09-11): the thirteen suites that link the changed modules all PASS (`krylov_and_dogleg` 309/0, `grid_and_gates` 187/1 with the one FAIL a WASP-52b SED file absent from the snapshot copy, `steady_species_rows` 195/0, `element_operator` 24/0, `carrier_retry` 122/0, `certification` 57/0, `attempted_step` 70/0, `carrier_returned_state_acceptance` 36/0, `species_masses` 9/0, `carrier_reference_scales` 14/0, `carrier_constraint_attribution` 10/0, `constrained_network_layout` 8/0, `steady_completion_flag` 3/0). The two fixtures re-run on that binary re-

produce P10's and P11's records: the HD 209458 b element reload, 12 passes, worst gated species row $2.61\text{e-}3 \rightarrow 5.47\text{e-}6$ with only the hydrodynamic mass row refusing ($8.893\text{e-}10$ against $3\text{e-}12$ at cell 3, $r = 1.0006$); the carrier reload, 3 passes, H2 $7.38\text{e-}2 \rightarrow 7.06\text{e-}2$ at cells 290 $\rightarrow$ 288, mass $8.12\text{e-}12 / 4.85\text{e-}12 / 4.96\text{e-}12$, momentum $3.22\text{e-}13 / 4.47\text{e-}13 / 3.54\text{e-}13$.

Second make check after P10 to P12 (advisor, 2026-09-11 15:15 KST, the shared build rebuilt from the merged tree): REGRESSION PASS, every DEFAULT_CASES output and the cold-start wasp_full_newton byte-identical to the goldens; no golden moved.

After P14 to P17, each count MEASURED by the item that owns the file and READ from its report: grid_and_gates **200/0** (P14, the second case of output_state_consistency.sh; 199/1 on the entry text), krylov_and_dogleg **321/0** after P15 and **334/0** after P17, certification **70/0** after P16 (57/0 on the entry driver), with steady_species_rows 195/0, adv_static_limit 53/0 and the other suites each item ran unchanged in both of its builds. The two rows that FAIL by design or by fixture are unchanged: residual_determinism's closure_spread_within_the_row_tolerance_atomic_elem_newton at $3.354\text{E}+01$ (N33, by design) and steady_selfconsistent_residual's handback_matches_the_accepted_iterate on a reload log that prints no iterate norm.

Third make check after P14 to P18 (advisor, 2026-09-11 21:10 KST, the shared build rebuilt from the final tree): REGRESSION PASS, every DEFAULT_CASES output and the cold-start wasp_full_newton byte-identical to the goldens; no golden moved.

Fourth make check after P21 and the terminology sweep (advisor, 2026-09-12 01:40 KST, the shared build rebuilt from the tree with P21, A1 to A4 and the closing edits): REGRESSION PASS, every DEFAULT_CASES output and the cold-start wasp_full_newton byte-identical to the goldens; no golden moved.

Reported by the workers and not fixed

- P1: relaxation_drift_covers_every_element in steady_species_rows builds a column whose species miss its own density by $1.660\text{E-}02$ (closed the same day by P8); relax_element_composition damps helium alone, which belongs to the outer composition control; element_diffusion_step's trace-metal branch re-seeds an exhausted cell and project_elements clamps grp_new at zero, the only place in the step that can create or destroy mixture mass, now covered by the closure test and not investigated further.
- P2: the carrier relaxation reaches into the steady residual for its advecting flux (carrier_rows_advect = .true.) while the element operator is given its flux by the caller, so the routine cannot be called at all unless someone else assembled a mass row; residual_row_scale in steady_residual.f90 carries a comment block that is the history of a change and not the physics; the fixed-point exit of the carrier relaxation is unreachable in production while the H2 front propagates, and loosening relax_tol to make it reachable is not a fix.
- P3: ~~distance_from_certification divides the hydrodynamic slot by the run's Resid_tol and not by each row's own certification tolerance, so the best_iterate ledger cannot see a mass row above $3\text{e-}12$ while $||R||$ is below $1\text{e-}8$, which is exactly the state D3 is about (it did no harm on the D3 run, where the returned iterate's mass row is $5.945\text{e-}12$ against the last iterate's $7.914\text{e-}12$, but the ranking does not guarantee it)~~ (fixed 2026-09-11, P10); steady_selfconsistent_residual assertion 2 (closed the same

- day by P8); `attempted_step/run.sh` and `carrier_returned_state_acceptance/run.sh` write a fallback `build_stamp.f90` into their own work directory, where the compile cannot find it, so a tree that has never been built fails with Cannot open module file 'build_stamp.mod' (fixed 2026-09-11, P12, in those two and in five sibling scripts).
- P4: the kind-generic rows do not form the equilibrium pressure term themselves (closed from `assemble_residual`, but the two places must stay one expression if the kind-generic rows are ever made to write their own face data back); the default path's PLM momentum scale carries a coordinate artifact, the geometric pressure term $(A^+ - A^-) p_j / dV$ entering the max separately although it is one physical term with the flux divergence, so at zero gravity with a uniform pressure the scale is $2 p / r$ where the physical force is zero (MEASURED: the well-balanced scale, which is the row's true content, sits 0.944 below the base scheme's for both Riemann solvers at every N; under WENO3 the two agree to $2E-13$). Fixing it moves the default momentum measure at every cell and therefore the acceptance of every recorded state, so it is a user decision. (fixed 2026-09-11, P11, on the user's instruction: the scale is now the largest of the three physical terms, the zero-gravity artifact reads 2.22507E-308 in place of 1.000, and the acceptance movement is measured, `wasp_full_newton` still CERTIFIED.)
 - P5: the same truncation-as-floor reasoning stood in three other documents (closed the same day by P8); the analysis document's own title is still stronger than its corrected conclusion; `docs/session_handoff_20260910.md` and `docs/steady_solver_design.md` carry statements of the same family, not read line by line; the experiment's private build is retained at `/tmp/exhale_partition_20260911.ZxXJEB` per its own section 9.
 - P8: `docs/certification_tolerance_anchoring_20260910.md` still opens with `cert_tol_element` and `cert_tol_carrier` at $1e-8$, which decision 22 (a) superseded, and whether that memo is a dated record or a live reference is a decision; `docs/code_status_20260910.md` section 3 described the stage-2 solver program as closed (this session's item P9 marks it superseded); `mass_closed_column` exists only in `src/tests/element_operator/element_operator_tests.f90` and two suites now build the same column by hand, so the recipe belongs in one place a test can call (fixed 2026-09-11, P12: `src/tests/test_columns.f90`).
 - P9: the Coupled carrier solve key comment in `src/modules/files_IO/input_read.f90` states that the alternation "was measured not to converge and not to be able to" (section 139). The first half is what P7 measured against: the alternation with the joint acceptance contract of P6 brings every elemental wind row of the atomic fixture inside $1e-5$ and moves the carrier front monotonically, so the comment's reading is at least narrower than it states. Not my file this increment; it needs the sentence rewritten to what section 139 actually measured (the old alternation, without the acceptance contract, and its sign error on the front's motion). (fixed 2026-09-11, advisor.)
 - P10: two rows of `src/tests/steady_species_rows/steady_species_rows_tests.f90` state the superseded semantics, feeding a raw `||R||` into the slot that now takes a distance and asserting that the hydrodynamic slot follows the run's `Resid tol` (MEASURED 193 PASS / 2 FAIL on the new objects) (fixed 2026-09-11, advisor); `dj` that the ledger ranks with is the accepted trial's own rows while the judged rows line prints the loop-top evaluation, so the reported best distance can sit below the minimum of the printed lines (`wasp_full_newton`: 8.469E-02 against a printed minimum of 1.178E+00), which is entry-text behavior and not a defect; in the

EXHALE_RESID_BRANCH_REPORT block one divisor for the whole class stays, because the block aggregates spreads across seeds whose maximum can sit in different rows.

- P11: ~~the kind-generic rows still hand back neither the momentum terms nor the face departures (hydrodynamic_rows.f90's store_the_interface_fluxes writes face_flux/face_p and nothing else), so under EXHALE_RESID_QUAD the well-balanced face departure and, on a reconstruction continuation with $0 < \lambda < 1$, which of the two pressure conventions the blended flux carries are read from the last RK_rhs evaluation instead of from the state just assembled; the shape of the fix is two weights in place of the use_plm branch, and the option is default off and a rounding control experiment nothing is adopted from.~~ (fixed 2026-09-11, P15, which was also the Codex review's second finding: the kind-generic rows store their own face departures, and the continuation combination is refused rather than weighted, because the two schemes reconstruct different faces and no single pair of departures satisfies the blended row identity. The momentum terms are still not handed back by the kind-generic rows, and they are not read from a previous evaluation either: the one expression of each is evaluated at the call site for the state just assembled, which is the P4 entry above.) The marching stop of mol_carrier reports a momentum measure of 1.979E+00 against a scale that is now the largest physical term in the layer, which is commensurable with the mass and energy entries in a way it was not; whether the marching stop should quote it at all is an acceptance question.
- P12: nothing further. The four sibling scripts carrying the same dead fallback and the work directories hardcoded with no override, which the P12 report listed as outside its own two files, are what P12b closed the same day.
- P14: ~~assert_written_state_is_the_accepted_one raises a false alarm whenever the flux spread reaches the rounding level (MEASURED on the partitioned carrier reload, 3.0982E-14 as written against 2.7487E-14 accepted on a state whose mass flux is flat to fourteen digits, while the same run at a spread of 1.7736E-03 passes the test); it needs an absolute floor beside the relative one, or the warning will keep accusing the best states the solver produces and three rows of the grid_and_gates suite write to a fixed directory under \$ROOT/build/tests/grid_and_gates and ignore EXHALE_TEST_OUT, so two concurrent runs of the suite collide there (both fixed 2026-09-11, advisor, the floor at 1e-12 and the three rows named in the paragraph above).~~ What is left is one line of documentation: src/tests/grid_and_gates/README.md gives one line for each test name, and the output_state row now asserts two things where the README says one.
- P15: quad_rows_calls and quad_rows_seconds are incremented in hydrodynamic_rows_in_quadruple_precision and public, and no file reads them (grep over src/), so the quadruple-precision rows' measured cost, which the module header calls a measured number and not an estimate, reaches no run summary; relatedly, nothing in run_log says whether EXHALE_RESID_QUAD took effect, so P15 had to establish that its quad run was a quad run indirectly, from the iteration-1 measure moving between the two binaries and from the sweep-kind reset READ in steady_newton.f90. A line in the setup report, or those two counters reaching it, would make a quad run self-documenting. And the continuation refusal the item added has no executed row: it ends in error stop, which cannot be asserted from inside the driver process, and no production path reaches it, the key being default off and the continuation controller disarming before the JFNK finish (READ).
- P16: the control build of the HD 209458 b reload ends with a state its own certification refuses and **exits 0**, because the outer loop's ending is the pass budget and no stationary claim is made. That is the run-mode contract

working as written, but a reader of the shell status cannot tell that case from a certified one.

- P17: ~~hydrodynamic_distance_from_certification has no production reader left and cannot be deleted without breaking the compile of a test file the item does not own (item P18, above).~~ Two fixture observations, no action taken: running `grid_and_gates` twice at once with distinct `EXHALE_TEST_OUT` makes the `flux_spread` sub-test report `mass_flux_spread_recomputed_from_output measured=no_dump_line`, so the two invocations share something outside `EXHALE_TEST_OUT` (each invocation on its own gives 200/0), which is worth a line in that suite's README for whoever next runs two builds side by side; and the log-driven block of `steady_species_rows` requires a `(species_jac_test)` line in the log, so the best judged `iterate <= min judged rows` seen invariant it states is checked only on logs made with the species Jacobian probe armed, and P17 had to apply that comparison by hand (0 of the 12 solves of its new build report a best above their own printed minimum, against 1 of 12 in the control). A fixture log carrying that probe would be worth pinning.
- P21: the archived experiment drivers under `docs/audit_20260905/partition_experiment_20260911/` (`transport_wind_experiment.f90` and the `driver_snapshot.f90` copies beside the runs) call `relax_photochemical_composition` with its old six-argument signature. They are not in the Makefile SRC list and no test `run.sh` builds them, so nothing compiles them; they are the record of the run they belong to and are left as they stand. The H2 front: diagnostic P21 reported beside it was fixed by the advisor (above).

9. PLAN_20260912: the seven findings of the stage-2 review (2026-09-12)

Plan of record: `docs/PLAN_20260912_review_fixes.md`. User instruction of 2026-09-12: analyze `docs/Update_EXHALE_stage2_review_20260912.md` against the code, and where its analysis is right, write the plan and start the corrections. All seven findings R1 to R7 were confirmed in the source before the plan was written (the confirming lines are listed at the head of the plan); the corrections are items Q1 to Q4, Q1 and Q4 by the advisor, Q2 and Q3 by workers under `docs/worker_rules.md`. Every number below is MEASURED on the tree of this section unless marked READ.

The state contract at a fixed hydrodynamic state (R2, R3)

The decision behind Q1: at a fixed hydrodynamic state the quantities the solve holds are the conserved variables $u = (\rho, \rho v, E)$. A composition update at fixed hydro keeps u and recomputes the pressure and the temperature from the unchanged thermal energy $\rho e = E - (\rho v)^2 / (2 \rho)$ through the caloric EOS of the new composition (`pressure_from_energy_density` after `get_species_densities`, then `comp_T_from_p`). A held pressure is not a state the solve knows: the review's 500-cell probe showed that keeping p while the main loop keeps E describes two molecular states $3.3e-3$ apart.

The items

- **Q1 (R1, R2, R3, R6, R7 and the outer progress measure), advisor.** `diffusive_photochemistry.f90` (+206/-204), the carrier branch and the summary of `steady_wind_with_element_diffusion` in `EXHALE_main.f90`, `src/tests/carrier_retry/carrier_retry.f90` (+195/-77) and its `run.sh`.

- R1: `photochemical_transport_step` takes an optional `trial` argument; a trial of the stationary relaxation returns before the history mark, the message and the stop, so a rejected trial (an uncovered interval, a chemistry that did not close, a displacement beyond the movement bound) leaves `carrier_history_certifiable_flag` as it found it. The marching path passes no trial and is unchanged.
- R2, R3: `relax_photochemical_composition(u, v, f_sp, p, T, heat, cool, eta, trust, drift, nstep, outcome)` now takes the conserved state; `p` and `T` are outputs. `pressure_and_temperature_at_fixed_conserved_state` is the one expression of the contract above, and `equilibrate_chemistry_at_fixed_conserved_state` cycles (`p, T` at fixed `u`; sweep; `p, T` again) until the largest relative change of `T` falls below $1e-6$ (at most five cycles), reads the sweep ledger (`n_nonfinite`) and a finite composition, and reports ok; a refresh that does not close refuses the trial (`carrier_relax_chemistry_refused`, a new named ending "the chemistry of the shortest admissible trial did not close") and the entry composition, background, `p, T`, heating, cooling and `eta` are restored. The caller in `EXHALE_main.f90` rebuilds `W` from the unchanged `u` and takes the returned composition; nothing is reconstructed from a held pressure any more.
- R6: the carrier retry suite's column is built by `column_carrying_its_own_density` of `src/tests/test_columns.f90` (the mass-closed constructor of the element suites; `run.sh` compiles it before the driver) with the ions moved out of the neutrals; a row asserts the column reconstructs its own density to $1e-13$ before any step. The consistency rows are absolute: the returned pressure is that of the unchanged thermal energy (0.0), the returned temperature is that of the returned composition (0.0), the background third-body density and temperature the sweep stored are those of the returned composition ($6.53e-7$ and $2.28e-7$ relative, allowance 10 ieq_res_tol), and one further sweep leaves the returned composition within the sweep tolerance ($8.83e-7$ against `ieq\_res\_tol`). Nine rows for R1: a rejected trial keeps the history certifiable, keeps no step, hands back the entry composition bit for bit and names the refused interval; a pass after the refusals keeps a step and stays eligible; a refused trial in phys mode reports exhausted, leaves the history certifiable and writes no composition. RED on the entry text (the trial marking the history, the refresh not closed): five rows fail (pressure $1.2e-7$, temperature $3.5e-7$, further sweep $1.2e-6$, the two R1 rows 0 of 1); GREEN on the tree: 138 PASS, 0 FAIL.
- R7 and the progress measure: the outer summary of `steady_wind_with_element_diffusion` ranks every entry by its binding distance (`dist_bind`, `row_at_bind`, `jbind` where the entry carries them), so the mass row's refusal is attributed to the cell that binds it and not to the largest row over another cell's tolerance; and the progress control halves `omega` and the carrier movement bound on the JOINT distance of the state (the largest entry over its own tolerance, `prog_worst`), not on the worst species row alone. An entry that is not finite or not available counts as an infinite distance.
- The review's 500-cell probe (`docs/audit_20260905/stage2_review_20260912/molecular_wind_state_probe.f90`, adapted to the new signature, hot-Uranus carrier reload, one hydro solve and one bounded relaxation at trust 0.01): returned temperature against the EOS of the returned state 0.0 (was $5.5997819e-9$), pressure of the unchanged thermal energy against the returned

pressure 0.0 (was 3.2993e-3), thermal-energy change a held pressure would need 6.9e-16 (was 3.3250e-3); mass closure 8.33e-15 and the five retained steps at drift 9.9955e-3 unchanged.

- Impact on the routes: the unbounded pass of P21 M1 repeated on the new contract (EXHALE_CARRIER_TRUST=1e6, cap 40) reaches the fixed wind's fixed point in 61 transport steps at drift 0.995 with the H2 front ($x_2 = 0.5$) at 3.079 R_p (P21: 57 steps, 2.237 R_p; both fully molecular fixed points, the front leaving the domain). The 12-pass production loop ends at H2 row 5.49e-2 at cell 279, front 1.0755 R_p (the 110-pass ladder of P16 at pass 12: 5.50e-2 at cell 279, 1.0765 R_p); no pass triggered the halving. atomic_elem_newton at three passes: pass lines identical to the control build. wasp_full_newton reload and the mol_carrier case (12000 steps, one thread): data rows identical to the control build and to the golden.
- **Q2 (R4 in the operator), worker (Sonnet).** binary_element_diffusion.f90 (+108/-43), src/tests/element_operator/element_operator_tests.f90 (+88/-1). element_diffusion_step judges its candidate on every call: the entry composition is kept, the four tests (solved, finite, X inside [0,1] to element_fraction_bound_tol, mass closure no worse than the entry's by element_mass_closure_tol) run whether or not status is present, and the entry composition is restored on a refusal. The outcome is always left in the new public element_step_last_status, and element_step_outcome_text names it. Two new rows repeat the ordinary and the NaN-residual steps with no status at the call site: RED on the entry text (the refused candidate with departure 0.86 adopted, the status left at accepted), GREEN after (element_operator 28/0). A second, independent defect found by the change: the Newton solve of the helium mass fraction (solve_mass_fraction) read its arithmetic floor only on a DESCENDING pass, so a step whose residual had already reached the floor and whose next pass found no improving trial reported unsolved (residuals 1e-12 to 1.1e-5 at the refusal, MEASURED on the closed-column relaxations of diffusion_tests T1a, T7c, T7d and T10, which went 35/0 to 27/8 under the unconditional judgment alone); the no-descent exit now reads the same floor, and newton_drop moved from 1e-6 to 2e-5 so that the ~1e-5 floor of the $K_{zz} = 2e12$ homopause column (T7), which the module header already documented, lies inside it (diffusion_tests 35/0 again). Cost on mol_diffusion at 300 steps: 54.4 s against 54.2 s, inside the noise. mol_diffusion and lower_profile at 300 steps: data rows identical to the control build. atomic_elem_newton: pass 1 identical; at pass 2 the judged rows are identical and omega differs (0.500 control, 0.250 Q2), attributable to the floor change turning a floor-level residual from unsolved to solved (READ from the Q2 report; the advisor's own three-pass run of the merged tree, above, has omega 0.250 at pass 2 on both builds because the control binary of that run already carried Q2).
- **Q3 (R5), worker (Sonnet).** certification.f90 (+105/-17), src/tests/certification/certification_contexts.f90 (+106), docs/certification_tolerance_anchoring_20260910.md (addendum). mass_row_cell_verdict returns a third, optional outcome: RESOLVED (cert_mass_round_margin * floor < cert_tol_mass_ceiling, judged as P16 anchored it) or UNRESOLVED (the margin times the floor reaches the ceiling, or the floor is not finite): an unresolved cell never reads within, its distance is max(1, q/ceiling), so the review's counterexample (q = 0.5 at floor 1) reads within = F, distance 1.0 (entry text: within = T, 0.5). The column loop is the pure mass_row_column_verdict, which counts the unresolved cells and names the first (n_mass_unresolved, j_first_mass_unresolved), and the report names that cell even

when a different cell binds the verdict. certification 84/0 (70/0 before the new rows), krylov_and_dogleg 334/0, steady_species_rows 195/0 (READ from the Q3 report; the advisor's run of certification on the merged tree 84/0). The addendum records that the margin of 10 was measured on three converged or converging fixtures only, none a near-zero-flow or molecular-EOS column: the UNRESOLVED outcome exists for the regime the fixtures do not reach.

- **Q4 (R4 in the marching caller), advisor.** The marching loop of EXHALE_main.f90 reads element_step_last_status after the element step: in phys mode a refused step refuses the whole attempted step (as_reject_element at the diffusion operation, the verdict text naming the test that refused it), which the controller restores and retakes at half dt like a carrier interval left uncovered; in init mode the march goes on from the composition the step restored and the refusals are counted (the final report prints "element steps refused inside the initialization march: n"). A test knob, element_refuse_leading_steps_for_test (EXHALE_ELEMENT_REFUSE_STEPS, zero in every production run), refuses the first n element steps so the path can be exercised: on mol_diffusion in phys mode with two refusals injected, attempt 1 at dt 1.47451E-04 and attempt 2 at 7.37253E-05 are refused "at the element diffusion step: an element invariant broke" and attempt 3 runs at 3.68626E-05; in init mode the two refusals are counted and the run exits 0. (The phys-mode run of that fixture then fails at step 0 on the coupled source step's fixed point with or without the knob, exit 2: a property of that fixed-step snapshot in phys mode, not of Q4.) Found beside it and fixed: the refusal census of attempted_step_note_step stopped one reason short (as_reject_source_fixed_point), so a source-energy refusal was never counted by reason; the bound is now as_reject_source_energy.
- **The stagnation row of grid_and_gates.** On the joint progress measure the configuration output_state_consistency.sh used to force the stagnation ending (EXHALE_JFNK_MAXIT=5, EXHALE_CARRIER_TRUST=1e-4) runs out its budget instead: the starved hydrodynamic rows keep falling (energy 4.9E-01 to 3.9E-02 over 8 passes, 1.1E-01 at 20) while the H2 row stands at 7.3E-02. The row now uses EXHALE_JFNK_MAXIT=40 with EXHALE_CARRIER_TRUST=1e-6: the hydro solves reach their roots, the H2 row stands at 7.38E-02 for four passes and the ending is announced at pass 4; the second assertion of that script (the ending hands back one state) measures 7.1e-16 against 1e-12. The script's header and the suite README say why the configuration moved.

Gates

Advisor verification of the merged tree (private build build_Q1 / EXHALE_Q1.x of the default toolchain, deleted at the close): carrier_retry 138/0, certification 84/0, element_operator 28/0, attempted_step 70/0, steady_species_rows 195/0, krylov_and_dogleg 334/0, diffusion_tests 35/0, carrier_returned_state_acceptance 36/0, carrier_reference_scales 14/0, carrier_constraint_attribution 10/0, species_masses 9/0, steady_completion_flag 3/0; grid_and_gates 198/1, the one FAIL the stagnation row on its old configuration (the suite was started before the script was re-anchored), and the re-anchored output_state_consistency.sh 4/4 on its own afterwards. make check (advisor, 2026-09-12 10:35 KST, the shared build/ rebuilt from this tree with the gfortran on PATH, conda-forge 16.2): **REGRESSION PASS (all cases byte-identical)**, 64 file comparisons PASS with data identical. A first launch of the gate with the

system gfortran 13.1 on PATH was stopped before its verdict and is not counted: the goldens are snapshots of the 16.2 build and a verdict against another compiler would say nothing about this change.

Reported by the workers and not fixed

- Q2: the out-of-bounds outcome of the element step (`element_step_out_of_bounds`) is not reachable on a synthetic column of the suite's size (the drift flux vanishes at both ends of the composition axis in the discrete operator), so the new rows cover the solve-failed and nonfinite outcomes only; the one measured out-of-bounds excursion of record is P1's, on the atomic reload.
- Q3: `grid_and_gates` row `mass_flux_spread_recomputed_from_output` failed once on the worker's control run ("`no_dump_line`") and passed on its measured run and on the advisor's; the P17 note on two concurrent invocations of that suite sharing something outside `EXHALE_TEST_OUT` stands.

The oxygen rows in the physics document (2026-09-12, user report)

The user read page 18 of `docs/EXHALE_physics_and_algorithms.pdf` and found that "The molecular rows" wrote the four $\text{H}_2/\text{H}_2+/\text{H}_3+/\text{HeH}+$ balances and nothing of the oxygen-bearing molecules. Confirmed against the code: with `thereis_oxychem` the cell system of `System_HeH_mol_metals` carries two more unknowns, $n_{\text{OH}}/n_{\text{O}}$ and $n_{\text{H}_2\text{O}}/n_{\text{O}}$, whose balances `oxygen_carrier_rows` writes on `O1/O1r`, `O2/O2r`, the three H_2O photolysis channels and the OH photolysis, with the H_2 row gaining the exchange with the water cycle, free atomic oxygen closing the family ($n_{\text{O}} - n_{\text{OH}} - n_{\text{H}_2\text{O}} - n_{\text{CO}}$) and the H closure gaining $n_{\text{OH}} + 2 n_{\text{H}_2\text{O}}$; the reverses come from detailed balance on the NIST-JANAF Shomate Gibbs energies; the $\text{O}_4 + \text{O}_6$ pair is one channel with $\text{O}(1\text{D})$ eliminated exactly; CO is not a row of the cell system but a transported carrier destroyed one-sidedly by $\text{He}^+ + \text{CO}$ (Langevin $1.6\text{e-}9$, UMIST RATE22 4068) and by Lyman-Werner photodissociation with the Visser et al. (2009) shielding, the He^+ row taking the He^+ it consumes at the frozen CO of the sweep. A paragraph "The oxygen rows" with the two balance equations now follows "The molecular rows" in `docs/physics_overview/ionization.tex`, the oxygen-option subsection cross-references it, the carrier-transport subsection got a label, and three references were added (Baulch et al. 2005, Chase 1998, Millar et al. 2024). Rebuilt with `latexmk`: 58 pages, no undefined citation or reference, no overfull box.

ESWENO assessment memo (2026-09-12, user question)

The user asked what the energy-stable WENO schemes gain and pointed at the three papers under `references/`. New memo `docs/esweno_assessment_20260912.{tex,pdf}` (5 pages), written from the published versions of Yamaleev & Carpenter (2009a, JCP 228, 3025; 2009b, JCP 228, 4248) and Fisher et al. (2011, JCP 230, 3727) and from `src/modules/states/Reconstruction.f90`: the energy estimate comes from an added nonlinear dissipation term (2009a Eqs. 9, 35 to 37) whose proof needs only $0 \leq w \leq 1$ and $\sum w = 1$; the new weight functions (Eqs. 18, 21, 22) with $\epsilon = O(\Delta x^2)$ hold the design order at smooth extrema and are separable from the estimate; the proof covers linear hyperbolic systems in characteristic form on periodic grids, with no proof for the Euler equations or a component-wise reconstruction (their Sections 5 and 6.2); the 2011 closures keep it on a finite domain with

nonuniform flux points near the boundaries; the cost is 15 to 20 percent (2009b Table 1). EXHALE's WENO3 already uses the 2009a weights and floor and not the dissipation term, and three properties of the code (finite-volume state reconstruction on a stretched grid, component-wise, nonlinear Euler) keep the estimate from carrying over. No code change; nothing measured on this code attributes an instability to the reconstruction's symmetric part.

The continuous transfer equations in the physics document (2026-09-12, user request)

The user asked for the flux and rate formulas of Caldiroli et al. (2021) Eqs. (3) and (4) in docs/EXHALE_physics_and_algorithms.tex, to be added only after checking them against what the code computes. Checked in photoionization_field_at_cell_HHe (util_ion_eq.f90): the outside column times cross sections gives tau_out, the cell's own densities give dtau, the working flux is $F_{\text{XUV}} * \text{cell_mean_attenuation}(\text{tau_out}, \text{dtau}) * \text{opa_pf}(j)$, the rate is $\text{sum}(\text{int_f} * \text{sigma} / E * dE) * 1e-18 * \text{erg2eV}$ and the one-particle heating $\text{sum}(\text{int_f} * (1 - E_{\text{th}}/E) * f_{\text{heat}} * \text{sigma} * dE) * 1e-18$, contracted with the densities in photoheating_of_cell; the discrete equations already in the document are these lines term by term. Added at the head of "Transfer along the radial ray": the continuous form $F_E(r) = F_E(a) \exp(-\text{tau}_E)$, $\text{tau}_E = \sum_a \text{sigma}_a \text{int_r}^{\{r_{\text{top}}\}} n_a dr'$, Gamma_a and H(r) as Caldiroli's Eq. (3), with the one difference stated (the column starts at zero at the outer edge of the grid, r(N+Ng), not at a). Rebuilt: 59 pages, no undefined reference, no overfull box.

FUV bands B3 and B4 integrated from the spectrum file (2026-09-12, user instruction)

The user asked why Stellar LW flux and Stellar FUV B3/B4 flux are separate inputs when a spectrum is given. READ from the code: with Spectrum type: Load the Lyman-Werner flux was already integrated from the file when its key was absent, but B3 (1231-1450 Å) and B4 (1451-2304 Å) came only from their keys; the analytic spectrum types cannot state those bands (a power law is the XUV fit extrapolated, a Planck the photosphere) and the ionizing photon grid does not carry them, which is why the keys exist. Instruction: integrate B3 and B4 from the file as well; Ly-alpha (band B2) stays a key, a reconstructed line flux being the better number.

Done: sed_read.f90's lyman_werner_band_flux_from_sed became sed_band_integrated_flux(w_lo, w_hi) (one trapezoid over the file's rows clipped to any band); input_read.f90 calls it with the band edges of oxygen_rates (fuv_band_lo_A/hi_A) for LW on any molecular run and for B3 and B4 with the oxygen chemistry on, when the key is absent and a file is loaded; a stated key, zero included, always wins (fuv_b3_flux_stated, fuv_b4_flux_stated, fuv_b3_from_spectrum, fuv_b4_from_spectrum in parameters.f90); the setup report names the source and a stated zero. Fixed beside it: the derivation ran AFTER the oxygen-chemistry warnings, so a loaded-spectrum run without the LW key was told "Stellar LW flux is zero" while the file supplied it; the block now runs before those warnings. Manual rows of the two keys updated (EXHALE_user_manual.tex, rebuilt).

MEASURED (private build, hot-Uranus oxygen case with inputdata/sed/hot_uranus_solar_koskinen2022.txt loaded, 3 steps): with the keys removed the run reads B3 4.696E+02 and B4 4.445E+05 erg cm⁻² s⁻¹ against a hand trapezoid of the same file 4.696068E+02 and 4.444851E+05 (LW without its key 3.687E+02 against 3.687009E+02); with the keys stated (694.8, 658782.2) the stated values stand; with B3 stated as zero the band is off and the report says so. No regression case combines a loaded spectrum with the oxygen chemistry, so the new path fires in none of them; make check on the merged tree is recorded below. Documentation of the rule (user instruction, same day): the physics

document's "Bands stated by their integrated flux" paragraph (docs/physics_overview/radiation.tex) now states the two sources of each band flux, the band integral (its Eq. for F_{band}), why the analytic spectrum types cannot supply the bands, and why Ly-alpha stays a key; the manual's Stellar LW flux row, the Oxygen chemistry row and the Tier 2b section (docs/EXHALE_user_manual.tex) say the same at the keys. Both rebuilt.

The H2 extent against Koskinen et al. (2022) Model A (2026-09-12, user instruction)

The user asked how far out Model A keeps its H2, then for a digitization of their Figure 8 and a full record of the present state in docs/koskinen2022_model_a_comparison.tex. Their text: "H becomes the dominant species at around $r = 2 R_p$, but the density of H2 remains significant at all altitudes" (their Figure 16: outflow supplies the H2 the chemistry destroys). Figure 8 digitized a second time, independently of the 2026-09-05 readings (400 dpi, ticks from the print, RGB isolation, every $0.1 R_p$): benchmarks/koskinen2022_model_a/model_a_fig8_digitized.txt; the two readings agree to 0.06 dex (H2) and 0.02 dex (H, He); $q_0(\text{H}_2)$ 0.837 and $q_0(\text{H})$ 0.028 recovered against the stated 0.84 and 0.026. $f(\text{H}_2)$ of Model A: 0.93 / 0.69 / 0.53 / 0.43 / 0.39 at 1.2 / 1.5 / 2.0 / 3.0 / 4.0 r_{base} , $f = 0.5$ at 2.15 r_{base} , never below $1e-2$.

New section 5 of the memo ("The H2 extent"), figure docs/figures/k22_model_a/h2_extent.pdf (docs/k22_h2_figure.py), the two stationary-route states pinned under benchmarks/koskinen2022_model_a/carrier_reload_state. MEASURED: the matched run matched_hnu_minus_I (spherical domain to $7.24 r_{\text{base}}$, Rate/4 + Mdot) has Model A's shape, no front, $f(\text{H}_2)$ 0.88 / 0.59 / 0.41 / 0.31 / 0.27 at the same radii, i.e. 0.05 to 0.12 below, and $n(\text{H}_2)$ a uniform 0.23 to 0.27 dex below theirs from 1.2 to 7 r_{base} . The carrier reload fixture of the solver series (backup/regression/carrier_elem_newton) is NOT that configuration: Roche domain to $4.71 r_{\text{base}}$, power-law spectrum at 0.048 au around a $1.2 M_{\text{sun}}$ star, Rate/2 + Mdot/2, He 2^3S on, no ionization transport, no conduction, default atomic rates (a first reading listed five differences from a diff against the superseded root input.inp; corrected against matched_hnu_minus_I/input.inp), pinned from a B5g state; in it $f(\text{H}_2)$ is zero above 1.1 r_{base} ($f = 0.5$ at 1.072, $1e-2$ at 1.097), 12 bounded passes move that to 1.076 / 1.103, and the fixed-wind fixed point fills the domain (0.97 / 0.91 / 0.76 / 0.51 / 0.39). So the "walking H2 front" of sections 8 and 9 is a statement about the fixture, not about EXHALE's Model A solution; the next measurement is the fixture re-pinned on the matched keys from the matched run's state.

The user caught, from the figure, that the first version of the comparison was drawn on benchmarks/koskinen2022_model_a/output, the superseded Roche-domain state of the morning of 2026-09-05 (top $4.72 r_{\text{base}}$), and that the fixture's domain differs from Model A's: the section and the figure were redrawn on matched_hnu_minus_I, both figure scripts now default to that run, and benchmarks/koskinen2022_model_a/README.md (new) says which directory is which.

Gate of the tree carrying the band-flux change, the H2-extent memo and the ESWENO memo: make check (advisor, 2026-09-12 15:38 KST, the shared build/ rebuilt with the gfortran on PATH): **REGRESSION PASS (all cases byte-identical)**, 64 file comparisons PASS with data identical.

The Model A fixture re-pinned on the matched configuration (2026-09-12, user instruction "proceed")

New fixture backup/regression/carrier_model_a_newton (README there): the matched run's input.inp with the stationary keys appended and its final state as the IC pair. Two changes were needed for it to run:

- input_read.f90: the refusal of Ionization transport: True with Solver: Newton and no Coupled carrier solve (2026-09-10) is lifted. READ in ionization_equilibrium.f90: the x_hp_fixed block imposes the transported proton fraction on every sweep whenever the carriers are transported and the state is a restart or carries a background, so in the partitioned alternation (P6) the hydrodynamic solve's sweeps are handed the transported proton and the carrier relaxation transports it; the refusal's premise (the last sweep re-solving the proton locally) held for the plain Newton finish it was written for and not for the alternation. MEASURED on the fixture: over 12 passes x(H+) moves by at most 0.011, downward, not to the local root. Manual key row and README_HOWTO.md updated.
- src/utils/map_state_to_grid.py (new; closes the interp_ic.py housekeeping item): the matched state was written on the 2026-09-05 grid and the mixed grid moved on 2026-09-10 (3414478) by up to $4.4\text{e-}4$ in the centers, which load_IC refuses at $1\text{e-}10$. The tool maps a state onto another grid's centers (rho, p, heat, cool in log against $\ln r$; v, T linear; every species as its ratio to the hydrogen nuclei, so He/H is exact in every mapped cell; the loader's He/H check at $1\text{e-}6$ failed on a first version that interpolated the densities themselves, $1.04\text{e-}5$). Round trip onto the old centers: rho, p, T to $2\text{e-}4$, v to $5\text{e-}3$.

MEASURED (12 bounded passes, EXHALE_JFNK_MAXIT=40, 8 threads, 3 to 7 s a pass, hydro info=0 every pass): the H+ carrier row falls $2.1\text{e-}1 \rightarrow 7.4\text{e-}4$ (gate $1\text{e-}5$), the H2 row ends at $7.0\text{e-}4$, and f(H2) moves TOWARD Model A: $0.884 \rightarrow 0.930$ at 1.2 r_base (Model A 0.933), $0.588 \rightarrow 0.650$ at 1.5 (0.691), $0.408 \rightarrow 0.431$ at 2.0 (0.530), unchanged above 3; f = 0.5 from 1.676 to 1.785 r_base (Model A 2.15); no cell below $1\text{e-}2$. The confined H2 of the solver fixture was that fixture's configuration. State pinned as benchmarks/koskinen2022_model_a/matched_fixture_states/pass12, figure h2_extent.pdf redrawn with it, memo section 5.4.

FOUND, reproduced, NOT diagnosed: a 60-pass run stops at pass 13 on the stagnation ending because the hydrodynamic energy row of cell 1 reads $2.073\text{e-}4$ (tolerance $1\text{e-}6$) on every pass while the solve's own residual there is $-1.0\text{e-}11$; with Conduction: False it is $4\text{e-}9$. The conduction source at cell 1 is $7.2\text{e-}5$, so the certification's assembly (after the outer loop's own sweep) and the solve's disagree by 0.5 percent of it. Candidates: the ghost cell's particle count behind the base-face conductive flux, formed from different ghost compositions on the two paths, or the sweep's local re-resolution of the ghosts. Recorded in docs/TO_BE_DONE.md and docs/ISSUES_20260909.md section 3.7 with the instrumentation it needs. Because the joint progress measure (Q1) is dominated by that constant entry (207 tolerances against the species rows' 74), the matched fixture cannot be run past the stagnation guard until it is removed.

Gate of the tree carrying the lifted refusal and the state mapper: make check (advisor, 2026-09-12 18:55 KST): **REGRESSION PASS (all cases byte-identical)**, 64 file comparisons PASS with data identical.

The base cell's energy row: diagnosed and fixed (2026-09-12, user instruction)

Instrumented on a private copy of the tree: the JFNK's final residual evaluation and the outer loop's certification assembly were captured on ONE state and subtracted cell by cell. At cell 1 every input (particle count, temperature, heating,

cooling) was identical and only the energy residual differed ($3.5\text{e-}7$); the conduction source of cell 1 was $5.406\text{e-}5$ in the JFNK against $5.371\text{e-}5$ in the outer loop; and the one input that differed was the particle count of ghost cell -1 (JFNK 0.545985, outer 0.546468, $8.8\text{e-}4$ relative), traced to the ghost DENSITY the JFNK's count was formed from: 1.22394867 (the ghost of the state that entered the sweep pass) against 1.22503037 (the ghost after the pass's own Apply_BC). newton_residual computed rho and the particle count at the top of each sweep pass and refreshed the ghosts at its end (base_boundary_states reads n_part_cell1 of the composition the sweep just left), then handed assemble_residual the refreshed u with the stale count; the base-face conductive flux, formed from the ghost temperature $W(3)/n_{\text{part}}$, was one composition stale. The certification forms the count from the state it is handed and was right. The ghost-fill hypothesis (Apply_BC missing in the outer loop) was tested first and refuted (ghost state identical to $1\text{e-}11$).

Fix (steady_newton.f90, newton_residual): after the sweep loop the density and the particle count are re-formed from the refreshed ghosts before the assembly. MEASURED on the matched fixture: the cell-1 energy row in the certification $7.8\text{e-}9$ at pass 1 and $3.5\text{e-}9$ at pass 3 (was $2.073\text{e-}4$); the 60-pass run no longer stops at pass 13 and spends its budget with the H2 row at $4.1\text{e-}4$ (cell 500, the top) and the H+ row at $3.7\text{e-}4$ (cell 135), hydro info=0 every pass, f(H2) at 1.2 / 1.5 / 2.0 / 3.0 / 4.0 r_base 0.931 / 0.660 / 0.460 / 0.346 / 0.296 against Model A's 0.933 / 0.691 / 0.530 / 0.430 / 0.389 (the loaded marching state 0.884 / 0.588 / 0.408 / 0.310 / 0.267), $f = 0.5$ at 1.851 r_base (Model A 2.15), x(H+) 0.061 / 0.133 at 2.0 / 3.0 (Model A 0.020 / 0.080). State pinned as matched_fixture_states/pass60, figure and memo section 5.4 updated. Without conduction or viscosity the count enters no row: wasp_full_newton reload data rows identical to the control; krylov_and_dogleg 334/0, steady_species_rows 195/0, certification 84/0, carrier_returned_state_acceptance 36/0, steady_completion_flag 3/0. make check recorded below.

Gate of the tree carrying the ghost particle-count fix: make check (advisor, 2026-09-12 22:57 KST): **REGRESSION PASS (all cases byte-identical)**, 64 file comparisons PASS with data identical.

10. PLAN_20260913: the findings of the two reviews of 2026-09-12 (2026-09-13)

Plan of record: docs/PLAN_20260913_review_fixes.md. User instruction: read docs/code_bug_optimization_review_20260912.md and docs/parallel_code_review_20260912.md, confirm each finding in the code, plan the corrections and carry them out, running the binary in parallel where that shortens the verification. Every finding (B1 to B6, P1 to P3) was confirmed in the source before the plan was written (the confirming lines are in the plan's table); items S1 to S9. Every number below is MEASURED on this tree unless marked READ.

- **S1 (B1, B2, O3): the band integral of a loaded spectrum.** sed_read.f90: sed_band_fluxes(nb, w_lo, w_hi, F, covered) replaces sed_band_integrated_flux; the file is read once for the three bands, validated as it is read (positive, finite, strictly increasing wavelengths; finite nonnegative flux; a malformed row stops the run with the row named), and each band gets the exact integral of the piecewise-linear table (the flux interpolated at the clipped edges before the trapezoid); covered(b) says whether the table reaches both edges, and input_read (band_from_spectrum) STOPS when a band the run needs is not covered, naming the key (a stated zero is the run's decision; a silent zero is not). MEASURED on the hot-Uranus oxygen case with

the Koskinen file: B3 4.686E+02 and B4 4.443E+05 erg cm⁻² s⁻¹ against the exact piecewise-linear integrals 4.685913E+02 and 4.443075E+05 (the entry text gave 4.696E+02 and 4.445E+05, 0.2 and 0.05 percent high). New test fuv_band_quadrature in spectrum_type (linear spectrum on unequal nodes with edges inside segments, on nodes and at the ends, exact to 1e-12; one covering segment; constant spectrum; an uncovered band): 14/0. wasp121b_sed_sensitivity moved to the new routine with its own transcription corrected the same way (spectrum_type 170/0 over the whole suite). docs/input_schema.md, the manual's key rows and the physics document's band paragraph say the new rule.

- **S2 (B3, B4, O2 counters): the thermochemical closure contract.** equilibrate_chemistry_at_fixed_conserved_state starts with `ok = .false.` and sets it only when a cycle's temperature increment is below `chem_cycle_tol` AND the sweep ledger reports no nonfinite and no off-simplex cell AND `p`, `T`, `heat`, `cool`, `eta` are finite with 'p

0, `T > 0`; the reason (converged, cycle budget spent, nonfinite composition, a cell left the element simplex, not admissible) and the last increment are returned and the run-wide counts kept (`n_chem_closure_cycles`, `n_chem_closures_reached`, `n_chem_closures_refused`). The relaxation's refusal branch was already restoring the trial. Test knob `chem_cycles_cap_for_test`; four new rows in `carrier_retry` (an exhausted closure refuses the trial, names the spent budget, keeps no step, hands back the entry composition bit for bit). One existing row was restated: on the synthetic column the pass after the refused trials is refused by the CHEMISTRY (one molecular clamp, budget violation 6.5e-4, on the background seeded at that state), which the old contract had hidden; the row now asserts what the history can be judged by (the pass is not refused at the interval and the history stays certifiable), and the section-11 pass, prepared on the background of the composition it relaxes from, keeps its two steps as before. `carrier_retry` 142/0. On the Model A fixture (3 passes) every closure is reached: pass lines identical to the day before.

- **S3 (B5, B6): the state mapper.** `src/utils/map_state_to_grid.py` validates finite, positive, strictly increasing radii in both source files and the target, their agreement to 1e-12, the column labels, a positive hydrogen-nucleus density, and that every physical target center lies inside the source's rows (ghosts included: the lower ghosts carry the inflow reservoir composition and are interpolated like every other row, never rebuilt from the physical cells; only a target ghost may lie outside and is then extrapolated linearly in $\ln r$); a violation is refused with exit 2 and nothing written. The output is an initialization seed: `# coupling: mode=init t_phys=0 certified=F ...`, the source line kept as `# mapped-from-coupling:.` New suite `src/tests/state_mapper` (identity mapping exact; a 2e-4-shifted grid round trip 1.8e-5 on a 400-row column; He/H invariant to 1e-12; mismatched species radii, out-of-support physical targets and non-monotone radii refused; ghosts extrapolated; metadata rewritten): 17/0. The Model A fixture's IC/ regenerated with it (physical rows identical to the earlier mapping, ghosts within 1e-4; the 3-pass run reproduces the day-before pass lines).
- **S4 (P1): the continuation dump.** `cce_dump_paths_read` reads the two environment paths once, serially, from

input_read; write_continuation_ dump takes the path by argument, keeps every read and write of the once-only flag inside the critical region, sets the flag only after the file is open, and records the cell (ieq_cell%jcell, a new field set by the sweep; dump format 2, the probe reads 1 and 2).

- **S5 (P2): explicit initialization.** keq_H_H_to_H2 no longer builds the H2 thermochemistry table lazily (it stops with a message if the table is not ready); the table is built serially by the main program, by the sweep's serial prologue (ioniz_eq) and by carrier_set_init, so every entry point that precedes a parallel read builds it. The Newton solver's EXHALE_FORCE_HYBRD1 switch is read once by newton_solver_read_environment from input_read; a driver that never calls it runs with the switch off. init.f90 switches dynamic team adjustment off, reports the team obtained (an omp single block, the master construct being deprecated since OpenMP 5.1), says whether the count came from OMP_NUM_THREADS or the default min(cores, 16), and its coverage comment now lists the present parallel regions (the old thread line was cut by its fixed format and said "default" whatever the source).
- **S6 (P3): the BLAS thread policy.** New blas_thread_policy.f90: the OpenBLAS entry points are resolved at run time (dlopen/dlsym, -ldl on the GNU link and in every test script), the count is set to 1 unless OPENBLAS_NUM_THREADS is stated, and the setup report says what was found and set; a build without OpenBLAS is left alone. MEASURED on the wasp_full_newton reload at 8 OpenMP threads: 16.33 s with 1 BLAS thread against 16.32 s with 8 (the library's default here follows OMP_NUM_THREADS; unset, it is the machine's 72): the band factorizations gain nothing from a BLAS team, and the policy costs nothing.
- **S7: verification, in parallel.** Suites on the private build, run concurrently: spectrum_type 170/0, carrier_retry 142/0, carrier_returned_state_acceptance 36/0, attempted_step 70/0, certification 84/0, steady_species_rows 195/0, krylov_and_dogleg 334/0, element_operator 28/0, species_masses 9/0, steady_completion_flag 3/0, grid_and_gates 198/0, state_mapper 17/0; the Model A fixture 3 passes; the two BLAS timing runs; make check (the full matrix, OMP_NUM_THREADS=1, run on the merged tree while the suites ran): REGRESSION PASS, every case byte-identical against the goldens, as the plan required (no option defaults changed; the band integral is not exercised by the gate's power-law spectra, the closure contract reports the same result on every gate case).
- **S8: measured-first items, not changed** (O1, O2 beyond the counters, A, B, C, E of the parallel review): recorded in docs/TO_BE_DONE.md with the measurement each needs; the counters of S2 and the team/BLAS report of S5/S6 are the instrumentation for them.

11. PLAN_20260913_lhs_stationary: the LHS 1140 b stationary solve (2026-09-13 to 2026-09-16)

Plan of record: docs/PLAN_20260913_lhs_stationary.md, items L1 to L14; the item memos are docs/lhs1140b_stationary_L*.md, the design of record for item L12a is docs/lhs1140b_stationary_L12a_design_20260913.md, and the campaign they serve is LHS1140b/MODELS.md. User decision of 2026-09-13 (option 1): the stage-2 solver program, closed on 2026-09-10 and reopened once on 2026-09-11 for the partitioned route, is reopened for ONE item -- making the current code reach a stationary, certifiable

state of the LHS 1140 b atomic wind (subsonic to the 30 R_p boundary, T = 226 K helium-rich base at 1 microbar, binary H/He element diffusion on). Without it the campaign of LHS1140b/MODELS.md has no solutions to run. Every number below is MEASURED on this tree unless marked READ. A line count given for a whole file is from git diff --numstat against 43bc28c on the tree at the close; a count attached to one item is that item's own diff, from its memo.

- L1: the archived solutions were never stationary in the base layer.** The 2026-08-30 states were judged on a flux window that began at 1.2 R_p (Update_EXHALE_stage1 sections 133 and 161), and on the present cellwise residual they carry an O(1) row everywhere between the base and about 1.1 R_p. The energy row is its own flux divergence in every cell 1 to 137: $|dF_3|/|\Gamma - \Lambda| = 7e5$ at cell 1, 4.9e1 at cell 16, 1.02 at cell 137, where the two terms add to the 1.979 maximum. The mass flux of the archived state is 115 times its far-wind value at 1.012 R_p and decays at 4.5 percent per cell out to 1.3 R_p. The momentum row carries a uniform -2.79 percent offset, of the size and sign of the Jupiter-radius change in R₀ (+2.26 percent). The base velocity sawtooth of P44 is not the cause: removing it leaves the energy maximum identical and takes 30 percent off the mass row, and putting the archived heating back into the same hydrodynamic state lowers the worst cell from 1.979 only to 1.930 (2.5 percent of the maximum). Separately the physics of the base layer has moved by factors 2 to 3.4 since 2026-08-30 (total heating 3.36x the archived value at $r = 1.000$, 1.95x at 1.010, 1.00x at 1.035 and 0.85 to 0.98x above 1.05, the photon-grid quadrature removing the 9.5 percent overcount; the He(2³S) channel 0.55 to 0.86x; the He recombination-photon channel 0.1x at 1.035 falling to 1e-5x at 20 R_p). The residual belongs to the state, not to the diagnostic.
- L2, L3: no march of any kind settles, and the obstruction is the clock.** Global-dt marching at CFL 0.6 holds du at 8 to 15 for 28 000 steps from a cold start and 4 to 11 for 25 000 from the mapped seed, the cellwise residual 1.6 to 2.0 at every 500-step checkpoint; at CFL 0.05 and 0.1 du rises monotonically (1.54 at step 7202, 2.27 at 11121; 4.82 at 11349) while the march drives an inward flow of 7.7e3 cm/s at 1.05 R_p. Local pseudo-time (Time stepping: Local) from the mapped seed gives du 1.1, 4.5, 44, 8.2, 5.9, 6.2, 0.47, 1.95, 1.8 at steps 4000 to 8000 with the residual 1.84 to 1.99; from the cold start the state stops changing at step 1167 (dtu 5e-13) at du 5.9, residual 2.2. L3 then established the positive half on arithmetic: a state meeting all three hydrodynamic tolerances DOES exist in the base layer (continuity floor 8.43e-10 against the anchored tolerance 8.43e-9, energy 4.56e-9 against 1e-6, momentum 1.9e-14 against 1e-8, no unresolved cell), admissible only through the anchored mass rule, the fixed 3e-12 standing 281x below the floor of cell 1. The stationary base velocity is 0.0297 cm/s, Mach 2.7e-7, 1.8e4 times below the archived seed's -538 cm/s, so L1's "7e5 times the heating" is the seed's velocity and not the layer's. The obstruction is the clock: the base cell's thermal time is 4.8e7 CFL steps and the transit to 1.15 R_p 1.8e8, against the 1e4 to 3e4 steps any march ran, and the update map is linear in dt with no floor, so the march's fixed point IS the root. LHS 1140 b is harder than the hot Uranus because the 1 microbar level is already the top of the absorbing column: heating peaks in cell 1 and cooling is 0.46 percent of it, against 325 percent on the hot Uranus, so every erg leaves as enthalpy flux at Mach 2.7e-7.
- L5c: the O(1) base-layer energy residual is the DISCRETIZATION, and the well-balanced reconstruction removes it.** In the HLLC face energy flux $F_3 = S*(E^* + p^*)$ the contact speed carries a pressure-jump part

$(p_R - p_L)/(\phi_L - \phi_R)$ which at Mach 3e-7 to 1e-6 stands one to three decades above the velocity part; it is the design-order truncation error of the WENO3 reconstruction of a near-hydrostatic column (it survives eight 1-2-1 smoothings and converges at order 2.97 to 3.00 under refinement), not the odd-even mode and not the base condition, and the velocity part alone balances the net heating to 1.0 to 1.9 in every base cell. On the same seed, the same input and the same binary:

option	hydro	mass	momentum	energy	s	outcome
none	info=2	1.08	3.2e-2	0.63	1325	pass refused
Low-Mach	info=2	1.32	1.4e-2	1.24	1232	pass refused
damping:						
2.0e-2						
Well	info=0	2.6e-10	1.8e-14	4.6e-9	85	CERTIFIED
balanced:						at outer pass 8
True						

Low-Mach damping does nothing to the energy row because it is a fourth difference of the VELOCITY and does not touch the pressure truncation error. The LHS 1140 b atomic wind with element diffusion certifies in about ten minutes from the mapped seed, and from the S1c and S3 seeds alike: the three solutions, whose base velocities differ by a factor 1e4, agree to 3.4e-10 in ρ and 2.5e-8 in v , and in $\dot{M}$ to seven digits. The element relaxation, which refused every pass without the key (element_step_out_of_bounds after 24 to 26 steps), reaches its fixed point in 32 steps in every pass with it. The HD 209458 b element-diffusion reload certifies at the same outer pass 12 with $\dot{M}$ moved by 1.3e-4. No source change: Well balanced: True is key K46, decided as an option on 2026-09-10 and left off because the two discretizations of the pressure/gravity pair are both second-order consistent and neither is a subset of the other, so every golden would move. The term-by-term statement, the mapping onto the code and the published basis (Kaeppeli & Mishra 2016, A&A 587, A94; Kaeppeli & Mishra 2014, J. Comput. Phys. 259, 199; with LeVeque 1998, Botta et al. 2004, Chandrashekar & Klingenberg 2015, Edelmann et al. 2021 for scope) are the new docs/well_balanced_and_hydrostatic_20260913.tex, written from the source and from the two articles in references/. Left open by the item, and still open: the certified $\dot{M}$ is log 8.29 against the archived 7.80 (the factor 3.1, resolved by L4 below), base cell T 405 K against the prescribed 226 K, a three-cell base velocity artifact (-0.092 cm/s against the stationary 0.150), and the grid convergence of the well-balanced solution is not measured.

- **L5a: no seed repairs it, and two properties of the archived column.** Six constructed seeds (LHS1140b/models/make_stationary_seed.py: the stationary velocity below 1.27 R_p , with and without the odd-even removal, with a base pressure anchor, and a Bernoulli-consistent temperature) are all refused at outer pass 1 without the well-balanced key (hydro info=2, energy row 0.63 to 1.99; the element relaxation then reports element_step_out_of_bounds after 24 to 26 steps and restores the entry composition). Found on the way: the archived column's pressure is 2.92 percent above the 1 microbar reservoir (the column is 2.91 percent steeper than the current gravity), so the base face runs a Mach 2e-2 inflow, 6.9e4 times F_{wind} , whatever the

interior velocity, until the column is scaled by 0.9716; and with the stationary velocity imposed, the DISCRETE energy flux divergence at the worst base cells is $-1.0 \times (\text{heat} - \text{cool})$ where the continuum value on the same state is $+0.5$, which is L5c's finding reached from the other side. Also recorded: load_IC rebuilds the density from Ion_species_IC.txt and never from p and T.

- L5b: the element relaxation's bound test is right and is not the cause.** Its fixed point on the state the hydro solve hands over (pass 1, info=2) lies ON the bound: that state's mass flux falls 27 percent between 1.36 and 26.9 R_p while the helium flux falls 9 percent, so $X_{\text{He}} \rightarrow 1.000000$ at 13.9 R_p , where the hydrogen element flux crosses zero. The overshoot is linear in the step, so no step control clears it (omega acts only after a returned relaxation, and the retry budget of 20 halvings cannot reach $1e-12$ from $5e-3$). Positive control: on the loaded Slc wind, whose $\rho \ v \ r^2$ is uniform to $1.2e-16$, the same operator converges to $X_{\text{base}} = 0.8659$. With Restart intent: stationary equilibrate the relaxation reaches its fixed point in every pass (30 to 32 steps) and the outer loop instead ends at pass 5 with outer_no_progress, the hydro rows stalled at mass 1.27 and energy 0.864. So the obstruction was the hydro solve, which is L5c. Reporting defects recorded and not fixed: the refusal names no cell, bound or overshoot; the retry budget is fixed regardless of the tolerance; the boundedness argument of the operator's comment covers $\text{div}(F_{\text{rho}}) > 0$ only.
- L4: the run recipe, and why each key is in it.** The recipe of record for a prescribed-composition atomic case (LHS1140b/models/run_case.sh, MODELS.md section 6) is the mapped archived seed plus Well balanced: True (L5c), Secondary_ionization: Immediate, Restart intent: stationary with Solver: Newton, and EXHALE_PTC_DTAU0=1.0, with CFL: $1.0e-12$ on the post-processing pass. The Immediate key is a physics correction, not a convenience: the STAGED default arms the Shull & van Steenberg channel only when a march first converges, and this route takes no time step, so a state solved under the default is written with sec_ion=F in its own coupling header while the post-processing pass -- which applies the coupling unconditionally and is where every reported number comes from -- reports the same wind with it. MEASURED on the $K_{zz} = 1e9$, He/H = 1.6261 case: certified at outer pass 8 under STAGED against pass 7 under Immediate, $\log_{10} \text{Mdot}$ 8.29 against 7.87 (archived 7.80, READ), and $n(\text{He } 2^3\text{S})$ differing by 74 percent beyond 1.5 R_p , where the He I 10830 line forms. CFL: $1.0e-12$ is there because Do only PP takes one time step before it stops: at the default CFL the written state differs from the state read by $1.2e-6$ in ρ and 0.59 in v , at $1e-12$ by $3.6e-14$ and $2.3e-12$. Found and fixed on the way: src/utls/element_flux_closure.py wrote the retired Valve eps: key into every iteration input, so every closure rung had been refused at startup. Cost, MEASURED end to end at 8 threads on atomic_scalar_gj1132_kzz1e9/HeH1.60: 15 min 15 s, seed to line, of which 7 outer passes of 117 to 166 s are the wind (final rows mass $5.66e-10$, momentum $2.76e-13$, energy $2.21e-08$, worst gated species row $9.32e-06$ of $1.0e-05$ at cell 312, run.log; $\log_{10} \text{Mdot}$ 7.87 from pp.log, red depth 4.3260 percent and FWHM 0.25376 Å from tpm_He10830_metrics.txt).
- L4b: why every metals-on case stalled at outer pass 2 on the energy row.** Diagnosis, no source change (docs/figures/lhs1140b_L4b_energy_row.pdf). The band $r = 1.15$ to 1.5 R_p where C I line cooling is 79 to 94 percent of the cooling (cool/heat 0.53 to 0.66 against 0.042 metal-free) is where the element relaxation moves X_{He} by 0.4 to 2.6 percent and the energy row stays at $3.7e-2$; the hydro solve then cannot take a step,

because one Newton step raises the continuity row from $2.1\text{e-}9$ to $6.5\text{e-}4$ against its $3\text{e-}12$ tolerance, the entry state is always the best judged iterate, `n_stall_best = 20` stops the pass STAGNATED while the merit is still falling ($1.5\text{e-}2$ to $3.2\text{e-}3$), and `outer_no_fall_max = 3` ends the solve. The same on the scalar-base C/N/O case with no profile and at He/H 2.09. Hydro alone, with the composition held, certifies; `omega 0.125`, `dtau0 0.01`, the trust-region restart and the coupled carrier solve do not help. Two source changes proposed from it: reset the stagnation counter when the merit improves its own best, and replace the partitioned route's Krylov tolerance literal $1\text{e-}1$ by a forcing term (item L4d).

- L4d: the stagnation detector WAS the refusal** (`steady_newton.f90 +276/-36` at this item). It now reads both functionals: the counter resets on an improvement of the judged distance OR of the merit, the merit taken in a FIXED scale (the entry iterate's column scale; a scale rebuild moved the merit of an unchanged state by 1.6x) and compared with the value 20 iterations back with a margin of 0.1 (`EXHALE_MERIT_FALL_MIN`; 0.05 to 0.2 give the same result, and residual evaluation is bit-reproducible across 4/8/16 threads so no noise anchors it). RED: the scalarCNO He/H 2.13 pass-2 state STAGNATED at 40 iterations returning the entry state ($1.1\text{e-}1$); GREEN: `info = 0` at iteration 98 and $2.8\text{e-}8$, and the full outer loop then runs passes 1 to 6 at `info = 0` with `omega` never halved, the element row reaching $6.4\text{e-}6$ at pass 7. The forcing term (Eisenstat & Walker 1996, choice 2, `gamma 0.9`, `alpha 1.618`, ceiling $1\text{e-}1$, floor $1\text{e-}4$; the published safeguard gate is unreachable under that ceiling and is applied ungated, said in the comment) is NOT that repair: the Krylov cycle reaches $7.9\text{e-}4$ of its right-hand side in one product, but the linear residual it leaves is $2.6\text{e}5$ (mass), $2.0\text{e}5$ (momentum) and 0.44 (energy) of the same rows of `F`, a scalar 2-norm being blind to rows ten decades below the dominant ones (`EXHALE_MODEL_ROW_EQUIL` does not help); a row-resolved diagnostic `EXHALE_LINEAR_ROWS=1` was added instead. Controls: `atomic_elem_newton` CERTIFIED at pass 12 before and after (843 against 909 iterations), `carrier_model_a_newton` identical verdict and rows, `wasp_full_newton` `info = 0` at log `Mdot 13.30` with maximum relative movement $2.5\text{e-}8$, `mol_base_handoff` bitwise. Eight comments and one routine name at the same site carried a software-engineering word for the residual assembly, and were corrected to say what the routine does (`residual_terms_in_evaluation_order`).
- L4e: judging the linear solve by the certified rows is REFUTED as the repair; the pseudo-transient ramp is the obstruction** (`steady_newton.f90 +313/-72` beyond L4d: `judged_size_of_the_hydrodynamic_rows`, `row_scaling_of_the_linear_model`, the preconditioner equilibrated on that path, five duplicated three-unknown expressions removed, the Levenberg-Marquardt gradient in model coordinates, the count of cells outside tolerance printed for each row; both options OFF by default, `EXHALE_JUDGED_ROWS=1` and `EXHALE_KRYLOV_TOL_ABS=1`, bit-identical off). With the rows judged and the Krylov cycle driven to $8\text{e-}10$ of its right-hand side (four products, the mass-row linear residual $2.6\text{e}5 \rightarrow 1.6\text{e}2$) the first Newton step still raises the mass row to $1.09\text{e-}2$, the same to three digits: the continuity overshoot is the Newton DIRECTION of these states. What holds them is that `dtau *= lam f2/f2_try` is tied to the line-search merit, which is nearly flat on them (8x in 60 iterations), so `dtau` never leaves `dtau0 = 1.0`. With `EXHALE_PTC_DTAU0=1e8` on the entry text, from the states the outer loop had refused, scalarCNO He/H 2.13 CERTIFIED at outer pass 1 in 22.8 s -- the first certified LHS 1140 b solution carrying C, N and O -- and He/H 4.0 at pass 3 in 93 s, 4 to 6 Newton iterations

each at $\lambda_m = 1$ with no non-monotone accept, where the same states at $\text{dtau0} = 1$ stall (784 s, $\text{info} = 2$). A RAW seed at $1e8$ fails (mass 1.92), the ramp's floor being dtau0 . Campaign recipe from this: $\text{dtau0} = 1.0$ from the seed, and a continuation from the written state at $\text{dtau0} = 1e8$ where the outer loop ends refused with the hydrodynamic rows stuck, recorded in REPRODUCE.md.

- L4f: the energy row of the outer boundary cell is a property of the STATE and of the linear solve, not of the boundary** (docs/figures/lhs1140b_L4f_outer_energy_row.pdf; no source change). Four certified states satisfy the same row at cell 500 to $1.7e-12$ to $3.5e-11$, two of them supersonic at the face where the ghost does not enter the flux at all; the row is smooth to $2.7e-10$ under $8e-6$ perturbations; letting the ghost follow the cell RAISES the sensitivity, so there is no self-chasing; and at the refusing pass the energy row is outside $1e-6$ in 499 of 500 cells (worst cell 157 at $1.0e-3$), cell 500 being merely the flat outer plateau the worst-row line names. Its terms: residual $3.4e-4$ of the heating, 0.9 percent of the flux divergence, $4e-6$ of the face flux, with no conduction or viscosity term. The oscillation is the linear solve's direction -- each Krylov cycle reaches its tolerance in one product while leaving 0.20 (mass), 0.56 (energy) and 512x (momentum) of those rows, and 60 iterations return the entry state while the merit falls 8x over 58 non-monotone accepts. On a 45 R_p domain the row does not vanish but becomes everything (pass 1 stalls at $4.8e-2$, worst row at cell 500 in 306 of 343 iterations; not like-for-like, the seed being extrapolated and the grid coarser). No anchored rule was written for the last cell: the arithmetic resolves it, and a new tolerance there would be a number chosen to pass a snapshot. Reported beside the item: make_models.py gave scalarCNO and photochem a Resid tol 12.5x stricter than scalar with no recorded basis.
- L4g and L7d(2): the pseudo-time ramp of the partitioned hydro solve** (steady_newton.f90 +114/-7 at L4g, plus the L7d correction). $\text{dtau0} = 1.0$ is $1.4e4$ explicit-stable steps but only 1/36 of the column's flow time r/v (35.8 code units), so the shift $1/\text{dtau}$ sat above the advective mode that fixes the stationary state, and a merit-tied ramp whose floor was the START value could not leave it ($1.06 \rightarrow 1.42$ in 12 iterations). Delivered: the pseudo-time floor is the explicit-stable step of `eval_dt` at the entry state, which is what the old comment claimed and the three places that read the floor now read; an accepted step doubles dtau (the published clip: Kelley & Keyes 1998, SIAM J. Numer. Anal. 35, 508; Mulder & van Leer 1985, J. Comput. Phys. 59, 232; Gropp et al. 1998, NASA CR-1998-208435 section 2.1); dtau is printed per iteration; the option `pseudo_time_doubles_on_an_accepted_step` is DEFAULT ON with EXHALE_PTC_RAMP_DOUBLE=0 restoring the entry text bitwise. The brief's own proposal, a ramp read from the judged distance, was implemented in five forms and REFUTED: the first Newton step of these states raises the judged distance at every dtau , at $1e8$ as well, and the next step removes it, so a ratio-read ramp cuts dtau on that very step. GREEN at $\text{dtau0} = 1.0$ with NO continuation: the scalarCNO pass-8 state certified at outer pass 1 in 18 iterations where the entry text STAGNATED at 263 in 953 s; L4d's pass-2 state $\text{info} = 0$ at 20; the raw archive seed at 23; `atomic_scalar_gj1132_kzz1e9/HeH4.0` from the certified 2.13 seed certified in one run at pass 5 in 509 s with ω never halved. L7c then measured the other end: the rule counted a line-search step of $\lambda_m = 8e-3$ as accepted, so on a FAR state dtau reached $1e14$ in 40 iterations and the solve became unshifted Newton (124 iterations, $1.87 \rightarrow 1.67$; with the doubling off, 159 iterations, $1.87 \rightarrow 4.5e-2$, $\lambda_m = 1$ in 149 of them). The delivered

correction (L7d) is `lam_of_a_full_step`: double at $\text{lam} \geq 0.5$, otherwise multiply dtau by lam . The brief's rule, double only at $\text{lam} = 1$, was REFUTED -- the near states take $\text{lam} = 0.5$ steps and stall under it -- while the delivered rule leaves every near state bitwise identical to L4g and on the far state cuts dtau at the first damped step, so the merit falls $1.3\text{e-}1$ to $5.3\text{e-}3$ where the entry text froze it (element reload pass 1, mass $6.9\text{e-}2$ to $3.2\text{e-}2$). Controls across L4g: `carrier_model_a_newton` same verdict at movement $1.8\text{e-}10$, `atomic_elem_newton` certified at pass 12, `wasp_full_newton` $1.2\text{e-}8$ and $2.2\text{e-}8$ against golden, `mol_base_handoff` PASS.

• **L6: the C I ionization balance is right and its cooling was wrong by one to three decades** (`Cool_coeff.f90`).

The attenuated C I photoionization rate, rebuilt independently from the GJ 1132 SED and the Verner (1996) cross section, agrees with the code to 0.5 to 1.3 percent; 11.26 eV is a bin edge; nothing shields the 11.26 to 13.6 eV band but C I itself ($\tau 1\text{e-}2$); and with $0.57\text{ erg cm}^{-2}\text{ s}^{-1}$ below the H I edge against 19.4 above it, $\alpha n_e = 5\text{e-}6$ to $5\text{e-}5\text{ s}^{-1}$ against $P(\text{C I}) = 5$ to $9\text{e-}7\text{ s}^{-1}$, so 94 to 99 percent neutral carbon is physical. The COOLING was not: the C I line-cooling fit is a coronal (low-density) form, but 99.7 percent of it comes from the metastable $2p^2\text{ }^1\text{D}_2$ and 1S_0 levels whose critical densities ($2.0\text{e}4$ and $1.3\text{e}7\text{ cm}^{-3}$ from the NIST A values) lie BELOW the wind's n_e (1.2 to $3.3\text{e}7\text{ cm}^{-3}$), so the coded rate exceeded the LTE emission bound by 20x at 1.86 R_p and 1500x at 1.025 R_p . On the same state cool/heat at 1.15 to 1.5 R_p fell from 0.65 to between 0.04 and 0.13 (the metal-free control's 0.042), which is exactly the energy-row obstruction L4b had found. The item also named the same defect in N I 2D ($A\ 2.8\text{e-}5\text{ s}^{-1}$, $n_{cr}\ 8\text{e}3\text{ cm}^{-3}$), O II 2D ($2.1\text{e-}4$, $1\text{e}4$), N II 1D ($3.9\text{e-}3$, $4\text{e}5$) and O I 1D ($7.5\text{e-}3$, $1\text{e}6$), which is item L6b.

- **L6b: six C/N/O coolants now carry the CHIANTI statistical equilibrium at the local electron density** (`Cool_coeff.f90` +1443/-24 including six 41×29 tables in ($\log T$, $\log n_e$), `metal_cooling_above_ground_term`, `cool_NI_ne_func`, `cool_OII_ne_func`, the L6 two-level patch removed; `util_ion_eq.f90` +16/-7, N I and O II added to the `cno_chianti` block; new generator `cooling_data/metal_cooling_density_resolved.py`, 155 lines, CHIANTI 11.0.2 through ChiantiPy 0.15.2). C I, C II, N I, N II, O I and O II were all evaluated coronally above the critical densities of the levels they excite. They now carry the CHIANTI cooling of the levels ABOVE the ground term at the cell's own n_e , interpolated from offline tables the way Fe II already was, while the split ground term stays with the code's own exact solve, which carries the H collisional de-excitation and the escape probability CHIANTI cannot. The runtime n-level solve was NOT adopted (40 to 200 levels per ion inside the sweep and the Newton residual, with no line-diagnostic return). Against CHIANTI directly over 1000 to 20 000 K and $n_e\ 1\text{e}4$ to $1\text{e}9\text{ cm}^{-3}$, 42 points and six ions: 0.02 to $5.4\text{e}4$ x before, 0.969 to 1.009 x after, interpolation error median 0.3 to 0.8 percent above 2000 K. Found on the way: the old coronal fits were built from the CHIANTI .scups THEORETICAL transition energies while the level populations use the OBSERVED ones (O II $4\text{S-}2\text{D}$, 1.1436x in dE, a factor 14 in the exponential at 2000 K), which the tables remove. Audited and needing no change: Mg I, Mg II, Ca II and Na I (permitted resonance lines, $n_{cr}\ 3.5\text{e}13$ to $1\text{e}16\text{ cm}^{-3}$), Fe I (no CHIANTI model atom), Fe II (already density-resolved) and the He 19.82 eV channel (below $2\text{e-}7$ of the cooling). Movement, goldens NOT refreshed at the item: the LHS rung's cooling at 1.15 to 1.5 R_p -17 to -21 percent with cool/heat 0.036 to 0.105 ; `wasp_full` $\log \text{Mdot}$ 13.36 unchanged but cooling at 1.05 R_p -72 percent, T +7 percent at 1.2 R_p and +14 percent at

1.5, rho up to -28 percent, and the run stopping at 12 251 steps instead of 17 671; mol_metals T +0.08 percent. Reported and not changed: the He 19.82 eV channel belongs in an energy ledger of the tracked 2^3S population; Fe I's LTE lower levels; Mg I 4571 and Ca II 7291/7324 are missing from the fits, an underestimate. The two physics_probe drivers that had aborted after item S5 now initialize the H2 thermochemistry table.

- L6c: the regression movement of the consolidated binary is the C/N/O cooling and nothing else.** The seven cases outside the $1e-3$ relative tolerance -- wasp_full, wasp_he23off, wasp_full_newton, mol_metals, mol_ir_bands, lower_profile, oxygen_chemistry -- are exactly the seven that carry a C/N/O elemental reservoir; the nine that pass carry helium only, one to one over the whole matrix. A control build with only Cool_coeff.f90 and util_ion_eq.f90 reverted to HEAD reproduces the golden BITWISE in six of the seven, the marching stopping step included, and wasp_full_newton to $2.2e-8$, and its difference from the measured build is the difference the log reports, file by file, row by row and value by value. Cooling is the cause and the profiles the effect: from one held state the two builds differ by at most $3.7e-3$ in T while the cooling differs by factors 0.02 to 0.78, and each case's movement is ordered by $|\Delta \text{cool}|/\text{heat}$ (lower_profile 0.51, the H2 front advancing ten cells from 3.16 to 3.43 R_p; the WASP-121 b cases 0.17 to 0.30 with T +14 percent, rho -27 percent and the converged wasp_full_newton mass flux +5.1 percent at log Mdot 13.30 -> 13.33; the molecular gates $3e-3$; oxygen_chemistry $4e-5$). DECISION recorded there: refresh those seven once at the end of the series and leave the other nine. Reported out of scope: Hydro_ioniz_adv.txt column 10 (adv_mass_row) holds exact zeros against $1e-14$ round-off, so the harness's relative comparison reports 1.0 on a physically identical file and needs an absolute floor there.
- L4c: no molecular case has a route, and a seed is the only opening** (diagnosis, NEGATIVE). The cold march stops at step 1200 to 1400 with a persistent non-root chemical equilibrium in ghost cell -1 (element violation $2.4e-4$ against $1e-6$ after 1000 consecutive sweeps, crossing the tolerance at step 984 and drifting up monotonically); moving q_H2_base away from its ceiling refutes the ceiling margin as the cause. The He-poor case stops at step 980 when four cells at $T = 7e4$ K have no energy bracket below the $5e4$ K ceiling of the H2 rovibrational table. Snapshots taken below both stops (du 9.6) put on the partitioned route end at outer pass 1 to 4 with hydro info = 2 (mass 1.0 to 1.4, energy 1.5 to 2.0 at 2.5 to 7.8 R_p), the same as an atomic cold-march snapshot; the atomic cases escape only through an archived solved seed, which no molecular group has, and the restart contract rightly refuses an atomic seed for a molecular run (mol and carrier decide the unknown count, K44).
- L7: the molecular seed from a certified atomic solution** (pilot, user-approved 2026-09-13 17:20; new src/modules/init/molecular_seed_from_atomic_state.f90, 573 lines, mode EXHALE_MOLECULAR_SEED=<atomic output dir> with ..._INVARIANT=p|T and ..._X2=<0..1>|local; load_IC.f90 lets the six molecular tokens differ only in a seed conversion; write_output.f90; docs/input_schema.md appendix D.3; new suite src/tests/molecular_seed). The seed exists and is exact (conversion identity $2.6e-15$, EOS closure $4e-16$, budgets to round-off), the loader accepts it, and on the certified He/H 1.60 and 2.13 states the chemistry has a root in every cell at every sweep -- the ghost reservoir failure of L4c is NOT reproduced from a stationary base -- while the hydrodynamics certifies with the local or the zero partition (the handoff x2 = 0.99998 partition gives no descent). What refuses is the H2 CARRIER

BALANCE row, exactly 1.000 at $r = 2.7 R_p$, where H_2 is $5.6e-27 \text{ cm}^{-3}$ against a gas of $1e6$: the carrier row is a relative measure with no absolute floor, where the element rows have $1e-20 \text{ rho } X_{\text{base}}$, so a physically absent species gated the solution. Two blockers were reproduced and diagnosed here: (a) the caloric EOS is not in fact limited to $5e4 \text{ K}$ (it continues the H_2 table linearly with a consistent (u, c_v) pair), $5e4 \text{ K}$ being only the energy BRACKET selected by the boolean $n(H_2) > 0$ where the guarded term is $1.9e-4$ of the internal energy; (b) the ghost reservoir pins $2 n(H_2)/n_H = 0.99998$ of the TOTAL hydrogen, infeasible once the march has ionized the ghost (the ionization balance asks $3.2e-4$ for H^+ , 19 times the room left), so the boundary may prescribe the partition of the NON-IONIZED hydrogen only, and nothing checked feasibility.

- L7b: all three are defects and are fixed** (diffusive_photochemistry.f90 +58/-9, certification.f90 +75/-13, energy_semi_implicit.f90 +26/-19, ionization_equilibrium.f90 +71/-2, EXHALE_main.f90). (1) A carrier below $1e-20$ of its element's free reservoir in a cell is reported and does not gate -- the element rows' own rule, now a named constant: the gate moves from a cell with $n(H_2)$ exactly zero to cell 218, where H_2 is real at $2.5e-6$ of n_H , and still refuses at 0.995, so an absent species no longer blocks but the molecular solution is not yet certified; carrier_model_a_newton rows identical before and after. (The x_{H_2} pin of ionization_equilibrium under transport is design, not a defect.) (2) One energy bracket ceiling for every cell, with the validity statement at the code site: the He-poor cold march that stopped at step 983 at $7.07e4 \text{ K}$ now runs 1200 steps with a maximum of $3.94e4 \text{ K}$, and a probe with only this change reverted stops again at step 982. (3) The base ghost's H_2 row is imposed as $x_2 (1 - x_{\text{ion}})$, the partition of the non-ionized hydrogen, with an infeasible prescription refused by cell, demand and room: the march that stopped at step 1240 with $x(1) + x(4) = 1.000307$ runs 3000 steps with $x(4)/(1 - x_{\text{ion}}) = x_2$ to eight digits. Movement on the seven mol_* regression cases, single thread, control against measured with only these five files differing: $3e-6$ to $2.7e-4$, all within $1e-3$.
- L7c: the refusing H_2 carrier row was catastrophic cancellation in the equilibrium fit, not a statement about H_2 .** $q_{h_2_equilibrium}(\text{lower_column.f90})$ evaluated $(A + t - \sqrt{t(2A + t)})/B$, whose two terms both grow like t while the difference falls like $A^2/(2Bt)$: above $t = A/\sqrt{2\epsilon} = 9.4e7$ the result is the rounding of t itself. On the LHS column 186 of 500 cells were off by more than 1 percent, 166 received exactly 0 where the value is positive, and the worst was $1.48e4$ times the true value; the local seed carried $x_2 = 5.04436e-6$ in four cells to twelve digits and 0 in their neighbours where the fit falls smoothly from $5.8e-10$ to $3.0e-10$. At the refusing cell 218 ($r = 1.204 R_p$, $T = 4940 \text{ K}$) the state held 3504 cm^{-3} of H_2 against an equilibrium $3.6e-2$ ($9.9e4x$), the dissociation time ($8.0e3 \text{ s}$) is below the crossing time ($1.8e4 \text{ s}$), three-body formation is five decades below, and the carrier row cannot relax a mixing-ratio spike (x_2 moved $3.9e-9$ relative in four passes). The tolerance is not the issue (round-off floor $1.4e-14$ against $1e-5$). The item also measured the ramp's other end, delivered as L7d(2) above, and found that the molecular input.inp files still carried the cold-march keys, so the molecular patch's wind pass would have discarded the seed.
- L7d: the three corrections L7c names.** (1) $q_{h_2_equilibrium}$ is written in the conjugate form with the square root as $\sqrt{t} \sqrt{2A + t}$ and the $u > 30$ branch removed: 186 of 500 cells off by more than 1 percent and 164 exact zeros before, none after (worst $2.1e-15$ against a 60-digit reference), and the local seed now carries

the fit's own smooth value. The same fit was duplicated in `src/utls/run_lower.py` with BOTH asymptotic guards inverted, and is fixed there too. Movement on the `mol_*` and `hp_*` cases: bitwise on four, below $1e-3$ on five, and `mol_sec_ion` $5.8e-3$ and TOWARD its golden. (2) the ramp rule, above. (3) Nine molecular cases regenerated with the seed keys, the runner's molecular branch in place (`make_models.py`, `run_case.sh`, `write_reproduce.py`; SEED_X2 default local, and SEED_GIVEN had been read before ATTEMPT existed, fixed), with 14 atomic campaign cases run through the swapped runner, all certified. Test T-L7-5: the hydro rows descend on this route for the first time (pass 2 `info` = 0, mass $7e-7$, energy $6e-6$; the continuation $1.2e-8$), and the block is now the PHYSICAL H2 carrier balance, 1.000 at cell 280 ($r = 1.60 R_p$, $T = 5986$ K) with H2 within 1.6 to 2.0x of the fit, the handoff partition returning its entry state. That is item L7e, approved by the user on 2026-09-14 15:05 as the first item after the campaign wrap-up and closed on 2026-09-15 (below).

- **L8: the outer boundary of a subsonic wind** (`docs/figures/lhs1140b_L8_outer_boundary.pdf`; `map_state_to_grid.py --extrapolate-beyond`). The certified 30 R_p solution is the subsonic part of a TRANSONIC escape whose critical point lies at 40.06 R_p , just outside the domain, and the isothermal hydrostatic ghost stands in for it, stating "a static isothermal atmosphere at cell N's temperature outside" ($p_{\text{inf}} = 0.134 p_N = 2.0e-10$ dyn cm⁻²; at the face Mach 0.52, so $v - c$ points inward and the ghost rule is an external condition, not an extrapolation). Solved again on 45 and 60 R_p domains, both certified, the critical point is inside (40.05, 40.08) and the two agree with each other to better than 0.1 percent everywhere: log10 Mdot 7.8737, 7.8740, 7.8735 at 30, 45 and 60 R_p ($5e-4$ dex), the He 2³S column within 0.23 percent, the equivalent width within 1.7 percent (1.1702, 1.1504, 1.1537 percent Å) and the FWHM not at all. The 30 R_p PROFILE is contaminated outside 10 R_p (T +25 percent, ρ +29 percent, v -22 percent at 28 R_p ; 1.9 percent at 8 R_p , 0.14 percent at 2). Conditions carried by every campaign number: Mdot is quotable as the flux the interior sets, the He 10830 numbers carry a 1.7 percent boundary systematic, nothing above 10 R_p is quoted from a 30 R_p run, and above 7.65 R_p -- the exobase of the extended solutions sitting at 29 R_p , below the critical point -- no radius is a validated continuum result. Found on the way: the `_adv` profiles diverge from the solution in the outer wind (429 K against 1513 K at 29 R_p), which is item L10.

- **L10: the `_adv` profiles in the outer wind, and one real defect in the upwind recursion** (`post_process_adv.f90` +56/-3, `src/utls/collisional_validity.py` +52/-10, `docs/collisional_validity.md`). The departing `_adv` temperature is NOT a post-processing defect: above 4 R_p the recombination and photoionization Damkoehler numbers fall below $1e-2$, the composition is frozen and advected, and the advection correction -- which satisfies its own stationary internal-energy equation to $1e-6$ -- then carries up to 5.9x the photoheating of the equilibrium composition, with T following it (1.45x at 10 R_p , 3.5x at 29). What that says is that the certified solution's photoionization-EQUILIBRIUM composition is wrong beyond 4 R_p . One real defect found and fixed: MINPACK `info` = 4 iterates whose residual was $1.5e-17$ to $5.2e-17$ of the row's largest term were discarded in three cells and the upwind recursion restarted above 32.6 R_p , which is where the 45 and 60 R_p `_adv` profiles disagreed; such an iterate is now adopted as the root (45 against 60 R_p : 0.751 -> 0.0019), with `wasp_full` (metals on, Brent path) bit-identical. `collisional_validity.py` now reads the SOLUTION by default, the critical point being a property of the

momentum-balanced state (32.5 R_p from `_adv`, 40.06 from the solution). THE LINE: the He I 10830 equivalent width is 1.1702 percent A with the `_adv` composition against 15.5 with the solution's composition (temperature is 0.15 percent of the EW, composition a factor 13), so the He 2³S column and the crossing He/H stand on the ionization closure and not on the hydrodynamics -- which is what opened item L12. Reported out of scope: the same info $\neq 1$ discard in the advected-ionization and metal re-solve branches, and the Brent window $[0.05, 4] \times T_{in}$ and the metals-on MINPACK band $[0.5, 2] \times T_{in}$, which above 16 to 33 R_p remove the correction altogether.

- L11: seeding a composition ladder from the certified solutions** (`map_state_to_grid.py --reservoir He/H, pick_seed.py, run_case.sh, write_reproduce.py, status.py`). The loader refuses a certified state whose reservoir He/H is not the run's (metadata rule, $1e-6$), so the mapper now rescales the helium species by new/old over the whole column -- the diffusion profile's shape kept and the base exactly at the new value, as `load_IC`'s own `He_diffusion` rule does -- rewrites `rho` and `T` so the file is self-consistent ($T = p/((n_{tot} + n_e) k_B)$, `rho` the species mass sum; 395 $\rightarrow$ 328 K at the first ghost for 1.60 $\rightarrow$ 2.13), rewrites the reservoir line and records the rescale in `# mapped:` and in `REPRODUCE.md`. MEASURED: `atomic_scalar_gj1132_kzz1e9/HeH2.13` seeded from the certified `HeH1.60` certifies in 4 passes and 13.6 min at `log Mdot 7.88` and EW 1.3934 percent A, where the archive seed succeeds at 1.60 and fails at 2.13 from the same loaded state (its base is the old sawtooth column at 226 K and $v = -643$ cm/s at cell 3, O(1) from the current root; whether the hydro solve climbs out is a basin question). `HeH4.0` from the certified 2.13 has pass 1 hydro info = 0, so the seed is in the basin, and stalls in the joint alternation, which was the L4d obstruction and not the seed. Addendum of 2026-09-14: `run_case.sh` continues a refused solve from its written state at `dtau0 = 1e8` (L4e) and, where the seed was `pick_seed`'s, walks down its ranked list (`SEED_ATTEMPTS`, default 3), keeping each failed attempt in `attempt_<n>/`; and `pick_seed.py` ranks every tier by ONE rule -- a composition step above 0.3 in $\log_{10}(\text{He}/\text{H})$ after every nearer candidate, whatever its tier -- and gained the tiers `tier2k` (a certified case of the same physics at another `K_zz`) and `tier4k` (the archive at another `K_zz`), because from the certified 3.5 state rescaled onto 0.55 (a factor 6.4) the element relaxation found no admissible advance while the certified `kzz1e10/HeH0.55` is the same composition one eddy coefficient away. --list N prints plain lines.
- L12a stage A: one routine imposes the transported ionization fractions.** The substitution that replaces an ionization balance row by the fraction the flow carries stood in two places (`System_HeH_mol`, `System_HeH_mol_metals`) and was absent from the other five systems that reach the same rows. It is now written once, as `impose_transport_ionization_fractions` in `ion_residual_core.f90`, and all seven systems (`System_H`, `System_HeH`, `System_HeH_TR`, `System_HeH_metals`, `System_HeH_TR_metals`, `System_HeH_mol`, `System_HeH_mol_metals`) call it once, last, after every term of their rows and after any scaling of them. The routine takes the cell state as an argument, so `ion_residual_core` stays free of state of its own; rows 1 to 3 are $x(\text{H II})$ per H nucleus and $x(\text{He II})$, $x(\text{He III})$ per He nucleus in every system that reaches them, and only row 1 is written, the cell state carrying no imposed helium fraction until stage D. The gate in `ionization_equilibrium.f90` that decides when a fraction is imposed is unchanged (stage C changes it), so this stage changes no behaviour.
- Byte identity, MEASURED.** A control binary of the tree before the change and a measured binary after it,

differing in these eight files and in nothing else (the source checksums of all 208 files were compared between the two builds). The four cases that carry Ionization transport: True or the carrier transport (hp_front, hp_zero_seed, hp_trace_seed, mol_carrier) give outputs identical byte for byte between the two binaries, in every output file, the run timestamp of the provenance header excepted. So is wasp_full, the atomic case with the key off, run unbounded to its own du stop: both binaries stop at count = 12251 with du = 9.7613E-04, their two 12518-line run logs differ in the execution time and nothing else, and all ten output files agree byte for byte. Against backup/regression/golden/ the four molecular cases are WITHIN 1e-3 and not bitwise, by 1e-7 to 5e-5 relative, and wasp_full EXCEEDS it at a velocity that changes sign at row 209 -- on the control binary exactly as on the measured one, to the last digit of every quoted value. Those goldens stand behind other changes in this working tree, not behind this one.

- **The new test suite**, src/tests/ionization_imposed_fractions, 31 assertions, all exact: the imposed row is $x(1) - x_{hp_fix}$ in each of the seven systems and in the routine itself; no other row moves; with nothing imposed the residual does not read the carried value at all; and the imposed row carries the same number in all seven. RED, MEASURED in a copy of src/: the substitution removed from the routine fails 9 assertions, the call removed from System_HeH_TR alone fails 2, which is the drift case the suite exists for. Suites re-run on the private build: carrier_retry 142/0, carrier_returned_state_acceptance 36/0, carrier_reference_scales 14/0, carrier_constraint_attribution 10/0, attempted_step 70/0, constrained_network_layout 8/0, certification 84/0, steady_species_rows 195/0 without the reload rows and 198/1 with them, the one failure standing on the control binary as well.
- **L13: the elemental-flux closure could not reload its own iterate, on two metadata fields.** Every closure rung stopped at iteration k01 with load_IC refusing the seed on the reservoir field, C/H off by 2.624e-5 against the 1e-6 threshold. The reason is physical: the photochemical column moves EVERY elemental ratio between iterations -- MEASURED on atomic_photochem_gj1132_kzzprofile/HeH9, k00 -> k01: He/H +1.020e-5, C/H +2.624e-5, N/H +2.624e-5, O/H -1.145e-4 -- while the seeding step carried helium alone. map_state_to_grid.py now takes --reservoir <El>/H <value> for any element of the # reservoir line, repeatable (one factor over that element's ionization stages, the mass at calc_rho's weights, the pressure held and T following the particle count, the field rewritten), replacing the helium-only option; and element_flux_closure.py (+217/-18) reads every $X_{<El>}$ of the profile by name (reservoir_ratios, profile_match_ratios) and passes the ones that moved in one call. A SECOND refusal on the same path, the grid field, was measured and closed the same day: with a lower-atmosphere profile in use, R0 is the profile's radius at the matching level times R_J, so it moves with the profile (k00 1.62166044189e-1 R_J, k01 1.62166043926e-1, 1.6e-9) while the cell centers in R_p do not, and the closure had handed the mapper the SEED's own file as the target grid. DECISION: the contract stays exact and the driver builds the target header. profile_values_at_match plus one multiply by RJ = 7.1492d9 reproduce input_read's R0 BITWISE (k00 -> 1.1593574831171966E+09, the string the k00 run wrote; k01 -> 1.1593574812390110E+09, the string the k01 run states; the match row's p equals p_match_bar, so value_at_pressure takes its exact-level branch and no interpolation enters), and write_target_grid_header writes <iter>/target_grid_Hydro_ioniz.txt

with only that field replaced, at meta_num's ES23.16. The mapper is called when the reservoir OR R0 moved. Reproduction: no field refused, the # grid line of the seed character- identical to the one the run writes, and the iteration solved at info = 0. Addendum of 2026-09-14 21:40: the same field refused the three atomic_photochem_gj1132x{0.15,0.10,0.01}_kzzprofile/HeH9.7 cases from the RUNNER, which had handed the mapper the shared current_grid_Hydro_ioniz.txt, whose R0 is the planet radius 1.1273716464000001E+09; the x0.33, x0.30 and x0.20 cases escaped only because their seeds are archived states with no metadata block to compare. The matching level, R0 and the metadata number format moved into the new src/utls/profile_match_level.py (168 lines), which the closure driver imports and run_case.sh runs, and a profile case now maps against its own target_grid_Hydro_ioniz.txt; write_reproduce.py records the file and the number. The three reloaded with no field refused (1.1592997812185061E+09, the string their refusal had named) and two then CERTIFIED (x0.15 at pass 9, log Mdot 7.04; x0.10 at pass 12, log Mdot 6.85).

- L14: the outer loop's progress control read a quantity that is not monotone while the composition is travelling** (src/EXHALE_main.f90 +165/-29, the progress control of steady_wind_with_element_diffusion). The five low-XUV cases the campaign of 2026-09-14 11:24 left uncertified were NOT refused by the cell-1 hydrodynamic energy row (that line is a note inside a JFNK solve which then returned info = 0 with mass 2.30E-08, momentum 3.39E-14 and energy 8.49E-07 of 1.0E-06) but by the GATED ELEMENTAL TRANSPORT He/H PARTITION row, 2.12E-02 of 1.0E-05 at $r = 2.50 R_p$, ended by the progress control itself: on the 0.10, 0.15 and 0.20 cases the element relaxation's own distance to its fixed point fell 6.62E-01 -> 4.24E-01 -> 3.39E-01 every pass while the gated row ROSE 1.28E-02 -> 2.12E-02, so the control halved omega, which takes the alternation's contraction factor from 2.00 to 1.14 per pass, and stopped at three strikes. The certified 0.30 case travelled the SAME distance (4.89E-01) and certified at pass 13 with omega at 0.500, the distance halving every pass. FIX: a pass counts as progress when the joint distance fell, OR when the state's distance from certification is carried by an entry other than a hydrodynamic row AND the composition's own distance fell; only when neither holds are the step lengths shortened and the consecutive-pass counter advanced, outer_no_fall_max = 3 being unchanged. The distance is read from the two most recent updates, so the first two passes of a solve carry no measurement and count as progress. TWO FORMS were written and the first was MEASURED to be wrong: an "every hydrodynamic row inside its tolerance" test halved omega twice on the first ladder rung (.L14/x005_HeH9.7 spent its whole budget at a gated row pinned at 2.64E-02 and an element distance creeping at 1.10 per pass, with the halvings at passes 2 and 10, whose hydrodynamic energy rows read 1.03E-06 and 1.32E-06 against a tolerance of 1.0E-06 while every other pass is inside it) -- it called a wind solved to three percent of its tolerance an unsolved wind. The delivered condition is which entry CARRIES the distance, hydro_worst_dist < prog_worst, which separates the two cases by five decades rather than by three percent (at dtau0 = 1 the energy row stands at a distance of 6.3E+05 against the elemental row's 5.8E+02, and in the continuation 1.03 against 1.9E+03). run_case.sh also gained EXHALE_OUTER_PASSES, exported with a default of 40 against the binary's own 20 (the certified 0.30 case had already spent 13 from a nearer entry and the 0.10 case of this item was accepted at pass 30 exactly); EXHALE_PTC_DTAU0 became a DEFAULT of 1.0 rather than a fixed assignment, so a caller may state the pseudo-time start of the FIRST solve, with

nothing about a run that does not state it changed; and the walk down `pick_seed.py`'s ranked list is skipped where the refused solve left every hydrodynamic row inside its tolerance, such a state being a stationary wind the composition refuses, for which a seed one further XUV step away can only make the wind worse (measured: those attempts spent their passes at $\lambda_{\text{m}} 1\text{e-}6$ and $\text{dtau} 1\text{e-}4$ without moving the residual, which is item L4h). RESULT: `atomic_scalar_gj1132x0.{20,15,10}_kzz1e9/HeH2.13` CERTIFIED at outer pass 19, 24 and 30, $|R|$ 1.228E-07, 7.112E-07 and 4.842E-07, log Mdot 7.14, 7.01 and 6.83, with zero omega halvings, and the $\text{dtau0} = 1$ stage of each reproduces the refused run of 11:24 to every printed digit. New spectra `LHS1140b/sed/lhs1140_sed_gj1132_at_b_xuv0p{07,05,03,02,015}.txt`, the flux column of the reference SED times the factor, written the way the existing grid was (regenerating `..._xuv0p10.txt` by that rule reproduces the file on the tree in all 29 992 rows). Regression cases that can reach this code: `mol_diffusion` PASSES within $1\text{e-}3$ (1.272e-07, 1.814e-06, 2.537e-05, 3.083e-06 on the four files); `lower_profile` FAILS IDENTICALLY ON THE CONTROL BINARY (du 3.1666E+00 at step 12000 and max rel 1.269e+00 at row 474 col 3 in both), so it is a golden the tree has moved past, reported and not acted on. THE 0.02 WALL, 2026-09-15: the three 0.01-XUV cases are reached by a ladder in `LHS1140b/models/.L14/` at 0.05, 0.03, 0.02 and 0.015, each rung solved at `EXHALE_PTC_DTAU0=1.0e8` because the $\text{dtau0} = 1$ solve of these columns falls into L4h. $\text{He/H} = 2.13$ certified at 0.05 (pass 20) and 0.03 (pass 19); $\text{He/H} = 9.7$ at 0.05 (pass 29) and 0.03 (pass 24); the lower-atmosphere-profile case at 0.05 (pass 29) and 0.03 (pass 25). ALL THREE chains then refuse at 0.02 in the same shape -- hydro info = 2 at every outer pass with the worst row within $0.005 R_p$ of the base and $\lambda_{\text{m}}/\text{dtau}$ walking down from a $1\text{e}8$ start -- independently of the helium abundance and of the lower-atmosphere treatment, and at a step (a factor 1.5) gentler than the $0.05 \rightarrow 0.03$ that certified on every chain. On the one rung that ran its full 40-pass budget the gated elemental row fell $2.95\text{E-}02 \rightarrow 4.58\text{E-}08$, four decades INSIDE its tolerance, with omega at 0.500 and no halving, so the composition is not what refuses these states and the block is in the wind half. The ladders reach 0.03; the three 0.01 cases are NOT CERTIFIED and are handed to item L4h.

- **L13 sections 12 and 13: the runner formed two restart-contract fields itself, and both now live in one module.** The grid refusal of the three XUV-scaled profile cases was not "the seed was a scalar case": `run_case.sh` handed `map_state_to_grid.py` the shared `models/current_grid_Hydro_ioniz.txt` as the target for EVERY case, and the mapper copies its target's # grid line, so every profile case whose seed carries a metadata block was refused whatever its seed ($x0.33$, $x0.30$, $x0.20$ escaped only because their seeds are archived states with no block to compare). The reservoir refusal that followed on 2026-09-16 00:42 is the same shape: `load_IC::compare_reservoir_field` walks the line ELEMENT BY ELEMENT at $\text{heh_dev_tol} = 1\text{e-}6$, while `run_case.sh` carried one element, He/H , read from `input.inp` - which is the whole composition only where the case states it there, a case with Lower atmosphere profile: taking every ratio from the profile's $X_{<\text{El}>}$ column at the matching level. The six XUV-scaled photochemical cases read their column from the flux-closed $\text{He/H} = 9.7$ rung, which the re-run had just re-solved, so C/H moved by $3.7\text{e-}6$ and all six were refused at load with their helium correctly carried. Both rules now live once in `src/utils/profile_match_level.py` -- `profile_match_ratios` (what the run will ask for), `reservoir_ratios` (what a state carries), `reservoir_moves` (what differs by more than the contract's own $1\text{e-}6$) and the command `--reservoir-options`, which prints them as the `--reservoir`

<El>/H <value> arguments the mapper takes -- with `run_case.sh` calling the command for a profile case and passing every option it prints, and `element_flux_closure.py`, which had grown its own copy of the same rule, importing it and computing nothing of its own. Verified: the command on the pre-L21 state of `atomic_photochem_gj1132x0.10_kzzprofile/HeH9.7` against its new column prints four options (C/H, He/H, N/H, O/H), the mapped seed then carries all four at the profile's values, and all six cases load and enter the solver; the nine closure rungs are unaffected, the routine being the one they were already using, moved. The grid refactor is behaviour-preserving: rebuilding the L13 reproduction seed through the moved helpers gives data rows byte-identical to the pair the earlier run loaded, the only difference being the target path the # mapped: provenance line records.

- L14 final form: the progress control reads a distance and not a displacement, and three questions no longer share one cell index** (items R2 and R3 of the review of 2026-09-15; `EXHALE_main.f90`). R2: the carrier half of `comp_drift` was `carrier_drift`, the change the relaxation KEPT, which is bounded by `trust_pass` -- the same quantity the outer loop shortens whenever a pass does not help. The loop shortened the bound, the displacement fell because the bound had fallen, and the control read "the composition is closing" precisely where it had stopped letting the composition move. The carrier half is now the carrier's own fixed-point residual (`carrier_steady_residual`, taken on the state the pass hands back in the isolated workspace the certification uses, so the frozen background, the photolysis rates and the row terms are the ones the next step would read); the element half was already a distance (`relax_element_composition` returns an undamped one, `comp_omega` damping what is applied and not what is measured). `EXHALE_CARRIER_DRIFT_IS_DISPLACEMENT=1` restores the old measure. MEASURED: the L14 fixture is atomic and the block is gated off there; on `carrier_elem_newton` (uncoupled) the two logs differ in THREE LINES and nothing else, three passes on which the old measure declared the composition not to be closing while the carrier's own residual was falling, so three false stagnation counts disappear and every number and the ending are unchanged; on `carrier_model_a_newton` the ending is the same (the budget of 20, two active equations refusing) and the iterate is not -- the old measure halves the bound at pass 7 and ends at `trust 2.5e-03` with a smaller worst carrier row (`5.75e-04` at cell 500) than the new one (`5.21e-03` at cell 248), which is not an argument for it: a carrier held still by its own bound perturbs its balance row less, and that circularity is what the correction removes. R3: the worst carrier-drift cell (`carrier_drift_location`), the cell holding the particle-count bound (`bound_last_j`) and the worst-residual cell (`carrier_steady_residual`) are logged apart every pass and are three DIFFERENT cells -- 333, 327 and 373 at pass 7 of `carrier_model_a_newton`, with the row that refuses the certification naming a fourth.
- L4h: the pseudo-time guard, delivered, with its published basis checked and found absent** (`steady_newton.f90 +186/-1`; `EXHALE_PTC_RAMP_GUARD` default ON, `EXHALE_PTC_RAMP_GROWTH_GATE` default OFF). DIAGNOSIS: the pin is the step-size rule and not the direction. At the floor the Krylov cycle reaches its tolerance in ONE product, because the shift $I/d\tau$ stands above every entry of the Jacobian, so the step is $dY = -d\tau F$, an explicit Euler step at the CFL interval, and a column that needs $O(1e5)$ of them cannot move in 200 iterations. The pseudo-time reaches that floor through ACCEPTED steps alone: $d\tau *= \lambda_m$ with λ_m unbounded below (`3.91e-03` measured) against a growth clipped at 2, eighteen decades in 41 iterations of pass

1 while the merit falls 47 percent; outer pass 39 spends 60 of its 69 iterations walking dtau from $1\text{e}+08$ to $7.93\text{e}-05$ and hands back the state it was given. (The $1\text{e}+14$ climb of the 0.01-XUV pair is a log of a binary that PREDATES the L7d rule.) RULES, default on: (1) the cut of an accepted step is clipped at ten, $\max(\text{lam}, 0.1)$; (2) it stops at $1\text{e}-3$ of $\text{dtau}_{\text{earned}}$, the largest pseudo-time at which the solve has taken a WHOLE step (dtau_0 alone does not serve: on the stationary restart it is 1.0 and the floor is then no protection, MEASURED), the explicit-stable interval staying the floor of the REJECTION cut alone; (4) eight consecutive damped steps return the solve to its best iterate with $\text{dtau}/10$, and two such returns with no full step between them stop it at its best iterate with $\text{info} = 2$. **Rule (3), a residual-progress growth gate, is OFF, and that is a measured negative result:** no such test separates the two regimes -- on the LHS 1140 b C/N/O near state the merit hovers at $1.28\text{e}-01$ to $1.42\text{e}-01$ against an entry $1.44\text{e}-01$ while $||R||$ falls $6.98\text{e}-01 \rightarrow 2.80\text{e}-01$, so the SER ratio leaves dtau at 1.13 and the 10-percent stagnation-window gate at 2, and the gated run finds no admissible step from iteration 6 and is at 1.959 by 14, worse than the state it was given. On the element reload with the delivered earned anchor the ratchet stops where the rule says: the guarded run holds dtau at $2.05 = 1\text{e}-3 \times$ the $2.05\text{e}+03$ of its last whole step while the unguarded one falls to the explicit floor $9.68\text{e}-05$; at iteration 27 the two stand at $||R||$ $4.035\text{e}-02$ / merit $9.71\text{e}-04$ against $1.543\text{e}+00$ / $1.11\text{e}-02$, and the four statements compose visibly (the floor holds dtau through six damped steps, the eighth fires the return to the best iterate, the full step after it takes $||R||$ from 1.62 to $3.899\text{e}-02$ in ONE iteration, a second return follows and the third run of eight stops the solve at its best iterate at iteration 62), ending pass 1 at $||R||$ $4.032\text{e}-02$ against the guard-off $8.336\text{e}-01$ at the 80-iteration cap, twenty times better on $||R||$ and the energy row in three quarters of the iterations. Controls: guard off is bitwise the entry text; the C/N/O near state reproduces iteration for iteration for 14 iterations; krylov_and_dogleg 334/0, attempted_step 70/0, certification 84/0, steady_species_rows 199/0 with the reload rows. **THE LITERATURE DOES NOT CARRY THESE RULES, and the item says so** (memo section 4.2, from the primary sources): Kelley & Keyes' "switched evolution relaxation" and Gropp et al.'s "successive evolution-relaxation" are the same rule from the same source, Mulder & van Leer (1985), whose own text carries NEITHER name; and what that paper prescribes is a target on the CHANGE OF THE STATE, $\text{dt}^n = \text{eps}/\text{RES}^n$, "Equation (4) guarantees that in the explicit case ($\alpha = 0$) the relative change of the state quantities per time-step will nowhere exceed epsilon", the successive-ratio form being a reading of it between two steps. Its own under-relaxation is applied to the OPERATOR ($\alpha \geq 1.2$ in $[I/\text{dt}^n + \alpha M^n] \Delta_t W = G^n$) and leaves the timestep rule untouched; their algorithm takes the FULL step and has no line search anywhere in it, so lam has no counterpart in any of the three papers and neither does a cut keyed on it. Only the CLIP ON THE GROWTH is supported explicitly (Coffey, Kelley & Keyes' phi caps and nothing else, and their section 3 reports limiting the increase "to a factor of 2"), and the explicit-stable interval as a floor has no counterpart either (in Kelley & Keyes the CFL enters as the scaling matrix V). **Statements (1), (2) and (4) are OURS and rest on the measurements of the memo's section 5 and on nothing in the literature**, which the code site now says. One thing the papers do bear on and this item met independently: Coffey et al. report that "the success of the method is relatively sensitive to the initial time step", which is the dtau_0 sensitivity L4g and L7d record. The guard does NOT solve the 0.02 rung: with it x002_HeH2.13 ends pass 1 at $||R||$ $2.081\text{e}-03$ against the control's $6.455\text{e}-04$, a factor 3.2 worse, the control buying that by letting dtau fall to $6.78\text{e}+01$ which the guard forbids, while the

guarded pseudo-time OSCILLATES on its floor ($1.00\text{e}+05$ to $4.00\text{e}+05$ from iteration 173) with $||R||$ drifting $6.7\text{e}-03 \rightarrow 2.7\text{e}-03$: an eighteen-decade collapse turned into a bounded oscillation, which is the change in kind it was written for. That rung is item L17.

- L15: the JFNK solve is not reproducible at eight OpenMP threads, and the residual evaluation is not where that enters** (OPEN; docs/lhs1140b_stationary_L15_20260916.md). Two runs of the same fixture (backup/regression/atomic_elem_newton_reload) at eight threads part after about 21 JFNK iterations and end pass 1 at mass rows $3.08\text{e}-02$ and $1.20\text{e}-02$; one thread is bitwise reproducible. FIRST STEP, MEASURED 2026-09-16 on the certified fiducial handed back to the binary as its own IC pair with Restart intent: stationary evaluate (ten runs, five at OMP_NUM_THREADS=1 and five at 8, each in its own directory): the data lines of Hydro_ioniz.txt have md5 8f15a045baa3 and those of Ion_species.txt 8a427fb6e00f in ALL TEN, the mass row $2.175\text{E}-09$ at cell 1, the momentum row $5.417\text{E}-13$ at cell 500, the energy row $2.126\text{E}-08$ at cell 1 and $||R||$ $2.1257\text{E}-08$ in all ten, and the whole run.log with timing and thread lines removed has one md5 within each thread count. Since Ion_species.txt carries the composition and Hydro_ioniz.txt the heating and the cooling, the ionization sweep, the radiative rates and the hydrodynamic row assembly are all inside that comparison. The ONE difference between the thread counts is the runtime's exit-time floating-point exception summary (IEEE_UNDERFLOW_FLAG added at eight), which is the union of the flags raised on the threads that ran and names no array. So the dependence enters at the Krylov stage or later -- the matrix-vector products, the norms and inner products, the banded preconditioner solve, or the line search. **This is explicitly NOT called harmless rounding**: a difference that appears only at eight threads is a race until proved otherwise, a reduction order is one explanation and a scratch array written by more than one thread is another, and the two are told apart by reading the parallel regions and not by the size of the difference. Next step, named in the memo: enumerate the parallel regions of the JFNK path and, for each, the arrays written, the variables carried across iterations and the reductions formed, and attribute only then. Every control/measured comparison of this series was taken single-threaded or on the same schedule; the campaign cases ran at eight threads and are certified states of the EQUATIONS, not of a path.
- L16: a recorded pass-1 triple that the current source does not reproduce** (OPEN, READ 2026-09-15 from the L4h memo, not chased). docs/lhs1140b_stationary_L4g_20260914.md section 7.3 records $2.16\text{e}-09 / 4.24\text{e}-13 / 1.55\text{e}-08$ on a fixture where the L4h control build gives $3.08\text{e}-02 / 1.90\text{e}-05 / 8.34\text{e}-01$. The item is to rerun that fixture on the L4g-era build (its md5 is in that memo) and on the current one, single-threaded, and say which change moved it; L15 is a candidate, the L4g numbers having been taken at eight threads.
- L17: the 0.02-XUV hydrodynamic solve, with the ramp excluded and the discretization named** (closed 2026-09-15 by the user, DIAGNOSED AND NOT SOLVED; docs/lhs1140b_stationary_L17_20260915.md, steady_newton.f90 only). With the L4h guard the pseudo-time makes the march the continuation is for and STAYS there taking full steps ($\times 002_{\text{HeH}2}$. 13 at iteration 175: $\text{lam } 1.00$, $\text{dtau } 1.05\text{e}+02$ and doubling, the Krylov cycle reaching its forcing term in 3 products of 40), so the ramp is not the cause; what refuses is the WIND half, every outer pass returning hydro info = 2 with the hydrodynamic rows flat while the gated elemental row falls ($\text{p}002_{\text{HeH}9}$. 7 passes 1 to 8 give mass $5.08\text{e}-02$ and momentum $2.86\text{e}-09$ IDENTICAL to every digit -- the

solve hands back its input -- while its elemental row falls $2.94\text{e-}02 \rightarrow 1.78\text{e-}02$). **The verdict: it is the Newton model of a wind that is subsonic to Mach $1\text{e-}8$ over its whole lower atmosphere, and not the anchor and not the finite-difference action.** The anchor is generous, not tight: at the binding cell the continuity row stands at $1.6189\text{e-}04$ of its own flux against a tolerance of $3.0\text{e-}07$, and the row's MEASURED response to one ulp of every cell density there is $7.65\text{e-}14$, so the row is $5.3\text{e+}02$ times its tolerance and $2.1\text{e+}09$ ulps of its own arithmetic away from where the certification asks it to be; and the base velocity at 0.02 is that of the CERTIFIED 0.03 state to three digits ($-1.28048\text{e-}01$ against $-1.27875\text{e-}01$ cm/s, Mach $8.595\text{e-}07$ against $8.589\text{e-}07$), so nothing collapsed between the rungs. The finite-difference action is rounding-limited at the arc the Krylov cycle uses (additivity defect $1.510\text{e-}02$ scaling as the arc to the power -0.976) and multiplying the probe arc by a thousand gives twelve times a better operator and the same stall in the same place. What fails is the linear model and the step built on it: the Krylov cycle reaches its forcing term and leaves the binding rows at 1.7 and 21 times their entry values (3.0 and 24 with a 200-vector subspace), and the scaled trust region, made available on this route by this item, predicts a three percent fall of the squared merit at its first radius, measures it nearly DOUBLING (ratio -26.8), cuts the radius twelve times to $2.3\text{e-}12$ and accepts no step at all. **The discretization is named, and the obvious repair was implemented and REVERTED.** The HLLC flux selects its branch on the SIGN of the contact speed and $S_{\text{star}} \sim v$ sits at $1\text{e-}09$ to $1\text{e-}02$ of the wave span over this whole domain, changing sign from one base cell to the next; the mass component of the star flux is $\rho * S_{\text{star}}$ with $\rho *_{\text{L}}$ and $\rho *_{\text{R}}$ differing by the face density jump, 4 percent per cell at this base, so the flux is continuous at $S_{\text{star}} = 0$ and its derivative is not and every base cell sits on that kink (nine tenths of the second difference of the residual along a probe direction in cells 1 and 2, ratios 1.06 and 1.33 where a smooth residual gives 4). A C1 blend of the two star fluxes over $|S_{\text{star}}| < b$ ($S_{\text{R}} - S_{\text{L}}$) at $b = 1\text{e-}3$ covers 409 of 500 cells out to $6.74 R_p$ and moves the residual of the same state from ($1.619\text{e-}04$, $8.045\text{e-}12$, $5.216\text{e-}04$) to ($3.309\text{e-}03$, $1.284\text{e-}07$, $1.370\text{e-}03$) -- not a regularization of this discretization but a different one, and narrowing b does not help because inside the band the perturbation is about half the face density jump INDEPENDENTLY of b . Five step controls at twenty outer iterations each leave the state where it was. Delivered anyway, default off and byte-identical off: the scaled trust region the hydrodynamic half of the partitioned route did not have, because it is what produced the measurement and not because it repairs the case. The three 0.01-XUV catalog cases stay recorded as NOT SOLVED; a smooth low-Mach flux would be a discretization change and is not opened.

- **L18: a certified state did not reproduce its certification when re-read, and the difference was in the mass density** (CLOSED, POSITIVE; load_IC.f90). A state file pair states the density twice -- the ρ column, and the species columns that weigh $\sum_i m_i n_i$ -- and the two agree only while the composition closes its own mass, $\sum_i f_i A_i = 1$. It does not: the closure of a state written at the end of a stationary solve drifts about $1\text{e-}14$ per outer pass, MONOTONICALLY, reaching $5.1\text{e-}13$ on .L14/x003_HeH2.13 after 19 passes and $6.2\text{e-}13$ on x003_HeH9.7 after 24, where a production state of models/ closes to $1\text{e-}15$. The loader took the species as the authority and rebuilt the density from them, so the evaluated state stood some 2000 ulp from the certified one -- and at a base whose energy row cancels its largest term by $6.2\text{e+}05$ that is the whole row. **FIXED:** the conserved density is read from its own column and the loaded species are projected onto it, one factor per cell, which leaves every element ratio and ionization split untouched, and the departure is reported on every restart.

.L14/x003_HeH2.13 now re-reads at mass 3.764e-08 (cell 7), momentum 1.814e-14 (cell 1), energy 8.255e-07 (cell 7) -- the rows it was written at -- and exits 0 where it exited 2; **all 120 certified states of LHS1140b/models/re-certify (exit 0)**, five of the seven ladder states reproduce their in-run row to every printed digit and two of the three the control refused now certify. NOT reproducible bitwise, and it cannot be: one ulp of the density moves the cellwise maximum of the energy row by 11 percent, so the mass row comes back within 0.78 to 1.71 of its in-run value (median 1.12) and the energy row within 0.89 to 1.93 (median 1.22), and the restart contract now SAYS so with those numbers (restart_contract_design_20260909.md section 6, write_reproduce.py); .L14/x003_HeH9.7, certified at 0.87 of its tolerance, re-reads at 1.18 and stays refused on that ground. Codex review applied: which half of the file states the density is decided by the loader's own INTENT -- each block that changes the composition says so -- and no longer by the SIZE of the disagreement, a restart of the same equations being allowed only rounding (1e-10 reported, above 1e-8 REFUSED as a damaged or mismatched pair). Opened by it: L19 and L20.

- **L19: nothing measured the mass closure of a state, and the sweep was ratcheting it** (CLOSED, POSITIVE; ionization_equilibrium.f90, certification.f90). The mass fractions are $f_i = n_i m_i / \rho$, so the sum is one by definition; the sweep holds the density fixed and returns a composition that weighs it only to the arithmetic of the cell solves, leaving a few units in the last place per sweep AND IN THE SAME DIRECTION, which is what ratcheted to 6.2e-13 over twenty-four outer passes. FIXED: the composition a sweep returns is projected onto the density it was given, one factor per cell, immediately after the sweep and before the heating and cooling are contracted with it -- every element ratio, ionization split and metal-to-hydrogen ratio survives a common factor, and it is the same rule load_IC applies to a restart file (L18), so a state written, read back and re-measured passes through one projection and not two. Default ON, EXHALE_MASS_PROJECTION=0 restores the old composition. REPORTED and gating nothing: certification_evaluate measures $\max_j |\sum_i f_i A_i - 1|$ from the same (u, f_sp) every row is measured on and prints it with its worst cell, its natural scale being nine decades below every tolerance in the inventory. MEASURED on the state a run writes: 4.151e-14 -> 4.196e-16 (hp_front, 100 steps) and 1.537e-14 -> 3.623e-16 (hydrostatic_column, 300 steps), the drift replaced by the rounding of the projection. Codex review applied: the projection is bounded and reports itself -- silent below 1e-12, reported to 1e-8, and above that NOT applied and the sweep reported as having failed its closure -- with finite and positive guards and the absolute element-total movement named.
- **L20: the cool column of a marching-written state, NEGATIVE as posed** (src/tests/grid_and_gates/output_state_consistency.sh). The 0.35-0.76 cooling ratio L18 reported was measured on backup/regression/wasp_full_newton/IC/, a state PINNED on 2026-09-08 from an earlier code generation and not anything the current code writes. MEASURED on every state the current code produces, the cool column and the total of Cooling_breakdown.txt -- which recomputes every channel from the written (T, rho, f_sp) through eval_cool -- agree to between 3.4e-15 and 3.0e-10 over the golden matrix (wasp_full_newton 2.98e-10, wasp_full 3.4e-15, the four short cases 2.1e-14 to 2.9e-14), and the heat column agrees with Heating_breakdown.txt to 2.4e-16; the residual is the documented one-sweep lag of the temperature. Rebuilding the pair at the final state was written, measured and REVERTED:

it takes one more sweep, moves every golden, and leaves the same lag in the other direction (hp_front 2.9e-14 -> 6.5e-12). WHAT WAS WRONG is that the invariant was asserted for heat and never for cool, which is how a pinned state 1.8x out could sit in the suite unnoticed: output_state_consistency.sh now carries cool_column_matches_breakdown_total at the same 1e-6, and the stationary evaluate route writes both breakdown files so the same statement can be taken on a re-measured state. For the user: backup/regression/wasp_full_newton/IC/ is stale, its own IC/README.md says it is refreshed only together with the golden, and backup/ is read-only for this work, so the refresh was not made.

- L21: the base boundary decided its entropy branch on a quantity that is not a flux, and its window sat inside this planet's operating point** (src/modules/states/base_boundary.f90). TWO defects, only one of them the window. (1) THE SIGN: the branch was keyed on the cell-centred product $\rho_1 v_1 r_1^2$ of the first cell, which at this base is not a flux but the collocated odd-even mode of P44 -- MEASURED on the certified atomic_scalar_gj1132_kzz1e9/HeH2.13, the Riemann face mass flux is +1.00000 F_{wind} at every one of the 500 faces INCLUDING the base face (largest departure 1.6e-9) while the cell-centred product reads -2.00 F_{wind} at cell 1 and -2.27 at cell 2, so the answer to "is $v(1) < 0$ a reversal" is NO. The boundary therefore ran an INFLOW face at $w_{\text{rev}} = 0.9737$, and the two branches are not close: $\rho_{\text{rev}}/\rho_{\text{res}} = 0.531$, exactly $T_i/T_{\text{res}} = 420.4 \text{ K} / 223.2 \text{ K} = 1.88$ at the common face pressure, so base.inp's 226 K entered the state with weight 0.026 and cell 1 sat at 417.6 K. No window repairs that (narrower takes the reversal branch MORE completely, $w_{\text{rev}} = 1.000$ at 1e-8; wider tends to 0.5). (2) THE WIDTH: the module's own note asks for a window far below the operating point and quotes the hot-Uranus gate at face Mach 6e-6, but this planet's base Mach is 4.6e-7, BELOW the old 1e-6, and even with the sign repaired the old width leaves $w_{\text{rev}} = 0.215$. FIXED: the branch is keyed on the conserved mass flux read where the cell-centred product IS the flux -- the mean of $\rho v r^2$ over the wind window $r \geq r_{\text{flux}}$ that flux_spread_of_state is already defined on -- mapped onto the face by the interior density, and the window is 1e-8, 46 times below this planet's operating point and 600 below the hot-Uranus one; on the same state the branch reads $M_{\text{branch}} = +4.02e-07$ and $w_{\text{rev}} = 0.000$. The face flux itself is NOT read, because it is this boundary's own output and reading it would be circular or would put history into the residual; the wind is consulted only where the window has standing to speak, a cold start's wind window carrying no wind. Old behaviour: EXHALE_BASE_BRANCH_ON_CELL1=1, EXHALE_BASE_MACH_BLEND=<value>. Two review rounds then reshaped it. FIRST: the handover is C1 and blends the two WEIGHTS rather than the two Mach numbers (blending the Machs is bounded only by $|M_i|/\text{blend}$ and was measured to swing the face density by 0.398 across two samples), and four assertions were added as src/tests/grid_and_gates/base_branch_discriminant.f90. SECOND (R1): the stationary/marching split that first round had introduced was WRONG and is REMOVED -- $R_{\text{stat}}(U^*) = 0$ does not imply $R_{\text{time}}(U^*) = 0$, and MEASURED on the certified fiducial the same state reads mass 2.17e-09 / energy 2.13e-08 under the flux-keyed face and 1.31 / 1.05 under the local one. There is ONE boundary now, keyed on have_F and the weight s and not on the route, and the certification LABEL ("of the stationary operator") is written into docs/restart_contract_design_20260909.md section 7. USER DECISION 2026-09-16, APPLIED: the two jobs base_face_mach_blend was doing are separated and the second is stated SCALE-FREE -- the wind window's standing in the branch is read off the window's OWN relative flux spread du_window

= (max - min)/|mean| of $\rho v r^2$ over $r \geq r_{\text{flux}}$, the functional the convergence gate is defined on, with $s_{\text{wind}} = 1$ at or below $\text{base_wind_window_spread} = 1e-2$ and 0 at or above twice it on the same cubic, while $\text{base_face_mach_blend}$ keeps $1e-8$ and one job. The threshold is the geometric middle of the MEASURED gap, chosen as $\text{flux_spread_th_default}$ was: 147 certified states read $1.51e-04$ to $2.95e-03$ and the tightest state that is NOT a wind reads $7.50e-02$, $\sqrt{2.95e-3 \times 7.50e-2} = 1.5e-2 \rightarrow 1e-2$. Old behaviour: EXHALE_BASE_WIND_SAY_ON_MACH=1, threshold EXHALE_BASE_WIND_SPREAD. MEASURED with it at one thread: `wasp_full` 300 steps, `hydrostatic_column` 300 and `hp_front` 100 are all IDENTICAL to the same binary carrying the pre-L21 boundary (the $3.954e-04$ the Mach-keyed form left on `wasp_full` is gone); the certified fiducial is BITWISE against the campaign binary and re-certifies (`du_window` $2.458e-04$, `s_wind` 1.000000, `w_rev` 0); and the L14 fixture $x0.20$, re-solved from its seed, reaches CERTIFIED at outer pass 35 with all 35 pass lines and both written state files identical to the Mach-keyed build. `base_branch` 6/6 PASS, handover ratio 0.5000118; `report_base_face_state` prints what the boundary did and the EXHALE_RESIDUAL profile carries the stored Riemann face mass flux, so neither has to be reconstructed by hand again. **AND THE CAMPAIGN HAD TO BE RE-SOLVED:** the certified states are not states of the corrected boundary -- re-entered, the mass row of the certified atomic case reads $9.41e-01$ against the $1.49e-09$ it certifies at under the old one. On the fiducial, re-solved from its own certified state, CERTIFIED at outer pass 3: the base ghost goes $413.7 \text{ K} \rightarrow 226.0 \text{ K}$, which is `base.inp's T_base`, cell 1 $417.6 \rightarrow 238.0 \text{ K}$ at $1.73x$ the density, the profiles cross near cell 21 and agree to 0.2 percent in the outer wind, and Mdot $7.5096e+07 \rightarrow 7.3767e+07 \text{ g/s}$ ($\log_{10}$ $7.8756 \rightarrow 7.8679$, **-1.77 percent**) -- far above the 0.1 percent rule, so every case of the planet was due.

- **L7e sections 11 to 23: the H2 carrier row is CLOSED, through four defects and one wrong seed** (`diffusive_photochemistry.f90`, `System_HeH_mol.f90`, `certification.f90`, `EXHALE_main.f90`; new suite `src/tests/carrier_boundary_jacobian`). **The measure of 1.000 was a row with one live term, and not a mask, a scale or a floor:** MEASURED at cell 280 ($r = 1.6044 R_p$, $T = 6032 \text{ K}$) the H_2 production is $3.598e-01 \text{ cm}^{-3} \text{ s}^{-1}$ against a total loss of $2.154e-06$ and a transport divergence of $3.2e-09$, so the residual IS the production; the sum-of-magnitudes scale reads 0.99999 and the floor is $6.2e-14$, twelve decades below the terms. All of that production is $\text{R9 H}_2^+ + \text{H} \rightarrow \text{H}_2 + \text{H}^+$, fed not by H_2 ($n(\text{H}_2^+)$ is $5.1e3$ times $n(\text{H}_2)$ there) but by $\text{He}(2^3\text{S}) + \text{H} \rightarrow \text{HeH}^+ + e$ and R19, and the row's root is $n(\text{H}_2) = 145 \text{ cm}^{-3}$, $1.7e5$ above the carried value. **The relaxation never reached it because it never took a transport step:** the chemistry closure refused every trial on `n_offsimplex` > 0 (12 cells, all molecular clamps, worst excursion $6.5e-04$), a count that describes the ROOT SEARCH and not the admissible state both clamp branches hand back, and behind it `chem_cycle_tol` $= 1e-6$ sat below the sweep's own far-wind noise (the traced closure contracts $7.6e-05 \rightarrow 1.1e-05 \rightarrow 2.8e-06$ and then holds a $1.7e-06$ to $3.4e-06$ band over thirty further cycles at 10 to 11 R_p). With both corrected (`chem_cycles_max` 5 $\rightarrow$ 12, `chem_cycle_tol` $1e-6 \rightarrow 1e-5$) the carrier transports for the first time on this route. Three refuted attempts and one wrong seed then followed, each measured: an e-fold clause on the movement bound was implemented and WITHDRAWN (it is looser exactly where x_j is LARGE, so it freed the base and the front instead of the wind cells, and the premise that the wind cells were held by the bound is refuted -- their whole required change, $1e-06$ in absolute units, was always far inside it); an implicit-chemistry prescription was REFUTED because the carrier substep is ALREADY backward Euler with an implicit source

(carrier_residual evaluates carrier_source at the trial, solve_carriers Newtons on it with the source Jacobian and a block-tridiagonal solve; newton 2 resid 1.26E-13 per substep); and the third gate was the SEED -- SEED_X2 = local imposed the thermochemical $q_{H2}(p, T)$ where the thermochemistry is not the balance that holds, seeding a MOLECULAR HUMP over the whole outer wind (13 to 21 percent of the gas in H2 from 5.9 to 29 R_p at 580 to 1570 K, 80 to 360 times the root of its own row) so that the relaxation kept ZERO steps on every pass and both refusals came from $r = 7 R_p$. SEED_X2 = local now means the smaller of the thermochemical fit and the root of that cell's own H2 carrier row, the root taken from carrier_source through the new carrier_h2_chemical_root so the chemistry is stated once (the root needs the molecular ions, so the fit is applied, ONE sweep is taken, the root is read and that sweep's composition is discarded before the revised partition is applied); cell 412 falls from $x_2 = 4.4e-01$ to $5.2e-03$, and MEASURED along the way, the fit is SATURATED at $x_2 = 1$ over BOTH ends of the column (cells 1-125 and 425-500), so the old local mode was the element-ratio ceiling there and not a chemical equilibrium. **WHAT WAS LIMITING WAS THE MOVEMENT BOUND, AND IT HAS BEEN REWRITTEN:** it is now the relative change of each cell's particle count $n_{tot} + n_e$ -- what the fixed wind responds to, since that and the mean molecular mass set the pressure at the conserved thermal energy -- at the same trust, with EXHALE_CARRIER_BOUND_FRACTION=1 restoring the carrier-fraction measure. The old one was refused on every pass at a base or front cell and NEVER at the cells the certification is about, whose change was four decades inside the allowance; the new one answers to its setting, the particle-count change going $9.9238e-03 \rightarrow 7.7835e-04$ and the carrier drift $1.6874e-02 \rightarrow 1.3440e-03$ as trust goes $1e-2 \rightarrow 1e-3$. **THE ROW IS CLOSED:** .L7e/fixed5/HeH2.13 certifies every species row -- H2 carrier balance **1.93e-08 of 1.0e-05, against the 1.000 this item opened on** -- and its momentum and energy rows, and was refused by ONE entry, the hydrodynamic mass row at cell 1 ($1.555e-07$ against $3.7e-09$), STALLED and not descending, whose decomposition names the base boundary: the handoff is imposed on the GHOSTS at $x_2 = 0.99998$ while cell 1, which it does not impose, stands at the network's root $x_2 = 0.3315$, so the mean molecular mass differs by 12.8 percent across one face and the density steps 26 percent. The certified ATOMIC case of the same planet and He/H has a mass row of $1.487e-09$ and a base density step of 4.2 percent with no composition discontinuity, so the molecular cell-1 row was 105 times the atomic case's worst and the difference was exactly that composition step -- which item L21 then removed from the other side (the same state on the corrected boundary is ACCEPTED at outer pass 11). **REVIEW ITEM R4, added 2026-09-16:** the carrier matrix was missing two boundary derivatives and carried one wrong factor -- the outward face of the last cell when the mass flux enters, the base face of cell 1 when the base composition is not imposed, and a spurious division by the carrier mass (measured factor 2.0088 for H2). New suite src/tests/carrier_boundary_jacobian (10 assertions, central difference of carrier_advective_divergence against the assembled coefficient) passes after the fix; **no refusal is attributed to it.** The row decomposition of the two refusing well-mixed states was then measured with the faces separated (EXHALE_CARRIER_ROW_TERMS=1 now writes both faces of every row, diagnostic, default off): on HeH0.083 cell 500 ($r = 29.0 R_p$, the domain edge) the chemistry is 3 percent of the residual and the row is an unbalanced diffusive inflow against two advective face fluxes that cancel to a third of it; on HeH0.55 cell 306 ($r = 1.95 R_p$, $x_2 = 2.7e-06$) it is diffusive supply against He+ + H2 destruction with advection at 7 percent and radiation at 0.09 percent of the loss. What holds BOTH columns is neither: every carrier relaxation of both runs

ends on the composition movement bound and never on its own residual, the bound is cut $5.0\text{e-}03 \rightarrow 2.5\text{e-}03 \rightarrow 1.25\text{e-}03 \rightarrow$ floor $1.0\text{e-}03$ with the drift falling in proportion, and the cell that ATTAINS the bound is the H2 front (cell 231, cell 225) and not the cell that refuses. HeH0.083 is STALLED (worst row flat $2.41\text{e-}02$ to $2.46\text{e-}02$ for ten passes) and HeH0.55 has NOT ARRIVED (its refusing cell migrates 233 $\rightarrow$ 306 at the pace of the $x_2 = 1\text{e-}02$ radius). Nothing was changed for that measurement and no cell was excluded from certification; the three well-mixed cases are recorded as NOT SOLVED with that reason, and the remedy is proposed as item L22. MEASURED and recorded for the next judgment: the relaxation's interval is the cell's crossing time and carries no chemical time ($dr/|v|$ is 21 to 76 times the H2 chemical time at cells 194/218/250/280) while the operator may subdivide it only 8 times.

- L7f: the 7-to-21 disagreement between the wind network and the handoff was the ROOT FORMULA, and three rate-table entries were wrong** (docs/lhs1140b_stationary_L7f_20260915.md; mol_rates.f90, molecular_reaction_heat.f90, write_output.f90). carrier_h2_chemical_root formed the row's root as production over loss RATE at the trial, on the stated ground that the production does not depend on $n(\text{H}_2)$. It does: the dominant formation channel is the three-body association R15, $k_{15} n(\text{H I})^2$, and carrier_source closes atomic hydrogen out of the element budget as $n_{\text{H,avail}} - 2 n(\text{H}_2) - \dots$, so the production FALLS with $n(\text{H}_2)$ and is exactly zero where every nucleus is bound. The row is quadratic in its unknown, and the seed evaluated the quotient immediately after applying a thermochemical fit SATURATED at $x_2 = 1$ over the whole base layer, i.e. at $n(\text{H I}) \sim 0$ where the production it divides has collapsed. With the root SOLVED (a bisection on $g(n_2) = \text{production} - \text{loss through carrier_source itself}$, so the chemistry is still stated once), MEASURED on the same state: x_2 at cell 1 goes $8.915\text{e-}02 \rightarrow$ **9.837e-01** against the handoff's $9.9998\text{e-}01$, and the layer ratio $7.24\text{-}21.00 \rightarrow$ **1.02 at the first cell, 5.87 at the top**. NEITHER CHEMISTRY IS WRONG; the H^- route and grain-surface formation, which the item had suspected, are not needed and are not adopted. **What destroys H2 in that layer, measured for the first time** (the row dump now writes the H2 row reaction by reaction): the helium-ion channels R17/R20/R23 at 20 to 39 percent with R18 behind them, the proton channels R10/R13 at 8 to 53 percent (most of the latter recycling through R6), Lyman-Werner photodissociation 2 to 4 percent with its attenuation verified ($N(\text{H}_2) = 5.70\text{e}18 \text{ cm}^{-2}$, $f_{\text{shield}} = 7.37\text{e-}04$ at cell 1), H2 photoionization 4 to 17 percent surviving to the base as X-rays, and **thermal dissociation R12 nine decades below the first cell's loss at 1023 K and twenty-two at 629 K**, nowhere reaching 0.4 percent of it. That put weight on three rate entries, and the user supplied the primary sources. **R20 ($\text{He}^+ + \text{H}_2 \rightarrow \text{HeH}^+ + \text{H}$, 4.2e-13) IS RETIRED**: Schauer et al. (1989), J. Chem. Phys. 91, 4593 (READ) measures the radiative and the dissociative charge transfer at 15-40 K and no HeH⁺ channel, states that the HeH⁺ channel "apparently does not become allowed until the collision energy approaches 9 eV", and its H3⁺ signal is the SUM of the radiative and HeH⁺ channels, which bounds the latter at $\leq 1.0\text{e-}14$, 42 times below the Koskinen Table 1 value; Boehringer & Arnold (1986) independently agree ("a very low upper limit could be derived for the cross section of reaction (1c), $\sigma < 6\text{e-}21 \text{ cm}^2$, at energies between 0.13 and 7.5 eV"). The HeH⁺ source the network now carries is the one Table 1 omits, $\text{H}_2^+ + \text{He} \rightarrow \text{HeH}^+ + \text{H}$ at $3.0\text{e-}10 \exp(-6717/T)$, read from the primary source (Black 1978, ApJ 222, 125, Eq. 12 and rate table) and default ON because without it the HeH⁺ row has no source at all. **R17 IS CORRECTED**: Boehringer & Arnold measure over 18-408 K a two-body total $k_2 = 1.1\text{e-}13 (300/T)^{0.24}$ and Johnsen, Chen & Biondi's

Fig. 3 carries it to $T_{\text{eff}} = 700$ K, while the Koskinen Arrhenius ALONE is $2e4$ below the measured rate at 300 K and 200 below at 400 K, the reaction having TWO mechanisms (tunnelling out of a long-lived He^+-H_2 complex at thermal energy, and an over-barrier branch above about 400 K); the code now carries their sum, $k(R17) = [1.1e-13 (300/T)^{0.24} - k(R23)] + 1e-9 \exp(-5700/T)$, the bracket being the measured two-body TOTAL less the radiative branch the network carries as R23, so the coded channels sum to the measured total and the branching is stated once -- verified against Johnsen's Fig. 3, INSIDE A FACTOR 2 from 300 to 700 K, with the three-body channel left out and its reason at the code site ($k_3 n = 6.0e-19$ at 1000 K and $n = 7e12$, five decades below) and the omission now GUARDED IN EACH CELL: $k_3(T) n / k_2$ is formed there and warns once with that cell's own n and T (no cell trips it on the certified state). **MEASURED effect of both rate repairs: almost none on the H2 budget and decades on the ion budgets** -- at the first cell with the ionization re-solved, retiring R20 takes $n(\text{He}^+) \times 1.11$ and $n(\text{HeH}^+) / 487$ at 1023 K ($\times 4.4$ and $/ 8.0e4$ at 629 K) while the total H2 loss moves $1.916e+03 \rightarrow 1.904e+03$, and the R17 correction takes the He^+ channels' H2 loss from 739.34 to 739.37 while $n(\text{He}^+)$ moves $512.5 \rightarrow 502.6$. The helium ions of that layer are SINK-LIMITED, so removing one He^+ sink raises $n(\text{He}^+)$ until the others carry the same flux: the 25-30 percent share R20 appeared to hold overstated it. The seed's base root is unmoved ($0.98372 \rightarrow 0.98323$). Black's reaction (14), $\text{H}^+ + \text{He} \rightarrow \text{HeH}^+ + \text{h}\nu$ at $1e-18$, is NOT adopted: it is a 1978 proposal argued from a level count, adopting an unmeasured estimate is how R20 got in, and MEASURED it would be the dominant HeH^+ source and would still change nothing observable. Still unobtained: Moses & Bass (2000), JGR 105, 7013 -- READ later, and it did NOT upgrade the basis above 700 K, its Table A2 row R596 carrying the reference column "Estimate, see text" and the 5700 K exponent being discussed there for a DIFFERENT reaction; a rising term is nevertheless REQUIRED by the data (Johnsen $1.05e-13$ at 300 K against $8.5e-14$ from the plateau alone), so it is carried in the source's own words. **PARTLY SUPERSEDED THE SAME DAY BY L21**, and the pointer is in the memo's header: every base-face and base-temperature number of this item was measured on the base boundary as it stood. Re-measured on the corrected boundary with the same seed and the same network: base ghost **226.02 K** against 873 K, cell 1 **808 K** against 1023, the cell-1 mass row **4.92e-09** against $1.68e-07$, the energy row $9.96e-09$ against $7.4e-07$, and the state ACCEPTED at outer pass 11 where it was refused after forty. The composition step across the base face is still there and the row certifies anyway, so the base-BC proposal of the memo's section 6 is no longer the open question it was; the root correction, the reaction record and the three rate repairs are independent of the boundary and stand.

- **L7g: the molecular base is hotter than the atomic one because of the molecular reaction heat, and the ledger does not double count** (OPEN in one named remainder; docs/lhs1140b_stationary_L7g_20260915.md). On the old boundary the two base ghost rows carried 879.1 and 873.0 K and cell 1 **1023.0 K** where the ATOMIC case of the same planet, He/H and boundary carried 419.3, 413.7 and 417.6 K, with base.inp stating 226 K. FIRST TEST, on the user's instruction: the infrared coolants are NOT the answer -- the same case continued with Base IR field: True and Molecular IR bands: True for 20 outer passes gives ghost 873.15 K against 873.0 and cell 1 **1022.83 K against 1023.0**. WHY, measured channel by channel from Heating_breakdown.txt and Cooling_breakdown.txt: at cell 1 the heating is **97.7 percent heat_mol_chem** ($5.205e-07$ of $5.327e-07 \text{ erg cm}^{-3} \text{ s}^{-1}$, heat_HeI 2.1 percent and

everything else below 0.2) against a total cooling of $1.101\text{e-}08$, so the base heating exceeds the base cooling by a factor 48 and turning both infrared options on raises the cooling only to $1.225\text{e-}08$. The arithmetic: R15 three-body association runs at $6.92\text{e+}04 \text{ cm}^{-3} \text{ s}^{-1}$ at cell 1 and releases 4.48 eV per H₂ formed, which is $4.95\text{e-}07 \text{ erg cm}^{-3} \text{ s}^{-1}$, essentially the whole of `heat_mol_chem`, while the H₂ infrared bands radiate $1.17\text{e-}09$, 0.2 percent of it. Found on the way: **H3+ cooling needs neither key** -- `util_ion_eq` calls it for every cell with a nonzero `n(H3+)` once `Molecular chemistry` is on, and `Base IR field` only switches it from the emission-only rate to the net rate that also absorbs the diluted `B_nu(T0)`. **VERDICT: THE LEDGER DOES NOT DOUBLE COUNT.** The construction forbids it -- every species carries ONE formation-plus-excitation energy from a single reference state (`species_formation_energy`), every reaction heat is a difference of those and is written nowhere else, and a photoionization deposits `hν - IP` and no more -- and it is VERIFIED FROM OUTSIDE THE TABLE: Schauer et al. (1989) print the two He⁺ + H₂ channels as `dH = -6.51` and `-9.16 eV` and the ledger gives `+6.5109 (R17)` and `+9.1615 (R23)`, neither fitted to the paper, with every forward/reverse pair an exact negative. The cycles close on the ionization potential ONCE: the helium loop (He + `hν`; R17 `+6.511`; H⁺ radiative recombination; R15 `+4.478`) gives the gas `hν - 13.598` with 13.598 leaving as the recombination photon and `6.511 + 4.478 = 10.989 = IP(He) - IP(H)`; the proton loop (H + `hν`; R13 `+4.349`; R6 `+9.250`) gives the gas the WHOLE photon, `4.349 + 9.250 = 13.599 = IP(H)`, because the H₃⁺ DISSOCIATIVE recombination returns the proton's binding energy as heat instead of radiating it. So `heat_mol_chem` is where the ionization energy of H⁺ and He⁺ is deposited when the molecular channels neutralize them, and that is physics. **REVIEW FIX (Codex sections 2 and 5, applied): closing the arithmetic is NECESSARY and not SUFFICIENT for assigning the heat RECIPIENT.** Audited channel by channel, exactly one of the seventeen is radiative -- R23, H₂ + He⁺ -> H₂⁺ + He + `hν` -- and it was depositing its own photon as heat. Boehringer & Arnold (1986) p. 1461 on Hopper's mechanism (READ, verbatim): "This process involves a radiative transition from the first formed excited state of the collision complex (He+.H2) to the ground state (He.H2+) which then decays into the products (1a). The wavelength of the emitted photon should be about 153 nm and H₂⁺ should preferentially be produced in a vibrationally excited state ($v = 2$)." 153 nm is 8.1035 of the 9.1615 eV and at 1530 Å is longward of the Lyman-Werner bands and the Lyman continuum, so nothing absorbs it where it is made; the ledger now deposits **1.0579 eV**, the reaction ENTHALPY is untouched so the `physics_probe` reaction-energy identity still holds, and the range is stated at the code site (no re-absorption modelled, the $v = 2 \sim 0.55$ eV deposited rather than followed, so the term overstates by at most that and understates nothing). MEASURED share on the certified state: the photon removed is $9.201\text{e-}11 \text{ erg cm}^{-3} \text{ s}^{-1}$ at cell 1 against a total heating of $1.128\text{e-}06$, **8.2e-05 of it** -- corrected because it is wrong, not because it matters here. No other channel is radiative. Comment strengths were corrected at the same time (Black's coefficient is "an approximation valid under an assumed thermal H₂⁺ vibrational distribution" and NOT a bound, used outside its assumption because the H₂⁺ lifetime against R5/R8/R9 is $\sim 1\text{e-}3$ s against ~ 1 s for radiative vibrational relaxation, so the error may run either way; R17 above 700 K is a model uncertainty). AND the base temperature is not the reservoir's BY CONSTRUCTION, which is what item L21 then fixed: `characteristic_base_face_state` formed the face density as `rho_b = (1 - w_rev) rho_res + w_rev rho_rev`, and at `lm_il` ~ 7 to $8\text{e-}7$ with the old $1\text{e-}6$ window `w_rev` was 0.95 to 0.98, so the face state was 95 to 98 percent the interior's own isentrope. ON THE CORRECTED BOUNDARY: the

ghost at $r = 1.0000$ carries **226.02 K at $p = 1.0001 p_0$** , base.inp's stated pair to four digits, cell 1 falls 1023 -> **808.3 K**, x_2 0.3315 -> 0.3553, the cell-1 mass row $1.68e-07$ -> **4.92e-09** and the state is ACCEPTED at outer pass 11. The first resolved cell is still 3.6 times the boundary it is fed from and the layer above is unmoved (1527 K at cell 25 against 1544), so the verdict stands, and **226, 418 and 808 K are three different things**: the stated reservoir, the ATOMIC case's solved base with no heat_mol_chem at all, and the solved base with that source on. The photochemical column cannot disagree, because it does not solve for temperature: READ from the photochem clone, T is mapped from a prescribed profile (map_press_temp_edd), there is no energy equation and no heating rate in its state, so q_H2_base was computed at whatever T that profile states. THE ONE QUESTION LEFT, and it is not the ledger: whether h2_vibrational_heat_fraction, which evaluates to very nearly 1 at this base, should hold back more of the 4.478 eV R15 releases -- the newly formed H2 is born vibrationally hot and a real atmosphere radiates much of it in the rovibrational lines. The validity of the present form is now stated where it is used: it is an EFFECTIVE thermalization fraction, Hollenbach & McKee's $v = 1$ expression applied to the 4.478 eV of three-body association which is not a $v = 1$ excitation, valid only where the collider density is far above n_{cr} of every level of the nascent cascade -- which this layer satisfies ($n(H_2) = 4e9$ to $5e13$ against $n_{cr}(v=1) = 2e6$ to $6e6$) and a shallower base would not.

- **L22: a carrier relaxation whose movement bound is one scalar over the whole column** (PROPOSED 2026-09-16, NOT IMPLEMENTED; the decision is the user's). What is measured is in the L7e bullet above: on both refusing well-mixed states every carrier relaxation of every pass ended on the composition movement bound and none on its own residual, the bound was cut $5.0e-03$ -> $1.0e-03$ with the pass movement falling in proportion, and the cell that ATTAINS the bound is the H2 front (cell 231 at $1.2565 R_p$, row measure $6.6e-03$; cell 225 at $1.2309 R_p$, $5.1e-03$) while the cell that REFUSES is four to five times worse and decades away in abundance (cell 500 at $29.0 R_p$, $2.46e-02$; cell 306 at $1.95 R_p$ at $x_2 = 2.7e-06$). The bound exists because the wind is held fixed while the carriers relax, so it is RIGHT that the front be limited; what is not established is that the far wind must be limited with it. What a proposal would have to answer before anything is written: over what region the linear response of the FIXED wind is one statement (the bound is on the particle count, which the wind feels through the pressure at fixed thermal energy, so the region is a physical one and not a box of cells); whether the right object is a bound per region, a bound weighted by each cell's own contribution to the wind's response, or a relaxation that advances the front and the wind together so that no bound of this kind is needed; and how any of the three is prevented from becoming a knob that certifies a state by loosening what holds it. Nothing in the physics path is to be changed for this item until that is decided.
- **The two Codex reviews of this series, and what each finding cost.** Both were checked against the source before anything was done, and the verification tables are docs/session_handoff_20260915_rev1.md sections 12 and 17. The FIRST (2026-09-15 21:13, docs/session_handoff_20260915_review.md), nine source findings, every one CONFIRMED: R23 depositing its whole 9.16 eV as gas heat (molecular_reaction_heat.f90 adds $k_{23} n(H_2) n(He^+) q(R_{23})$ to gamma_chem with no photon term) -> reopened under L7g, every radiative channel of the network audited, the molecular cases re-solved after the fix and an atomic case checked byte-identical; the L21 switch between the two discriminants being discontinuous (base_boundary.f90 selects

M_wind only above the window) -> a C1 blend of the two weights, four tests, the LHS branch unchanged and verified bitwise; the L18 density authority decided by the SIZE of the discrepancy (load_IC.f90) -> authority by INTENT, with a stated rounding allowance and a refusal above it; the L19 projection unbounded and unreported (ionization_equilibrium.f90) -> maximum correction measured and reported, warning above rounding, sweep failure above $1e-8$; the rate commentary overstating its own strength (mol_rates.f90) -> wording to the stated assumptions and the three-body omission turned into a guard on the local n and T ; the pseudo-time CFL comment stated as general (steady_newton.f90) -> reworded as the measured control it is; L15 labelled nonblocking -> relabelled a prerequisite; the R15 heat fraction lacking a formation-state model -> named as the L7g remainder; and three reporting errors of the handoff itself (a case-row sum of 101, "log Mdot -1.8 percent" for a -1.8 percent Mdot and -0.0077 dex, sections read as live instructions). A frozen source manifest was written to accompany the binary md5, and the review's suggestion that a golden refresh follow attribution alone was NOT taken: the standing rule is one deliberate refresh at the end of a series with the movement reported, and the user decides. The SECOND (2026-09-16 07:09, docs/session_handoff_20260915_review1.md), R1 to R7 plus four wording items: R1, the stationary and physical-time routes using different base-boundary operators (confirmed, and it had been a deliberate choice to keep the marching goldens bitwise) -> one boundary for both routes, measured on one saved certified state; R2, composition_closing reading a bounded displacement -> the carrier's own fixed-point defect; R3, one global particle-count budget and three different cell indices under one label -> the three logged separately and measured distinct; R4, the carrier matrix omitting the copied-ghost derivative at $j = N$ when $F\rho(N) < 0$ and at the inner ghost when no base composition is imposed -> both derived from the same imposed-versus-copied ghost rule as the residual, with directional-derivative probes for both signs and both branches, and NOT claimed as the cause of any failure; R5, the outer-boundary reading of the well-mixed failures exceeding the evidence -> the wording withdrawn and the row decomposition taken instead; R6, R23's post-photon remainder deposited at once -> open with the R15 formation-state question, an effective thermalization fraction only with its timescale conditions stated; R7, "order-dependent reduction" too specific -> the wording corrected and the frozen-state residual test made the first step of L15. The review's own order of experiments was adopted as the order of the next work, under the standing rule that no pass budget is raised, no tolerance loosened, no boundary row suppressed and no rate changed to obtain convergence.

- **The campaign re-run on the corrected base boundary, twice** (docs/lhs1140b_rerun_20260915.md, three-column table by LHS1140b/models/compare_trees.py, which reads both trees with the readers of status.py). The catalog was re-solved on db87b88d1ce5 (built 2026-09-15 20:38: L4h, L7e, L7f, L18, L19, L21), which measures the boundary correction itself, and then AGAIN on c2e9c9990b9f (built 2026-09-16 10:43), which carries the review fixes too, so that every case of record stands on one md5. The 2026-09-14 results are preserved unchanged in LHS1140b/models_20260914_preL21/ and the db87 ones in LHS1140b/models_20260915_db87/ (83 cases, 413 MB), and both are the SEED of what follows: pick_seed.py gained a first tier, a case's own most recent certified state, so each case continues from the state that already solves every equation but the base row. WHAT MOVED, against 2026-09-14: the first physical cell falls by up to 72 percent in temperature (median 46) and its density rises by up to a factor 3.40 (median 1.83), while the ghost carries base.inp's T_{base} to the digit it is given in and nothing else in the column moves by that much (the profiles cross near cell 21 and

agree to a fraction of a percent in the outer wind); **the mass-loss rate falls by about 2 percent everywhere**, 0 to -4.0 percent with a median of -2.0 and NO case rising, which is a colder, denser base making a smaller wind; the He I 10830 red-pair equivalent width moves by a median -2.0 percent with a range of -82 to +4, the large negatives being the low-helium rungs where the metastable column is small and sits low ($kzz0/HeH0.55$ 0.1682 $\rightarrow$ 0.0305 percent A as its first cell goes 304.7 $\rightarrow$ 234.0 K), which shows that the 2^3S population of a cool, helium-poor base is not a robust number; and the crossings move by 1 to 3 percent in He/H except the two least diffusive ladders, $kzz0$ 2.7169 $\rightarrow$ 3.1010 and $kzz1e5$ 2.7074 $\rightarrow$ 3.0706, because their rungs sit nearest the measured line and a 2 percent shift in equivalent width walks a long way along a shallow ladder. A CONSISTENCY CHECK fell out of the GJ 699 ladder: four of its six rungs are BIT-IDENTICAL across the correction, because on that weak spectrum the interior isentrope at the base was already the reservoir, so the branch the old boundary took wrongly happened to give the right state; only HeH1000, whose base ran at 472.9 K, moves (to 260.9 K, $\dot{M}$ -1.69 percent). **THE FINAL PASS, and what it proves:** the 79 cases that certified on db87b88d1ce5 were solved again on c2e9c9990b9f, each from its OWN db87 state, and **all 79 certified, each in ONE outer pass** -- 67 prescribed cases in five minutes of wall clock (about 24 s apiece), the 9 closure rungs in forty minutes, the 3 XUV-scaled photochemical cases in under a minute -- which is what a seed that is already the root costs. The two generations agree far inside the tolerances the certification asserts: the red-pair DEPTH is bit-identical in all 79, the mass-loss rate in 52 of 79 and to 8.7 decimal digits at worst ($1.8e-9$), the first cell's temperature and density to 8.3 digits and the equivalent width to 7.6, with the largest departures all in the flux-closure rungs, whose Photochem column is re-solved and whose alternation stops at a marginally different point of the same fixed point. The cost of a case that was NOT near its root is the exception and it is a solver cost, not a seed problem: at 0.10 to 0.33 of the fiducial spectrum the first solve spends its forty outer passes with the hydrodynamic rows refusing while the gated species row is already at $1e-12$, and thirteen such cases were re-solved at EXHALE_PTC_DTAU0= $1.0e8$ (models/run_lowxuv_dtau0.sh, the recipe of item L14, the stopped attempts kept in models/.stopped/), MEASURED: $x0.30/HeH2.13$ 3 outer passes in 2 h 18 refusing $\rightarrow$ CERTIFIED in 7 min, $x0.33$ 4 passes in 1 h 02 $\rightarrow$ 6 min, $x0.10$ 8 passes in 3 h 21 $\rightarrow$ 16 min, $x0.20$ 4 passes in 6 h 22 $\rightarrow$ 17 min, $x0.25$ 3 passes in 4 h 00 $\rightarrow$ 18 min. Nine of the thirteen certified that way within the hour; FOUR did not and keep the same behaviour AT $dtau0 = 1e8$, the ramp falling back to $dtau$ of order 10 with the cell-1 mass row holding $||R||$ near 1.5, so raising the start is not what they need either. A defect this re-run found is the reservoir half of the L13 bullet above. STATE OF THE CATALOG, MEASURED by reading MODELS.md section 7 at 2026-09-16 11:32 (three molecular cases were still running): 95 rows, **81 certified** -- 79 of the 86 atomic cases and 2 of the 9 molecular ones. The seven atomic cases not solved are the three at 0.01 XUV (item L17) and atomic_scalar_gj1132 $x0.10$ _kzz1e9/HeH9.7 with the photochemical 0.15, 0.20 and 0.25, whose hydrodynamic rows refuse under a composition that is still travelling. Of the nine molecular cases, kzz1e9/HeH2.13 and HeH9.7 are certified, kzz1e9/HeH0.55 sits at a marching stop, kzz1e9/HeH0.083 is info = 0 but uncertified, the three well-mixed ones are the L22 case, and three were still running; the two molecular_photochem cases were redefined on the re-run closure rungs' own converged columns (He/H 2.0924 and 9.0103) instead of the stored 2026-08-30 profiles, so that each sits on the same column as its atomic counterpart and can be seeded from its certified state. The crossings of the re-solved

catalog are 0.4044 (scalar_gj1132_wellmixed), 0.0526 (scalar_gj699_wellmixed), 3.1010 (kzz0), 3.0706 (kzz1e5), 2.8930 (kzz1e6), 2.5365 (kzz1e7), 1.9729 (kzz1e8), 1.5161 (kzz1e9), 1.1579 (kzz1e10), 0.8927 (kzz1e11) and, for the first time in a molecular group, 1.7300 (molecular_scalar_gj1132_kzz1e9, between the rungs 0.55 and 2.13); the flux-closed ladder still has no bracket, every one of its rungs sitting above the measurement, and its composition converges as before (He/H at the matching level 2.0924, 3.0034, 5.0057, 7.0080, 8.0091, 9.0103, 9.7111, 10.011, 12.014).

- **The final binary, its manifest, and where the work ran.** EXHALE.x md5 c2e9c9990b9f14f1be8cd77abca68945, built 2026-09-16 10:43:25 with bare make (the conda-forge gfortran 16.2 on PATH and the OpenBLAS of its own prefix), HEAD 43bc28c plus the uncommitted work of this series. Its manifest LHS1140b/models/BINARY_MANIFEST_c2e9c9990b9f.txt carries the md5 of every one of the 256 source files at build time together with the `git diff --stat` of the tree it was built from, so the binary a case names in its REPRODUCE.md can be tied to source text and not only to a hash; the earlier generation has its own, BINARY_MANIFEST_db87b88d1ce5.txt, written the same way. The campaign ran on both machines of the NFS mount and each REPRODUCE.md names its host: the 68 prescribed atomic cases on lart4 at 6 jobs x 8 threads, the 9 elemental-flux closure rungs and the 6 XUV-scaled photochemical cases on lart3 at 4 x 8, the 13 raised-pseudo-time re-solves on both, and the final pass on both at the same shapes. models/archive_results.sh <tag> is what preserves a generation of results before the next one overwrites it.
- **The runners and the state tools.** src/utlis/map_state_to_grid.py (+528/-14) carries the reservoir rescale of any element (L11, L13), the --extrapolate-beyond option of L8 and the seed metadata rules; src/utlis/element_flux_closure.py (+174/-18) the reservoir handoff of every element that moved, the target grid header and the removal of the retired Valve eps: key; src/utlis/profile_match_level.py (new, 242 lines) is the one implementation of the matching level, R0, the elemental reservoirs a profile states and the metadata number format, for both callers (L13 sections 12 and 13); src/utlis/collisional_validity.py (+52/-10) reads the solution rather than the _adv profile; src/utlis/run_lower.py (+15/-6) carries L7d's fit correction. Under LHS1140b/models/: make_models.py writes every case's input from one table; pick_seed.py ranks five tiers by one composition rule; run_case.sh runs the four passes of a prescribed-composition case (seed, wind, _adv profiles, line) and grew the molecular seed branch, the continuation at $\text{dtau0} = 1\text{e8}$, the seed walk with SEED_ATTEMPTS, EXHALE_OUTER_PASSES, the EXHALE_PTC_DTAU0 default and the target grid header a profile case needs of its own; run_closure.sh runs one elemental-flux rung; run_campaign.sh and run_rungs.sh run the tree JOBS at a time (defaults 8 x 8 and 4 x 8 of the 72 cores), skipping any case that already carries tpm_He10830_metrics.txt; write_reproduce.py writes each case's REPRODUCE.md (binary and md5, the seed and its mapping command, every command with its environment, the outer passes, the certification, the line measures and the clock); status.py reads every case back and writes MODELS.md sections 7 and 8; make_stationary_seed.py built the six L5a seeds; compare_archive.py overlays a solved case on its 2026-08-30 counterpart (docs/figures/lhs1140b_wb_vs_archive_kzz1e9.pdf). Three more arrived with the re-run: compare_trees.py (355 lines) reads two whole trees with the readers of

status.py and writes the three-column table and the agreement block of docs/lhs1140b_rerun_20260915.md; archive_results.sh preserves one generation of results under its own tag before the next overwrites it (models_20260914_preL21/, models_20260915_db87/); and run_lowxuv_dtau0.sh re-solves a named case at EXHALE_PTC_DTAU0=1.0e8 with SEED_ATTEMPTS=1, keeping the stopped attempt in models/.stopped/ and naming it in the new REPRODUCE.md. pick_seed.py also gained the first tier the re-run needed, a case's own most recent certified state.

- **The test suites, MEASURED by running each run.sh on the final binary** (2026-09-16, OMP_NUM_THREADS=2):

suite	pass / fail
physics_probe	1491/0
krylov_and_dogleg	334/0
steady_species_rows	195/0
grid_and_gates	202/4
spectrum_type	170/0
carrier_retry	143/0
certification	84/0
state_mapper	76/0
attempted_step	70/0
adv_static_limit	53/0
energy_update	53/0
carrier_returned_state_acceptance	36/0
ionization_imposed_fractions	31/0
coupled_source_step	30/0
run_mode	31/0
element_operator	28/0
fuv_band_ledger	24/0
acceptance_classes	22/0
transit_census	21/0
molecular_seed	25/0
carrier_reference_scales	14/0
carrier_boundary_jacobian	10/0
carrier_constraint_attribution	10/0
species_masses	9/0
constrained_network_layout	8/0
residual_determinism	5/1
species_face_flux	1/0

Three suites are new with this series and three grew inside it: src/tests/molecular_seed (17 assertions with L7, then 25 as L7e and L7f redefined the local seed partition and added a row assert-

ing that the returned root balances the H2 carrier row), src/tests/ionization_imposed_fractions (L12a, 31) and src/tests/carrier_boundary_jacobian (review item R4, 10 assertions, a central difference of carrier_advective_divergence against the assembled coefficient); src/tests/state_mapper grew from 17 to 76 over L11 and L13, carrier_retry to 143, and grid_and_gates to 206 rows as L21's base_branch_discriminant.f90 and L20's cool_column_matches_breakdown_total were added. The run_mode failure standing at the start of the series is gone (31/0). The five failing rows that remain are pre-existing and fail identically on the control binary: grid_and_gates four -- outer_iteration_ending_is_the_stagnation_one, whose fixture reaches the pass budget where its comment records a stagnation ending, and the three stationary_evaluate_* rows, which read the pinned backup/regression/wasp_full_newton/IC/ state item L20 found stale -- and residual_determinism one (closure_spread_within_the_row_tolerance_atomic_elem_newton). steady_completion_flag and steady_selfconsistent_residual assert nothing without EXHALE_STEADY_RUNLOGS and report that, so they are not counted here.

- The campaign of 2026-09-14, on the pre-L21 base boundary** (the record of that generation, preserved case by case in LHS1140b/models_20260914_preL21/). 95 cases were written under LHS1140b/models/ and 83 were CERTIFIED, the 12 that were not being the three 0.01-XUV atomic cases blocked on L4h and the nine molecular cases blocked on L7e. Ten groups bracketed the He I 10830 equivalent width of the measurement (1.108 percent A) and reported a crossing He/H by the log-log chord between the two rungs that bracket it: 0.4012 (scalar_gj1132_wellmixed), 0.0526 (scalar_gj699_wellmixed), 2.7169 (kzz0), 2.7074 (kzz1e5), 2.6334 (kzz1e6), 2.4844 (kzz1e7), 1.9330 (kzz1e8), 1.4922 (kzz1e9), 1.1413 (kzz1e10), 0.8784 (kzz1e11). The nine elemental-flux closure rungs all converged, eps_H $8.0\text{e-}3$ to $4.7\text{e-}2$ and eps_He $2.0\text{e-}2$ to $4.6\text{e-}2$ against the tolerance $5\text{e-}2$, at $k = 4$ (eight rungs) or $k = 5$ (He/H 2.09), every iterate exhale_info = 0. The XUV grid, added after the ladders, spans 0.10 to 0.33 of the reference spectrum at He/H 2.13 and 9.7 with the scalar K_zz and at 9.7 with the profile: log10 Mdot 6.83, 7.01, 7.14, 7.24 to 7.25, 7.33, 7.37 for the scalar cases and 6.85, 7.04, 7.18, 7.28, 7.36, 7.40 for the profile ones, against 7.88 and 7.87 (scalar, He/H 2.13 and 9.7) and 7.91 (profile) at the full spectrum. Against the archived 2026-08-30 solution of the reference case ($K_{zz} = 1\text{e}9$, He/H = 1.6261), overlaid by models/compare_archive.py in docs/figures/lhs1140b_wb_vs_archive_kzz1e9.pdf: red-pair equivalent width 1.17576 against 1.11635 percent A, red depth 4.34388 against 4.17649 percent, FWHM 0.2543 against 0.2507 A and log10 Mdot 7.87 against 7.80 (READ) -- +5.3, +4.0 and +1.4 percent and +0.07 dex, measured on 2026-09-13 before the L6b cooling tables landed. Part of that difference is the archived curve's pre-2026-09-03 transit reader ghost rows. **Every number of this bullet was superseded by the base-boundary correction of item L21;** what stands now is the re-run bullet above.
- Open at the close of the series.** (a) **L15**, the JFNK stationary solve is not reproducible at eight OpenMP threads: the frozen-state residual evaluation is excluded (bitwise at 1 and 8 threads over ten runs) and the next step is to enumerate the parallel regions of the Krylov stage and what each writes, before attributing to a reduction order or to a shared write. Every control of this series was single-threaded for that reason. (b) **L16**, the pass-1 triple recorded in the L4g memo that the current source does not reproduce on the same fixture; to be rerun on the

L4g-era build and on the current one, single-threaded. (c) **L22**, the carrier relaxation's movement bound as one scalar over a whole column, PROPOSED and not implemented, with the three questions a proposal must answer first and the standing instruction that nothing in the physics path changes for it until the user decides; the three well-mixed molecular cases wait on it. (d) The **L7g remainder**: whether `h2_vibrational_heat_fraction`, very nearly 1 at this base, should hold back more of the 4.478 eV that R15 releases, the newly formed H2 being born vibrationally hot; the present form's validity condition is now stated where it is used, and the H2 infrared bands radiate 0.2 percent of that energy. (e) **L9**, a read-only diagnostic and output path: Do only PP exits the time loop AFTER a step and its output, and CFL: $1.0\text{e-}12$ suppresses that step's movement ($3.6\text{e-}14$ in ρ , $2.3\text{e-}12$ in v) but is a workaround and not an identity; until an explicit path exists that evaluates the held state with no time update, every REPRODUCE.md says the pass was taken at CFL $1\text{e-}12$ and the runner rechecks the written state against the certified one. (f) **The seven atomic and seven molecular cases that are not solved**, each with its reason: the three at 0.01 XUV are item L17 (diagnosed, not solved: the Newton model of a wind subsonic to Mach $1\text{e-}8$, with the HLLC contact kink named and the obvious regularization measured and reverted); `atomic_scalar_gj1132x0.10_kzz1e9/HeH9.7` and the photochemical 0.15, 0.20 and 0.25 hold the cell-1 mass row at $||R||$ about 1.5 on the colder, denser base, the same at $\text{dtau0} = 1\text{e}8$ as at 1, which is a base obstruction of the L17 kind at higher XUV and is not yet measured; and the molecular remainder is L22 and the still-running cases. (g) **L4e's two proposals**, not implemented: scale the linear system by the judged size `tol_k s_k(j)` (the option exists, is off by default and was measured not to be the repair on its own, the continuity overshoot being the Newton direction) and a line search that descends the judged distance the ledger ranks. (h) `src/tests/grid_and_gates/output_state_consistency.sh`, the row `outer_iteration_ending_is_the_stagnation_one`, FAILS ON THE CONTROL BUILD as well as on this one: its fixture reaches the pass budget where the comment records the stagnation ending measured on 2026-09-12, and the test itself says what to do, pick a setting that reaches the ending again rather than loosen the assertion. Its three `stationary_evaluate_*` neighbours fail on the pinned `backup/regression/wasp_full_newton/IC/` fixture that item L20 found stale; `backup/` is read-only for this work, so neither was refreshed. (i) The three coolant notes L6b reported and did not act on: the He 19.82 eV channel belongs in an energy ledger of the tracked 2^3S population rather than in the cooling sum, Fe I's lower levels are treated as LTE with no CHIANTI model atom to replace them, and Mg I 4571 and Ca II 7291/7324 are missing from the fits, which makes those two coolants an underestimate. (j) Named and left as proposals by item L21: separating the preconditioner's boundary block, and the base face's missing eddy-diffusive flux that item L7f's section 6 proposed before the boundary correction removed the row it was for.

The golden refresh of 2026-09-16

The matrix was run once against the delivered binary EXHALE.x md5 `c2e9c9990b9f14f1be8cd77abca68945` (manifest `LHS1140b/models/BINARY_MANIFEST_c2e9c9990b9f.txt`), single-threaded through REGRESSION_EXE on a fixed copy, and **all sixteen cases failed against the goldens of 2026-09-10**. This is the one refresh of the series: the goldens were read and never written by any item, each item measured its own movement against a control build instead, and the refresh is taken here, once, with the movement attributed case by case.

What the comparator's numbers are. compare_within_tolerance.py reports $|a-b| / \max(|a|, |b|)$ with no floor, so a column whose two values are both physically zero can read 1.000e+00, and a velocity that changes sign reads almost 2. The table below therefore carries the worst entry the harness prints AND, beside it, what moved physically: the peak relative movement of rho, of T, and of the velocity WHERE THE WIND IS A WIND (|v| above one per cent of that state's own |v| maximum), and the worst composition column holding more than 1 cm⁻³.

case	harness worst	rho	T	v (in the wind)	worst species above 1 cm ⁻³	attributed to
wasp_full	1.976 v r=1.064	2.7e-01	1.7e-01	2.1e-01	Na I 8.2e-01 (907 vs 4.95e+03)	L6b
wasp_he23off	1.992 v r=1.065	2.7e-01	1.7e-01	2.1e-01	Na I 8.2e-01	L6b
wasp_full_newton	0.653 cool ghost	1.7e-01	1.2e-01	1.3e-01	Ca I 5.7e-01	L6b
mol_base_handoff	0.000 HeHp r=1.081	2.1e-02	6.5e-03	3.9e-03	H3+ 9.1e-01 (1.02 vs 10.9)	L7f
mol_metals	1.000 HeHp r=1.079	2.4e-02	7.2e-03	5.2e-03	H3+ 9.9e-01	L7f , cool L6b
mol_lyman_werner	0.000 HeHp r=1.057	4.7e-04	3.5e-04	4.2e-04	H3+ 3.6e-01	L7f
mol_diffusion	1.000 HeHp r=1.081	2.1e-02	6.5e-03	3.9e-03	H3+ 9.1e-01	L7f
mol_ir_bands	1.000 HeHp r=1.079	2.4e-02	7.2e-03	5.2e-03	H3+ 9.9e-01	L7f , cool L6b
mol_sec_ion	1.000 HeHp r=1.084	1.1e-02	3.6e-03	8.8e-03	H3+ 8.5e-01	L7f on L7b 's cell
mol_carrier	1.000 HeHp r=1.081	6.4e-03	3.9e-03	2.1e-04	H3+ 8.1e-01	L7f
lower_profile	1.269 v r=3.210	9.0e-01	8.3e-01	1.3e+00	O I 9.9e-01	L6b , amplified (du = 3.17)
hydrostatic_column	0.000 HeIII ghost	6.7e-15	1.3e-14	1.3e-14	none above 2.2e-14	L19 , PROVED
oxygen_chemistry	0.982 heat r=1.018	7.5e-01	5.9e-01	1.9e+00	O II 1.0 (569 vs 5.6e-05)	L19 , amplified (du = 4.34)
hp_zero_seed	1.000 HeHp r=1.097	2.7e-05	3.4e-05	1.3e-06	H3+ 9.8e-01	L7f
hp_trace_seed	1.000 HeHp r=1.097	2.7e-05	3.4e-05	1.3e-06	H3+ 9.8e-01	L7f
hp_front	1.000 HeHp r=1.086	4.3e-07	7.7e-07	7.7e-07	H3+ 6.9e-01	L7f

L6b, the density-resolved C/N/O and metal cooling, moves the seven cases that carry metals -- the three WASP-121 b cases, `mol_metals`, `mol_ir_bands`, `oxygen_chemistry` and `lower_profile`, which carries its elemental reservoirs in the profile rather than in a `metals.inp`. The size is what its own memo measured on `wasp_full` before any golden was touched: the cooling below 1.1 R_p falls by up to 72 per cent and the temperature above 1.2 R_p rises 7 to 14 per cent, with $\log_{10} \dot{M}$ unchanged at 13.36. Na I and Ca I follow the temperature, which is why a neutral trace metal, not a hydrogen or helium column, carries the worst composition entry in those cases.

The cool entries L6b carries in the two molecular-plus-metals cases read the same way and are worth stating, because the harness's number and the physics point in opposite directions. At `mol_metals` row 32 ($r = 1.006$) the cooling falls from $1.77\text{e-}08$ to $4.72\text{e-}09$ $\text{erg cm}^{-3} \text{ s}^{-1}$, a factor 3.7 at the cell that holds 85 per cent of that column's peak cooling -- and the TEMPERATURE there moves by $1\text{e-}04$ and the heating by $1\text{e-}03$, because the cooling is 2 per cent of the heating in that cell. A C/N/O cooling coefficient changed by a factor nearly four and the state did not notice: which is what L6b's own memo means by "`mol_metals` barely moves (T to $8\text{e-}4$) ... at 890 to 2650 K the C/N/O channels are in the exponential tail and the layer is cooled by H3+ and the ground-term fine structure".

L7f, the molecular rate repairs, carries the worst `Ion_species` entry in ten of the eleven cases that run the molecular network (the eleventh, `oxygen_chemistry`, is the one case below where nothing can be separated). Its signature is unambiguous and is the physics: the R20 channel $\text{He+} + \text{H2} \rightarrow \text{HeH+} + \text{H}$ is GONE from `mol_rates.f90` (HeH+ is formed only through $\text{H2+} + \text{He} \rightarrow \text{HeH+} + \text{H}$, Black 1978), so **HeH+ falls by four decades** -- $1\text{e-}06$ to $4\text{e-}05$ against a golden 0.02 to 0.15 -- in every one of the ten; H3+ and H2+ follow through the chain; and He+ and He++ rise by a factor 2.2 because a helium sink was removed. Item L7f measured exactly those ratios against its own control (HeH+ 9.998e-01, He+ 2.166, He++ 2.203, H3+ 9.871e-01, H2+ 9.519e-01) and recorded that the hydrodynamic state moves 1 to 3 per cent, which is the ρ and T columns above.

The goldens are older than L6b, which is why the metal cases move most: they were taken 2026-09-10 and L6b landed 2026-09-13, so no comparison made against a control build of this series could see it -- `wasp_full` at 300 steps moves by $2.9\text{e-}11$ between the tree as delivered to item L18 (922dec0fa183, which already carried L6b) and the delivered binary. The golden is the only reference that predates it.

L19, the mass projection of each ionization sweep, is the whole of two cases and nothing anywhere else. Bounded by experiment where it is not the mover: on `wasp_full` at 300 steps, the projection alone moves `Hydro_ioniz` by **2.902e-11** and `Ion_species` by $7.882\text{e-}05$ (an Fe I density of $2\text{e-}12$), so none of that case's 27 per cent is its. On `hydrostatic_column` this is PROVED and not inferred: re-running it with `EXHALE_MASS_PROJECTION=0` reproduces the 2026-09-10 golden **bit for bit in all four files**, so the $9.589\text{e-}08$ in cool and the ghost-cell entry below are that projection and nothing else. On `oxygen_chemistry` item L19 measured $1.926\text{e+}00$ and $1.000\text{e+}00$ against its own control -- the entire failure -- for the reason the fixture states: it is a 1000-step snapshot stopped at $du = 4.34$, still changing by tens of per cent per step, so a difference in the last bit has a thousand steps to grow in. That case carries metals, the molecular carriers AND the oxygen carriers, so L6b and L7f act in it as well and every one of its columns has moved by order unity (H2 by $1.08\text{e+}11 \rightarrow 3.85\text{e+}08$, O I by 907 $\rightarrow 2.75\text{e+}08$, O II by $5.6\text{e-}05 \rightarrow 569$); **it is the one case of the sixteen whose movement cannot be apportioned**, and the reason is that it is a relaxation snapshot and not a solution. Nothing is claimed about it beyond that.

L21, the base-boundary correction, moves no regression case at all. That was measured directly rather than as-

sumed: wasp_full at 300 steps, hydrostatic_column at 300 and hp_front at 100 are bit for bit identical between the delivered boundary and the same binary carrying the pre-L21 one (EXHALE_BASE_BRANCH_ON_CELL1=1 EXHALE_BASE_MACH_BLEND=1.0e-6). **L14 and L4h** changed the pseudo-time start a caller may ask for and the ramp's rule; no matrix case sets either, and wasp_full_newton's movement is L6b's in its composition, not the ramp's.

The two signatures the harness flags that are not movements.

(a) hydrostatic_column, Ion_species row 1 column 6. Row 1 is the first ghost cell and column 6 is **He III**. The two values are 1.264e-180 (new) against 1.164e-170 (golden), in cm^{-3} , in a cell whose H I is 9.29e+13 and whose He I is 7.37e+12. Both numbers are zero as densities -- 170 decades below one particle per cubic centimetre, and the cell's own He II is already 6.48e-90 -- and they are the underflow tail of the equilibrium chain, in which He III is suppressed by $\exp(-E/kT)$ at 200 K. A relative change of 1.2e-09 in He II moves that tail by ten decades; the comparator, which has no floor, reads it as 1.000e+00. **Attributed by experiment:** with the projection off, this column is the golden value exactly. The physical cells of this case move by 1e-14.

(b) the columns the comparator reads as exactly 1.000e+00 in the _adv files. Those are not physical columns at all. Hydro_ioniz_adv.txt carries adv_T_status (col 8), adv_comp_status (col 9) and adv_mass_row (col 10); Ion_species_adv.txt carries the same two status fields after its species, at col 35 (34-species cases), 39 (with the four molecular carriers) or 42 (with the three oxygen carriers as well). The status fields are the integer classification write_output.f90 defines: **0** adv_corrected (the steady advective correction was solved and adopted), **1** adv_retained (the run's own value kept -- the cell's mass row stands above adv_conditional_tol, or the thermal Damkohler number is above one, or the gas enters the cell), **2** adv_failed, **3** adv_unsupported (the closure omits the molecular and oxygen carriers and this cell's particle count is dominated by them), **4** adv_not_evaluated (a ghost the upwind difference cannot reach). They are dimensionless enumerations whose smallest possible change is one unit, so 1 vs 0 and 3 vs 0 both read 1.000e+00 by construction. Every such entry in this matrix is one cell changing category, and it follows the case's own physics. MEASURED, three of them:

- mol_base_handoff row 167, $r = 1.058$, adv_T_status 0 -> 3. The unsupported test is whether the omitted carriers hold more of the particle count than the species the closure models, and at that cell the ratio $(\text{H2} + \text{H2}^+ + \text{H3}^+ + \text{HeH}^+) / (\text{H I} + \text{H II} + \text{He I} + \text{He II} + \text{He III})$ is 0.9977 in the golden and 1.0025 now. The cell sat a quarter of a per cent below the threshold and L7f's chain -- H3+ 768 -> 719, HeH+ 1.49e-02 -> 6.91e-06 -- pushed it across. Nothing about the post-process changed.
- mol_metals row 197, 0 -> 1. Its omitted-carrier ratio is 0.11 and falls, so this is not that test but one of the three retained conditions.
- wasp_full row 290, 0 -> 1, in a case with no molecular carriers at all: L6b's temperature moved either the cell's mass row past adv_conditional_tol or its thermal Damkohler number past one. adv_mass_row (col 10) IS a real number: the cell's own mass row, the DIMENSIONLESS fraction by which the face mass flux changes across the cell, which is what adv_conditional_tol is compared with. wasp_full_newton's entry is 0 against 3.69e-15 -- the rounding floor of that fraction, twelve decades below the tolerance it is measured against -- and hp_zero_seed's and hp_trace_seed's are 6.3e-04 against 2.0e-05 at cell 5, a small difference of two small fractions at a front.

The comparator is left as it is. A floor in it would hide the case it was written for -- a golden value of zero that becomes non-zero IS a full difference -- and the reading above is what the table's physical columns are for.

The refresh. backup/regression/golden/ of 2026-09-10 00:30 is preserved entire, with its roundtrip directory, as backup/regression/golden_blockJ_20260910/ (a NOTE.txt there says what it is and where the attribution lives); run_check.sh golden then snapshotted the sixteen case directories the matrix had just written, each final: count= line matching the matrix log. The two pinned fixtures were refreshed with it, as their own rule requires:

- wasp_full_newton/IC/ from the new golden of that case (certified=T, sec_ion_step 2402 -> 2429). The old pair was a 2026-09-08 state whose cool column was 1.3 to 2.9 times the cooling of its own state and which no longer re-certified, so the three grid_and_gates rows that read it (stationary_evaluate_rebuilt_within_sweep_budget, stationary_evaluate_certification_unchanged, stationary_evaluate_exit_status) failed on EVERY binary back to 2026-09-14. wasp_full_newton/IC/README.md records the refresh, as it instructs.
- atomic_elem_newton/IC/ **was NOT replaced**, and the attempt is worth more than the pin would have been. That case is not in DEFAULT_CASES, has no golden, and no suite row that reads its pair fails; its pair is the state its own run hands back, and its documented value is being the FIXED 2026-09-09 reference that items N7, N26b and L15 quote their measurements against. It was re-run with the delivered binary and stopped before it ended, so the 2026-09-09 pair, run.log and output/ stand (the stopped run's log is kept as run.log.20260916_partial, and README.md there records all of this).

WHAT THE STOPPED RUN MEASURED: on 2026-09-09 this case stood still -- zero dogleg step, zero trust radius, $||R||$ 1.892 to every printed digit for eleven iterations, info = 2. With the 2026-09-16 binary it DESCENDS, the trust region accepting steps at ratio 0.99 and $||R||$ falling 1.892 to **9.186e-02**, one and a third decades -- and then STALLS there: iterations 60 and 61 report the same $||R||$ to four digits with the full Krylov budget spent (gm = 40) and the pseudo-time collapsing (dtau 9.60e-05 then 3.91e-04). **That is item L4h's ramp collapse, not the zero-dogleg obstruction item N7 pinned this fixture for:** the obstruction N7 recorded is gone and a different one now holds the same case. Re-pinning would have replaced the reference those items quote, so it is left for whoever means to move it.

run_check.sh check was then re-run with the same binary, single-threaded: **REGRESSION PASS, all sixteen cases byte-identical**, 64 of 64 files data identical, no case falling back to the tolerance branch.

The three grid_and_gates rows that read the refreshed pair went with it: stationary_evaluate_rebuilt_within_sweep_budget from 6.509639e-01 to **3.825e-10** (Ca I at row 252, against a tolerance of 1e-08), stationary_evaluate_certification_unchanged from 1 to **0** (the writer and the re-evaluation both certified=T, sec_ion_step 2429 on both sides), and stationary_evaluate_exit_status from 2 to **0**. The suite now stands at **205 PASS / 1 FAIL**, the one being outer_iteration_ending_is_the_stagnation_one, which reads a different fixture, has failed on the control build since before this series, and is recorded above under item L7b (e).

roundtrip was refreshed too (user decision, 2026-09-16), by its own invocation, because it is out of DEFAULT_CASES: its second half is not a bitwise comparison. Its golden was the oldest of all, 2026-09-06, and it was stale in two

different ways. MEASURED against it with the delivered binary: the state files moved $1.565\text{e}+00$ (v at row 147, $r = 1.044 -- 3.3\text{e}-04$ of that state's own lv maximum, the base mode again; the physical columns there move $1\text{e}-02$ in T and $3\text{e}-05$ in ρ) and $9.998\text{e}-01$ (HeH^+ at row 207, $9.53\text{e}-06$ against $3.94\text{e}-02$, L7f's signature once more; above 1 cm^{-3} the worst columns are $\text{H I } 4.4\text{e}-01$, $\text{He I } 5.9\text{e}-01$ and $\text{H2 } 6.1\text{e}-01$ at 1.3 to 2.7 R_p , which is a 40-step snapshot at $du = 2.58$ and not a solution). **And its two `_adv` goldens no longer had the SHAPE of the files the code writes** -- 7 columns against 10 and 38 against 40 -- because they predate the `adv_T_status`, `adv_comp_status` and `adv_mass_row` columns, so `check_roundtrip` could not have passed whatever the physics did. After `check_roundtrip / golden_roundtrip / check_roundtrip` it is byte-identical in all four files, and the 2026-09-06 state stands in `golden_blockJ_20260910/roundtrip/`.

The round trip's second half FAILS, and not on the golden: the restart drops one provenance field. `roundtrip_check.sh` hands the stage-A outputs back as the `_IC` pair, reloads with `EXHALE_DUMP_IC=1` and requires the `# coupling: header` to be preserved. It is not:

```
ref coupling: ... certified=F cert_reason=no_stationary_claim mode=init
dump coupling: ... certified=F mode=init
```

`parse_coupling_header (load_IC.f90)` restores `sec_ion`, `sec_ion_step`, `valve`, `recon`, `iontrans`, `certified`, `mode`, `t_phys` and `fluxconst` from the header it reads -- but NOT `cert_reason`, and nothing else sets `state_certification_reason` on a reload, so the re-dump writes the state with its reason blank. The two headers then say different things about the same state: `certified=F` with a reason means a certification ran and declined, `certified=F` with none means no certification was made at all, and the reloaded file claims the second where the first is true. This is NOT new and not this refresh's: the same failure, in the same field, comes out of the campaign binary `db87b88d1ce53facf1d61084fa535ca5`. The fix is one case (`'cert_reason'`) in that parser plus restoring it through `set_state_certified`; it is NOT made here, because it would change the binary the sixteen goldens above were just taken with. Reported for the next build.

- **The sixteen backup/regression/ case directories that carried the noun "arm" were re-named** (`PLAN_20260916_rev3` section 10; mapping in `docs/named_case_audit.md` section 6): the He/H ladder is now `heh_0p3/heh_3/heh_10/heh_30` and the $\text{He/H} = 1$ family `heh_1_{12k,40k,lw_12k,lw_40k,newton,lw_newton,newton_bigstack,x2matched}`, and the four quarantined cases are named for the key each exercises. None of them is in `DEFAULT_CASES` or has a golden, so no reference data and no number moved; inputs and stored outputs were not touched.
- **Restart intent: stationary evaluate now produces every output of a run from the state it measured, and the runner's CFL $1\text{e}-12$ step is gone** (item L9, `docs/lhs1140b_stationary_L9_20260916.md`). Three states are named and each product says which one it describes: the loaded state, kept untouched; the work state, which holds the loaded conserved density, momentum and energy and derives p and T from the refreshed composition through the caloric equation of state, and which reports its own chemical and thermal closure defect; and the advection-derived composition, whose `_adv` files now carry `# derived_from: provenance` in place of the `# coupling: header` they used to copy from the run state and certify nothing of their own.

The evaluation reports the file's claim and the work state's verdict as two answers. MEASURED on the certified atomic_scalar_gj1132_kzz1e9/HeH2.13: the mass row the post-process weighs its own rows by falls from 1.573 to 1.734e-09 at cell 1 and from 3.08e-05 to 4.00e-14 at cell 500, because the replaced step assembled it with PLM on a WENO3 solution; the _adv temperature moves by up to 58 per cent at cell 5 and by less than 1.3e-05 above $r = 1.20$, the He I 10830 equivalent width by 5.1e-06, and log10 Mdot not at all (7.87).

- **The molecular network's energy goes where the measurements say it goes, and the three-body channel runs at the rate of its own third body** (item L7g, docs/lhs1140b_stationary_L7g_model_20260916.md). Five corrections, none of which touches a reaction enthalpy: R5 and R16 leave one hydrogen atom in $n = 2$, so the gas is charged 0.749 and 1.554 eV instead of 10.948 and 11.753 and the 10.199 eV reaches the $H(n=2)$ population where the run carries one and leaves as radiation where it does not (Takagi 2002; Giusti-Suzor et al. 1983; Guberman 1994); R6 leaves its H2 fragment at $v = 5-6$, so 2.480 eV of its 9.250 is internal and branches like R15's (Kokoouline et al. 2001; Strasser et al. 2001); the third body of the R12/R15 pair becomes the collider $\sum k_1(H_2) n(H_2) + k_1(H) n(H) + k_1(Ar) n(He)$ in place of $k_1(H_2)$ on the total density, with the pair kept an exact detailed balance collider by collider (Cohen & Westberg 1983, p. 559); the heat fraction's radiative rate becomes the all-level maximum of the code's own 302-level ladder, $5.5939e-06 \text{ s}^{-1}$, 6.56 times the $v = 1$ value it used, reduced from the line list at initialization and agreeing to $1.1e-04$ with the complete quadrupole set of Wolniewicz et al. (1998); and its collisional side becomes three published quantum calculations, one per collider, in place of two 1979 fits and an absent helium term (Lique 2015 for H, Jozwiak et al. 2024 for He, Le Bourlot et al. 1999 for H2 -- the fits were 3.47 and 107 times those calculations at 300 and 808 K). MEASURED on the certified LHS 1140 b molecular base held as it stands: heat_mol_chem falls 21.6 per cent at cell 1 and 9.4 per cent at cell 25, essentially all of it the 22.2 per cent fall in the R15 rate, and the thermalized fraction stays 1 to a part in $1e6$. EXHALE_REACTION_HEAT_RECIPIENTS=0 restores the recipients and the old third body together for a control run; it does not restore the collider coefficients. The atomic case wasp_he23off is byte-identical over 300 matched steps, so none of this executes without the molecular network. TWO THINGS THE ITEM DOES NOT REACH: the chemistry solver, which still hoists the total-density third body because a collider-resolved coefficient is a function of its own unknowns; and the certified base state itself, which is no longer a root of the corrected equations -- a bounded re-solve leaves an energy residual of $4.27e-01$ through two outer passes with every other row at its tolerance, so re-solving it is a campaign and not a pass.
- **L22 step 2c: one outer ghost rule for the carrier column, and its outer boundary stated.** The certification and the fixed-wind relaxation evaluated the carrier row of one state through two different outer ghosts (step 2b). Both ghost rules of the column now live in carrier_face_mass_fraction, which every evaluation of the advective term goes through, and the carrier state array has one carrier_outflow_ghost in place of three inline copies. The rule: the ghost composition is the interior continued, $X_{ghost} = X_N$, in both directions of the face mass flux, because the outer face is an outflow face of a transported scalar and the hydrodynamic closure at the same face states no reservoir either; and NO diffusive, eddy or settling flux crosses either end of the column, at the outer face because the gas there is not collisional (bulk Knudsen number 1.4-2.4 at the 30 R_p top, exobase at 18.8-25.4 R_p, READ from docs/collisional_validity.md) and at the inner face because the handoff composition al-

ready enters on the inflowing base face. Forming the outer face instead would leave the inward settling drift of H2 alone crossing it, thirty times the largest term of the row it enters. The two step 2b keys that measured those variants are removed. A new test row compares the two evaluations of one row at tolerance zero: 2.1724e-02 before, exactly 0.0 after. The three frozen LHS 1140 b states are bitwise unchanged, measured against a control built from the same snapshot of the tree, because their species vector already held the outflow copy in the outer ghost; what the step removes is a certified carrier row's dependence on that accident. `mol_carrier`, `mol_base_handoff` and `mol_diffusion` are byte-identical over 12000 steps in every output file except the # provenance: wall-clock line, and no golden was refreshed. Memo: docs/lhs1140b_stationary_L22_20260916.md section "Step 2c".

- **One three-body rate for the H2 chemistry and for its heat, the helium collider at every call site, and the Le Bourlot table corrected against the published paper** (item L7h). The three-body association R15 and its reverse R12 run on a collider sum, $k_1(\text{H}_2) n(\text{H}_2) + k_1(\text{H}) n(\text{H}) + k_1(\text{Ar}) n(\text{He})$; the energy ledger formed it and the composition solver still hoisted the total-density coefficient once per cell, so the H2 abundance was set by one rate and its heat charged at another. The collider sum is now formed inside the balance rows from the routine that owns it, both directions carrying the same density so the pair stays an exact detailed balance, with the control key restoring the total density in the chemistry as well. MEASURED on the certified LHS 1140 b molecular base, held: the collider sum is 0.778 of the total density at cell 1 and 2.487 at cell 243, because above the H2 front the third body is atomic hydrogen, whose coefficient is temperature independent while $k_1(\text{H}_2)$ falls as $T^{-0.6}$; the R15 rate therefore falls 22.2 per cent at the base and rises 149 per cent in the wind, and the H2 carrier balance of that archived state goes from 5.645e-07, within its 1.0e-05, to 7.817e-01. On `mol_base_handoff` and `mol_sec_ion` at matched 500 steps the front moves outward: $n(\text{H}_2)$ at $r = 1.11$ rises by a factor 6.5 and 5.8 and $n(\text{H}3+)$ by 9.5 and 13, T by at most 4.3e-02 relative and ρ by 2.5e-02, with $\log_{10} \text{Mdot}$ 7.93 for both binaries. `h2_vibrational_heat_fraction` is passed the ground-singlet neutral helium at the three sites that had dropped it. And the Le Bourlot, Pineau des Forets & Flower (1999) H2-H2 coefficient is corrected: the paper states that ONE set exists, computed for a para-H2 ($J = 0$) perturber and applied unchanged to an ortho-H2 one, while the module was mixing the two files the Cloudy distribution carries; the para file holds the paper's own 51 levels and reproduces its Table 1 critical densities to ratios 1.59, 1.09 and 0.72 where the other file is low by 1.4 to 3.3, and `gamma_10_H2` is now 12 to 30 per cent below the mixture. New test src/tests/physics_probe/molecular_third_body_in_the_chemistry.f90, 9 assertions including the three Jacobian entries by central difference, RED 5 of 9 against an entry text. Memo: the "Follow-up" section of docs/lhs1140b_stationary_L7g_model_20260916.md.
- **The coupled block, its coupling measured and its entry named (item L22 step 3, increments I1 and I2, 2026-09-17).** The block the design of docs/lhs1140b_stationary_L22_step3_design_20260916.md settled on is the existing Coupled carrier solve: True route, so the first question is whether its matrix-free action is a usable difference quotient at the width the refusing cases have. EXHALE_COUPLED_JAC_ACTION=<file> (with EXHALE_COUPLED_JAC_CELLS and EXHALE_COUPLED_JAC_DIR, default off) writes that action against a CENTRAL difference of the full residual at two step lengths and against the banded preconditioner's row of

the same direction, row by row for every unknown kind, and stops the run without a step. MEASURED on the frozen molecular_scalar_gj1132_wellmixed/HeH0.55 state at `nvar_jac = 4`, direction on cells 300-312: the action reproduces the central difference to a mean relative error of $3.692\text{e-}04$ (carrier direction, 40 moved rows) and $3.792\text{e-}03$ (energy direction, 21 rows), against the $1.542\text{e-}03$ step 2b measured for the carrier operator's own action; no species component is blocked, the fraction-to-the-boundary length is 13.8 times the direction, and every preconditioner color resolved, so neither of the two ways the coupled route was set aside before appears on this state. Both cross terms are live (the smallest energy row under a carrier direction $4.472\text{e-}09$, the smallest carrier row under an energy direction $1.721\text{e-}11$), the band holds all thirteen two-cell reconstruction entries $d_{\text{res_c}(j+2)}/d_{\text{f_c}(j)}$ that the carrier relaxation's block-tridiagonal matrix drops, the species rows carry the certification's own `tscale_code`, and the outer ghost rule is bitwise the same in both evaluations, which is step 2c holding. Coupled carrier solve: is now three-valued: False, True and On stall, the last running the alternation first and handing the state to the block at the pass the alternation gives up on today (the joint distance has not fallen in three consecutive passes AND the carrier relaxation of this pass and of the two before it ended on the composition movement bound), with the reason printed; a word the key has no meaning for now stops the run instead of falling through to False. The acceptance is unchanged, and with the key absent or False every path is bit for bit the alternation. New suite `src/tests/coupled_block_jacobian/` (8 rows), and `grid_and_gates/coupled_carrier_h2` grew five rows.

- The numerical flux of the seven low-XUV atomic cases of the LHS 1140 b catalog (item L25, decided by the user 2026-09-17).** Those seven are solved with Numerical flux: ROE on the 500-cell catalog grid and the rest of the catalog keeps HLLC, because the HLLC flux does not reach the stationary root there at all while the Roe flux does: with the catalog recipe otherwise unchanged three of the seven certify and a fourth reaches a wind whose three hydrodynamic rows are all within tolerance, refused only by the gated elemental He/H partition at $10.85 R_p$. The obstruction is the grid and not the wind, MEASURED: the unmodified HLLC flux certifies the same $x_{0.10}/\text{HeH}9.7$ case on a grid of twice the cells and the state it reaches is the 500-cell Roe state (Mdot +1.0 per cent, equivalent width -0.4 per cent, T and rho within 0.1 to 3 per cent above $1.005 R_p$). Where both fluxes solve, Roe against HLLC lowers Mdot and the He 10830 equivalent width by 0.3 to 0.6 per cent and leaves the first cell 6 to 7 per cent colder and 6 to 8 per cent denser with the mass-flux spread unchanged, and that systematic is now stated wherever those cases are quoted. The flux is a property of the case in `LHS1140b/models/make_models.py` (ROE_CASES, default HLLC), `status.py` prints it as a flux column, `write_reproduce.py` records it with the systematic, and `LHS1140b/MODELS.md` and `docs/lhs1140b_exhale_vs_pwinds.tex` carry the decision. The catalog's states for these cases are made by the campaign re-run on the final binary; no output of the measurement directories is copied into a case directory. Memo: `docs/lhs1140b_stationary_L25_20260916.md`, step 3.
- L22 step 3, increments I3 to I5: a route is not an equation set, and the block measured against the alternation.** The `carrier_newton` token of a state file's # options line adds no row and no column: it says only whether the transported balances are unknowns of the Newton vector or are relaxed at a held wind, and the balances, the columns and the tolerances are the same either way, so under the L23 metadata contract it is metadata OF a state and not a statement of which equations the state solves. A difference in it is now admis-

sible at a reload with nothing named (opt_is_route beside opt_changes_layout), reported as the restart changes the ROUTE and not the equations and written as a # route_change line the rungs after it inherit; the refusal stands for the tokens that do change the equation set. The token a state is written with now states the route that produced it, the handover of Coupled carrier solve: On stall included. Five new rows in grid_and_gates/restart_option_change, four RED against the control. MEASURED on lart3, both routes started together on the certified molecular_scalar_gj1132_kzz1e9/HeH2.13 state reloaded: over the same wall clock the alternation ran twelve outer passes, its worst gated carrier row falling from 7.82e-01 to 8.18e-06, and CERTIFIED at pass 12, while the coupled block reached trust-region iteration 14 and moved ||R|| by 0.6 per cent with all thirteen of its linear solves ending with the Krylov subspace exhausted, the L24 failure mode. The certified state beside the archived one: the He I 10830 equivalent width moves by one per cent (1.5599 against 1.5754 %A), log10 Mdot by 7.9024 against 7.9091, the H2 front by one cell and the base temperature by 3.7 per cent (778.79 K against 808.32). No handover fired in the three refusing cases within the passes they reached. The entry certification of all four reloaded states puts the hydrodynamic energy row five to six decades outside its tolerance, so none of them is a state whose carrier row alone refuses it and the two questions the increments were opened for are not answered by these runs. The three Newton fixtures with the key absent print the same number at every place both binaries printed one. No golden refreshed, no default changed.

- L26: the wind window's say is an amplitude times a shape, and the zero window is an ordinary point of the base boundary.** The branch weight of characteristic_base_face_state mixed the local weight w_i with the window's weight through a scale-free gate on $(\max - \min)/|\text{mean}|$ of $\rho v r^2$ over $r \geq r_{\text{flux}}$, and that has two defects the probes of item L26 measured: the gate is unchanged when the whole window flux is scaled, so the face density has two different limits at the zero window (MEASURED 37.407 and 37.468 per cent of ρ_b on the two 0.03 and 0.02 LHS 1140 b states, 4.188 and 3.794 on the certified 0.10 state and the fiducial, 0.411 on a cold start), and the range is not differentiable where the cell carrying an extremum changes (MEASURED one-sided slopes $-2.422039\text{E}+02$ and $+1.279177\text{E}-02$, a factor $1.9\text{E}4$, converged at three step sizes). The weight is now $s_{\text{wind}} = A(M_{\text{wind}}) C(d_{\text{window}})$ with d_{window} the relative STANDARD DEVIATION of the same flux and A a cubic smoothstep that is zero at a zero window flux and one at $|M_{\text{wind}}| \geq 1\text{e}-12$: A vanishes quadratically, so s_{wind} and its derivative go to zero along every direction into the zero window, and the second moment has no argmax. The threshold is recalibrated on the new functional the way `flux_spread_th_default` was, the geometric middle of the measured gap to one digit: 302 certified states sit at $4.06\text{e}-05$ to $1.82\text{e}-04$ and the tightest state that is not a wind at $1.39\text{e}-02$, so $2\text{e}-3$. The three LOCAL closures the plan preferred were measured first and all three are refused by the states themselves: on the four LHS 1140 b wind states every reading of the base mass flux built from cells 1 to k is negative (-4.2 to -294 of the window flux) while the branch those states carry is inflow, because the base artifact carries twenty to three hundred times the wind's own flux and the wind's base Mach number is $1\text{e}-6$. Nothing the code meets moves: ρ_b , both ghost densities, w_{rev} and s_{wind} are bit for bit the old ones on all five probed states, the two certified LHS 1140 b states re-evaluate to outputs that differ only in the provenance line, and the fiducial resolved under the new boundary returns `info = 0`, CERTIFIED, and reproduces the catalog profile to $2\text{e}-12$ in velocity. EXHALE_BASE_WIND_SAY_ON_MACH is removed (the amplitude weight is that statement, at its own scale)

and EXHALE_BASE_WIND_AMPLITUDE is added. The probe is now the suite: 32 PASS|FAIL rows on each of five states. Memo: docs/lhs1140b_stationary_L26_20260916.md, section "Repair".

The golden refresh of 2026-09-17

The tree was built once with bare make (conda-forge gfortran 16.2.0, the OpenBLAS of the same prefix, -O3 -fopenmp) and the matrix was run against that binary, EXHALE.x md5 59bfdb3fc4d0104fc2e9c3734596d2f6 (manifest LHS1140b/models/BINARY_MANIFEST_59bfdb3fc4d0.txt), single-threaded by the script itself. make -q returned 0 after the build, so the binary is the one the sources make. **Eleven of the sixteen cases failed against the goldens of 2026-09-16, and the seventeenth directory, roundtrip, failed with them; the five cases that run no molecular network are byte-identical in all four files.** This is the one refresh of the 2026-09-16/17 series.

The verdict case by case, MEASURED. The harness worst column is what compare_within_tolerance.py prints, the same measure $|a-b| / \max(|a|, |b|)$ with no floor as in the refresh before this one; the three columns beside it are the peak relative movement of rho, of T and of the H2 number density with the cell they sit in.

case	harness worst	rho	T	n(H2)	attributed to
wasp_full	data identical	-	-	no H2	nothing moves it
wasp_he23off	data identical	-	-	no H2	nothing moves it
wasp_full_newton	data identical	-	-	no H2	nothing moves it
lower_profile	data identical	-	-	no H2	nothing moves it
hydrostatic_column	data identical	-	-	no H2	nothing moves it
mol_base_handoff	9.398e-01 H3p r=1.094	1.049e-01 r=4.210	9.246e-02 r=4.210	9.123e-01 r=1.095 (1.633e+10 vs 1.432e+09)	L7g, L7h
mol_metals	9.875e-01 H3p r=1.093	1.109e-01 r=4.172	9.266e-02 r=4.172	9.189e-01 r=1.094 (1.803e+10 vs 1.461e+09)	L7g, L7h
mol_lyman_werner	8.110e-01 H2 r=1.078	2.048e-01 r=4.249	1.664e-01 r=4.249	8.110e-01 r=1.078 (1.032e+10 vs 1.951e+09)	L7g, L7h
mol_diffusion	9.400e-01 H3p r=1.094	1.049e-01 r=4.210	9.245e-02 r=4.210	9.124e-01 r=1.095 (1.631e+10 vs 1.429e+09)	L7g, L7h
mol_ir_bands	9.875e-01 H3p r=1.093	1.109e-01 r=4.172	9.252e-02 r=4.172	9.189e-01 r=1.094 (1.798e+10 vs 1.458e+09)	L7g, L7h
mol_sec_ion	1.895e+00 v r=1.073	1.478e-01 r=4.134	1.231e-01 r=4.134	8.663e-01 r=1.099 (1.140e+10 vs 1.524e+09)	L7g, L7h

case	harness worst	rho	T	n(H2)	attributed to
mol_carrier	9.974e-01 H3p r=1.099	9.411e-02 r=4.289	8.330e-02 r=4.289	9.947e-01 r=1.099 (3.016e+10 vs 1.586e+08)	L7g, L7h
oxygen_chemistry	1.954e+00 v r=2.325	9.270e-01 r=1.017	7.336e-01 r=1.015	1.000e+00 r=4.058	L7g, L7h, amplified (du = 4.51)
hp_zero_seed	6.675e-03 HII r=1.085	5.061e-07 r=1.084	4.336e-05 r=1.850	5.276e-04 r=1.149	L7g, L7h
hp_trace_seed	4.034e-03 HII r=1.084	1.291e-06 r=1.083	4.336e-05 r=1.850	5.276e-04 r=1.149	L7g, L7h
hp_front	1.616e-02 heat r=1.086	3.612e-07 r=1.097	1.250e-04 r=4.535	5.268e-04 r=1.149	L7g, L7h
roundtrip	1.509e+00 v r=1.042	1.417e-02 r=1.117	3.417e-02 r=1.110	9.545e-01 r=1.114 (1.630e+10 vs 7.413e+08)	L7g, L7h

The mover is the molecular network and nothing else. Every case that runs it moved and every case that does not is byte-identical, which is the gate on `thereis_mol` that item L7g asserted and measured on `wasp_he23off` over 300 matched steps. `lower_profile` is in the second group: it hands the lower atmosphere over as a profile with solar C/N/O, and it runs no molecular chemistry, so the corrected rates never execute in it.

The signature is the H2 front, moved outward. In the seven `mol_*` cases `n(H2)` at $r = 1.08$ to 1.10 stands 5.3 to 190 times its golden value, and `H3+` follows it through the chain; the movement of `rho` and `T` is an order of magnitude smaller than that and sits far outside, at $r = 4.1$ to 4.3 , where the relaxation snapshot is furthest from a solution. That is the direction the corrected rate table requires and item L7h measured on the same two cases at 500 matched steps: above the H2 front the third body is atomic hydrogen, whose coefficient is temperature independent while `k1(H2)` falls as $T^{-0.6}$, so the collider sum stands 2.5 times the total density there and the three-body association runs that much faster than the total-density coefficient ran it.

The three `hp_*` cases move by 4e-3 to 1.6e-2 and no more, because they are 100-step restarts from a state the corrected rates have had a hundred steps to act on, and `oxygen_chemistry` moves by order unity in every column, because it is a 1000-step snapshot stopped at `du = 4.51` that carries the molecular and the oxygen carriers together; as in the refresh before this one, **its movement cannot be apportioned** and nothing is claimed about it beyond the attribution of the rates.

The `_adv` status columns read 1.000e+00 in nine cases and are not movements: `adv_T_status` and `adv_comp_status` are the integer classification `write_output.f90` defines, whose smallest possible change is one unit, and every entry here is 0 → 3 (`adv_unsupported`, the omitted molecular carriers holding more of the particle count than the species the closure models) or 0 → 1 at one cell that the moved H2 front pushed across its own threshold. The reading is the one recorded under the 2026-09-16 refresh and the comparator is left as it is.

What moved no case. L9 (the no-step stationary evaluation and the `# derived_from:` provenance of the `_adv` files)

changes a header, and the comparison excludes # lines. L22 steps 1, 2c and 3 default to the route the matrix runs, and Coupled carrier solve is absent from every case. L25's Low Mach velocity jump and the coupled-block diagnostic keys default off. L26's base-face wind weight is bit for bit the old one on the five states its own probes measured, and the matrix agrees: the five cases with no molecular network are byte-identical through it. L12 adds an acceptance suite and no module. L15's instrumentation enters nothing unless a key is set.

L23 moves no data line and fixes the round trip. cert_reason is restored on a reload, so the certification pair of a restart header round-trips: backup/regression/roundtrip/roundtrip_check.sh on this binary reports coupling header preserved: ... certified=F cert_reason=no_stationary_claim mode=init and restart round trip is the identity to 1e-12. On the previous binary that row FAILED, with cert_reason dropped from the re-dump, and the 2026-09-16 block reported it for the next build. It is closed here.

The refresh. backup/regression/golden/ of 2026-09-16 is preserved entire, with its roundtrip directory, as backup/regression/golden_L7g_20260917/, a NOTE.txt there saying what moved it item by item; run_check.sh golden then snapshotted the seventeen case directories the matrix had just written, each final: count= line matching the matrix log. run_check.sh check was re-run with the same binary, single-threaded: **REGRESSION PASS, all sixteen cases byte-identical**, 64 of 64 files data identical, and check roundtrip byte-identical in all four of its files as well. No pinned fixture was refreshed with this snapshot.

12. PLAN_20260917: the corrections that follow the review of the 2026-09-17 handoff (2026-09-17 to 2026-09-18)

Plan of record: docs/PLAN_20260917.md, items L27 to L36, written after docs/session_handoff_20260917_review.md (the Codex review of the first handoff text) and the revised handoff docs/session_handoff_20260917_rev1.md, whose sections 0 and 4 carry what the review established and what was verified in the source before the plan was written. The item numbers continue docs/PLAN_20260913_lhs_stationary.md, whose last item is L26. The code the plan stands on is EXHALE.x md5 59bfdb3fc4d0104fc2e9c3734596d2f6, repository HEAD 3c73905ca8a7fe2af92a5c2b014c225b2feedff3, with the seventeen regression goldens as the 2026-09-17 refresh of section 11 left them. Every number below is READ from the item's own report or from the memo named with it; nothing was measured for this section.

What the series is. Three items repair diagnostics the review found unsound: a probe that evaluated PLM on states solved under WENO3 (L27), a log reader that joined the records of two different solves (L28), and a set of archived certified=T headers that had never been re-measured under the current evaluate route (L29). Two items correct a real conservation defect of the transport operators, where diffusion and advection divided by two different cell volumes (L30, L30b). One gives the molecular approximations a validity contract, an energy audit and an uncertainty record (L31). Two are read-only verdicts on states that refuse to converge (L32, L33). One re-runs the atomic half of the LHS 1140 b catalog on the binary of record and closes the seven low-XUV cases (L34a). Two build ionization stage transport and put the H II row of a run on the element operator's own nucleus flux (L36, L36c). Items L27, L28, L29, L30 and L30b wrote no separate memo under docs/; their record is named with each below.

L27: the diagnostics evaluate the production stationary operator, and the certification report prints a face-flux budget

What changed. New `src/modules/states/stationary_operator.f90` (200 lines) holds `select_stationary_reconstruction`, `restore_reconstruction_selection` and `stationary_face_mass_flux`, with one line added to `SRC.EXHALE_main.f90` (+48/-33) replaces the nine literal WENO3 assignments of the stationary and residual routes by one call, and the evaluate route passes the budget to the report; the marching PLM and the residual probe's save and restore are untouched. `certification.f90` (+59/-3) gains an OPTIONAL third argument on `certification_report_write` and prints one block from it: no tolerance was given a number, no entry of the inventory reads it, and `rep%certified` is formed before the block. `base_boundary_continuity_probe.f90` (+118/-33) is rewritten on the new entry point; `stationary_evaluate_products.sh` (+24/-3) gains a stage that starts a Reconstruction scheme: PLM input under Restart intent: stationary evaluate. The budget travels as an argument rather than as saved module state, so no report can print a budget measured on a different state. The selection also disarms the `recon_lambda_on_blend`, which was already disarmed at all nine sites.

Tests. On `LHS1140b/models/.L26/fid_resolve/output/` the probe was RED at 32 PASS, 0 FAIL with a base face flux of 32.376280 of the window mean and no operator identity printed; GREEN at 35 PASS, 0 FAIL with the base face at 0.9999843726323676, the spread over the faces 1.78e-09 and the off-set from the window mean 1.56e-05. On the catalog state `atomic_scalar_gj1132x0.10_kzz1e9/HeH2.13` the RED reading was 336.551906 and the GREEN one 0.9999748281788378, spread 5.04e-08, off-set 2.52e-05. The new stage of `stationary_evaluate_products.sh` was RED at 21 PASS, 2 FAIL (`plm_input_evaluates_the_stationary_operator`, `plm_input_face_flux_budget_printed`) and GREEN at 23 PASS, 0 FAIL. The production report and the probe return the SAME BITS for the base face, the minimum, the maximum and the face indices on both states.

Movement. Byte-identical on the data lines of `wasp_full` (12251 steps), `mol_base_handoff` and `lower_profile` (12000 steps each), with the same final du on both builds; the whole-file difference is the two provenance lines. The 32.376 and 336.552 readings survive only on an explicitly labeled row that asserts nothing.

Record. Two head notes in `docs/lhs1140b_stationary_L26_20260916.md` sections R3 and R9 saying that the face-flux readings there are PLM evaluations of WENO3 states, pointing at the revised handoff's section 4.3 and at this item; the rest of that memo is left as the record it was.

L28: a stationary record names its solve, and the reader reads one completed solve

What changed. `steady_newton.f90` (+26/-8, print statements only): a module counter incremented on entry to `solve_steady_ptc` and to `solve_steady_jfnk` and appended as `solve=<n>` to the five records the module prints. `src/tests/steady_selfconsistent_residual/run.sh` (+204/-25) takes the iterate record, the certification block and the completion line of the LAST COMPLETED solve and of no other, by the token where the records carry it and by the span between completion lines where they do not, and prints `incomplete_evidence` where the records of that solve are not all present. Four synthetic fixture logs were added beside it.

Tests. On the real log `LHS1140b/models/.L22/i3_alt/run.log` the entry-text reader was RED,

handback_matches_the_accepted_iterate at 2.83103e-08 and exit 1; the delivered reader is GREEN at 0.842987, which is 1.208E-08 over 1.433E-08 from that log's own lines. On the four fixtures the entry-text reader gave PASS, FAIL, FAIL, PASS and the delivered reader gives PASS, PASS, PASS and one deliberate FAIL on incomplete_evidence with the reason records_after_the_last_completion_line. Full suite on the real log: 7 PASS, 0 FAIL. steady_species_rows 195 PASS, 0 FAIL. No interval or tolerance was changed.

Movement. All 437 archived run.log files under LHS1140b/models/ were re-read with both readers: 156 carry no stationary solve, 142 keep the same ratio, 7 change, 41 are runs stopped inside a solve whose old ratio was built from the PREVIOUS solve and are now refused, 16 name a solve with no iterate record and are now refused, and 75 carry no iterate record anywhere. Of the seven that change, five move from failing to passing the existing [0.5, 2] interval: .L22/i3_alt 2.83103e-08 to 0.842987, .L22/i4_kz0083 4.21785e-08 to 0.796117, .L25/roe_s010_HeH9.7 2.67982e-07 to 1, .L7f/HeH2.13_ir 1.88841e-07 to 1.10715, molecular_scalar_gj1132_kzz1e9/HeH0.55 1.10606e-04 to 1.25829; .L14/x003_HeH2.13 moves 0.770056 to 1 and .L8/r45 0.661335 to 1.00047. Against a control built with only steady_newton.f90 at its entry text, carrier_model_a_newton gives byte-identical data lines in all four output files and run logs that differ only in the wall clock once the token is removed; wasp_full carried to its own end gives byte-identical data lines in all ten output files.

The record L32 added to this reader. A solve accepted at its loop top printed a completion line and no iterate record at all, because the iteration line is printed at the END of the iteration body while the acceptance test sits at the top; on 16 archived logs the last completed solve is of that kind, and the entry-text reader had answered the assertion from an earlier solve. Item L32 step 0 adds one record at the loop-top stop, in the iteration line's format and carrying this item's solve=<n> token. RED, a control-format log: FAIL ... measured=incomplete_evidence, reason no_iterate_record_in_that_solve; GREEN with the record: PASS at measured=1, read from the identified scope of that solve. Print only; carrier_model_a_newton byte-identical on the data lines of Hydro_ioniz.txt and Ion_species.txt.

Record. A head note at "Step 3, increments I3 to I5" of docs/lhs1140b_stationary_L22_20260916.md saying that its 2.83103e-08 reading was a reader defect and that the records of the one completed solve give 0.843.

L29: every archived certification claim reclassified under the binary of record

What was done. 119 archived states were re-measured with EXHALE.x md5 59bfdb3fc4d0104fc2e9c3734596d2f6 through the no-step evaluate route, on scratch copies, single-threaded, nothing written into any case directory. The counts:

group	states	reproduced	refused	no claim	no state	not evaluable
catalog,	77	70	0	0	7	0
prescribed-						
composition						
atomic						

group	states	reproduced	refused	no claim	no state	not evaluable
catalog, flux-closure rungs (last rung of each)	9	9	0	0	0	0
catalog, molecular	9	0	3	5	0	1
memo .L14	2	0	1	1	0	0
memo .L25 ladder rungs	18	18	0	0	0	0
memo .L25 direct solves	3	3	0	0	0	0
memo .L26/fid_resolve	1	1	0	0	0	0
total	119	101	4	6	7	1

no claim is a file that states none (certified=F), not evaluable a run that never reached the evaluation and said why. The operator identity column is empty in every row because that binary prints none; item L27 adds it.

The four refused claims, with the worst entry and the binary each names in its own REPRODUCE.md: .L14/x003_HeH2.13, hydrodynamic mass row 9.984E-01 at cell 2 against 3.2E-10, on a11038c245050d4f11852af829e13fc7; molecular_scalar_gj1132_kzz1e9/HeH0.55, energy row 4.634E-01 at cell 257 against 1.0E-06; /HeH2.13, energy row 4.267E-01 at cell 203; /HeH9.7, energy row 3.597E-01 at cell 1, the three molecular ones on c2e9c9990b9f14f1be8cd77abca68945, which carries the molecular physics of before item L7g. No cause is attributed to any of them. The one state the route could not evaluate is molecular_scalar_gj1132_wellmixed/HeH2.13, whose mapped seed carries he_diff: T into a run that does not diffuse elements.

The x003 check was run first: the residual block of this item's log and of the review's retained docs/audit_20260905/handoff_20260917_x003_evaluate.log are line-identical, original claim: NOT REPRODUCED, certified=F cert_reason=failing_entries, exit status 2. Of the 95 catalog rows, 79 reproduced, 5 no claim, 7 no state, 3 refused and 1 not evaluable.

Record. LHS1140b/models/CLAIMS_59bfdb3fc4d0.md (the head and the 119-row table), the script LHS1140b/models/reclassify_claims.sh, which refuses to run on any other binary, and one new column in LHS1140b/MODELS.md section 7, whose removal reproduces the old section line for line so that no other value moved.

L30: one spherical geometry for all transport, and the element face flux exposed

The defect and the correction. Diffusion and advection divided by two different cell volumes in both the elemental and the carrier transport: the advective half used the exact shell $(r_+^3 - r_-^3)/3$ and the diffusive half $r_j^2 \max(dr_j, 1 \text{ cm})$, which agree only to $O(dr^2)$ because r_{edg} is the arithmetic midpoint of the centers.

The element and carrier operators now divide every term of one row by the same exact shell volume and multiply every face term by the same face area $A(f) = r_{\text{edg}}(f)^2$, both defined once in `grid_construction` as `spherical_face_area_and_cell_volume`. The element row is the divergence of a single face flux, and that flux is one public object, `element_nucleus_face_flux`.

What changed. `define_grid.f90` (+79/-6): the new routine, evaluated in the factored form $(r_+ - r_-)(r_+^2 + r_+ r_- + r_-^2)/3$ because the difference of cubes loses about three decimal digits at the dr/r of $2e-4$ of the production base cells; filled from the current `r_edg` at every call rather than stored, because several acceptance drivers write `r_edg` without calling `define_grid`; a cell of nonpositive volume and a non-ordered face pair are refused, and the 1 cm floor on the cell width is gone. `binary_element_diffusion.f90` (+341): the four diffusion sites, the advective divergence, the new public `element_nucleus_face_flux`, the diagnostic helium mass on the same volumes, the dead geometry dummies removed, and the DISCRETIZATION and BOUNDARY NUCLEUS FLUX paragraphs of the header. `diffusive_photochemistry.f90` (+85): the carrier advective divergence and face coefficients, the carrier diffusive residual and the block-tridiagonal assembly, the geometry filled before the OpenMP regions so the assembled blocks stay bitwise the sequential ones. `carrier_mass_fractions` now states the outer ghost rule in both directions of the face mass flux ($X_{\text{ghost}} = X_{\text{N}}$) and what it does to the limiter, which is the section 11 bookkeeping item of the plan folded into this one. One line was added to `DIFT_SRC` in the `Makefile`, a file the item does not own, because `diffusion_tests.x` could otherwise no longer link.

Tests. The RED numbers were measured on a build in which ONLY the six diffusion denominators were reverted, so they isolate the geometry:

row	RED	GREEN	tolerance
<code>element_row_is_the_divergence_of_the_face_flux</code>	2.40865e-14 PASS	0.00000e+00 PASS	1e-14
<code>element_column_sum_is_the_difference_of_the_boundary_fluxes</code>	1.038185e-09 PASS	2.03150e-19 PASS	1e-13

Both are relative to the operator's own summed row scale, the measure every other relative row of this operator uses. The second is the conservation defect itself: under the entry geometry the column sum of $V_j R_j$ misses the difference of the two boundary face fluxes by $1.4e-9$ of the size of the terms, and after the correction by $2e-19$, which is the state's own mass closure and not a property of the grid. The rows that CANNOT be RED are reported as such: a manufactured smooth profile cannot separate two weights that are both second-order accurate (the refinement rows hold at order 2.0058 and 2.0029), and a constant-flux or static column gives a zero row under any single volume weight. Three existing conservation rows of `src/tests/diffusion_tests.f90` had to follow the operator, because `helium_mass_in_column` summed with the old diffusive weight; after that, `diffusion_tests` is 35 passed, 0 failed on both builds. Every suite whose driver links the changed objects was run on an isolated build and passes. Not delivered as written: the carrier DIFFUSIVE weight has no unit row, because the only public route to `carrier_residual` needs a full molecular state that no light driver builds; it is established by reading the two sites and by the regression movement below.

Movement. Zero-step Restart intent: stationary evaluate on three catalog states, control against measured: `.L26/fid_resolve` elemental transport row 6.671e-06 to 6.679e-06 at cell 217 against a $1e-5$ tolerance, CERTIFIED

both ways; `atomic_scalar_gj1132x0.10_kzz1e9/HeH2.13` 8.378e-06 at cell 258 to 7.496e-06 at cell 257, CERTIFIED both ways; `molecular_scalar_gj1132_kzz1e9/HeH2.13` 5.725e-09 at cell 217 to 7.245e-08 at cell 315, refused on both builds for the same three rows, none of which moved. **No certification verdict flips.** On the regression cases, `mol_base_handoff` and `wasp_full` are byte-identical in their data rows because neither enters a changed routine, `mol_diffusion` moves by 1.566e-05 (v) and 1.026e-04 (H3p), `mol_carrier` by 8.658e-08 and 8.039e-06, and `lower_profile` by 1.247e-07 and 4.421e-07, with `log10 Mdot` unchanged at 10.57, 10.57 and 8.82.

Record. No separate memo; the statement of record is the DISCRETIZATION and BOUNDARY NUCLEUS FLUX text of `binary_element_diffusion.f90`'s header, the header of `spherical_face_area_and_cell_volume`, and the two new rows of `src/tests/element_operator/`.

L30b: one spelling of the spherical volume, the failing carrier budget row, and the element flux writer

The follow-up of the three findings L30 reported outside its own scope.

The volume is now one expression. The eleven sites that still formed the shell volume themselves (`species_face_flux.f90`, four in `RK_rhs.f90`, two in `hydrodynamic_rows_body.inc`, `Reconstruction.f90`, three in `steady_residual.f90`) read `grid_construction's spherical_cell_volume(j)`, a new public scalar function (+51/-6 in `define_grid.f90`) that the array form now fills its `cell_volume` from, so there is one expression and the array form is that expression over the physical column. A scalar function was needed because seven of the eleven sites assemble rows over the padded ghost range the array does not cover and two more are scalar functions called once for each cell. The one place the expression is still written out is the kind-generic `hydrodynamic_rows_body.inc`, where taking one factor of a quadruple-precision row from a double routine would leave that factor rounded at 1e-16; in double it still reproduces the production routines bit for bit, which the four `generic_double_*_is_the_operator_bitwise` rows read as exactly 0.000000E+00. The face AREA was already one spelling and was left alone.

The failing carrier budget row was the fixture. `h2_nucleus_total_changes_only_by_boundary_flux` built its own face composition beside the operator and got the outer ghost rule wrong: it continued its own cell N+1 value past the outflow face where `carrier_mass_fractions` states $X_{ghost} = X_N$, so the boundary term carried half a limited slope the operator's limiter had already flattened. The arithmetic of the fixture's own profile gives 8.6e-9 against the 9.9166e-9 the row reported. With the fixture reading `carrier_mass_fractions` and `species_face_fraction`, and on the UNCHANGED production objects, the row is RED at 9.9166E-09 and GREEN at 1.3739E-15 against a 1.0E-12 tolerance. **No production operator was changed for it:** the Runge-Kutta species advection telescopes. The row beside it, `helium_nucleus_total_changes_only_by_boundary_flux`, passed all along at 4.9361E-17 because it already called the production face routine, and that contrast is what identifies the defect.

The element flux writer. `write_element_flux_profile` now takes the state and calls `element_nucleus_face_flux` for both halves, so the advective half carries the reconstructed, limited, upwinded face mass fraction instead of a one-sided pick of a cell value; the call moved to after `project_elements`, so the file is the flux of a state that exists; F_H is $F_{rho} - F_{adv}$, so $F_H + F_{He} = Mdot_{face}$ holds to the last bit; X_{face} is the face mass fraction that carries the flux. On `mol_diffusion` the columns move by 1.30e-05 (X_{face}) to 5.65e-05 (F_{He}). How much of that is the writer and how much the state cannot be separated on that case, and the report says why: the helium mass fraction of that column is uniform to 1.39e-3 across all 499 faces, so the writer's change is a change of the RULE whose size on a

column with a real helium gradient is not measured there.

One fixture outside the item's list had to follow the operator. `src/tests/grid_and_gates/hydrostatic_residual.f90` spelled the volume out at six places: `zero_gravity_momentum_scale_is_the_rows_own_terms` was RED at $6.44697\text{E-}13$ against $1\text{e-}15$ on two grids and is GREEN at exactly $0.00000\text{E+}00$ with the fixture reading the shared function. $6.4\text{e-}13$ is the measured size of the two spellings' difference at the base of that refined grid.

Movement. Two measures are given for each file, because the largest $|a-b|/\max(|a|, |b|)$ over a file is dominated by a species that is a numerical zero in a given configuration: `hydrostatic_column` (300 steps) moves by $8.46\text{e-}15$ of the column scale in T, `wasp_full` carried to its own du gate at 12251 steps on both builds by $4.57\text{e-}13$ in v and $7.75\text{e-}12$ in CaI, `mol_base_handoff` by $1.19\text{e-}05$ and $4.68\text{e-}06$, and `mol_diffusion` by $1.60\text{e-}05$ and $6.05\text{e-}06$, with $1.07\text{e-}04$ entry-relative on H2. Every compared column of every case is below the $1\text{e-}3$ harness tolerance and not below $1\text{e-}6$. The two molecular gates are 12000-step relaxation snapshots of a breathing base, so a $4\text{e-}13$ seed at the base cells is carried for 12000 steps; `hydrostatic_column` and `wasp_full` are the cases that show the movement without that amplification. Over all 12251 matched steps of `wasp_full` the two builds' du agrees between $1.1\text{e-}15$ and $1.7\text{e-}11$, so the trajectory does not amplify the change at all. Cost, measured on alternating runs: `hydrostatic_column` 5000 steps at a ratio of 0.998 and `lower_profile` 400 steps at 0.999, both inside the run-to-run scatter.

Reported from this item and repaired under bookkeeping. `src/utis/fortdep.py` scanned module and use and not include, so `hydrodynamic_rows.o` did not depend on the file whose text it is compiled from and editing the kind-generic rows alone did not trigger a rebuild.

Record. No separate memo; the statements of record are the corrected headers of `species_face_flux.f90` and `hydrodynamic_rows_body.inc`, the header of `spherical_cell_volume`, and the comment of the repaired fixture row, which now says that its exactness holds only while the fixture divides by the same volume.

L31: the molecular approximations get a validity contract, a reaction-cycle energy audit and an uncertainty record

Verdict: nothing is proposed for change, and no molecular energy share is counted twice. Files: `molecular_reaction_heat.f90` (+151/-23, comments plus one new module procedure and its call), `h2_vibrational_relaxation.f90` (+58/-22, comments only), `mol_rates.f90` (+57/-25, comments only), `src/tests/physics_probe/molecular_energy_recipients.f90` (+433/-12, 62 assertions to 81), the new suite `src/tests/h2_level_ladder/` (an 818-line model, a 32-assertion driver, `run.sh` and a README), and one row of `docs/input_schema.md`.

The energy audit. Five closed cycles, with the gas share of every step measured from the production `molecular_chemical_heating` rather than restated, close to $2.20\text{e-}13$ eV or better; the one the review named, the nascent H2+ vibrational excitation of R23, is deposited once, at R23, and R5 is charged the ground-state enthalpy and adds none of it back. RED: a deliberate $+0.55$ eV on R5's gas share, exactly the $v = 2$ energy the R23 comment names, is caught by three rows at $5.5000000000000007\text{E-}01$, $5.500000000002210\text{E-}01$ and $5.5000000000000007\text{E-}01$ against a $1\text{e-}9$ tolerance; GREEN on the revert. The audit followed the excitation into files the item does not own and found no double count there either.

The four prescriptions now carry validity and uncertainty text, and every claim the review called unsupported is

removed. The R5/R16 $n = 2$ recipient is a low-state recipient approximation, with the released-energy identity in place of the withdrawn claim that the share is an upper bound, and a new `reaction_heat_recipient_validity_report` prints three lines once a run when the recipients are in force, including that this code's H_2^+ lives about $1.0E-03$ s against R5/R8/R9 and relaxes vibrationally in about $1.0E+00$ s, so it is vibrationally hot. The R6 internal energy is a model estimate from the position of a published peak, with the peak-to-mean distance unquantified and no figure digitized. The quench fraction is a first-event branching model, not a bound over the ladder. The helium third body is an argon-based estimate over 77 to 5000 K with an uncertainty of ± 0.3 in log, the distance from argon to helium explicitly outside it. The stale paragraph claiming the photoelectric and Lyman-Werner callers omit helium is corrected against the three call sites that do pass it, which is the section 11 bookkeeping item of the plan folded into this one.

The reduced ladder and its bracket. The new suite solves the statistical equilibrium of 54 rovibrational levels of X with $v \leq 3$ (H_2 -H from Lique 2015, H_2 -He from Jozwiak et al. 2024, energies, weights and A values from Roueff et al. 2019, everything above $v = 3$ as ONE explicitly uncertain group), 32 of 32 assertions GREEN: the all-level $A_{\max}$ agrees with the production module to $9.4e-06$ relative, the collision-only bath departs from Boltzmann by $2.97e-14$, $2.92e-15$ and $6.12e-15$ at 808, 2128 and 5083 K, the steady injection closes to $3.7e-13$ and one backward-Euler step to $3.4e-16$. RED, two deliberate faults each reverted from a copy: dropping the degeneracy ratio from the detailed-balance factor takes the Boltzmann departure to 6.61 and fails 9 rows; counting the group's collisional disposal in both the heat and the radiation fails 8 rows. The bracket on the radiated share:

cell	T [K]	1 - f scalar	1 - f ladder bracket	ratio
molecular depth	808.32	6.793198e-07	1.20e-08 to 5.140654e-08	13.21
H2 front	2128.37	1.006850e-05	2.76e-07 to 2.004984e-06	5.02
dilute upper column	5083.44	5.963722e-04	6.99e-06 to 2.146044e-04	2.78

The scalar form radiates 2.8 to 13.2 times what the most radiative member of the bracket does, so it keeps more energy out of the gas than any member and replacing it could only add heat: **the scalar form stays the default.**

The argon-helium extrapolation is negligible, measured. R15 forms H_2 out of ATOMIC hydrogen, so the H_2 abundance does not bound it. On the certified molecular fiducial `molecular_scalar_gj1132_kzz1e9/HeH2.13`, 76 of 504 cells lie above 5000 K (5007.83 to 5870.73 K); the largest R15 source there is $1.6465e-05 \text{ cm}^{-3} \text{ s}^{-1}$ against $4.6848e+00$ from the other three H_2 formation channels at the same cell, a ratio of $2.3404e-08$ to $3.5145e-06$ over the 76 cells, and the balance the pair holds is at most $3.9053e-11$ in $2 \cdot n(H_2)/n_H$. On the hottest atomic wind of the catalog, `atomic_scalar_gj699_wellmixed/HeH1000` at a maximum 9517.55 K, 184 of 504 cells are outside the range, the largest R15 source is $1.3357e-10 \text{ cm}^{-3} \text{ s}^{-1}$ and the largest balance $1.6745e-13$. Because R12 and R15 carry ONE collider sum, their ratio is $K_{\text{eq}}(T)$ with no rate coefficient in it, so the extrapolation can move the rate of relaxation towards the balance but never the balance.

The uncertainty record. Five quantities varied separately at three cells of the certified fiducial, each as an exact difference against the production assembly, as fractions of the nominal molecular chemical heat:

cell	T [K]	recipient	R6 branch	R6 E_int	quench	helium efficiency
molecular depth	808.32	2.1e-07	1.3e-09	1.9e-10	6.6e-07	+41% / -21%
H2 front	2128.37	+4.6%	4.5e-07	6.8e-08	4.5e-07	-0.20% / +0.10%
dilute upper	5083.44	+130%	5.0e-10	7.5e-11	5.0e-10	-0.065% / +0.032%

The ordering is the result: at the base the uncertainty of the heat is the helium third body and nothing else, four to six orders above every other approximation, because the base heat is dominated by the R15 and R12 terms the collider sum multiplies; higher up the recipient takes over, and the two internal shares carry no uncertainty in this layer at all, because they reach the gas only through $1 - f = 6.8e-07$. The collider sum was checked bitwise in the three places that form it: the chemistry against `h2_association_collider_density` exactly 0.0, the carrier row's R15 channel exactly 0.0, and the heat ledger to $1e-14$ relative on a cell whose only open channels are R12 and R15.

Movement. Byte-identical. Against a control built from the entry text of the three source files, every output file of `mol_base_handoff` at 400 steps is identical with the provenance line excluded; the only difference in standard output is the three validity lines. `physics_probe` moves from 1585 PASS, 0 FAIL to 1604 PASS, 0 FAIL.

Left as a request, not a change. A published helium three-body recombination coefficient for $H + H + He \rightarrow H_2 + He$ over 200 to 5000 K. The only helium calculation in hand, Paolini, Ohlinger & Forrey (2011), Phys. Rev. A 83, 042713, plots a total over 0 to 350 K and tabulates nothing, and the table above makes this the largest single uncertainty in the molecular base heat.

Record. `docs/lhs1140b_stationary_L31_energy_cycles_20260917.md`.

L32: one frozen rejected coupled Newton system, measured before any preconditioner is designed

Verdict: the directional Jacobian action is not a linear map of its direction, so the Krylov solve has no theory to stand on, and neither a wider subspace nor a preconditioner can repair it. That a wider subspace does not repair it is measured; that no preconditioner does is an argument from what a preconditioner is, since right preconditioning replaces $Ax = b$ by $A\hat{M}^{-1}y = b$ and where A is not linear neither system is one a Krylov method has a theory for. The checkpoint is the block route's ENTRY state on the certified `molecular_scalar_gj1132_kzz1e9/HeH2.13` state, 500 cells, 5 unknowns in each cell, 2500 rows, $\|R\| = 4.2666e-01$, $\text{arc sqrt}(\text{eps})(1+\|Y\|) = 3.1541e-07$, the pseudo-time shift of the leg zero. Two corrections to the text the plan and the handoff carry: the $2.597e-01$ is the FIRST linear solve's reduced residual and the largest of the seventeen legs, and trust-region iteration 14 was an ACCEPTED step at $1.337e-01$, the rejected ones being 13 and 16.

The seven measurements:

1. **One operator is being sampled.** $F(Y)$ twice, $F(Y)$ with an unrelated state evaluated in between, $A(v)$ twice and $A(v)$ with a state in between are all bitwise identical, against a noise floor of $1.6e-12$ in row-scale units. Nothing that follows is a reproducibility defect.

2. **The perturbation ladder, by row family.** Along the scaled right-hand side, four of the five families behave as a difference quotient should, the error falling to $3.5\text{e-}07$ to $3.6\text{e-}07$ at the route's own arc and rising at ten times it; the momentum rows do not, standing at $8.16\text{e-}02$ at that arc on 131 of the moved rows, and the elemental rows do not along the carrier and the energy directions ($4.71\text{e+}00$ and $3.55\text{e+}01$). The mean relative error of $3.692\text{e-}04$ measured on another state does not bound these. Worst cells 11 to 14, 42, 123 to 130 and 216 to 223.
3. **Additivity fails and homogeneity holds.** On the first two Arnoldi directions $||A(v_1+v_2) - A(v_1) - A(v_2)||$ is $1.867\text{e+}02$ of $||A(v_1)+A(v_2)||$, and on two directions off the basis $1.871\text{e-}03$; the defect scales as the arc to the power 2.291 and 1.958, so it is curvature and not a floor, and no component is blocked by the feasible set. Nine tenths of the squared defect sits in cells 1 and 2 and 87 per cent of it in the momentum rows. Meanwhile $A(3v)$ against $3A(v)$ is $1.400\text{e-}14$, the worst Arnoldi image column gap over 40 columns is $3.523\text{e-}16$ and the loss of orthogonality $2.164\text{e-}13$: each column is a correct product on its own direction, and the sum over the columns is not the product on the sum.
4. **The reported residual is not the residual of the step.** The default cycle reports $2.589\text{e-}01$ while the true $||b - Ax||/||b||$ of its own iterate is $5.113\text{e-}01$, and the Arnoldi image the trust region reads its predicted decrease from is $4.210\text{e-}01$ away from the operator's image of that step; choosing the step by its true residual gives $3.189\text{e-}01$ with a zero image gap. Factor 1.98 between the two, the same shape as N27's 15 per cent.
5. **Orthogonality and the chemistry tolerance are not it.** Orthogonality $3.269\text{e-}12$, four decades below the level at which a second pass is worth its inner products, and orthogonalizing twice moves the fifth digit. EXHALE_RESID_EQ_TOL from $1\text{e-}6$ to $1\text{e-}12$ leaves the attainable floor at $3.189\text{e-}01$ to four digits.
6. **Eight times the subspace buys nothing.** On one operator and one right-hand side, a subspace of 40 spends 40 products and reaches $3.18872\text{e-}01$, and 80, 160 and 320 all stop at 70 products at $2.81563\text{e-}01$, returning the same step to six digits. The cycle stops because the true residual of its iterate has risen at two consecutive checks: $3.730\text{e-}01$ at product 20, $3.189\text{e-}01$ at 30, $5.113\text{e-}01$ at 40, $2.816\text{e-}01$ at 50, $3.454\text{e-}01$ at 60, which a GMRES iterate on a linear operator cannot do.
7. **An assembled Jacobian and a direct solve on 50 cells.** The direct solve solves what it was given to $9.408\text{e-}12$, and the map it was assembled from does not agree with it: $||A(x) - (\text{assembled } A) x||/||(\text{assembled } A) x||$ is $1.196\text{e+}04$, the same at unit step length, with additivity $1.719\text{e-}01$ and homogeneity $1.170\text{e-}16$ on that grid. The memo states the caveat that a column probe displaces one unknown in its own units while a combined direction spreads the arc over every unknown, so this measures the departure from ANY single linear map.

The banded model omits no long-range coupling, so the third hypothesis is refuted: over six sampled carrier columns the part lying outside the band is exactly zero and the band's own entries are 0.09 to 3.5 per cent off, which is all of the $2.8\text{e-}02$ to $3.9\text{e-}02$ difference. **A base-only block is not a cure**: the modes that break the linear model are confined to cells 1 and 2, but the modes the solve is held by are carrier rows spread over cells 223 to 287 beside 1 and 2, and the refusing row is carrier H2 of cell 243 at $r = 1.317$. Every "relative residual reached X" in the L22 and L24 records is a reduced least-squares residual, not the residual of a step.

No production numerical change, no default moved, no preconditioner designed; the one source change is the print-only loop-top iterate record reported under L28 above.

Record. docs/lhs1140b_stationary_L32_20260917.md, with the frozen checkpoint, the identities and every run

log under docs/audit_20260905/L32_20260917/.

L33: slow outer relaxation or a stalled mode, read from the existing logs

Read-only; no solve was run and no source file was edited. The three runs stopped on 2026-09-17 are classified:

run	case	classification
i4_wm0083	wellmixed/HeH0.083	front relaxation. The $x_2 = 1e-2$ contour entered the domain at pass 7 and walked inward from 28.5489 to 9.13.
i4_wm055	wellmixed/HeH0.55	front relaxation, unambiguous: the refusing cell walks outward through 315 to 445, that is 2.1136 to 11.7471.
i4_kz0083	kzz1e9/HeH0.083	neither. The cell is fixed at 499 from pass 9 to 40 and both H2 contours stand still, which reads as a mode, but

The corrected progress factors. The revised handoff's three figures are confirmed: 0.95273 a pass over passes 26 to 37 of i4_wm0083 (4.73 per cent), 0.91232 over passes 13 to 16 of i4_wm055 (8.77 per cent), and 1.14, 1.04 and 0.88 per cent a pass over passes 20 to 40, 23 to 40 and 30 to 40 of i4_kz0083; the conditional extrapolations to $1.0e-05$ are 175.0 and 86.7 further passes and 384, 420 and 497. What the single-window figures leave out changes the reading: i4_wm0083 ROSE from $2.65e-02$ to $9.20e-02$ over passes 1 to 22 at a factor 1.0611 a pass, so 0.953 is the second half of a curve with a maximum; and the rate follows the movement bound, i4_kz0083 falling 2.49 per cent a pass at a bound of $2.5e-03$ and 1.04 per cent at $1.25e-03$, so halving the bound took the decrement to 0.42 of itself against 0.50 for the bound. The bound is near its floor of $1.0e-3$ and only ever ratchets down within a run.

The control that makes the reading possible is i3_alt, the run that certified kzz1e9/HeH2.13 at outer pass 12 and the only one of the four in which the carrier relaxation reached the fixed point of its own operator (carrier residual $2.87e-09$ to $4.70e-09$ at every pass, 11 of 11 passes ending "the carriers stopped moving at the fixed wind"). In the three refusing runs every relaxation of every pass was cut off by the movement bound, 37 of 37, 16 of 16 and 39 of 39, and the composition handed on stood $6.6e-02$ to $2.0e-01$, $1.6e-01$ to $3.0e-01$ and $1.72e-02$ to $1.86e-02$ from that fixed point. None of the three entered the coupled block, so L32's verdict applies to none of them.

The handover was never exercised. It needs outer_no_progress, three CONSECUTIVE passes without progress, together with three carrier relaxations ending on the movement bound. The second was met in every pass of all three runs; the longest consecutive run of no-progress passes is 1, 2 and 1, so the criterion was never reached. This corrects the wording "neither of the handover's two counts reached three": the bound-ending count reached three in all three runs. The handover is an unexercised route and nothing more.

One bounded continuation experiment was designed and NOT run: five rungs uniform in q_{H2_base} from 0.190110258 to 0.476178639 (He/H 2.1300 to 0.5500) seeded from i3_alt's certified state, because the base H2 fraction and not K_{zz} is what sets the front (at the same base fraction the eddy coefficient moves the $x_2 = 0.5$ front only from 1.2565 to 1.3705 R_p , while q_{H2_base} moves the $x_2 = 1e-2$ contour from 1.1626 R_p to beyond the 30 R_p boundary).

Record. docs/lhs1140b_stationary_L33_20260917.md and the reader LHS1140b/models/.L22/outer_pass_history.py.

L34a: the atomic catalog re-run, 79 reproduced and the seven low-XUV Roe cases

Verdict: 79 reproduced, 0 refused, 0 re-solved, and all seven low-XUV Roe cases certified. Every archived atomic state of the catalog, the 70 prescribed-composition cases that carry a state and the 9 last rungs of the flux-closure

ladders, reproduces its own stationary claim under EXHALE_3146d11b.x through the no-step evaluate route, so the re-solve branch never ran and no pass was spent on one. The verdicts match L29's on 59bfbdb3fc4d0 in all 79 rows.

The states did not move. Mass density bit-identical in every cell of every state; velocity moves in 32 of 79 by at most $2.219\text{e-}16$ relative; temperature by a median $5.836\text{e-}14$ and at most $3.807\text{e-}12$, which is the composition refresh of the work state. $4 \pi \rho v r^2$ of the outermost physical cell is unchanged to the last bit in all 79, and the Log10 of steady-state $\dot{M}$ line is unchanged to its two decimals in all 79. The He I 10830 equivalent width moves by a median $5.4\text{e-}06$ and at most $3.6\text{e-}05$, which is the size item L9 measured on the fiducial for the change of post-processing route ($5.1\text{e-}06$) and comes from the `_adv` profiles. The gated elemental He/H partition row of the 64 states that run element diffusion moved by a median 0.9 per cent and at most 16.2 per cent between the value each case's own solve accepted on c2e9c9990b9f and the value this evaluation measures, with the same refusing cell in 56 of the 64; every state still certifies, and the movement is NOT attributed to L30 alone, because the two numbers come from two binaries and two operations and the intermediate binary is not on disk. The L27 face-flux budget over the 79: base face within $3.5\text{e-}05$ of the wind-window mean of $\rho v r^2$ (0.9999650 to 1.0000210), face to face spread at most $5.05\text{e-}08$ with a median $1.96\text{e-}09$, operator identity WEN03, numerical flux: HLLC, well balanced: T on all 79. **The seven Roe cases**, each seeded from its L25 state, each `info = 0`, CERTIFIED and ACCEPTED at outer pass 1:

case	seed	R	log10 $\dot{M}$ dot	$\dot{M}$ dot now / L25 [g/s]	EW now / L25 [percent A]
atomic_scalar_gj1132x0.01_kzzprofile/HeH9.7	1132x0.01_kzzprofile/HeH9.7	1.16E-07	5.79	6.1644e+05 / 6.1609e+05	1.3699e-06 / 1.370e-06
atomic_scalar_gj1132x0.01_kzzprofile/HeH9.7	1132x0.01_kzzprofile/HeH9.7	1.16E-07	5.78	6.0017e+05 / 6.0016e+05	3.2459e-05 / 3.2456e-05
atomic_scalar_gj1132x0.01_kzzprofile/HeH9.7	1132x0.01_kzzprofile/HeH9.7	1.16E-07	6.81	6.4138e+06 / 6.4102e+06	0.23532 / 0.23532
atomic_photochem_gj1252x0.01_kzzprofile/HeH9.7	1252x0.01_kzzprofile/HeH9.7	1.08E-07	5.77	5.8235e+05 / 5.8235e+05	3.3576e-05 / 3.3576e-05
atomic_photochem_gj1252x0.01_kzzprofile/HeH9.7	1252x0.01_kzzprofile/HeH9.7	1.08E-07	7.02	1.0543e+07 / 1.0543e+07	0.48218 / 0.48218
atomic_photochem_gj1252x0.01_kzzprofile/HeH9.7	1252x0.01_kzzprofile/HeH9.7	1.08E-07	7.16	1.4347e+07 / 1.4347e+07	0.67111 / 0.67112
atomic_photochem_gj1252x0.01_kzzprofile/HeH9.7	1252x0.01_kzzprofile/HeH9.7	1.08E-07	7.26	1.8194e+07 / 1.8194e+07	0.84287 / 0.84287

The mass-loss rate agrees with L25 to at most $5.6\text{e-}04$ relative and the equivalent width to at most $8.3\text{e-}05$, over states re-solved on a binary carrying L27, L28, L30, L30b and L31 from mapped seeds; the face-flux budget of the seven puts the base face at 0.9999710 to 0.9999793 of the window mean with a spread of at most $2.25\text{e-}07$, operator identity WEN03, numerical flux: ROE, well balanced: T. Both 0.01 cases keep an equivalent width of a few times $1\text{e-}05$ percent A, so L25's statement that the line is gone at 0.01 of the fiducial spectrum stands on the catalog states too. atomic_photochem_gj1132x0.01_kzzprofile/HeH9.7 is the one whose history moves: its direct Roe solve was

refused by L25 on the gated elemental He/H row (2.234e-04 against 1.0e-05 at cell 440), the L4 ladder certified the same wind by continuation, and seeded from that ladder state the catalog case certifies at pass 1.

What was written. Each of the 86 case and rung directories carries its previous output/ as output_pre_L34/, created once and never overwritten, with that case's previous pp.log and He 10830 products inside it; the seven Roe cases had no output/ and so have no backup. Nine input.inp were repaired, the last rung of each flux-closure ladder, where the old post-processing pass had left Do only PP: True and no Solver line so that the file did not reproduce its own state; Solver: Newton and Restart intent: stationary are back and Do only PP is False, each recorded in that rung's REPRODUCE.md. 86 REPRODUCE.md name md5 3146d11b4090306dcea75bb9718edd22, and LHS1140b/MODELS.md sections 7 and 8 were regenerated. Nothing under backup/regression/, src/ or any molecular case directory was touched.

Record. docs/lhs1140b_stationary_L34a_20260918.md.

L36 and L36c: ionization stage transport

L36, stage B: the module exists, tested, with no caller in a run. New src/modules/functions/ionization_stage_transport.f90 (518 lines) holds the stage face flux on the element operator's own element_nucleus_face_flux, its divergence on the L30 geometry, its tridiagonal Jacobian rows and the simplex projection, with one line in SRC. The suite src/tests/ionization_stage_flux/ grew from 15 rows to 25, **25 passed, 0 failed**, the 15 rows of L12b reproducing their values exactly. The seven new GREEN rows include production_stage_flux_sums_to_the_element_flux at 2.0245e-16, stage_divergence_sums_to_the_element_nucleus_divergence at 4.4998e-16, stage_divergence_column_sum_is_the_boundary_flux_difference at 1.2421e-17 against 1e-13, and stage_row_jacobian_matches_a_central_difference at 1.8490e-11 against 1e-7. Three rows are RED by construction and must exceed their floor: a cell-centered volume breaks the column sum at 8.0694e-08, a full mixing-ratio eddy term in every stage breaks the identity at 1.4761e-04 (which is the derivation's central point measured directly, the sum left off the element flux by exactly one copy of the element's own eddy flux), and bounding the closing face value breaks it at 3.6208e-04. Stages C and E were NOT built: the ionization rows are solved inside diffusive_photochemistry.f90 and the configuration is refused in input_read.f90, neither of which the item owned, so the module was linked and exercised by the suite with no caller in a run. Stage D's promotion rule has no precondition for the same reason; the systematic it would be judged against was measured, the He 2³S radial column being 4.9292e11 cm⁻² from the solved state against 8.0114e10 from the _adv profile, a factor 6.15.

The one-velocity approximation is narrow, and the number is now at the site. On the certified atomic fiducial, |w/v| crosses 0.1 at 1.220 R_p for hydrogen and 1.243 R_p for helium with no resonant charge exchange, at 1.231 and 1.273 R_p with it, and the largest |w/v| outside 1.2 R_p is 0.625 and 0.430 without it and 0.518 and 0.336 with it. The momentum transfer is fast, t_{coll}/t_{flow} crossing 0.1 only at 14.51 and 8.25 R_p, and the stages nonetheless drift, because the drift is set by the ambipolar field against the friction and not by the friction alone.

Stage F, the transit side. EXHALE_TRANSIT_STATE selects the input PAIR, adv (the unchanged default) or solution, both files always from one selection, and every saved product carries state_selection, state_files and state_difference in its header. On the certified atomic fiducial the He I 10830 equivalent width is 1.5544 per-

cent A from the `_adv` pair and 20.3950 percent A from the solution pair, a **ratio of 13.12** (L10 reported 13.3 on a different model), with the measured distance between the two states over the 500 rows $7.366\text{e-}1$ in T, $6.313\text{e-}1$ in n(H I) and $9.862\text{e-}1$ in n(He 2³S). That factor is the size of the thing stages C and D move, not the answer they give: on this run the solve's composition is still the local ionization equilibrium. `post_process_adv` is not retired; it stays the independent discretization the solution is measured against.

L36c: the H II row of a run is now the stage row. The unknown is $x(\text{H II}) = n(\text{H II})/n(\text{H nuclei})$, a fraction per hydrogen nucleus, solved as a row of the transport-chemistry operator with the face flux $F(f) = x(f) N_{\text{H}}(f) - n_{\text{H}}(f) K(f) [x(j+1) - x(j)]/dr(f)$, where $N_{\text{H}}(f)$ is the hydrogen nucleus face flux of the element operator; nothing rebuilds a face coefficient of that operator. The bulk-mass `ic_Hp` spelling is retired. `diffusive_photochemistry.f90` (+492/-31) carries the row, its time term, block-Jacobian diagonal, chemistry-column chain rule, limiter and write-back; `binary_element_diffusion.f90` exports `element_nucleus_counts` so that the cell density a row divides by and the face density its flux divides by come from one count; `input_read.f90` (+40/-19) restates the two existing refusals with the real reason and adds two (a mixture with no helium, and Coupled carrier solve: True, whose row registry carries every carrier as a species mass fraction while a stage unknown is a fraction per element nucleus); `certification.f90` (+42/-1) gains the entry `ionization stage nucleus sum`, reported and not gating; `write_setup_report.f90` echoes the carried set. Which halves of N_{H} a run carries is stated in the header: the advective half exactly where the row itself carries the material advection, the element's diffusive half only where the run transports the elements against each other, and the stage's own eddy term in every configuration.

Tests. `ionization_stage_flux` 27 passed, 0 failed, with two new rows: `boundary_stage_rows_fold_the_ghost_onto_the_diagonal` GREEN at $1.8552\text{e-}15$ against $1\text{e-}7$, and `dropping_the_end_face_entry_changes_the_end_rows` RED by construction at $9.0722\text{e-}01$, which is the whole material advection through the boundary face that a purely diffusive end-face rule would drop. A private build with `-fcheck=bounds,do,mem -ffpe-trap=invalid,zero,overflow` runs the two key-on configurations for 40 steps with no runtime trap. Every suite whose driver links a changed object passes, with one RED row, section below.

Movement. Key off, byte-identical: `wasp_full` (400 steps), `mol_diffusion` and `mol_carrier` (12000 steps each) give byte-identical data rows in `Hydro_ioniz.txt` and `Ion_species.txt` against a control built from the entry text, and the two builds agree on every printed step line. Key on, on the three `hp_*` cases the largest movement is $4.2160\text{e-}03$ and $5.2914\text{e-}03$ (`hp_zero_seed`), $3.0318\text{e-}03$ and $3.5450\text{e-}03$ (`hp_trace_seed`) and $3.0201\text{e-}05$ and $4.1478\text{e-}05$ (`hp_front`); $x(\text{H II})$ itself moves by $1.2851\text{e-}10$ to $1.4259\text{e-}07$, and what moves is the helium partition and the neutral hydrogen at the one or two cells that carry the H2 and ionization front. Those goldens were ALREADY stale against the entry text by $3.2967\text{e-}03$ and $3.9439\text{e-}03$ (`hp_zero_seed`) and $2.7650\text{e-}03$ and $3.3068\text{e-}03$ (`hp_trace_seed`), so the refresh those cases need is one refresh and not two.

The key-on stationary solve. `carrier_model_a_newton` is the one fixture that carries Ionization transport: True with Restart intent: stationary, Solver: Newton and Coupled carrier solve: False. **The stage sum identity holds to the rounding of the sums on its solved states, 2.268e-16 on the loaded restart and 2.196e-16 on the state the first JFNK handed back**, against $2.0245\text{e-}16$ on the suite's synthetic column and against the $1.4761\text{e-}04$ to $3.6208\text{e-}04$ the RED rows read when the construction is broken. The state is NOT certified, and it was not on the control either: the carrier balance H+ row stands at $6.432\text{e-}03$ on the measured build against $6.392\text{e-}03$ on the

control, two and a half decades above its $1e-5$ gate in both spellings. No tolerance was changed. The proposed anchor for the new certification entry is $1e-13$, about 500 times the largest rounding measured and nine decades below the smallest break the RED rows produce; proposed, not set.

What remains blocked. $x(\text{He II})$ and $x(\text{He III})$ are not carried and an atomic gas is still refused, both for one reason in three files the item did not own: two flags beside `x_hp_fixed` in `ion_cell_state.f90`, two lines in `impose_transported_ionization_fractions` of `ion_residual_core.f90`, and the gate of the imposed-fraction block in `ionization_equilibrium.f90`. Without the third, the next equilibrium sweep of an atomic run puts the proton back on its local root and the transport leaves no trace, which is why the refusal is correct today and was kept. With those three in place the carrier operator takes the helium rows by adding two indices and three lines, the module already carrying an arbitrary number of stages. Consequently a solve of the atomic fiducial with the key on was not possible, so the He I 10830 equivalent width from a certified transported composition could not be measured and stage D's promotion rule still has no precondition. One RED row is left standing in `src/tests/carrier_retry/:one_further_sweep_leaves_the_returned_composition_within_the_sweep_tolerance` reads $1.034627e-06$ against $1e-6$ on the measured build and $2.685237e-07$ on the control, bound by HeH^+ at $1.33e-12$ of the gas in the base cell, which the fixture counts as present because its threshold is $1e-20$. A diagnostic line naming the species and the cell was added; the tolerance was not touched, and the choice between raising the fixture's threshold and accepting the row as a statement about the sweep's own noise is open.

Record. `docs/lhs1140b_stationary_L36_20260918.md` and `docs/lhs1140b_stationary_L36c_20260918.md`; the key's documentation is the rewritten K15f row of `docs/input_schema.md`.

L36d: both elements' ionization stages carried, and the anchor the stage rows already had

What changed. The carried set becomes $x(\text{H II})$ per hydrogen nucleus and $x(\text{He II})$, $x(\text{He III})$ per helium nucleus, the three fractions of L12a. Each is a transported row of the transport-chemistry operator on its own element's nucleus face flux, each is handed to every ionization sweep, the two helium stages share one simplex and one closing stage, and the certification carries one stage-sum entry per element. `ion_cell_state.f90` (+22/-0) gains `x_heii_fixed`, `x_heiii_fixed`, `x_heii_fix` and `x_heiii_fix` beside the proton's; `ion_residual_core.f90` (+6/-3) writes rows 2 and 3 of `impose_transported_ionization_fractions`; `constrained_chemical_equilibrium.f90` (+32/-1) fixes the two helium species and puts the continuation's seed on the constraint surface instead of on the local root, so that the system it solves is the system it is judged against; `ionization_equilibrium.f90` (+43/-18) moves the imposed-stage block out of the molecular carrier condition onto `ionization_transport` .and. (`bg_ready` .or. `do_load_IC`). `diffusive_photochemistry.f90` takes `n_carrier_max` from 5 to 7 with `ic_HeII` and `ic_HeIII`, gives every stage routine its element (`carrier_stage_element`, `carrier_stage_nucleus_density`), writes `ionization_stage_nucleus_sum` per element with all that element's stages together, and projects the two helium stages in one simplex; `certification.f90` gains ionization stage nucleus sum H and ... He, both reported and not gating; `input_read.f90` restates the three kept refusals and `write_setup_report.f90` echoes the carried set, the flux each rides on and the closing stage of each element. K15f of `docs/input_schema.md` was rewritten in the five places this item makes false. `src/EXHALE_main.f90`, `steady_newton.f90` and `binary_element_diffusion.f90` were not touched.

Tests. ionization_stage_flux **38 passed, 0 failed** (27 on the entry text) and ionization_imposed_fractions **48 passed, 0 failed** (31 on the entry text). None of the new rows can be run against the entry text at all, where ic_HeII, ic_HeIII, x_heii_fixed and x_heiii_fixed do not exist and both suites fail to compile; what stands in their place are the two rows that are RED by construction and keep measuring the two mistakes the construction avoids. operator_stage_rows_ride_on_their_own_element_flux 0.0000 bitwise against 1e-14, operator_stage_sum_is_the_hydrogen_nucleus_flux 2.2113e-16 and operator_stage_sum_is_the_helium_nucleus_flux 3.3525e-16 against 1e-13, helium_stages_are_returned_to_one_shared_simplex 0.0000 against 1e-15; against them charging_helium_to_the_hydrogen_flux_breaks_the_sum reads 1.2614e+00 and projecting_each_helium_stage_alone_leaves_the_sum 4.0000e-01, both of which must exceed 1e-3, the second being a cell holding 1.4 of its helium nuclei in ionized stages. Every other suite that links a changed object passes: carrier_reference_scales 16, steady_species_rows 197, element_operator 40, carrier_returned_state_acceptance 36, certification 84, attempted_step 70, physics_probe 1604, krylov_and_dogleg 334, state_mapper 76, each with 0 failures. carrier_retry keeps one RED row, below; four grid_and_gates route_change_* rows fail on the ENTRY TEXT's refusal and not on anything this item changed (item L36e repairs the fixture), and residual_determinism's closure_spread_within_the_row_tolerance_atomic_elem_newton reads the same 1.715e+04 on the measured and on the control build.

The anchoring measurement L12a section 2.4 and L36c owed, and what it returns. It is made as a row of the stage-flux suite, so it is repeated whenever the suite runs: a manufactured ionization column over [1, 2] whose faces are the same two radii at every resolution, with $n_{el}(r) = 1e10 \exp[-4(r-1)] \text{ cm}^{-3}$, a divergence-free nucleus flux $N_{el}(r) = 1e13/r^2 \text{ nuclei cm}^{-2} \text{ s}^{-1}$, $K(r) = 1e12 r^2 \text{ cm}^2 \text{ s}^{-1}$ and a monotone quintic $x(r)$ with $x' = x'' = 0$ at both ends, so the limiter never fires and the operator's own zero-eddy boundary rule is exact there. The row measure over the interior cells reads 7.2833e-06 at 250 cells ($dr = 4.000e-3 R_p$), 1.0135e-06 at 500 and 1.3327e-07 at 1000, observed order 2.845 / 2.927 / 2.886, the reconstruction's own order, with the advective and eddy halves of the source within a factor 2.38 of each other. Extrapolated with that order to the $1.8447e-3 R_p$ spacing on which cert_tol_carrier_wind = 1e-5 was anchored for a smooth ELEMENT column (docs/certification_tolerance_anchoring_20260910.md section 2 (3)) the stage column reads 7.80e-7, and a decade above it is 7.8e-6. **The rule of the anchoring memo, applied to a measurement of an IONIZATION row, returns the number the stage rows already carry, so 1e-5 is anchored for them and no tolerance was set or changed.** Two readings belong beside it: on the fixture's own Mixed grid the spacing at the gating radius (cert_regime_wind_r = 1.20) is 3.5215e-3 R_p , where the same column extrapolates to 5.04e-6 and the rule would give 5e-5, LOOSER than the gate in place, so the tighter reading stands; and a smooth column is a lower bound on the production discretization, as the anchoring memo says of its own. The stage-sum entry takes no tolerance and is left non-gating, the identity being exact in exact arithmetic: 2.21e-16 (H) and 3.35e-16 (He) on the suite's synthetic 500-cell column and 2.218e-16 and 4.266e-16 on the solved state, against 1.48e-4 to 3.62e-4 when the construction is broken. L36c's proposal of 1e-13 stands as a proposal.

Movement. Against a control built from the entry text of every file of this item (md5 91a0dca3daa3e72cf6ea5b61a16c52d6, the binary L36c delivered; measured build

cf3d2941d1ebe7864f14bd411d5bdfb9), key off and single-threaded: wasp_full at 400 steps, mol_diffusion and mol_carrier at 12000 steps each give byte-identical data rows in both output files. hp_front carries the key and moves by 1.600e-02 in Hydro_ioniz.txt and 1.561e-01 in Ion_species.txt over 100 steps, and the movement is this item's physics: He III 1.561e-01 at cell 205 (2.471972e+00 to 2.086097e+00 cm⁻³), He 2³S and HeH+ 4.368e-02, He II 4.271e-02, H2+ 4.025e-02, H3+ 3.201e-02, He I 2.563e-02, cooling 1.600e-02, heating 3.341e-03, and every other column, H I and H II among them, below 1e-3. The helium ionization partition is what the flow accumulated instead of the local root, and the molecular ions that are functions of it follow. The state stays inside the simplex on the final hp_front state: no species column negative anywhere, x(He II) spanning 5.10e-12 to 0.6500, x(He III) 4.88e-16 to 0.02451 with their sum at most 0.6745, and x(H II) 1.53e-08 to 0.7716. The write-back moves charge and not nuclei: over the 500 physical cells n(He)/n(H) is 0.0793000000 with a spread of 2.945e-12 on the measured build against 2.895e-12 on the control.

The refusals that were kept, and why. The atomic gas is still refused, and the blocker is NOT the imposition gate L36c named, which this item lifts. What remains, all read from the source, is that the operator has no caller in an atomic run (fourteen gate sites on thereis_mol .and. carrier_transport in src/EXHALE_main.f90, the marching operator-split step, the fixed-wind relaxation of the stationary route, the pass cap and eleven diagnostics and progress-control sites), no frozen background (bg_cell and bg_ready are filled only inside the molecular cell state, while carrier_state reads bg_cell(j)%ntot and %T_K for every cell), and no source outside the molecular network (carrier_source takes every stage's chemistry from mol_heh_rows). The refusal message now states the first of the three. Item L36e removes all three.

The one key-on solve, and the row that refuses it. backup/regression/carrier_model_a_newton is the only fixture that admits the key with Restart intent: stationary, Solver: Newton and Coupled carrier solve: False. The state is NOT certified on either build, and the refusing row is carrier balance He+, at 9.99e-1 on pass 1 falling to 5.09e-1 by pass 20 and standing at 7.74e-1 at pass 25, while the control build's carrier balance H+ falls 2.17e-01 to 5.71e-04 over the same passes, every pass hydro info=0 and every pass binding at cell 248, r= 1.2013, the first gated cell. The final certification of the measured build reads carrier balance H2 5.334e-06 (within), carrier balance H+ 1.172e-02 at cell 124 and 2.256e-04 gated, carrier balance He+ 1.000e+00 at cell 107 and 7.736e-01 gated, carrier balance He++ 5.767e-03 and 1.795e-04 gated, ionization stage nucleus sum H 2.218e-16 and ... He 4.266e-16, and every hydrodynamic row within (4.617e-11 / 5.442e-13 / 3.355e-09); the control's carrier balance H+ is 5.165e-02 with 5.708e-04 gated, so **the proton row is BETTER on the measured build at the gated cell and the He+ row alone refuses**. Read term by term from the row record, the He+ row is carried entirely by its chemistry, whose net source is 68 times the stage flux divergence at the gated cell and 5.0e4 times it in the shielded layer, while the two builds' helium compositions agree to 0.20 per cent at cell 248 (1.2951e+03 against 1.2977e+03 cm⁻³) and 0.23 per cent at cell 107. The same operator evaluating the same rows on the CONTROL's own state, so that only the state differs, gives carrier balance He+ 4.151e-02 against 7.736e-01, a factor 19 lower and still three and a half decades above its 1e-5 gate: the row is not broken, and on this fixture no composition either route produces certifies it. The scale carrier_residual divides by is built from the NET source, and for a stage whose production and loss cancel to three decades the net is the same size as the residual. That is the one thing that decides whether the He+ row can ever certify, it is a change to a shared acceptance interface with its own anchoring, and it was not made

here. It is item D3 of docs/PLAN_20260918_rev1.md.

The carrier_retry RED row, diagnosed. one_further_sweep_leaves_the_returned_composition_within_the_sweep_tolerance reads 1.044879e-06 against $\text{ieq_res_tol} = 1\text{e-}6$ on the measured build, 1.034627e-06 on the entry text and 2.685237e-07 one increment before it, the binding species being HeH^+ at $1.33\text{e-}12$ of the gas in the base cell. The verdict is **a seed that has not settled, in a self-coupled trace-ion chain, not the imposition of the transported fractions and not the sweep's own noise**, from four diagnostic measurements added to the fixture with no assertion and no tolerance touched. (1) At the binding cell the further sweep moves H I by $1.32\text{e-}11$, H II by $1.39\text{e-}16$, He I by $2.30\text{e-}16$, He II by $2.13\text{e-}16$ and H2 by $1.86\text{e-}16$, while H2+ moves $1.47\text{e-}08$, H3+ $1.05\text{e-}07$ and HeH^+ $1.04\text{e-}06$, a chain with a factor 7 to 10 per link whose every parent moves at $1\text{e-}11$ or below. (2) The chain closes on itself: with $n_e = 1.197\text{e+}05$, $\text{H}^+ = 5.636\text{e+}04$, $\text{H2}^+ = 3.450\text{e+}01$, $\text{H3}^+ = 6.312\text{e+}04$ and $\text{HeH}^+ = 1.226\text{e-}02\text{ cm}^3$, H3+ carries 53 per cent of the electrons and the imposed proton the rest, so re-solving the trace molecular ions moves the electron density their own balances read. (3) It settles: a THIRD sweep moves the composition by $1.0085\text{e-}10$, four decades below the second, where arithmetic noise would repeat. (4) It is not the imposition: the same sweep with the transported fractions NOT imposed moves the composition by $1.5067\text{e+}08$, its worst being He III at cell 4, the local root of the helium ionization there being eight decades from the transported value. The tolerance and the $1\text{e-}20$ presence threshold were not touched; the returned-state closure contract this row reads on is item D4 of docs/PLAN_20260918_rev1.md.

One defect fixed beside the stage work. carrier_ionization_stage_projection wrote into an unallocated cell record (pct_cell_constrained, which only limit_to_element_budget allocated), so a caller that projected on its own segfaulted; the routine now allocates it. Found by the new acceptance row that calls it directly.

Record. docs/lhs1140b_stationary_L36d_20260918.md. Goldens were not refreshed and EXHALE.x was not written; hp_front's golden is now stale by $1.6\text{e-}1$ in the helium columns and joins the two L36c reported.

L36e: the atomic run carries the three stages, and the certified state answers stages D and F

Verdict: the atomic run carries the three ionization stages, and the state it converges is CERTIFIED. On the certified atomic fiducial LHS1140b/models/.L26/fid_resolve with Ionization transport: True through the partitioned stationary route, the outer iteration accepted at **pass 56** with $\text{info} = 0$, the three stage rows at $8.620\text{e-}07$ (H^+), $7.147\text{e-}07$ (He^+) and $2.360\text{e-}06$ (He^{++}) against $1\text{e-}5$ and the two stage-sum entries at $2.161\text{e-}16$ and $3.078\text{e-}16$.

What changed. ionization_equilibrium.f90 (+124/-88) gains the public predicate transported_rows_exist(), "the transport-chemistry operator has a row to solve", and puts the frozen background $\text{bg_cell}(j) = \text{ieq_cell}$ and bg_ready on it instead of on thereis_mol , with the four keep, adopt and install routines of the stationary background; the cell temperature and gas particle density are set for any gas and the ionization-stage flags and imposed fractions move out of the molecular cell state. src/EXHALE_main.f90 (+24/-18) replaces thirteen gate sites on thereis_mol .and. carrier_transport by that predicate, two sites keeping the molecular condition and now saying why. diffusive_photochemistry.f90 (+135/-31) states the split in carrier_set_init (molecular carriers where the network exists AND is transported, the three stages on their own key), puts the operator's three entry conditions on the predicate, takes the atomic H and He rows from heh_tr_rows , adds the He/H charge exchange in both gases, and makes the row record print the scale the acceptance divides by. input_read.f90 (+42/-33) lifts the atomic refusal and keeps, with its real reason, the refusal for a molecular gas whose carriers are not transported; write_setup_report.f90

(+34/-20) moves the ionization-transport block out of the molecular one and names the source of a stage row in each gas. Four files outside the item's list were required and are reported as their own item: certification.f90 (+6/-5, four gate sites, without which an atomic key-on run's certification would print the three carrier balance entries as "the run does not carry this species", skip both stage-sum entries, and let the progress control call a state certified on the hydrodynamic rows alone), load_IC.f90 (+21/-13, iontrans removed from opt_changes_layout), carrier_retry/carrier_retry.f90 (+26/-6) and the carrier_returned_state_acceptance fixture (+9/-2).

Tests. The new src/tests/grid_and_gates/ionization_transport_atomic.sh, registered as iontrans_atomic, is GREEN on the measured build and RED on the control on its first four rows: ionization_transport_atomic_runs PASS at rc = 0 with no refusal against FAIL at rc = 1 refused in input_read; ..._setup_names_rows, ..._stage_sum (2 entries against 0) and ..._rows_measured (3 rows evaluated against 0) likewise. The fifth row, ionization_transport_molecular_local_refused, passes on both builds, which is the refusal that is KEPT. **The four grid_and_gates route_change_* rows L36d left failing are GREEN on BOTH builds:** their block tests the route token and nothing about the ionization stages, and it built its rungs from carrier_model_a_newton, which carries Ionization transport: True, so rung B's Coupled carrier solve: True hit L36c's refusal; the block now runs on carrier_elem_newton, which admits both routes and carries no ionization key. That the rows pass on the control build too is how one can tell the fixture was the defect; the refusal was not lifted. Every suite whose driver links a changed object passes: ionization_stage_flux 38, ionization_imposed_fractions 48, steady_species_rows 201, element_operator 40, certification 84, attempted_step 70, state_mapper 76, grid_and_gates (restart_option_change + iontrans_atomic + carrier_transport_inert) 29, steady_completion_flag 3 and steady_selfconsistent_residual 7 on the certified run's own log, each 0 failed. carrier_retry keeps its one RED row and coupled_block_jacobian two, both on the measured and on the control build.

Movement. Control cf3d2941d1ebe7864f14bd411d5bdfb9 (the binary L36d delivered), measured 6a36529386102b37a38cb40fde0a611c. Key off, byte-identical: wasp_full at 400 steps and mol_diffusion and mol_carrier at 12000 each give byte-identical data rows in both output files and identical step lines. The one key-on fixture, carrier_model_a_newton, moves by 4.2725e-05 in Hydro_ioniz.txt and 1.2479e-04 in Ion_species.txt, and that movement is the He/H charge exchange (Huang et al. 2023 Table 4 group B) now entering rows (1) and (2) of the stage sources: the largest movers are n(He II) 1.2479e-04, n(H I) 2.5916e-05, n(He III) 1.6337e-05 and n(HeH+) 1.1289e-05, exactly the species that pair couples.

The 25-pass front. Twenty-five outer passes, the bound the brief named, are not enough, and the reason is a relaxation front and not a refusing row. Over passes 1 to 25 the worst gated species row falls only 3.51e-02 to 1.76e-02 while the cell that binds it marches OUTWARD from 217 to 491, about 3.3 cells a pass: the restart's local-equilibrium ionization is being replaced by the transported partition and the front has to cross the column first. It reaches the outer boundary at pass 26, and from there the measure falls monotonically, 1.75e-02 at 26, 1.27e-02 at 40, 8.73e-03 at 45, 2.03e-03 at 50 and 8.68e-06 at 56, where the state is accepted. At 25 passes the refusing rows are carrier balance H+ 9.209e-03 at cell 445, He+ 6.622e-03 at 464, He++ 1.760e-02 at 491 and elemental transport He/H partition 2.256e-05 at 339; the transport divergence stands 13 to 29 times the net chemical source at those cells and the residual IS that divergence, while on the certified state the two balance to eight digits. **A campaign that judges this configuration at 25 passes reads a front and not a solution.** On this configuration the row scale that refuses L36d's hot Uranus does

not bite: the He+ row certifies four decades under its gate.

The certified state. carrier balance H+ 8.620e-07 at cell 150, He+ 7.147e-07 at 177, He++ 2.360e-06 at 183, ionization stage nucleus sum H 2.161e-16 at 294 and ... He 3.078e-16 at 166, elemental transport He/H partition 8.679e-06 at 500, level balance He 2³S 7.263e-21 at 442, eliminated-species closure System_HeH_TR 5.043e-15 at 495, and the hydrodynamic rows at 5.670e-10 / 9.788e-12 / 1.045e-08. The stage-sum identity holds to roundoff on a solved state, against the 1.5e-4 to 3.6e-4 the suite's RED rows read when the construction is broken.

Stage D is closed, on the negative side. On the certified state, the first state whose three ionization fractions meet their own gate, the He 2³S radial column over 1 to 29.03 R_p reads 8.0846e+10 cm⁻² from the solution against 8.0949e+10 from its _adv profile, a difference of **0.127 per cent**, twenty times below the 5 per cent promotion rule of L12a section 3: **the He 2³S level is NOT promoted to the carried set.** For contrast the same column of the local-equilibrium closure reads 4.9257e+11 against its _adv 8.0109e+10, the factor 6.15 L36 measured, reproduced here.

Stage F, and what the factor 13.12 was. The He I 10830 equivalent width, by the definition that reproduces L36's two numbers to four figures:

state	EW [percent A]	max depth
local-equilibrium solution (key off)	20.3950	66.1194
its _adv profile	1.5544	5.1959
transported solution at 25 passes (not certified)	2.9823	10.4172
transported solution, certified	1.3678	4.5745
its own _adv profile	1.4053	4.7338

Behind them, $x(\text{H II})$ reaches 0.0583 on the certified transported state against 0.6657 on the local-equilibrium one, and $x(\text{He II}) + x(\text{He III})$ reaches 0.1888 against 0.9117. **L36's factor 13.12 between the solution and its _adv profile is a property of the local-equilibrium closure, and carrying the stages removes it:** the transported solution and its own _adv profile differ by 2.7 per cent.

The validity range, restated beside the result. The ambipolar drift of an ion against its own neutral stays below 0.1 of the bulk velocity only inside 1.220 R_p (H) and 1.243 R_p (He) without the resonant charge-exchange channel and 1.231 and 1.273 R_p with it; over $r \geq 1.2$ R_p it reaches 0.625 and 0.430, and 0.518 and 0.336 with the resonant channel. The momentum transfer is fast ($t_{\text{coll}}/t_{\text{flow}} < 0.1$ inside 14.51 R_p for H and 8.25 R_p for He), so the stages are collisionally locked in TIME SCALE and still drift; above about 13 R_p the continuum equation is itself unvalidated by the run's own Knudsen measure (max Kn 0.88). The certified state therefore solves an equation whose one-velocity closure is exact only inside about 1.2 R_p, with a neglected drift of order 0.02 to 0.6 of the advective stage flux over the range the stages are carried.

Six defects found and fixed, a separate item from the stage work.

1. **An out-of-bounds write in the update-map diagnostic** (EXHALE_main.f90): res_um_all and terms_um_all were allocated (1:N,4) and handed to a dummy of (1:N,n_carrier_max=7) whose first statement zeroes the whole dummy, 3N doubles past the end of both arrays, wrong since the carrier set grew past four. Allocated by

n_carrier_max.

2. **The gas particle density was zero in every atomic run** (ionization_equilibrium.f90): calc_ntot was called only under thereis_mol, so an atomic run's frozen background carried $ntot = 0$, the pressure its sweep retry forms was the electron pressure alone, and the operator's $wfac = \rho n_0 / \max(n_{tot}, 1e-99)$ would be of order $1e99$ for any row that read it. The stage rows do not read it, which is why nothing measured here moves. The particle density is now formed in every gas from the one stoichiometric sum, and wasp_full (atomic, helium, He 2³S and metals) is BYTE-IDENTICAL across the change, so no existing run moves.
3. **The carrier row record did not print the scale the acceptance divides by**: the writer rebuilt a scale from the flux DIVERGENCE while carrier_residual builds it from the two face fluxes by their magnitudes. At the H+ binding cell of the 25-pass state the record read 0.934 where the certification read $9.209e-03$ for the same row of the same state, a factor 101 from the cancellation of the two faces. The record now carries row_terms_phys itself and reproduces the certification's three gated numbers exactly. (L36d's diagnosis table quotes that record's old, divergence-based measure; its gated numbers from the certification are unaffected.)
4. **The proton's record invented no split and printed none**: in an atomic gas heh_tr_rows returns the rows and not their halves, so the H+ production and loss were both written as zero while the row's source was not. The proton now carries its split where mol_heh_rows hands one out and its net on the side its own sign falls where none exists.
5. **iontrans was classified as a layout-changing restart token** (load_IC.f90), which refused the restart in both directions and made the two notes printed beside it unreachable. It is not one: n(H II), n(He II) and n(He III) have a column in every state file this code writes. The four tokens that DO change the columns (metals, mol, oxychem, carrier) are unchanged. Without this the fiducial could not have been restarted with the key on at all.
6. **The He/H charge exchange was missing from the stage row sources**, while every ionization system carrying helium adds it to rows (1) and (2) and it is on by default. Fixed in both gases; this is the $1.2479e-04$ movement of carrier_model_a_newton. Three statements naming the proton a molecular carrier were corrected in place with it.

What is still absent, and where it belongs. The metal charge exchange is absent from the stage row sources: cx_add_to_fvec writes group A (metal + H/H+) into row 1 and groups C and D into rows 2 and 3 of every metals system, and carrier_source does not call it, so in a run WITH metals the stage sources and the sweep's rows differ by those terms. The fiducial is metal-free, so nothing measured here is affected; adding it needs the frozen metal densities per stage and the cx_metal_base layout of the system being judged, neither of which the transport operator holds. **It is item D7 of docs/PLAN_20260918_rev1.md.**

Four key-on goldens are stale: hp_front, hp_zero_seed and hp_trace_seed from L36c and L36d, and now carrier_model_a_newton by $1.2e-04$. One refresh at the close of the series; none taken here.

Record. docs/lhs1140b_stationary_L36e_20260918.md; the key's documentation is the K15f row of docs/input_schema.md, rewritten again around the atomic configuration, the row source in each gas, the refusal that is kept and iontrans as a nameable restart token.

L37: the round trip of the molecular base mass row, and the probe that can be pointed at a state

Verdict: the molecular base's re-entry band is a decade above the atomic one, and it is a transient of the re-entry and not a property of the state. On the six certified molecular states in hand the cell-1 hydrodynamic mass row comes back from a single no-step re-entry at **2.39 to 10.09 times** the value the run printed for the same cell, and at 16.0 to 31.2 times that cell's own rounding floor, where the tolerance is ten floors. On three certified ATOMIC states of the same planet, the same grid and the same K_{zz} the ratio is **0.68 to 1.35** and the row stands at 0.21 to 1.52 floors, which is L18's 0.78 to 1.71 band met again. Every measurement was taken with LHS1140b/models/EXHALE_3146d11b.x, md5 3146d11b4090306dcea75bb9718edd22, single-threaded on scratch copies; no tolerance, anchor or key default was changed.

state	cell	in-run	E1, one re-entry	E4, settled	floor	E1/in-run	E1/floor	E4/floor
heh213_new	1	5.213e-09	1.2475e-08	5.5734e-09	7.8046e-10	2.39	15.98	7.14
heh213_pre	1	3.616e-09	1.7031e-08	3.0823e-09	7.6854e-10	4.71	22.16	4.01
heh97_new	1	2.319e-09	2.3398e-08	1.7066e-09	7.4894e-10	10.09	31.24	2.28
heh97_pre	1	2.330e-09	1.6456e-08	3.7884e-09	7.4264e-10	7.06	22.16	5.10
heh055_new	1	1.065e-08	2.5454e-08	5.0843e-09	1.2867e-09	2.39	19.78	3.95
i3_alt	1	3.249e-09	1.4863e-08	6.3734e-11	7.8046e-10	4.57	19.04	0.08
at213	1	1.189e-09	1.2016e-09	1.1852e-09	7.9044e-10	1.01	1.52	1.50
at97	2	1.481e-09	1.9924e-09	1.9924e-09	7.9345e-10	1.35	2.51	2.51
at055	1	7.414e-10	5.0212e-10	5.0212e-10	6.7147e-10	0.68	0.75	0.75

The settled readings. Feeding the evaluation its own written state again and again, the row settles: four of the six molecular states reach a reading that repeats exactly by the third or fourth re-entry, and at that reading cell 1 stands at 0.08 to 7.5 floors, **inside the existing anchor on every one of the six**. heh213_new goes 1.2475e-08, 7.0724e-09, 5.5734e-09, 5.5734e-09; heh97_new 2.3398e-08, 2.3132e-09, 1.7066e-09, 1.7066e-09; i3_alt 1.4863e-08, 1.1231e-09, 6.3734e-11, 6.3734e-11. Two do not settle at cells the verdict does not take, heh213_pre and heh055_new. The atomic states are at their settled reading on the FIRST re-entry, at97 and at055 reproducing their own output exactly at every one of the ten cells and at213 alternating between two values 1.4 per cent apart. The settled molecular reading also agrees with what the run itself printed, 5.5734e-09 against 5.213e-09 on the reference solution, a factor 1.07. The

measurement also identifies what L34b's third reading was: EXHALE_RESIDUAL=1 equilibrates the loaded composition before measuring, so with EXHALE_RELOAD_EQ=0 it is the evaluate route to every printed digit and with the default it is the SECOND re-entry, 7.0724e-09 on heh213_new, the 7.072e-09 L34b reported. Anything quoting the residual diagnostic beside the evaluate route should say which of the two it ran.

The mechanism: the base ghost composition and the caloric equation of state. The physical cells round-trip exactly, ρ , v and p of cells 1 and 2 coming back to all 17 digits and the composition of cell 1 to 1.7e-16. What does not come back is the pair of inner GHOST rows: ghost -1 moves by 2.90e-11 in v , 1.82e-11 in p and 9.89e-12 in T , ghost 0 by 2.90e-11 in v , and in the composition file ghost -1 by 9.139e-06 (the He 2³S column) and ghost 0 by 4.835e-08 (H I). Exchanging the two halves shows they are the carrier: giving the settled state the ARCHIVED ghost composition rows reproduces the archived reading 1.2475e-08 exactly, and giving the archived state the settled ghost composition gives 7.0724e-09. Scaling one ghost column by $1 + \epsilon$ makes the response linear with **slope 0.30 to 0.33** of the row's own scale per relative unit of the ghost H2 (3.3171e-07 at $\epsilon = 1e-06$ and 3.5562e-08 at $1e-07$, against 5.5734e-09 unperturbed), so a ghost H2 count off by $1e-08$ is four rounding floors of the cell-1 mass row and the measured ghost drift is of that size. The atomic response is unmoved at the printed precision up to $\epsilon = 1e-06$, a hundred times the molecular perturbation that matters, and the ghost pressure and velocity columns are inert in both, the base boundary rebuilding them. The reason is in the pressure map: where caloric_mixture_active is false the map is $(\gamma_{ad} - 1) \rho e$ and carries no composition, while the inner ghost of these molecular columns is $2 n(H_2)/n_H = 0.99998$, so its pressure, its sound speed and therefore the base face flux of the HLLC solve are functions of its H2 count; at a base Mach number of $3e-05$ a part in $1e8$ of the face state is a part in $1e3$ of the difference of two face fluxes. The Mach number is not what separates the two groups (molecular base cells run at $2.7e-05$ to $3.1e-05$ and atomic ones at $1.8e-06$ to $1.1e-05$, so ϵ/Mach is if anything smaller in the molecular base, and the rounding floors agree within a factor 2 across all nine states); what differs is what a base face flux is a function of. The composition refresh of the physical cells is innocent: no key skips the one equilibrium sweep, and the sweep moves the physical particle count by $1.295e-13$, which at the amplification above is $4e-14$ of the row, five decades below the $6.9e-09$ the re-entry moves it.

c_round = 100, proposed for the record and not adopted. Applying anchor (6) of docs/certification_tolerance_anchoring_20260910.md with the re-entry as the perturbation, the largest ratio is 31.24 and P16's own margin of 3.7 above it gives 116, a round **100** where the code carries 10. **The rule does not make that choice unambiguous**, because it fixes the margin above a measured ratio but not which perturbation the ratio is measured under: one ulp of the density is a perturbation the arithmetic forces on any evaluation and no re-evaluation removes it (P16 measured 1.5931, 1.8857, 2.6806 and 1.7733 on four states), while the restart round trip is not forced and two further re-entries of the same state remove it. Anchoring on the second would widen a gate to admit a restart that does not reproduce its own base ghost, which the standing rule forbids, so the number is reported and NOT proposed for adoption. The three ways out stated for the user are to make the restart carry its base ghost (at which the existing $c_{\text{round}} = 10$ passes every molecular state measured here with a margin of 1.3 to 125), to make the certification's input the route's own fixed point (a contract change, two extra no-step evaluations at about 7 s each), or to re-derive c_{round} on the round-trip perturbation. The contract question the measurement leaves is not about the sweep but **what a restart pair must carry so that the base face flux of a molecular column is the same flux after a write and a read**, and it is item **D5** of docs/PLAN_20260918_rev1.md. Two certifications inside one run, of one

conserved state, already differ by a factor 4.3 at the base cells while agreeing to four digits at the verdict cell, so the re-entry is where the movement is measured and not where it begins.

The probe can now be pointed at a state. `src/tests/physics_probe/molecular_energy_recipients.f90` (+745/-2 against the last commit, of which the reader is this item's) held its three representative cells as `real*8`, parameter arrays transcribed by hand, so every re-solve of the molecular fiducial made the L31 uncertainty bracket stale. It now reads them: `EXHALE_PROBE_STATE` names a directory and the pair read is `{Hydro_ioniz_IC.txt, Ion_species_IC.txt}`, the state a run hands back; unset, the compiled values stand and nothing is read, so the suite runs with no state in the tree. The species columns are taken BY NAME from the file's own # columns header, and the electron density and particle count are formed as the code's mass policy forms them. The three cells are a rule: molecular depth is physical cell 1, the H2 front is the first cell where $2 \cdot n(\text{H}_2)/n_{\text{H}}$ has fallen to half its value at cell 1 (L31 section 5.4's own definition), and the dilute upper column is the cell nearest `r_dilute_upper_column = 1.8422600916216840 R_p`, which is the choice itself, no threshold on the profile picking that cell because $2 \cdot n(\text{H}_2)/n_{\text{H}}$ is flat and rising outward there (3.43e-06 at cell 296 to 3.88e-06 at cell 303). Pointed at `HeH2.13/output_pre_L34` the reader reproduces the compiled values at **0.00000E+00 over all 33 numbers**, cells 1, 178, 299 at 808.319, 2128.367 and 5083.438 K; pointed at the L34b solution it gives cells 1, 177, 299 at 778.79, 2116.61 and 5061.59 K, a distance of 3.41525E-01 from the compiled values, and reproduces L34b block G (the helium third body at the depth +42.05 / -21.07 per cent of a nominal heat of 1.04863e-06, against L34b's +42.0 / -21.1; the R5/R16 recipient at the front +4.37 per cent of 8.60218e-09, against +4.4). Pointed at an atomic state the driver stops with FAIL `probe_state_has_no_column_H2` and at a directory with no state pair with FAIL `probe_state_file_unreadable`, rather than reading a zero. Suite with the variable unset: `src/tests/physics_probe/run.sh` **1604 PASS, 0 FAIL**, this driver contributing the 81 assertions L34b counted; the driver alone, pointed at each of the two molecular states, 81 PASS, 0 FAIL. No control run of the entry text was taken, because the file in the tree already stood ahead of the last commit through other items' work.

Record. `docs/lhs1140b_stationary_L37_20260918.md`, which carries every table in full. `docs/lhs1140b_stationary_L31_energy_cycles_20260917.md` section 5.4 now states that no rule drawn from the profile picks the dilute upper column and that the radius is the choice.

L34b: the molecular reference solution, certified at pass 12

Verdict: `molecular_scalar_gj1132_kzz1e9/HeH2.13` **CERTIFIED at outer pass 12**, `info = 0, \|\R\|\| = 1.417e-08`, in 40 m 35 s at 8 threads, seeded from its own archived certified state, which the current binary refuses on the energy row at 4.267e-01 at cell 203 exactly as L29 records. `HeH9.7` certified the same way at **pass 10** in 25 m 35 s. Binary 3146d11b4090306dcea75bb9718edd22 for every run; no source file was edited, no tolerance weakened, nothing under backup/regression/ touched. Of the other four cases of the step, `kzz1e9/HeH0.55` ran 6 passes in 3 h 35 m and did not solve, `photochem/HeH9` one pass at each of two pseudo-time starts in 1 h 04 m, `photochem/HeH2.09` no completed pass in 1 h 55 m, and the three `molecular_scalar_gj1132_wellmixed` cases were not started, the L33 ladder they need being five rungs at 25 passes each. Each case that did not solve carries a `not_solved.md` in the catalog's form.

The acceptance table of the reference solution.

acceptance item	answer
info = 0	yes, $ R = 1.417\text{e-}08$ at pass 12
CERTIFIED on every gated row at the existing tolerances	yes: mass at binding cell 188, 0.919 of its anchor; momentum $5.588\text{e-}12$ of $1\text{e-}08$; energy
the L27 face-flux budget	printed on the solved state: WENO3, HLLC, well balanced; window mean $5.73832\text{e-}07$
the energy balance closed cell by cell	yes, by the code's own operator with no step taken: cell-wise maximum $1.4464\text{e-}08$ at cell
the same balance assembled outside the binary	from Heating_breakdown.txt, Cooling_breakdown.txt and the hydro columns through
the L31 collider sum on the solved state	yes: chemistry against the routine 0.0 exactly, the carrier row's R15 channel 0.0 exactly,
the L31 uncertainty bracket on three of its cells	yes, on the solved state's molecular depth (778.79 K), H2 front (2116.61 K) and dilute u
the probe taking a state	it does not. The three cells were compile-time arrays, so the blocks were run through a
the He I 10830 EW through run_case.sh	1.5603 percent A over 10832.60 to 10834.20 A (air), R = 68 000 WINERED HIRES-Y
passes and wall clock	12 outer passes, 40 m 35 s at 8 threads

The comparison, archived against new, with the L31 uncertainty beside it.

molecular_scalar_gj1132_kzz1e9/HeH2.13:

quantity	archived (c2e9c9990b9f)	this item (3146d11b4090)	change	L31 uncertainty
log10 Mdot [g s ⁻¹]	7.90907	7.90239	-1.53 per cent in Mdot	--
He I 10830 red-pair EW [percent A]	1.57583	1.56032	-0.99 per cent	--
red-pair depth [per cent]	5.62817	5.57130	-1.01 per cent	--
FWHM [A]	0.262836	0.262836	0.00 per cent	--
base T [K]	808.319	778.792	-3.65 per cent	set by the base heat below
base heating rate [erg cm ⁻³ s ⁻¹]	1.12791e-06	1.06477e-06	-5.60 per cent	+42.0 / -21.1 per cent (helium third body)
2 n(H2)/n_H at cell 1	0.355303	0.335357	-5.61 per cent	the same
H2 front, x2 = 1e-2 [R_p]	1.16256 (cell 205)	1.15973 (cell 204)	one cell	--

molecular_scalar_gj1132_kzz1e9/HeH9.7:

quantity	archived	this item	change	L31 uncertainty
log10 Mdot [g s ⁻¹]	7.94648	7.94282	-0.84 per cent in Mdot	--
He I 10830 red-pair EW [percent A]	2.42118	2.40606	-0.62 per cent	--
red-pair depth [per cent]	7.68280	7.62903	-0.70 per cent	--
FWHM [A]	0.295123	0.295627	+0.17 per cent	--

quantity	archived	this item	change	L31 uncertainty
base T [K]	567.526	522.193	-7.99 per cent	set by the base heat below
base heating rate [erg cm ⁻³ s ⁻¹]	6.31976e-07	5.51060e-07	-12.80 per cent	+42.0 / -21.1 per cent
2 n(H2)/n_H at cell 1	0.0779932	0.0675867	-13.34 per cent	the same
H2 front [R_p]	1.05626 (cell 145)	1.05333 (cell 142)	three cells	--

Two readings of the pair. **The L7g corrections land in the base and the colder the base the larger they are**, -3.7 per cent in base temperature at He/H = 2.13 and -8.0 per cent at 9.7, against -1.5 and -0.8 per cent in the mass-loss rate and -1.0 and -0.6 per cent in the equivalent width: the line forms far above the layer that moved. And **the uncertainty of the base chemical heat is eight times the change the re-solve made in it**, the helium third body's +/-0.3 dex moving that heat by +42 and -21 per cent against the re-solve's -5.6 per cent; that is what the memo quotes beside every molecular number. The reference solution also reproduces the L22 I3 solution on the predecessor binary to four digits (base T 778.79 against 778.792, front 1.1597 R_p at cell 204 against 1.15973 at cell 204, log10 Mdot 7.9024 against 7.90239, EW 1.5599 against 1.56032), the largest difference being 0.03 per cent in the equivalent width.

The one acceptance item that does not come back clean. The no-step evaluate route `run_case.sh` runs after the solve REFUSES the state on one entry of the seven: the hydrodynamic mass row at cell 1, 1.248e-08 against that cell's own rounding anchor 7.8e-09, the accepted pass having printed 5.213e-09 there, a factor 2.39 across the re-entry against the 0.78 to 1.71 band L18 measured over 120 certified ATOMIC states. Six of the seven come back where they were, and HeH9.7 behaves the same way (2.340e-08 against 7.5e-09). Cell 1 carries $v = -1.9 \text{ cm s}^{-1}$ against a wind of $1.2 \times 10^5 \text{ cm s}^{-1}$ and lies far below the $r \geq 1.20$ window every gated species row is judged in, so this is the cancellation of the base and not a second opinion about the wind. Item L37 measures the band and its mechanism; the contract question is item **D5** of docs/PLAN_20260918_rev1.md.

What was written. `models/status.py` --write regenerated MODELS.md sections 7 and 8; `LHS1140b/make_memo_figures.py` regenerated the twenty docs/figures/lhs1140b_*.pdf, of which twelve changed when compared page by page, most of them drawing ATOMIC cases whose state files were rewritten in the same hours by the L34 steps 1 and 2 worker, so those changes are not claimed here; `latexmk -pdf docs/lhs1140b_exhale_vs_pwinds.tex` built cleanly at 80 pages, the same count as before. Also written: three `not_solved.md` files, two `REPRODUCE.md` written by their own runs and naming `md5 3146d11b4090306dcea75bb9718edd22`, `output_pre_L34/` beside each of the five cases touched, and `docs/figures/lhs1140b_L34b_reference.png`.

Record. docs/lhs1140b_stationary_L34b_20260918.md.

L34c: the remaining molecular cases, and the two seeding routes the restart contract refuses

Verdict: none of the six cases certified, one new certified solution was produced as a seed, and two of the three seeding routes the item was offered are inadmissible by the restart contract, which is the substantive finding. Binary 3146d11b4090306dcea75bb9718edd22, verified before the first run; the only source file changed is

models/status.py.

case	seed	passes	wall clock	verdict and where it stopped
molecular_scalar_gj1102_kzz1e9/HeH0.55	6	6	6 h 01 m	not solved; carrier row 8.40e-01 to 7.09e-02 at pass 5 and back to 1.19e-01, hydrodynamic solve info = 2 at every pass; stopped at the ceiling
molecular_photochem_gj1102_kzz1e9/HeH0.55	6	6	37 m 41 s	not solved; REFUSED, the element relaxation found no admissible advance and restored its entry composition; mass row 2.848e-01 at cell 2, energy 3.167e-01 at cell 43
molecular_photochem_gj1102_kzz1e9/HeH2.13	6	6	6 h 00 m	not solved; carrier row a band, 1.85e-01 to 4.42e-01, ending 3.07e-01; stopped at the ceiling
molecular_scalar_gj1102_wellmixed/HeH2.13	7	7	6 h 01 m	not solved; wind converged from pass 3, carrier row flat at 4.29e-01 from pass 5; stopped at the ceiling
molecular_scalar_gj1102_conversion/HeH0.55	31	31	6 h 01 m	not solved; wind converged from pass 5, carrier row to 3.76e-02 at pass 11, up to 1.56e-01 at 21, down to 6.48e-02 at 31; stopped at the ceiling

case	seed	passes	wall clock	verdict and where it stopped
molecular_scalar_gj1132_wellmixed/HeH0.08340 + 40	the certified atomic pair	2 + 4	5 h 19 m	not solved; REFUSED on its own terms, both budgets spent. Wind within every tolerance (mass 2.005e-08 of 2.8e-08, momentum 5.509e-13, energy 3.110e-08 of 1.0e-06), carrier balance H2 9.264e-02 of 1.0e-05
.L34/rung1, rung 1 of the L33 ladder	the certified reference, --reservoir He/H 1.411074554655	2 + 4	6 h 00 m	did not certify, which ENDS the ladder by its own acceptance; rungs 2 to 4 were created and not run
.L34/atomic_wm2.13	atomic_scalar_gj1132_wellmixed/HeH1, reservoir carried		2 m 44 s	CERTIFIED , info = 0, R = 2.476e-08, log10 Mdot 7.86, He I 10830 red-pair EW 1.9533 percent A

The two routes that are not admissible. (1) **A metal-free scalar molecular state cannot seed a photochem case.** load_IC refuses it: "metadata field reservoir: this run carries a reservoir for C/H (2.778025321E-04) and the restart files do not". A scalar case is metal-free and states one ratio, a photochem case states He/H, C/H, N/H and O/H at the matching level, and no factor makes a carbon reservoir out of a state that has none; both profile cases were therefore seeded the way MODELS.md section 3 defines for that group, from the certified wind of their own atomic closure rung. (2) **A diffused molecular state cannot seed a well-mixed run**, so the L33 ladder's element-treatment step is impossible on a molecular state. With Restart option change: he_diff in place, so that the option difference itself was admitted, load_IC still refuses: "the restart file does not carry the He/H the input asks for, and the state carries HeH+, whose nucleus of each element cannot be rescaled by one factor", the reference state's He/H running from 2.13 at the base to 6.042513123E-01 at cell 500. The same clause forbids a composition step INSIDE the well-mixed family, so the only entry to a molecular well-mixed case is the atomic-to-molecular conversion of an atomic well-mixed wind at that composition. The catalog's atomic ladder has no He/H = 2.13 rung, so **one was solved for the purpose, .L34/atomic_wm2.13, certified**, and wellmixed/HeH2.13, which MODELS.md recorded as never started, was started for the first time. The measured bound on how far a well-mixed molecular state can be carried at all: map_state_to_grid.py --reservoir He/H accepts the 2026-09-16 wellmixed/HeH0.55 state carried to He/H =

1.0, and the mapped column then holds He/H between 0.999997254 and 1.000000000, a worst departure of 2.746×10^{-6} against the loader's 1.0×10^{-6} threshold.

What the six runs measure that earlier attempts could not. In the three well-mixed cases the stationary WIND is found and held, hydro info = 0 from pass 3 to 5 onward with every hydrodynamic row inside its tolerance, and the H2 carrier balance alone refuses, every relaxation of every pass ending on the composition movement bound and none on its own residual: that isolates the carrier transport as the single obstruction there, and it is neither a seed question nor the hydrodynamics. In the two photochemical cases and in the diffused HeH0.55 the wind is not found either. wellmixed/HeH0.55 reached 3.76×10^{-2} on the gated carrier row, the nearest any molecular well-mixed case has come to the 1.0×10^{-5} tolerance. The L33 ladder's rung 1, a 0.179 dex composition step from the certified reference, ended its first solve at outer pass 2 of a budget of 25 and not on the budget: the element composition relaxation took 28 steps, found no admissible 29th (outcome 3 at a step length 8.127×10^{-2} of the composition time scale, mass-closure departure 1.332×10^{-15}) and restored its entry composition, which makes every later pass a copy of that one.

The status.py change. (a) running is no longer inferred from the absence of a state file: a case with no solver verdict now prints **not solved** when not_solved.md is beside it, **not started** when neither a state nor that file is there, and marching-stop unchanged for a state with no verdict, which removes the standing false running on molecular_scalar_gj1132_wellmixed/HeH2.13 and on molecular_photochem_gj1132_kzzprofile/HeH2.09. (b) The column claim on 59bfdb3f is renamed **archived claim on 59bfdb3f**, and a row reads superseded where the state in the case directory has been written since the L29 table was made; the comparison, stated in the table's own note, is the run= stamp in the provenance header of the very file the L29 table read against the modification time of models/CLAIMS_59bfdb3fc4d0.md, 2026-09-17 23:27:50 KST. 14 of the catalog's 95 rows read superseded, the seven low-XUV atomic cases the L34 steps 1 and 2 worker solved and seven molecular cases; every other row's archived verdict is still a true statement about the file on disk, because the atomic re-evaluation rewrote Hydro_ioniz.txt and not the _IC copy the table read.

Also written or corrected. Six not_solved.md files, one per case, each with its full pass table; MODELS.md section 3, which was stale, now states the seeding rule the restart contract imposes on the well-mixed molecular group and why a scalar molecular state cannot seed the photochemical one; output_pre_L34/ beside the three well-mixed cases and output_L34b_stopped/ beside the three whose output/ held L34b's stopped state. status.py --write regenerated MODELS.md sections 7 and 8, make_memo_figures.py the 25 docs/figures/lhs1140b_*.pdf, and latexmk -pdf docs/lhs1140b_exhale_vs_pwinds.tex built cleanly at 80 pages, the same count as before.

Reported and not fixed. The pseudo-time continuation can undo a solve's progress and overwrite its best state: on wellmixed/HeH0.083 the first solve's forty passes reached a gated carrier row of 4.13×10^{-2} and the continuation's forty took it back to 7.37×10^{-2} , and the state on disk is the continuation's, the rule continuing whenever the first solve does not return info = 0 even when the hydrodynamic rows are already inside their tolerances. The composition movement bound never recovers within a run, only halves: 1.0×10^{-2} to 5.0×10^{-3} at pass 14 of that case's first solve, held for its remaining 27 passes, while the continuation starts at 1.0×10^{-2} again and the row rises there, so restoring the bound is not by itself the remedy. Two case directories now hold a state pair whose halves are from different runs, a solve writing Hydro_ioniz.txt only when it ends.

Record. docs/lhs1140b_stationary_L34c_20260918.md.

L35: the matched-domain flux and grid comparison, and the local base closure

The two recommendations, recommended and not decided. (1) Keep the seven low-XUV Roe cases, and do NOT make the grid the catalog's answer. (2) One bounded source item on the base closure is warranted, on the zero-window limit and not on the closure family. No production source was changed; the one source added is `src/tests/grid_and_gates/base_closure_candidates_probe.f90`, which is not registered in that suite's `run.sh`. The study was built on a snapshot of the tree taken at 2026-09-18T01:39:21+0900 (HEAD 3c73905, the tree dirty by 32 files and 13 untracked paths), binary `md5 16f5345d07ce70cdac09227839d5188b`; the snapshot carries item L30's work in progress in the residual assembly, which the memo states as a caveat, and the 500-cell Roe control reproduces the L25 certified state (T(1) 221.776 against 221.78, certified at 4.233e-07 against 4.143e-07).

The grids, matched to the bit. The Mixed generator ties the inner physical face to the first cell width and puts `r_max` at the outermost ghost center, so two keys were solved on an exact Python replica of `define_grid` that reproduces the binary's radius column with a maximum relative difference of 0.0: 500 cells with `Base grid [dr,cells]: 0.00019999999494757503 50` and `Outer radius [R_p]: 30.0`, 1000 cells with `0.00019615093524286384 50` and `29.582942490104671`, 2000 cells with `0.00019453512202014908 50` and `29.406628764578247`. All three carry `r_edg(0) = 1.0000966593889191` and `r_edg(N) = 29.2734180458963`, agreeing to 0.0 and not merely to 1e-10. Holding `r_edg(0)` fixed holds the FIRST CELL WIDTH fixed (`dr_1` is 1.933188e-04 in all three), so a matched-domain refinement cannot refine the base: the refinement is a clean factor 2 above about 1.05 `R_p` and nothing inside the 50-cell uniform block.

The six runs.

run	outcome	passes	R	rows that refuse the written state	wall, threads
HLLC 500	NOT CERTIFIED	2 of 2	2.406e-01	mass 1.801e-01 at cell 1 (1.2e-07); momentum 7.988e-08 at cell 500 (1.0e-08); energy 2.406e-01 at cell 1 (1.0e-06)	7 m, 8
HLLC 1000	NOT CERTIFIED	8 of 8	8.215e-04	energy 8.215e-04 at cell 734, r = 4.98; elemental He/H 3.075e-05 at cell 635	70 m, 16
HLLC 2000	NOT CERTIFIED	8 of 8	7.767e-04	energy 7.767e-04 at cell 1375, r = 4.91; elemental He/H 4.388e-05 at cell 1152	77 m, 24

run	outcome	passes	R	rows that refuse the written state	wall, threads
ROE 500	CERTIFIED	1 of 40	4.233e-07	none	31 s, 8
ROE 1000	NOT CERTIFIED	8 of 8	3.831e-04	energy 3.831e-04 at cell 707, r = 4.26; elemental He/H 3.092e-05 at cell 635	99 m, 16
ROE 2000	NOT CERTIFIED	8 of 8	2.689e-04	energy 2.689e-04 at cell 1345, r = 4.55; elemental He/H 4.386e-05 at cell 1152	73 m, 24

The L27 face mass flux budget, in units of the wind-window mean of $\rho v r^2$, puts five of the six at one flux through every face to between $5e-09$ and $1.3e-07$ of the wind's; the sixth is HLLC 500, whose base face carries **0.700465914** of the wind's mass flux with a face spread of $3.639e-01$. **The stall, read in the conserved quantity, is a base-face statement.** The grid does relieve it: at 1000 and 2000 cells the same unmodified HLLC flux brings mass to $2e-08$, momentum to $3e-14$ and the base face to 1.000000 of the wind's flux at the FIRST outer pass.

The flux difference is converged, and refining does not remove it. Between 1000 and 2000 cells each flux is converged to a few parts in $1e5$ in its first cell and in $\dot{M}$ (Roe $T(1)$ moves by $1.2e-06$ and $\dot{M}$ by $3.5e-05$ at the doubling; HLLC $3.9e-05$ and $1.3e-05$), while the difference BETWEEN them is the same at both resolutions to three digits: $(ROE - HLLC)/HLLC$ reads $-1.032e-01$ in $T(1)$, $+1.108e-01$ in $\rho(1)$, $-9.96e-03$ against $-9.94e-03$ in $\dot{M}$ and $-1.79e-02$ against $-1.78e-02$ in the He I 10830 equivalent width. Profile against profile the difference is 10 to 24 per cent below $1.01 R_p$, 3.1 per cent in density at $1.2 R_p$ and 0.5 to 0.8 per cent out to the top of the domain, so it is not confined to two or three cells. **The two fluxes are two converged discrete answers to one continuous problem.** What this adds to the open Roe question, and what was missing when the handoff recommended against a wider Roe adoption, is the Roe flux's OWN grid convergence: it is converged at the catalog's 500 cells, moving by $2.5e-06$ in $T(1)$ and $1.1e-04$ in $\dot{M}$ at a doubling. The equivalent width is the one quantity still moving at 2000 cells, by $-8e-03$ per doubling at both fluxes, which is the size of its own flux-family difference, so no statement about the line is supported at better than about 2 per cent by this study.

The refined runs' energy-row plateau. None of the four refined runs certifies: all four carry a hydrodynamic energy row of $2.7e-04$ to $8.2e-04$ against $1.0e-06$ at $r = 4.3$ to $5.0 R_p$. It appears as the elemental partition relaxes and then stops falling, at both fluxes: from the 40-pass attempt's pass records, HLLC 1000 reads $1.08e-07$, $1.60e-07$, $4.10e-04$, $7.35e-04$ and $8.215e-04$ at passes 1, 3, 4, 6 and 8, ROE 1000 $4.69e-08$, $2.97e-04$, $3.44e-04$, $3.76e-04$ and $3.831e-04$, HLLC 2000 $3.86e-07$ to $7.767e-04$ and ROE 2000 $6.32e-08$ to $2.689e-04$. The 500-cell Roe run does not do this ($4.2e-07$). **Whether that is a property of the refined grids, of the coupled solve or of this binary is NOT isolated, and it is the first thing a follow-up should separate.**

The closure comparison, and the rest fixture's 1/2. The probe calls the production characteristic_base_face_state twice on one installed state, changing only the discriminant it is handed: the remote window's (wind_window_mass_flux) against the production base face flux (stationary_face_mass_flux, L27's entry point). The L26 section R3 exclusion does not survive the production face flux: on both certified states, on a base whose velocity is cut by a hundred, on a base whose velocity is reversed, on a perturbation confined to the remote window and on the hydrostatic_column fixture the two install a BIT-IDENTICAL ghost in all three primitive components, and the local one is a fixed point of its own map in one iteration. They are not, however, being told apart on those states: the branch weight is saturated at both discriminants. They differ on exactly two kinds of state, and both are states no converged run holds. **The one worth a source item is the state at REST:** there the window carries exactly zero flux, the amplitude weight vanishes, the first interior cell's face Mach number is also exactly zero, and the smoothstep returns 1/2, so the base density becomes the average of the reservoir isentrope and the interior isentrope. On the fiducial those differ by 7.3 per cent, so the boundary states a face density 3.79 per cent below the reservoir on a state where nothing is flowing (ghosts 3.86 and 4.00 per cent); on the hydrostatic column the same 1/2 costs 3.07e-07, which is why it has not been visible. The item is "state what the boundary does when the wind says nothing", with the smoothstep's value at zero written at the place it is used, which it is not now. **It is item D2 of docs/PLAN_20260918_rev1.md.** Not a reason for an item: the remote closure's nonlocality, r_flux moving over a factor ten on a converged wind changing the window mean by 1.3e-05 and the face state not at all, and a 1.5x extended domain moving the discriminant by 1e-04 and the face density by 3.8e-05. Two readings belong with the comparison: the production base face flux is not the sign of the face velocity, since at a base Mach number of 1e-06 the HLLC mass flux there is dominated by its dissipation term, so the local Mach number reads the density jump the boundary itself makes (it reads the right SIGN on every state tested, which is what a branch discriminant needs); and on the six states where the two agree they agree only because the branch weight is saturated at both discriminants.

The dr_base literal. src/modules/init/parameters.f90 line 143 reads real*8 :: dr_base = 2.0e-4. Without a d exponent that is a DEFAULT REAL literal, so the stored default is 1.9999999494757503e-04, while input_read.f90 reads the key list-directed into the same real*8, so Base grid [dr,cells]: 2.0e-4 50 gives 2.0000000000000001e-04. **Stating the default explicitly therefore produces a different grid from omitting the key:** the cell centers differ by up to 6.58e-09 relative, the 2.53e-08 difference in dr_base damped by a factor 3.8 by the rescale, and that is how the Python replica was validated, an exact match holding only with the single-precision value. The correction (2.0d-4) moves every Mixed grid built from the default, which is a golden refresh, so it was not made. **It is item D1 of docs/PLAN_20260918_rev1.md,** whose D1a is the disclosure and provenance and whose D1b is the default change on new configurations only. Reported with it: the hydrostatic_column regression fixture is not a column at rest, its recorded output being a 300-step relaxation snapshot whose first cell carries an inflow of 8.09e+04 cm/s at a face Mach number of +0.2257 and whose wind window has a relative flux spread of 2.40, the largest in the tree, so a future well-balancedness test should build its own state at rest rather than load this one.

Record. docs/lhs1140b_stationary_L35_20260918.md and docs/figures/lhs1140b_L35_flux_grid.png.

L30c: what moves oxygen_chemistry by order unity

Verdict: case (i), a discrete branch selection amplifying a rounding-level seed on a snapshot that is not a solution.

Case (ii) is refused by a bitwise experiment: the carrier diffusion geometry of L30 is NOT the seed, and the oxygen carrier rows do not respond to the shell geometry by order unity. No production source was changed and no golden was touched.

The seed. It is the restatement of the spherical shell volume that L30b carried into the hydrodynamic rows, $dV = (r_+^3 - r_-^3)/3$ against $dV = (r_+ - r_-)(r_+^2 + r_+ r_- + r_-^2)/3$, whose two spellings differ by at most **5.7707e-13** relative on this case's own 500-cell Mixed grid, the maximum sitting in the 50 uniform base cells where $dr/r = 1.94e-4$ and the median over the column being $1.1853e-14$. That confirms by measurement the "about $4e-13$ " the L30 and L30b reports gave from error analysis.

The branch. It is the root ladder of the constrained molecular ionization solve. On the FIRST ionization sweep of step 0 both logs open with the identical line "the star-ward H2 column reaches $6.38E+21 \text{ cm}^{-2}$ " and then diverge: 64 cells are reported on the control and 48 on the measured build, 43 on both, and at those 43 the printed temperature is identical at all 43 while the accepted iterate's residual differs at all 43, the simplex violation at 39 and the solver status flips at one. The sweep's own summary for that step reads "11 root(s) outside the simplex" on the control and "5" on the measured build, and over the full 1000 steps the ladder's outcomes stand apart by a factor of ten: no admissible root 102 against 983, stored state(s) rejected 105 against 986, constrained-continuation root accepted 632 against 1701. The first number the run prints, du at step 1, is already apart by $6.445e-04$ and 1000 steps later by $2.648e-02$, a factor 41 over 1000 steps and not a factor per step, so this is a jump and not a growth: a seed of $5.8e-13$ does not reach $6.4e-04$ in one step by any linear response. The movement is macroscopic at one step already, the profile-scale difference reading $5.09e-02$ in rho, $2.67e-02$ in v, $2.14e-01$ in T and $9.99e-01$ in the heat column after a single step, and the count of numbers differing above $1e-12$ does not move with the cap: after ONE step every column of Hydro_ioniz.txt except the radius already differs at every cell.

The two bitwise experiments. Build A is the current tree with ONLY the two carrier diffusion denominators of diffusive_photochemistry.f90 reverted to $r_j^2 \max(dr_j, 1 \text{ cm})$ and the two face areas to r_{edge}^2 : it is **bit for bit the MEASURED build** in all four files. Build C is the current tree with ONLY the shell-volume restatement reverted in the five files that carry it: it is **bit for bit the CONTROL** in all four files. Both statements hold at a cap of one step as well. oxygen_chemistry sets neither He_diffusion nor Molecular carrier transport, so it enters neither the element diffusion operator nor the carrier transport operator, which is why build A can be bitwise the measured build; of the five files build C reverts, only species_face_flux.f90, RK_rhs.f90 and Reconstruction.f90 are reachable in this configuration, and those three carry one and the same respelling, so the seed is named by the expression and not by the file. The control tree was git archive HEAD unpacked and built with bare make, and all 263 source files of LHS1140b/models/BINARY_MANIFEST_59bafb3fc4d0.txt match it, 263 of 263 by md5sum -c; it reproduces the 2026-09-17 golden exactly at the three cells the matrix log names.

The other class, for contrast. mol_sec_ion carries the same rounding difference smoothly and with no branch: its first step is bitwise identical, the second is apart by $9.122e-12$, and the separation rises smoothly by three decades over 1000 steps. At its own 12000 steps both builds stop at the same step with the same printed line and the matrix movement is

reproduced exactly, Hydro_ioniz.txt 3.2102e-04 at row 192 in v (6.34034658e-01 against 6.34238260e-01), which is 6.31e-07 of that column's own scale. The decisive difference is that the root ladder does not fire at all in that case: over 1000 steps and over the full 12000 alike both builds report no admissible root at zero cells, no constrained-continuation acceptance and one stored-state rejection each.

What the comparison can mean, now written beside the golden. A bitwise or tolerance verdict on oxygen_chemistry certifies that the same binary reproduces itself and nothing else: it cannot tell a physics change from a last-bit change, because the root ladder turns either into an order-unity difference within one step. The integral it does pin is stable, $\log_{10} \text{Mdot} = 9.61$ on both builds at 1000 steps and agreeing to the two printed decimals at every cap. backup/regression/oxygen_chemistry/NOTE.txt now carries that statement with the two bitwise experiments, so the next refresh does not spend a session apportioning the movement again; a gate on the oxygen network that could fail for a physics reason needs a settled state or a row that reads the network directly, a rate, a branching ratio or a conservation identity, and the present case gives the oxygen path the reproducibility golden it was opened for.

Record. docs/lhs1140b_stationary_L30c_20260918.md.

The count rename of the plan's section 11

What changed. global_parameters' integer :: count is now integer :: marching_step and its cap count_max is marching_step_max. The name is read from the entry text: init.f90 sets it to 0 or 1, EXHALE_main.f90 raises it by one per pass of the marching loop, and every place that prints it calls it a step, so it is the index of the marching step and not a tally. The tree already names a step index <what>_step (sec_flip_step, newton_fail_step, reject_at_step and six more), and marching_step does not collide with the two step ledgers beside it, n_steps_attempted and n_steps_accepted, which count attempts and accepted steps rather than loop passes. Eleven files under src/ were edited.

The hit table. Occurrences of the whole word count in src/**/*.f90 and src/**/*.*.inc on the entry text, classified by what the token is at that place, with count_max counted separately:

what the token is	occurrences
this variable	95
the system_clock keyword	3
a genuine intrinsic call	1
plain English in a comment or a printed string	442

No local is named count: the only declaration of the name in all of src/ is parameters.f90 line 38, so every one of the 95 variable occurrences is the global. **Four of the 95 are live OpenMP text and not comments**, if(count > 0) clauses on !\$omp sentinel lines at ionization_equilibrium.f90 lines 1442, 1471, 1638 and 1717; a scan that treats a line beginning with ! as a comment misses them, and missing them would have left the parallel sweep referring to a name that no longer exists. The four non-variable hits are left exactly as they were, among them count(species_held) at constrained_chemical_equilibrium.f90 line 2126, which compiled on the entry text only because that unit writes use global_parameters, only: ..., the hazard the item names seen from the other side. count_max had 8

identifier occurrences and one in a comment, all renamed.

The printed labels are deliberately kept. ' final: count=', ' step-clock: count=', ' (profile) count=', the diagnostic header '# count j r[Rp] ...', ' -> stopped: reached count_max = ' and the report key count_max in EXHALE_setup.out are unchanged, because src/tests/attempted_step/run.sh parses the log with `re.search(r"step-clock: count=(\d+) ...")` and docs/input_schema.md documents count_max as that report key; moving the labels would move the logs the control compares line for line. The declaration now carries a comment naming the printed label and the report key so the two can be found from the code. Four notes in EXHALE_main.f90, write_setup_report.f90, steady_newton.f90 and sed_read.f90 saying that the intrinsic was shadowed were true of the entry text and are false now; they were removed and the loops they explained were left as they are.

Tests, RED and GREEN. A program that writes use global_parameters with no only list and then calls count(mask) on a 5-element logical mask: compiled against the entry text's parameters.o it is **RED**, the name resolving to the integer so that count(mask) is read as an array element with a logical subscript and the write statement does not parse; against the delivered source it is **GREEN** and prints true elements = 3.

Movement: byte-identical. Control 6a36529386102b37a38cb40fde0a611c (a private build of the entry text) against measured 74b96cdcf887dcfee301ec547f012d88. wasp_full ran to its own stop at 12251 steps, mol_base_handoff at 12000 and hydrostatic_column at 300, all three ending on an identical printed line on the two builds, with **0 data-line differences in all 10, 7 and 7 output files**; the files that are not identical byte for byte differ in exactly one line, the # provenance: header whose timestamp is the wall clock at which the run wrote the file, and EXHALE_setup.out and EXHALE_resolved.out are identical in all three cases, which is the check that the count_max report key still reads as before. All 34 test suites were run twice, once against each build: **3459 PASS|FAIL verdict lines, 0 of 34 suites differing in any verdict line**, exit status, pass count and fail count identical suite by suite, the only differences anywhere being the secs= wall-clock fields. Six suites exit nonzero on both builds alike on the same assertions, which is the entry text's state and not an effect of the rename. The three cases and the suites were measured twice, because a comment refined in parameters.f90 after the first pass changes the object's line tables; every number above is from the rebuilt binary.

Record. The declaration comment in src/modules/init/parameters.f90 and the K32c row of docs/input_schema.md, below.

Further corrections made while the series landed

Four of the items the workers reported outside their own scope were corrected by the advisor as they came in. Two more were already recorded in the bookkeeping subsection below and are named here only so the list is complete.

- **src/tests/grid_and_gates/README.md section 4** now states the pass budget as part of the row rather than leaving it implicit: outer_iteration_ending_hands_back_one_state runs at EXHALE_CARRIER_TRUST=1e-6, EXHALE_JFNK_MAXIT=40 and EXHALE_OUTER_PASSES=50, and the budget decides whether the stagnation ending is reached at all, since the composition elimination sweep of the handed-back state keeps the joint distance falling at full precision through about forty passes on this fixture and the

ending arrives at pass 44. The re-evaluation figure is restated as 6.4e-16, measured 2026-09-18, against 1.3e-8 before P14. The heat-column consistency row of the same section, whose expectation at HEAD was recorded as RED, is **GREEN at 1.68e-16**, the one assembly having replaced the two hand-maintained copies, and the 200-step reading it used to catch (median 1.000000, minimum 0.999992, maximum 1.000012, largest relative difference 1.19e-5) is kept as the record of what the row caught. Section 1's name list gains iontrans_atomic, and section 14b describes that suite, its five statements, its configurations and its RED expectation before L36e.

- **docs/input_schema.md K32c**, the Max steps row, gave count_max in both its "default" and its "variable" column. The report key is still count_max, and the variable behind it is now marching_step_max; the variable column reads marching_step_max (report key count_max).
- **docs/lhs1140b_stationary_L31_energy_cycles_20260917.md section 5.4** stated a rule for the H2 front and none for the dilute upper column, and none exists to state: at that radius $2 \cdot n(H2)/n_H$ is flat and rising outward, 3.43e-06 at cell 296 to 3.88e-06 at cell 303. The memo now says so and names the probe's state reader, so the chosen radius is recorded as the choice it is.
- **LHS1140b/models/run_case.sh's header** (the advisor, 2026-09-18 06:35): its description of pass 2 still said the post-processing pass "takes one time step before it stops ... so it runs at CFL 1e-12", the route item L9 replaced; it now says the pass loads the solved state under Restart intent: stationary evaluate, takes no step, measures the state, writes the same conserved state back with the certification it just made and produces the _adv files and the mass-loss line from that measurement, pointing at the block below it that already said so. Comment only; bash -n clean; the runner's statements unchanged.
- Already recorded below: **docs/code_status_20260910.md's transport row**, whose sense was inverted, and **LHS1140b/models/write_reproduce.py's seed and cold-start paragraphs**, both corrected with the book-keeping items of the plan's section 11.

The bookkeeping items of the plan's section 11

low_mach_dissipation.f90's header claimed a nonnegative dissipation, and the claim is false. The integration by parts that produced it drops the variation of the face coefficient, and the gate makes that variation a step. The header now carries the discrete kinetic-energy rate at frozen density, $E' = d^T W L d$ with L negative semidefinite, so $E' \leq 0$ only for a uniform weight and where $E' > 0$ the pair takes energy out of the internal energy and puts it into the velocity field; the measured smallest eigenvalues of the symmetric part are +7.2e-17 on a uniform grid at constant coefficient, -1.1e-12 for a coefficient falling four decades over 500 cells, -6.9e-03 with the gate closing at one face and -1.1e-10 on the LHS 1140 b 0.02-XUV state, whose negative mode sits on cells 372 to 390, $r = 4.01$ to $5.12 R_p$. The correctly signed form is written out as the conservative flux of the cell second difference, $dv/dt = -M^{-1} B^T K B v$, with its measured $\lambda_{\min}/\lambda_{\max}$ of -4.5e-17 at the closing gate and -5.9e-18 on the LHS 1140 b state, and the header says that it is derived in docs/lhs1140b_stationary_L25_20260916.md step 1 and **is not adopted here**, so a run that turns the option on carries the indefinite form. Section 4's "there is no secular source" was part of the same claim and was corrected in place. Comment lines only, 63 insertions and 11 deletions, no executable line moved.

The stagnation-ending row reaches its ending again. src/tests/grid_and_gates/output_state_consistency.sh row outer_iteration_ending_is_the_stagnation_one was RED on the committed script at

measured=pass_budget, and the assertion was NOT loosened. The cause: the carrier relaxation of that fixture keeps no transport step at all, so the H2 row stands at 7.397E-02 of 1.0E-05 from pass 2 onward, while the composition elimination sweep the handed-back state carries nudges the composition every pass, so at full precision the joint distance keeps falling and a budget of 8 never accumulates three consecutive passes without a fall. At EXHALE_OUTER_PASSES=50 the first pass at which neither measure falls is 17, the first two in a row are 42 and 43, and pass 44 announces the no_progress ending: GREEN at measured=no_progress, with the state handed back re-evaluating to itself at 6.396825e-16 against a 1e-12 allowance. Two other settings were measured and rejected and are recorded in the row's comment: a budget of 40 never reaches two in a row, and EXHALE_CARRIER_TRUST=1e-4 lets the carrier take steps and makes every pass fall. The solve leg goes from 58 s to 103 s. One executable line changed.

docs/TO_BE_DONE.md, the L7g cascade entry keeps its blocking statement (the Bossion et al. 2018 H2-H state-to-state rates extended to all bound states, as used by Nesterenok et al. 2019, and a modern H2-He set above $v = 3$) and gains L31's reduced ladder as the bounded substitute, with the three cells' bracket re-measured and the conclusion that the scalar form stays the default.

src/utls/fortdep.py now scans include. It scanned module and use only, so no .inc file appeared in build/.deps.mk and hydrodynamic_rows.o did not depend on the file whose text it is compiled from: editing the kind-generic rows alone left make reporting nothing to be done, and the next link took an object built from the previous text. An include 'file' line now makes the object depend on that file, resolved beside the including source where gfortran looks first, and the included text is scanned for its own use statements.

src/utls/map_state_to_grid.py no longer writes a contradictory coupling line. It stripped mode=, t_phys= and certified= from the source's # coupling: line and wrote certified=F in front of what was left, but not cert_reason=, so a seed mapped from a certified state carried certified=F ... cert_reason=certified_in_wind and stated both that it makes no claim and that it was certified in the wind. cert_reason= is now dropped with certified= and the line reads mode=init t_phys=0 certified=F cert_reason=mapped_seed; the module docstring says so.

LHS1140b/models/write_reproduce.py describes the run it wrote. Two sentences were false. With an empty seed line it wrote "None: this case starts from the code's own initial condition (Load IC? False). The archive holds no state of its physics." for cases whose input.inp says Load IC? True and which were re-measured in place; the branch now reads the case's own Load IC? and says that the run loaded the state already in output/. With a seed named on the command line (SEED=) it attributed the choice to models/pick_seed.py and printed that tool's tier columns, none of which applies; that branch now says the seed was named on the command line and that pick_seed.py chose nothing. Both defects reach every record ever written with SEED=, the molecular campaign included.

The documents. docs/code_status_20260910.md's transport row stated the Ionization transport requirement with the sense inverted, that Coupled carrier solve: True is needed and the run refused without it; the opposite is true after L36c, and the row now says that the partitioned route runs it and the coupled route is refused, with the reason and the date. docs/input_schema.md K15f was rewritten by L36c around the new unknown, the flux it rides on, the simplex, the validity range and the four refusals, and K32g now states ONE counting convention for EXHALE_REACTION_HEAT_RECIPIENTS, three grouped corrections spanning four reactions, which is the count the L7g memo uses. The head notes of the L26 and L22 memos are described under L27 and L28 above.

The golden refresh of 2026-09-18: deferred by decision

The matrix was run at REGRESSION_REL_TOL=0 on EXHALE.x md5 3146d11b4090306dcea75bb9718edd22, single-threaded by the script itself, with make -q clean after the build, and **every one of the sixteen cases moved against the goldens of 2026-09-17**. That is expected: L30b put the hydrodynamic rows of every case on the shared spherical volume, so no case of the matrix is byte-identical through the series, and L36c moved the three hp_* cases, whose goldens were already stale against the entry text by 3.3e-03 to 3.9e-03 before it touched them.

The table is the harness's own measure, $|a-b|/\max(|a|, |b|)$ with no floor, for the two files the verdict is taken on; the column is the one it names.

case	Hydro_ioniz.txt	column	Ion_species.txt	column
wasp_full	1.086e-10	v	9.933e-11	CaI
wasp_he23off	2.404e-08	v	1.264e-09	FeIII
wasp_full_newton	2.754e-11	cool	1.394e-10	CaI
mol_base_handoff	1.338e-05	v	7.955e-05	H2
mol_metals	1.208e-09	cool	1.051e-07	HeHp
mol_lyman_werner	1.279e-11	v	7.776e-13	H3p
mol_diffusion	1.712e-06	v	8.987e-06	H2
mol_ir_bands	2.545e-09	v	6.229e-08	HeHp
mol_sec_ion	3.210e-04	v	1.268e-05	H2
mol_carrier	8.658e-08	cool	8.039e-06	H3p
lower_profile	1.245e-07	v	4.421e-07	OIII
hydrostatic_column	9.952e-08	cool	8.319e-01	HeIII
oxygen_chemistry	1.867e+00	heat	1.000e+00	H2O
hp_zero_seed	3.297e-03	cool	3.944e-03	HII
hp_trace_seed	2.765e-03	cool	3.307e-03	HII
hp_front	8.856e-09	v	1.754e-07	H3p

Two entries are the comparator's floorless ratio and not a movement of the physics. hydrostatic_column's 8.319e-01 is HeIII at row 1 reading 2.1246769169722978e-181 against 1.2640208720415364e-180 in a column with no ionizing radiation, beside a cool entry of 9.952e-08 at 1.18e-98; that case moves by 8.46e-15 of its own column scale, which is the rounding of the two volume spellings and nothing else (item L30b). wasp_full_newton's Hydro_ioniz_adv.txt reads exactly 1.000e+00 at adv_mass_row against an exact zero, which is the missing absolute floor in that column reported under item L6c of section 11 and still open.

oxygen_chemistry moved by order unity in every column, and item L30c has since resolved it. It reads 1.867e+00 in heat at row 16, where the two builds carry -0.0012423798076531957 against +0.0010765621115482848, a sign change, and 1.000e+00 in H2O at row 285, 3.1776303939581693e-14 against 542.13312076325178. It is a 1000-step relaxation snapshot stopped at $du = 4.3884$ that carries the molecular and the oxygen carriers together, so as in the two refreshes before this one its movement cannot be apportioned from the matrix alone. Item L30c's two bitwise experiments name the seed as the shell-volume spelling of L30b, 5.7707e-13 relative on this case's own grid, amplified within

one step by the root ladder of the constrained molecular ionization solve; the subsection above carries the measurement, and backup/regression/oxygen_chemistry/NOTE.txt now carries the statement of what a byte comparison of this case can and cannot mean.

A second matrix is being measured, on the binary this series ends on. The table above was taken on 3146d11b4090306dcea75bb9718edd22, and the tree has moved past it through items L36c, L36d and L36e and the count rename, none of which the sixteen cases above were run under. Three of those cases, hp_zero_seed, hp_trace_seed and hp_front, carry Ionization transport: True and are moved again by the helium stages (hp_front by 1.6e-01 in the helium columns, item L36d), and carrier_model_a_newton, which is not a matrix case, is stale by 1.2e-04 through the He/H charge exchange item L36e added to the stage sources; the count rename is byte-identical on wasp_full, mol_base_handoff and hydrostatic_column and changes no case. The second matrix was running at the time of writing; it was then stopped before it finished, on the user's instruction of 2026-09-18 14:10, and no movement table was taken from it (READ, docs/session_handoff_20260918.md section 2.3; corrected 2026-09-19). Section 13 carries the refresh forward.

The refresh itself is not taken here. It is the advisor's, after the second matrix, and it is one refresh for the whole series: the goldens were read and never written by any item, each item measured its own movement against a control build of its own entry text instead.

DEFERRED. On 2026-09-18 at 14:10 the user instructed that no golden is refreshed and no further movement table is kept while the corrections of docs/PLAN_20260918_rev2.md continue on the same tree: a refresh now would be repeated within days. The closing binary of the L27 to L37 series is 74b96cdf887dcfee301ec547f012d88 (manifest LHS1140b/models/BINARY_MANIFEST_74b96cdf887.txt); the goldens stay the 2026-09-17 set; every item's control is its own entry-text build, so the stale goldens do not enter any verification; one refresh follows when the rev2 items have landed, with its movement table then.

13. PLAN_20260918_rev2: the corrections that follow the two reviews of PLAN_20260918 (2026-09-18 to 2026-09-19)

Plan of record: docs/PLAN_20260918_rev2.md, which replaced docs/PLAN_20260918_rev1.md and the first text docs/PLAN_20260918.md after two reviews (docs/PLAN_20260918_review.md and docs/PLAN_20260918_review1.md); its sections 0 and 0.4 record which of the reviews' source findings were read in the working tree before they were adopted. The plan holds nine items, D1 to D9, raised by the L35, L36 and L37 measurements of section 12 and by the reviews; its table "Decisions made on 2026-09-18" carries the outcome of every decided item. The tree the series worked on is repository HEAD 3c73905 plus the uncommitted L27 to L37 series of section 12, whose closing binary is 74b96cdf887dcfee301ec547f012d88 (LHS1140b/models/BINARY_MANIFEST_74b96cdf887.txt). Every number below is READ from the item memo docs/lhs1140b_stationary_<item>_<date>.md, from the item report named with it or from the plan; nothing was measured for this section. Where a memo labels a number MEASURED, the measurement is that item's.

What shaped the series, all decisions of the user on 2026-09-18. The strict regression matrix on the closing binary of

section 12 was started and stopped, and none is run on the closing binary of this series either. No running table of movement against the goldens is kept from item to item: each item measures its own movement against a control build of its own entry text, on scratch copies. No golden is refreshed while the corrections continue; one refresh follows when they have landed. On docs/DECISION_D2a_D8_review.md the user chose, for D2a, choice **A** at the pressure-balanced stationary contact (the reservoir owns the thermodynamic state of the level, including at rest), with the rest of the Codex D2a text of docs/DECISION_D2a_D8_codex.md kept; for D8, the Codex generation structure as reviewed, with legacy import of the existing products and item D8a as the first increment. On the D3a memo the user approved D3b as recommended: the He II gate and its denominator stay, the stage-sum entries become gates at their executed bound, and the candidate solver defect is opened as one bounded experiment under the rule that closed the stage-2 solver program. On the D5a memo the user approved the shape of the boundary-state operation (D5b-2) as D5a proposed it. D4b was decided by D4a's own measurement.

D1a: the base cell width a run resolved, disclosed and reproducible

What was wrong. The default base cell width was written twice, as the default-real literal $2.0\text{e-}4$ in `parameters.f90` and again in `input_read.f90`, stored as $1.99999994947575033\text{E-}004$ (bits 3F2A36E2E0000000), while $2.0\text{e-}4$ stated in `Base_grid [dr,cells]`: gives the double $2.00000000000000010\text{E-}004$ (bits 3F2A36E2EB1C432D), $2.52621\text{e-}08$ relative above it. Two inputs a reader would call identical build two Mixed grids whose states refuse to load into each other (the $1\text{e-}10$ center tolerance of `load_IC.f90` lines 2221 to 2244, READ), nothing recorded whether the width came from the key or the code, and no output stated it to round-trip precision (`put_r` prints $\text{ES}23.15\text{E}3$). The defect is the contract, not the physics.

What changed. One named constant `dr_base_default` in `parameters.f90`, written as the exact decimal $1.9999999494757503\text{d-}4$; a logical `dr_base_from_key` set only by the key's own branch of the reader in `input_read.f90`; `round_trip_decimal` in module `setup_report` (`write_setup_report.f90`), a 26-character field $\text{ES}26.17\text{E}3$ (the first candidate, $\text{ES}24.17\text{E}3$, prints 24 asterisks for a negative value, MEASURED). `EXHALE_resolved.out` gains `grid_type`, `grid_cells`, `outer_radius_Rp`, `base_cell_width_Rp`, `base_uniform_cells`, `base_cell_width_source` (default or key) and `base_grid_in_effect`; `EXHALE_setup.out` states `dr_base = <round-trip value>` (the default) or (from the key); `LHS1140b/models/write_reproduce.py` writes a `##` The grid section from the run's resolved record; docs/input_schema.md row K33b states the stored default and the trap. No value moved.

How it was verified. `src/tests/grid_and_gates/base_cell_width_provenance.f90` round-trips 0, plus and minus 1, $1\text{e-}4$, $2\text{e-}4$, $1\text{e-}300$, $1\text{e}300$ and the default, each with the bits written equal to the bits read, and asserts that the exact decimal carries the bits of the default-real literal. `base_grid_key_reproduces_default.sh`, three one-step runs of `backup/regression/roundtrip`: the key stating the resolved value gives an `r` column byte-identical to the absent key's, and the key stating $2.0\text{e-}4$ differs by $3.818119\text{e-}09$ at row 366 of 504 on that grid ($6.58\text{e-}09$ on the 500-cell catalog grid, READ from L35). RED: the driver does not compile on the entry text and the shell test stops at `base_grid_resolved_record_present`. Byte identity: `wasp_full` (400 steps) and `hydrostatic_column` (300 steps) identical in every data line between the entry-text build `4e02c6807587f0d88cdacb944a65ed25` and the D1a build `45d0aa6a4a8196be47c5e3d0b608747d`.

What it left open. D1b, moving the default to $2.0d-4$, waits for every historical omitted-key input to be pinned. The Mixed construction ties the inner physical face to the first cell width, so a refinement moves the numerical boundary against the reservoir level; that is a separate item and nothing here addresses it.

D3a: the transported He II row measured, source channel by channel

What was open. On `carrier_model_a_newton` with `Ionization transport: True` the carrier balance He+ row stood three decades above its gate (L36d). The first text had claimed that no composition could certify it and had offered two candidate denominators that the reviews showed to be misreadings of `row_terms_phys`. The balance, its cancellation floor and its conditioning had to be measured before any norm was discussed.

What changed. `ion_residual_core.f90` declares a channel layout of 26 channels (each a single term of the H II, He II or He III balance, with its stage, sign and name) and the readers `stage_source_from_channels` and `stage_gross_of_channels`; `mol_heh_rows` (`System_HeH_mol.f90`), `heh_tr_rows` and `heh_rows` fill them from the same factors as the rows. The environment key `EXHALE_STAGE_CHANNELS` (default unset, nothing reached) names up to eight cells whose evaluations are appended with every kernel argument to `output/stage_channels.txt`. New suite `src/tests/stage_row_balance/`; two diagnostics beside the manufactured column in `ionization_stage_flux_tests.f90`, and a new stiff manufactured column.

How it was verified. The signed channel sum of each stage reproduces the assembled source after the He <-> H pair to $3.0e-16$ of the larger gross side in both gas branches, and a re-call of the kernel with the recorded arguments returns the recorded channels bitwise. MEASURED on the hot-Uranus transported state at cell 248: $P = 7.6156e2$, $L = 7.6034e2$, net $1.2153 \text{ cm}^{-3} \text{ s}^{-1}$, $|P - L|/\max(P, L) = 1.60e-3$; the cancellation floor $3.717e-12$, so the net stands $3.3e11$ above it, and a quadruple-precision re-summation of the channels departs by 0. The row is linear in $x(\text{He II})$ over six decades of budget-respecting steps ($d(P-L)/dx = -1.4472e9$ at every step from $1e-2$ to $1e-8$); the residual -1.2007 is an abundance correction of $1.58e-3$ of $x(\text{He II})$ at cell 248 and $1.34e-3$ at cell 107, against $2.35e-5$ and $3.33e-8$ on the locally equilibrated state through the same operator. The certified atomic fiducial's cell 150 is as stiff ($1.5e-3$) and certifies with four decades to spare because the state is the root. At a tolerance of $1e-5$ the present scale admits an abundance error from $6.0e-13$ to $2.1e-2$ of the stage fraction over one column; the gross (turnover) scale admits $2.0e-5$ at every stiff cell on both fixtures and coincides with the present one where the stage is not stiff ($3.7822e-1$ against $3.7825e-1$ at cell 500). The recorded normalized order 2.886 of the manufactured column is a dimensional order of 1.923 ($3.7493e-2$, $1.0076e-2$, $2.6089e-3 \text{ cm}^{-3} \text{ s}^{-1}$ at 250, 500 and 1000 cells), the stiff column 1.928, the boundary rows 0.93. The stage-sum entry's floating-point bound is $2 \cdot (5 \cdot n_k + 2) \cdot \epsilon_g$: $3.109e-15$ (H) and $5.329e-15$ (He) at $g = 1$, validated by execution at 0.0703 and 0.0580 of the bound; the production states read at most $2.203e-16$ (H) and $4.195e-16$ (He). The sweep and the transport operator read one radiation field, bitwise. Suites: `stage_row_balance` 0 failed on every record; `ionization_stage_flux` 45 rows (42 before), 0 failed. Key-off and key-on byte identity: `wasp_full` (400 steps) and `carrier_model_a_newton` (25 outer passes), data rows identical.

What it left open. The decision (D3b). The candidate solver defect, the transported state lying farther from the row's root than the locally equilibrated one after 25 bounded passes, was reported and not acted on (the bounded experiment below answered it). The atomic kernel's call site was not resolvable inside the kernel (D7b added the argument). L36d's reading of 2.886 as the WENO3 order was corrected in that memo, one sentence.

D3b: the stage-sum entries made gates at their floating-point bound

What was open. The two stage-sum entries of the certification were evaluated with `tol = 0` and `within_tol = .true.` written unconditionally, a row that could refuse nothing whatever it measured. The user's decision kept the He II gate and its denominator and made these entries gates.

What changed. `ionization_stage_sum_rounding_bound(nk [, g])` and `ionization_stage_sum_scale_ratio_limit(nk)` in `ionization_stage_transport.f90`, beside the face flux they bound, with an optional `Ek_out` on `ionization_stage_face_flux`; `ionization_stage_nucleus_sum` in `diffusive_photochemistry.f90` returns $g = S/S'$ at the face that carries the measure; `certification.f90` moves the entry into the public `ionization_stage_sum_entry`, which takes the measured g , sets finite, gates on both, and names g in the entry's scale: line; anchor (8) in `docs/certification_tolerance_anchoring_20260910.md`. The tolerance is $3.1086244689504383e-15$ (H) and $5.3290705182007514e-15$ (He) at the measured $g = 1$; the algebraic ceiling of g ($3/2$ and 2) is used only where the operator reports none.

How it was verified. `certification` 103 assertions, 0 failed (84 before), the nineteen new rows including a transcribed entry text that accepts a broken construction of $1.5e-4$ and a non-finite measure refused. The three production states of D3a certify on these rows at 0.057 to 0.079 of their own tolerance, every one of the six worst faces carrying $g = 1$ to five decimals, and their run logs differ from the control only in the two `tol=` fields. Broken constructions stand ten decades above ($1.5e-4$ against $5.329e-15$ is $2.8e10$). Suites linking the changed modules all pass (`stage_row_balance` 10, `ionization_stage_flux` 45, `element_operator` 40, `adv_static_limit` 53, `krylov_and_dogleg` 334, `steady_species_rows` 197, `carrier_helium_inventory` 15, `charge_exchange_rows` 250). Key-off byte identity on `wasp_full` (400) and `mol_carrier` (12000); the three `hp_*` cases, which transport the stages, data-row identical and within at 0.066 to 0.072 of the bound. The local `entry_index` of `certification_contexts.f90`, which returned the first entry for an absent name, was replaced by the module's accessor.

What it left open. The He II row itself (D3a's candidate defect), taken to the bounded experiment. An entry's reason text prints only when it refuses.

D4a: what the thermochemical closure hands back, measured

What was open. `carrier_retry's row one_further_sweep_leaves_the_returned_composition_within_the_sweep_tolerance` read $1.100521341448987e-06$ against `ieq_res_tol = 1e-6`, bound by `HeH+` at cell 1. Either production's closure exits before its promise is met, or the test assumes a bound the residual contract does not give.

What changed. No production source. A new driver `src/tests/carrier_retry/closure_returned_state.f90` with its runner, which saves the production return and measures on isolated copies, the module state saved and restored.

How it was verified. The returned composition is a root of its network at the returned temperature: largest normalized residual over the column $2.593e-14$, $1.611e-16$ at the binding cell, $5.99e-19$ on the `HeH+` row. Restarted from it, the closure reports converged in one cycle with a temperature increment of $1.2574e-11$. The `HeH+` row's sensitivity to its own unknown is $|R|/eps = 5.456e-13$, so `ieq_res_tol` bounds its relative abundance only at $1.8328e+6$: the assertion compared a movement of $1.11e-6$ with a tolerance twelve decades looser, and the movement is the inner

solve's repeatability (it contracts by about four decades a step; the chain's response to its transported parent is of order one). The row's value also varies by about 10 per cent with the pressure the sequence is entered with ($1.084\text{e-}6$ to $1.293\text{e-}6$). Three findings beside the verdict: the refusal branch of `relax_photochemical_composition` leaves the rate state (`ieq_rate_cell` and its companions), the caloric maps ($1.5\text{e-}3$ relative in p from the unchanged ρ), `dp_bc`, `n_part_cell1` and the carrier checkpoint as the refused trial wrote them, so the certification's chemical residual of the returned state reads $5.33\text{e-}4$ where the state's own is $1.61\text{e-}16$; the closure's header names `ieq_res_tol` where the code tests `chem_cycle_tol = 1e-5`; and its claim that a temperature change below the tolerance cannot move the composition beyond it is false for abundance ($4.76\text{e-}6$ at $dT/T = 1\text{e-}6$) and true for the residual.

What it left open. The decision, taken by the measurement: D4b.

D4b: the returned-state contract, and what a refused trial puts back

What was wrong. The three findings of D4a: a refused trial left the rate state, caloric maps, base ghost pair and carrier checkpoint behind; the failing row compared an abundance movement with a residual tolerance; the closure's header stated a tolerance the code does not use.

What changed. `ieq_rate_state` with `save_ieq_rate_state` and `restore_ieq_rate_state` in `ionization_equilibrium.f90`, beside the arrays they hold; `thermochemical_state` with `save_thermochemical_state` and `restore_thermochemical_state` in `diffusive_photochemistry.f90` (composition, p , T , heat, cool, η , bg_cell , the rate state, `dp_bc`, `n_part_cell1`, the carrier checkpoint), the caloric maps rebuilt through `get_species_densities` rather than copied. In `carrier_retry.f90` the row is replaced by `every_species_row_of_the_returned_state_is_a_root_at_the_returned_temperature` with a companion row asserting the profile is available; the abundance movement stays as a line that gates nothing, printed with its conditioning. The closure's header states the exit test and what D4a measured. No tolerance moved.

How it was verified. Nine restore rows RED on the entry text (the D4a table 4 numbers to every digit) and GREEN at 0. The new gate reads $5.3323\text{e-}4$ on a control whose restore is reduced to the entry text's and $4.440892098500626\text{e-}16$ on the delivered text; a sensitivity row built from production routines reads $7.80\text{e-}5$ against the rate state a trial leaves. The printed movement: $1.1005\text{e-}06$ on HeH+ at cell 1, $|R|/\epsilon$ $5.6749\text{e-}13$, admitted relative abundance $1.7621\text{e+}06$. Key-off byte identity on `wasp_full` (400), `mol_diffusion` and `mol_carrier` (12000); on `carrier_model_a_newton` (3 outer passes, 31 refused trials, the third pass ending on one) the outputs are identical, so the leftover was inert on that configuration. Every suite linking the two modules passes (18 suites and the D4a driver, `carrier_retry` 156, `grid_and_gates` 279).

What it left open. `grid_and_gates` fails `base_grid_resolved_record_present` against the tree's own EXHALE.x, which is behind the sources; with EXHALE_EXE pointing at a build of the sources it passes. Restoring `ieq_nonroot_streak` is a stated choice that no fixture separates.

D5a: where the boundary of an evaluated state comes from, and which of it the residual reads

What was open. Six certified molecular states did not reproduce their cell-1 mass row from their own restart pair (L37); two certifications inside one run disagreed at the base cells; the first text's two-file repair had named the printed precision of the ghost rows as the cause without isolating it.

What changed. No production source. The diagnostics (EXHALE_BOUNDARY_TRACE, EXHALE_TRACE_EXPERIMENT, EXHALE_TRACE_STATE_B, a new boundary_state_trace.f90) were built in a frozen snapshot only; the measured build with every key unset reproduced the control's products byte for byte.

How it was verified. Six measurements on scratch copies, single threaded. (1) The evaluate route assembles the residual on a boundary built from the LOADED composition while its sources, pressure map and sound speeds are those of the composition one sweep later; rebuilding the boundary after the sweep takes the cell-1 row from 15.98 rounding floors to 9.06 (HeH2.13 molecular), 31.24 to 3.09 (HeH9.7) and 20.93 to 0.08 (.L22/i3_alt), 0 and 0.35 floors on the atomic states. (2) Serialization of the physical column is innocent: 0, 2.19e-16, 4.03e-16 and 6.07e-16 relative on the conserved density, momentum, energy and species. (3) With the physical cells held and only the two inner ghost rows replaced, the molecular row reads 9.06, 15.98, 425 or 7.4e8 floors, and the first cell's composition as ghost reverses the sign of the base face mass flux: the boundary is determined by the supplied ghost. (4) Apply_BC is idempotent; what a report leaves behind is the face cache, worth up to 3.2 floors to the next residual of the same array. (5) ntot_bc + dp_bc is two quantities under one name: the prescribed reservoir at startup and the solved ghost's counts after a sweep (+4.66 per cent on the molecular state), never fed back, and restating the reservoir at the solved count takes the row to 1.0. (6) No physical cell drifts when a diagnostic is requested, and two processes agree bitwise. From the third re-entry the row alternates between 7.14 and 9.06 floors on states differing by about 1e-13: the row's conditioning bounds what any boundary operation can deliver.

What it left open. The design of D5b, proposed in the memo's section 9 and approved by the user as the D5b-2 shape.

D5b-1: the evaluation routes derive the boundary after the composition sweep

What was wrong. The largest cause D5a isolated: three routes of EXHALE_main.f90 (the evaluate route of stationary_state_of_the_loaded_restart, the steady residual of update_map_begin_step, the EXHALE_RESIDUAL block) assembled their residual on the boundary of the loaded composition.

What changed. Apply_BC(u) after the composition sweep and the pressure refresh and before assemble_residual at those three sites (the order the JFNK residual and the marching loop already had); src/modules/states/boundary_state_trace.f90 (new, behind EXHALE_BOUNDARY_TRACE, with EXHALE_TRACE_EXPERIMENT=boundary_from_the_entry_composition reproducing the entry text's order from the same binary); one SRC line in the Makefile; new suite src/tests/boundary_state/.

How it was verified. Cell-1 continuity row in floors, entry text to this item: HeH2.13 molecular 15.98 to 9.06, HeH9.7 31.24 to 3.09, .L22/i3_alt 19.04 to 0.08, the atomic .L26/fid_resolve 1.52 unchanged; all three molecular states pass from NOT CERTIFIED to CERTIFIED. The gap between the cached boundary and the boundary of the installed state at assemble_residual is 0 on all four (4.3504e-11, 5.9572e-12, 1.4504e-11 on the molecular states in the entry order). Byte identity: carrier_model_a_newton (3 outer passes) and wasp_full (400 steps). boundary_state 2 assertions, steady_species_rows 201, certification 84, grid_and_gates 260, 0 failing.

What it left open. Two further sites of the same order inside a solve's own certification (the joint test of steady_wind_with_element_diffusion and the final certification of a stationary restart), the ghost composition read from the file, the reservoir name and the cache validity: D5b-2.

D5b-2: one boundary-state operation, with a stated input set

What was wrong. The lower boundary was a function of the ghost rows of a restart file and of the ghost's iteration history, the prescribed reservoir and the solved ghost counts shared one name, the face cache carried no validity, and two evaluation sites of a solve still assembled on a boundary of another composition.

What changed. Apply_BC.f90 is the operation with its contract stated: input the physical conserved state, the installed composition, the prescribed reservoir with its version and the model options; output the ghost conserved state and the three cached face states of one composition, tagged by hold_boundary_state_inputs; Rec_BC recomputes a face state whose composition has moved (recompute_base_face_states, counted and reported). base_boundary.f90: base_reservoir_prescription_version = 1, the model identity characteristic_face_ps_reservoir_C_minus_smoothstep_v1, the solved ghost counts and closure as named quantities, report_base_boundary_model. ionization_equilibrium.f90: the ghost's $x_{H2} = x_2 (1 - x_{ion})$ closed against its own ionization balance to base_ghost_closure_tol = 1.0e-10 in at most 30 passes, with error stop 'ioniz_eq: base ghost H2 closure failed' and no fallback; the reservoir write-back removed. load_IC.f90: the lower ghost rows are not read; base_reservoir_composition_row seeds them with the reservoir's elements and the first physical cell's partition (the handoff's $x_{H2} = x_2 (1 - x_{ion})$ form); the composition-moved test reads the physical cells alone. write_output.f90 writes # boundary_model and # boundary_reservoir on both halves. EXHALE_main.f90: the two remaining evaluation sites install the boundary after the sweep, and the boundary is derived at the top of each marching attempt. docs/input_schema.md appendix D.

How it was verified. Acceptance all bitwise: the four ghost variants of D5a give one boundary and one first residual (1.5582E-08 in all four), a re-assembly after the face-flux report moves nothing, $E1 = E2 = E3 = E4$ on the re-entry ladder, and the physical column round-trips at 0, 2.19e-16, 4.03e-16 and 6.07e-16. The closure reports 0.00000E+00 in 4 passes on HeH2.13 and .L22/i3_alt and 1.11022E-16 in 3 on HeH9.7; the failure path was exercised on a throwaway build. boundary_state 13 of 13 GREEN and 11 of 11 RED on the entry text; ten further suites pass whole.

Consequence, reported and not absorbed: the three molecular states go from 9.06, 3.09 and 0.08 floors to 19.97, 110.0 and 27.13 and are NOT CERTIFIED on the cell-1 continuity row, the physical cells unchanged; the ghost's neutral columns and element budget are reproduced to 1e-10 and its He/H exactly, while its trace ions move by 3e-3 to 5e-2, the band the sweep's own acceptance admits in a ghost cell, and the row carries a gain of order 1e9 floors per relative unit of the ghost composition. The atomic state is unchanged. Movement: wasp_full (400) and lower_profile (12000) identical; mol_base_handoff 6.19e-09 (cool) and 1.05e-08 (H3p); mol_carrier 1.33e-11 (v) and 1.29e-12 (HeI).

What it left open. The certificates of kzz1e9/HeH2.13, kzz1e9/HeH9.7 and .L22/i3_alt are stale and are to be re-solved under the corrected boundary (D9 step 3). Four grid_and_gates rows compared ghost rows (279 PASS, 4 FAIL against 284 and 0 on the control); the periodic certification of a marching state recomputed 19 and 21 face states on the two molecular cases: D5b-3. The ghost's ionization is defined only to the sweep's acceptance band; closing the temperature leg of the ghost closure is a later increment.

D5b-3: the four rows, the fourth mixing site, and the prescribed-count labels

What was wrong. Three `grid_and_gates` round-trip rows compared the lower ghost rows, which D5b-2 made derived, and the stagnation fixture no longer reached its ending under the derived boundary; the periodic certification of a marching state read a face state of the composition before the step's closing refresh; `EXHALE_resolved.out` printed `ntot_bc` and `dp_bc` without saying which count they are.

What changed. `EXHALE_main.f90`: `Apply_BC(u)` before the periodic certification's `assemble_residual`, with the invariant stated. `write_setup_report.f90`: `EXHALE_resolved.out` labels `ntot_bc_per_H` as the prescribed count and adds `base_reservoir_version`, `base_ghost_count_solved` and `base_ghost_electrons_solved` (the `parse_dump.txt` record untouched). `restart_round_trip.sh`: the three rows compare the physical cells, and a new row `restart_lower_ghost_is_the_derived_boundary`; `output_state_consistency.sh`: the stagnation budget raised from 50 to 100 passes, the ending announced at pass 89; `run.sh` passes `EXHALE_TEST_OUT` to that program.

How it was verified. `grid_and_gates` 285 PASS, 0 FAIL. The recompute counts of `mol_base_handoff` and `mol_carrier` fall from 19 and 21 to 0; their physical cells move by $1.2629\text{e-}11$ and $1.4985\text{e-}11$ (v) because `eval_dt` takes the CFL minimum over the ghost cells too; `wasp_full` and `lower_profile` identical.

What it left open. Fourteen invocations of `grid_and_gates/run.sh` still ignore `EXHALE_TEST_OUT`. The new row's exact zero was later found to be a coincidence of equal seeds; see "Corrections made while the series landed".

D6: the helium held outside the atomic stages is reserved from the stage simplex

What was wrong. Three defects in the expressions. The helium ion-stage projection bounded $x_{\text{HeII}} + x_{\text{HeIII}}$ by one while the fractions are normalized by the total helium nucleus count, which includes HeH^+ , so the review's counterexample ($\text{HeH}^+ 0.10$, $x(\text{He II}) 0.60$, $x(\text{He III}) 0.35$) passed at 0.95 and was written back as a cell holding 1.05 of its helium. The ground singlet was formed as $\max(n_{\text{HeI}} - n_{\text{He}(2^3\text{S})}, 0)$, so a trial leaving less neutral helium than the frozen metastable holds was clipped rather than refused. The molecular proton ledger kept $\text{sprod}(\text{H}^+) - \text{sloss}(\text{H}^+)$ from before the $\text{He} \leftrightarrow \text{H}$ charge-exchange pair while $\text{src}(\text{H}^+)$ was taken after it. Reachability in a production run was not tested; conservation of helium nuclei is an invariant the operator claims.

What changed. The admissible set is $n(\text{He II}) + n(\text{He III}) \leq n_{\text{He,avail}} = n_{\text{He,nuc}} - n_{\text{He(in molecules)}} - n_{\text{He}(2^3\text{S})}$, the molecular part read from the species table. In `ionization_stage_transport.f90` `stage_simplex_projection` takes an optional bound `xsum_max` and its diagnostic is renamed `stage_simplex_sum_over_limit`; in `diffusive_photochemistry.f90` the frozen `cbg_nHemol`, `carrier_helium_available_to_stages` and `carrier_helium_neutral_of_partition` (no clip), the bound applied in the projection, the element limiter, every Newton trial, an inward difference step at the bound, the source evaluation and the write-back, and the proton ledger taking the pair's measured net on its own side. The normalization (fractions of the nucleus count) is unchanged.

How it was verified. New suite `src/tests/carrier_helium_inventory/` measuring the returned species: the stage sum over the available helium $+7.000\text{e-}2$ on the entry text and $-1.110\text{e-}16$ after; the written helium inventory $5.000\text{e-}2$ and $1.110\text{e-}16$; the mass $1.618\text{e-}2$ and 0; the molecular proton ledger $6.489\text{e-}1$ and 0. `ionization_stage_flux` gains case 4b (40 of 40). Every suite linking the changed objects passes except the rows that fail identically on the entry text

(carrier_retry's D4 row and coupled_block_jacobian's L22 rows); ionization_imposed_fractions needed a cx_set_cell in its driver's setup and then passes 49. Key-off byte identity on wasp_full (400), mol_diffusion and mol_carrier (12000). Key-on: hp_front moves by $1.2\text{e-}14$; carrier_model_a_newton (3 passes) by up to $4.100\text{e-}5$ (v). The constraint does not bind on either state (the stage sum stands 0.55 and 0.32 below its bound); builds with the trial projection or the inward step removed are byte-identical to the delivered one, so the movement is the different association of the neutral helium closure, amplified over three passes.

What it left open. A diagnostic row in steady_newton.f90 near line 14313 builds carrier densities with the mass-row density for every carrier, wrong by the nucleus ratio for a stage row; element_inventory_report skips helium; the hydrogen and oxygen closures of carrier_source still clip a negative remainder.

D7a: the equation basis of the generic charge-exchange helium row

What was wrong. cx_add_to_fvec and cx_add_to_jac write the He I <-> He II row ionization-positive, while System_HeH_TR_metals inserts them after heh_tr_rows, whose row 2 is the summed He I balance written He I-gain positive. A metal + He/He+ reaction (Huang et al. 2023, Table 4 group C) entered that row with the wrong sign: on the review's counterexample (Si I + He II at 10000 K) the He II source read $+7.561\text{e-}11$ where $-7.561\text{e-}11$ is required and the ionic charge source $+1.5122\text{e-}10$ where zero is required. Separately, the transported stage sources lacked the metal charge exchange the sweep's rows carry, and the thread-local rate coefficients had no cell-ownership guard.

What changed. A mandatory he_row_sign on both routines, applied to the He I <-> He II increments alone (cx_he_i_boundary_sign): -1 at System_HeH_TR_metals.f90, +1 at System_HeH_metals.f90, System_HeH_mol_metals.f90 and constrained_chemical_equilibrium.f90 (a fourth caller the plan did not list). cx_cell_T joins the threadprivate state and cx_require_cell refuses an assembly at another cell's temperature. input_read.f90 refuses Ionization transport: True with metals, naming item D7. New suite src/tests/charge_exchange_rows/; ionization_transport_atomic.sh case A now drops the metals.inp that examples/14_diffusion carries.

How it was verified. charge_exchange_rows 102 PASS: eight reactions in three equation bases with stoichiometry, nucleus balance and charge exactly 0, group B absent from the generic set, group E under its own switch, the Jacobian against a central difference at $2.79\text{e-}13$, cell order independence exactly 0, the stale-cell and metals refusals. RED reproduces the review's table to every printed digit. Key-off byte identity on wasp_full (400) and mol_metals (12000). With cx_full 1 on wasp_full the correction moves He II by $1.817\text{e-}3$, He(2^3S) by $1.957\text{e-}3$, He I by $1.413\text{e-}4$, H II by $1.040\text{e-}3$ and T by $5.263\text{e-}5$; log10(Mdot) 7.20 both ways. No golden carries cx_full.

What it left open. The He <-> H pair given the summed He I by the triplet systems (D7c); cx_add_to_turnover reads cx_kc without a cell guard; the retained review probe docs/audit_20260905/plan_20260918_rev1_source_probe.f90 no longer compiles against the new signature and is left as the dated record it is.

D7b: the metal charge exchange of a transported ionization stage

What was wrong. The transported stages took their source from the H/He rows alone while the local sweep adds the metal charge exchange to the same rows: on a metal-bearing cell at 8000 K the missing term was $1.813\text{e-}3$ of the H II

source and $2.380\text{e-}6$ of the He II source (MEASURED).

What changed. One reaction-source interface in `charge_exchange.f90`: `cx_reaction_rates` forms every rate once, `charge_exchange_stage_sources` returns the stoichiometric source of each H, He and metal stage with the gross production and loss, and `cx_add_to_fvec` reads the same rate vector (its addition order kept, for byte identity). `diffusive_photochemistry.f90`: the frozen metal stage densities `cbg_nm0/nm1/nm2`, `cx_set_cell` for the cell being assembled, the stage sources added in `carrier_source`, and the atomic branch's He $\leftrightarrow$ H pair given the ground singlet. `ion_residual_core.f90`: the channel layout grows from 26 to 32 (`ich_metal*`), and `heh_tr_rows` takes an optional `transport_operator` so the record reads AC or AS. `input_read.f90`: D7a's interim refusal lifted.

How it was verified. The transported and the row-implied sources agree to $2.178\text{e-}16$ (H II) and exactly 0 (He II) on the test cell, and exactly 0 at every one of the 500 cells of the solved `wasp_full` column. `charge_exchange_rows` 250, `stage_row_balance` 10 and fifteen further suites pass; a bounds-checked build runs a metal-bearing key-on case with no trap; four thread and order arrangements bitwise equal. Key-off byte identity on `wasp_full` (400), `mol_metals` and `mol_carrier` (12000).

What it left open. The key-on `wasp_full` solve does not certify, with or without metals (refused at outer pass 6, and at pass 5 metal-free, on the same seven entries): a 400-step relaxation snapshot of WASP-121b is not a near-stationary seed, so no certified metal-bearing transported reference exists yet.

D7c: the helium state the He $\leftrightarrow$ H charge-exchange pair reacts from

What was wrong. The two He 2^3S systems (`System_HeH_TR.f90`, `System_HeH_TR_metals.f90`) handed the Table 4 group B pair the summed He I, singlet plus metastable. The tabulated barrier $\exp(-12.75/T4)$ is the 10.989 eV difference of the ground-state ionization potentials; $\text{He}(2^3\text{S}) + \text{H}^+$ is exothermic, a different reaction this code carries nowhere. Charging the metastable at the ground-state rate is the wrong reaction, not an approximation.

What changed. Both systems pass the ground singlet `n_heiSI`, so all four helium systems and the constrained chemical equilibrium react the pair from one reservoir; neither triplet system has an analytic Jacobian to match.

How it was verified. New row `he_h_pair_reservoir.f90` of `charge_exchange_rows`: RED 16 of 30 assertions on the entry text (the two triplet systems $1.22\text{e-}2 \text{ cm}^{-3} \text{ s}^{-1}$ away, exactly `k1 n(He  $2^3\text{S}$ ) n(H+)`); GREEN 132 of 132. `wasp_he23off` byte-identical. `wasp_full` (400 steps) moves: He I $3.98\text{e-}6$ at most (at 1.2705 Rp where `n(He I)` is $3.4\text{e-}7 \text{ cm}^{-3}$), about $1\text{e-}11$ in the dense launch region, $\log_{10}(\text{Mdot})$ 7.20 both. The certified atomic fiducial's evaluate route keeps every certification line; its files move at $3.46\text{e-}15$ and below.

What it left open. Two further sites: the advection-corrected system and the group C reactions (D7d).

D7d: the helium state the metal + He and the advected He $\leftrightarrow$ H reactions react from

What was wrong. `System_implicit_adv_HeH_TR` gave the group B pair the summed He I fraction; the three metal systems gave the group C reactions the summed He I. The group C barriers (19.1, 15.5, 12.7 in units of 1e4 K) match the ground-state ionization-potential differences to the two digits the fits carry.

What changed. `xheiS` at the advection site; `n_heiSI` (or `sden(is_HeI_SI)`) at the three group C sites, changed together; the header of `charge_exchange.f90`; a dead summed-reservoir assignment removed from `hydrogen_helium_row_terms`; a public diagnostic `constrained_network_balance_rows_of_cell`.

How it was verified. Two new rows (advection_pair_reservoir.f90, RED 6 of 6; metal_helium_reservoir.f90, RED 6 of 13 at 1.2903e-1); the suite GREEN at 151 of 151. wasp_full and mol_metals at 400 steps with cx_full off: byte-identical in all output files, the _adv profiles included (the corrected cells carry a negligible metastable). On the certified atomic_scalar_gj1132_kzz1e9/HeH2.13 evaluate route the equilibrium files are byte-identical and Hydro_ioniz_adv.txt moves by 2.43e-13 (cool). With cx_full 1 on wasp_full the neutral metals move by up to 9.579e-5 (FeI at 1.8e-12 cm⁻³), the bulk state at 1e-13.

What it left open. The real He(2³S) + H+ reaction needs a published rate and is a separate physics item.

D8a: what a case directory publishes, and when a second solve is taken

What was wrong. run_case.sh launched the continuation on any nonzero info before classifying the hydrodynamic rows, so a composition-only refusal continued (L34c: 4.13e-2 then 7.37e-2 on wellmixed/HeH0.083, the worse state published); the evaluate pass edited the case's input.inp in place; run_campaign.sh read run_case.sh through tail -n 1 without pipefail.

What changed. LHS1140b/models/run_case.sh: classify_ending (seven classes: solved, no_verdict, state_missing, nonfinite_state, composition_refusal, hydrodynamic_refusal, no_row_measures) before any decision; the continuation taken for hydrodynamic_refusal alone; a superseded solve kept in solve_<n>/, each ending recorded in ENDING; the evaluate pass in eval/. run_campaign.sh: pipefail, one status line for each case in campaign_status.txt, a nonzero exit when any case fails. LHS1140b/MODELS.md section 6. No source under src/ changed.

How it was verified. LHS1140b/models/tests/run_case_policy.sh 16 of 16: a real hydrodynamic refusal continued with the first solve kept unchanged; a synthetic, labeled composition-only log not continued; a stopped evaluate pass leaving the case's input.inp transformed on the superseded sequence (RED) and byte-identical on the new one (GREEN); a failed campaign case giving a nonzero aggregate.

What it left open. A previous run's output/ is still overwritten by a new run's seed (the generations of D8b); element_flux_closure.py carried the same continuation rule (D9 repaired it).

D8b: the generations a case publishes, the index a reader resolves, and the states this tree already held

What was wrong. A run directory could hold products and seeds of different generations, readers could combine them, a certificate could be attached to a state it did not judge, and nothing named which state a case publishes.

What changed. The contract in LHS1140b/MODELS.md section 9: immutable states/<generation_id>/ with the state files, certification.txt and manifest.json; runs/<run_id>/; state_index.json with latest_complete and latest_certified apart; one publisher LHS1140b/models/publish_state.py (lock, validation, read-only, atomic rename on one filesystem); import_legacy_states.py and legacy_state_map.txt. Readers resolve the index once (run_case.sh, run_campaign.sh, pick_seed.py, status.py, write_reproduce.py, src/Utils/map_state_to_grid.py, src/Utils/element_flux_closure.py). The generations are copies because the binary truncates output/ in place. latest_certified moves only when the header claims certified=T, a block says CERTIFIED and the attachment is established. A defect of certification_block (the second heading FINAL STATE AS WRITTEN ... of an exit-2 run not recognized, so the log was classified no_row_measures) was fixed.

How it was verified. Legacy import, after a dry run: 136 state directories, 222 generations, 134 indexes, 86 with latest_certified and 48 with none, +132 MB; a second run publishes nothing. The certificate attachment is a surrogate (the header's second plus agreement of verdicts) and every manifest says so: 135 of 222 attached, 84 of the 87 others being output_pre_L34/ states with no log of their own. kzz1e9/HeH2.13 and HeH9.7 lose the certified status their REPRODUCE.md states (the pair on disk is the L34 evaluate pass's certified=F) and carry stale_under: boundary_model_v1. The two catalog defects of the bounded experiment are exposed: kzz1e9/HeH0.083's first generation is the seed its forty passes started from (md5 1229b961323d7ac710146650da72454a) and the second, the state they left, is now latest_complete; wellmixed/HeH0.083 has its ending classified from the log. tests/state_generations.sh 27 of 27 (a publisher stopped by PID before the rename leaves the previous index; a mixed pair refused by every reader); the four D8a and D9 fixtures pass (16, 12, 10, 11).

What it left open. The retention policy for the 84 output_pre_L34/ generations with no verdict (the user's). No crash durability across a server failure is claimed, no exact algorithmic continuation, no best reference. 81 of the 134 published states were written by the evaluate pass over a solve not on disk. atomic_scalar_gj1132_kzz1e9/HeH2.13's _IC file is the earlier run's seed, not its solved state; which state L34 was handed is worth checking.

The bounded experiment of D8 step 8: the movement bound held at 1e-2

What was open. D3a's candidate solver defect (25 bounded passes leaving the transported state 1.58e-3 of x(He II) from the row's root) and L33's reading of the monotone halving of the composition movement bound as a throttle. The user's D3b decision opened one bounded experiment under the rule that closed the stage-2 solver program: its result decides whether an item exists.

What changed. The diagnostic environment key EXHALE_CARRIER_TRUST_HOLD (default unset; a positive value holds the bound and takes the halving branch out) in EXHALE_main.f90, and the row carrier_bound_hold (src/tests/grid_and_gates/carrier_movement_bound_hold.sh). Nothing in the acceptance interface moved.

How it was verified. Key absent: carrier_model_a_newton byte-identical to its entry text. The row: 5 PASS on the delivered build, carrier_bound_key_holds failing on the entry text. The three cases:

- carrier_model_a_newton, ten passes from the 25-pass transported state: held and halved runs **bitwise equal**; every relaxation ends on "the carriers stopped moving" with displacements 9.43e-4 to 4.71e-4, a decade below either bound; the He II distance from the row's root goes from 1.579137e-3 to -7.222846e-4, falling 7.52 per cent a pass and crossing zero, while the gated measure falls 1.87 per cent a pass. D3a's "moved away" was one side of a crossing.
- molecular_scalar_gj1132_wellmixed/HeH0.083: the bound binds at every pass, and held at 1.0e-2 the gated H2 row RISES monotonically, 7.37e-2 to 9.35e-2; in the one pass the runs differ, the halved step makes it fall (to 8.30e-2).
- molecular_scalar_gj1132_kzz1e9/HeH0.083, from .L22/i4_kz0083's exit state (the catalog publishes that run's seed): with the bound held at 1.0e-2 the row goes 8.15e-04, 6.29e-03, 1.93e-03 and pass 3 returns info = 2; the ten passes were not reached (a JFNK line-search collapse, lam near 1e-6, a pass costing 3660 s). The monotone descent L33 saw is a property of the small bound.

Result: no item on the movement bound, and no default recommended for change.

What it left open. The two catalog defects (taken by D8b). A JFNK line-search collapse costs the whole iteration cap before the outer loop can judge the pass. On a stiff stage the gated measure and the distance from the root move at different rates.

D9, steps 1 and 2: the budget of a solve by configuration, and seed compatibility before distance

What was wrong. Rev1 had turned one recorded history into a rule (3.3 cells a pass, 40 passes before a row could fall), which the second review withdrew; pick_seed.py's Physics.matches accepted two inputs differing only in the ionization-transport key, and tier 0 took the case's own state with no comparison; element_flux_closure.py continued on any nonzero info.

What changed. run_campaign.sh: a pass and a wall-clock ceiling for each configuration class, each an empirical allowance with its source (atomic_prescribed 40 passes and 30 min, atomic_closure_rung 20 for each EXHALE solve and 45 min for the rung, molecular_alteration 40 and 6 h, transported_ionization 90 and 6 h, the last wall clock not a measurement of its class); overrides in models/budget_overrides.txt refused without an authorized: sentence; a case at a ceiling recorded UNCERTIFIED INCOMPLETE. pick_seed.py: compatibility before distance in every tier (spectrum, K_{zz}, lower boundary, reservoir inventory, metals, mol, oxychem, carrier, he_diff, iontrans), the configuration read in three layers, an iontrans transition offered only when the case names it and ranked last, no certificate transferred. element_flux_closure.py sources the policy block of run_case.sh (RUN_CASE_POLICY_ONLY=1) instead of restating it; run_dtau0_first.log is gone. LHS1140b/MODELS.md sections 3 and 6.

How it was verified. tests/campaign_budget.sh 12 of 12 (stub runner), tests/seed_compatibility.sh 10 of 10 (RED on the tool before the item: the local-equilibrium state offered as tier1; six real cases choose the seeds they chose before), tests/closure_continuation.sh 10 of 10, and D8a's fixture 16 of 16 again. A first draft that compared only the tokens both sides stated readmitted 262 archived states to a molecular case; the layering is the repair.

What it left open. Step 3, the reference refresh; no declared matrix of seeds, resolutions and regimes was run; no bounded repeat of the L36e fiducial.

D2b: the lower boundary as a contact, upwinded after the acoustic matching

What was wrong. At zero and reversed flow the base face density was the midpoint of a smoothstep blend of the reservoir and interior isentropes, a model choice with no stated physics (L35); the user's decision A asked for a stated contact law, the direction decided after the acoustic matching, the transport operators with their own base conditions and any width a declared regularization with a convergence test.

What changed. base_boundary.f90, characteristic_base_face_state: acoustic matching by C- with the reservoir's pressure first ($v_b = v_i + (p_b - p_i)/(\rho_i c_i)$), moved ahead of the branch); the contact's direction read after it; entropy and composition upwinded with w_rev in {0,1} and no width. The reservoir owns the level at inflow and at a pressure-balanced contact at rest (the lower atmosphere as a heat bath on the time scales of a stationary solution, stated with its validity range at the head of the module); a reverse flow carries the interior's trace out with only the pressure imposed. The smoothstep keeps one use, the supersonic-outflow transition at M_i

= -1, as the declared regularization `base_face_mach_blend` (default $1e-8$). `EXHALE_BASE_BRANCH_ON_CELL1` is removed. Model identity `characteristic_face_ps_reservoir_C_minus_contact_upwind_v2`, reservoir prescription version 1. `viscous_conduction.f90`: `conduction_base_level_T`, the level's own temperature (`base_reservoir_temperature_at`), with the budget entry `conduction_base_heat_flux`. `binary_element_diffusion.f90`: the Dirichlet reservoir composition stated as a condition that drives a flux either way, and `element_base_diffusive_flux` recording face 0. `diffusive_photochemistry.f90` (comment): the carrier carries zero diffusive flux across the base face and its advection is upwinded on the face mass flux. `docs/input_schema.md` (`T_base` is the temperature of the level; the two model identities) and `LHS1140b/MODELS.md` section 4.

The one departure from the decision text, for the user. The direction is read from the wind window's mass flux where the window has standing (the existing thresholds $|M_{\text{wind}}| \geq 1e-12$ and $d_{\text{window}} \leq 2e-3$, now thresholds rather than blend weights) and from the matched face velocity v_b elsewhere, with a rounding floor $v_{\text{floor}} = 4 \text{ eps_mach} [(|p_i| + |p_b|) / (\rho_i c_i) + |v_i|]$, about $1e-15$ of the sound speed, at which the reservoir owns the contact. The reason is MEASURED: on the three certified catalog states v_b is -7.254, -2.098 and -3.415 cm/s while the base face carries +0.9995324 to +0.9999844 of the window flux; v_b there is set by the $1e-4$ pressure residual a converged state leaves at the base, two orders of magnitude above the flow's own velocity, so a bare sign of v_b would put every certified state on the reverse branch. The decision text said the window must not override a local reverse flow; on these states the window and the local face flux agree, and no state has been measured where the window has standing and the local face flux reverses.

How it was verified. `boundary_state` `contact_law` 22 assertions (seven FAIL on the entry text, three rows not compiling there), the face state bit-identical when the width moves from $1e-10$ to $1e-6$ ($1.50e-06$ and $1.80e-06$ on the entry text); `grid_and_gates` `base_branch` 11 (five FAIL on the entry text); `boundary_state` 35 of 35; the `grid_and_gates` rows 39, `hydrostatic_residual` 60 and `base_continuity` 35 all pass; the whole `grid_and_gates` 290 of 290 on the advisor's verification (READ, the plan's D2a row). Movement: `wasp_full` identical in every number; `mol_base_handoff` $1.185e-05$ (v, cell 203) and $2.403e-06$ (H2); `mol_carrier` $5.226e-06$ and $5.595e-06$; `lower_profile` $3.312e-04$ (cool, cell 495 near the outer edge); on the two molecular marching cases and `lower_profile` the window is silent and v_b reads reverse where the first cell read inflow. `carrier_model_a_newton` (30 marching steps, conduction on) $2.596e-05$, the conduction operator now reading the level's temperature, 4.7 per cent from the ghost's `p/n_part`. The three catalog states' evaluate routes are identical, with the same verdicts. On L35's rest fixture the midpoint is gone: three states take the reservoir and the atomic fiducial at rest takes the interior's trace ($v_b = -6.667$ cm/s, a resolved reverse flow).

What it left open. `write_setup_report.f90`'s label for `base_face_mach_blend` (corrected by the advisor; see below). `element_base_diffusive_flux` is recorded and not printed, because face 0 in `element_flux_profile.txt` would change a gated product. The ghost's `p/n_part` and the level's temperature remain two numbers 4.7 per cent apart on a molecular base.

What the user decides next

- **The direction rule of D2b.** Whether the one departure stated in the D2b subsection stands: the direction read from the wind window where it has standing, because a bare sign of v_b on a converged state is the sign of the base's $1e-4$ pressure residual (-7.3 , -2.1 and -3.4 cm/s on the three certified states against $+0.9995$ to $+0.9998$ of the window flux through the face).
- **D1b**, the default base cell width $2.0d-4$, only after every historical omitted-key input is pinned from the D1a record.
- **The D8 retention policy** for the 84 output_pre_L34/ generations that carry no verdict (nothing deletes them today).
- **D9 step 3:** re-solving the reference states under boundary model v2, including the three molecular states whose certificates are stale since D5b-2 (molecular_scalar_gj1132_kzz1e9/HeH2.13, HeH9.7 and .L22/i3_alt), and then the one golden refresh that follows.

Corrections made while the series landed

Corrected by the advisor as the items came in.

- **src/tests/stage_row_balance/stage_row_balance_tests.f90**, the header of stage_sum_rounding_bound, gave the bound at $g = 1$ as $1.55e-15$ and $2.66e-15$, which is $(5 \text{ nk} + 2) \text{ eps}$ without the factor 2 of its own derivation; it now reads $3.109e-15$ and $5.329e-15$, which the routine's DIAGNOSTIC line prints (READ, D3b section 8). The executed rows always used the correct expression.
- **src/modules/time_step/certification.f90**: the element-inventory comment named the statement made cell by cell with a hyphenated "per" compound, which this tree's wording rules forbid; it now reads "the cell-by-cell statement".
- **src/modules/files_I0/write_setup_report.f90** printed entropy-branch blend window, face Mach = before base_face_mach_blend, which after D2b no longer sets an entropy-branch window; the label is now supersonic-outflow regularization width, face Mach =.
- **src/tests/grid_and_gates/restart_round_trip.sh**, the row restart_lower_ghost_is_the_derived_boundary that D5b-3 added at an exact zero: the zero was a coincidence of equal seeds, since the ghost's solve closes to base_ghost_closure_tol and its last bits follow the seed the loader takes from the file's first cell; after D2b the seeds differ and the row reads $4.0e-16$, so it now asserts the round-trip rounding band $1e-14$, and whether the file's ghost rows are read is decided by the four ghost variants of src/tests/boundary_state/ (reason in docs/lhs1140b_stationary_D5b2_20260918.md section 11).
- **docs/lhs1140b_stationary_L33_20260917.md** section 6.3 gained an "Executed 2026-09-18" paragraph stating the bounded experiment's result on the case it designed.
- **The EXHALE.x in backup/regression/wasp_full/**, dated 13:27 on 2026-09-18 (md5 74b96cdcf887dcfee301ec547f012d88, the closing binary of section 12), was first reported as a stray. It is not one: run_check.sh copies the binary under test into every case directory before running it, so each case directory holds the binary of its last check (corrected 2026-09-19).

The golden refresh: still deferred by decision

No golden was refreshed in this series and no item wrote under backup/regression/golden/. The goldens are the 2026-09-17 set, and they no longer match the tree on the hp_* cases and on every case D7c, D5b-2, D5b-3 and D2b moved. The movements the memos measured, each against the item's own entry-text control (not against the goldens):

item	case	largest relative move
before this series (L36c to L36e, READ from D9)	hp_front, hp_zero_seed, hp_trace_seed, carrier_model_a_newton	1.6e-1 (helium column)
D6	hp_front (100 steps)	1.2e-14
D6	carrier_model_a_newton (3 passes)	4.100e-5 (v)
D7c	wasp_full (400 steps)	3.98e-6 (HeI where)
D5b-2	mol_base_handoff (12000)	6.19e-09 (cool), 1.0
D5b-2	mol_carrier (12000)	1.33e-11 (v), 1.29e-
D5b-3	mol_base_handoff	1.2629e-11 (v), 1.80
D5b-3	mol_carrier	1.4985e-11 (v), 5.64
D2b	mol_base_handoff	1.185e-05 (v), 2.403
D2b	mol_carrier	5.226e-06 (v), 5.595
D2b	lower_profile (12000)	3.312e-04 (cool), 2
D2b	carrier_model_a_newton (30 marching steps)	2.596e-05 (v), 5.212

Measured identical by every item that ran it: wasp_full under D1a, D3a, D3b, D4b, D5b-1, D5b-2, D5b-3, D6, D7a, D7b, D7d and D2b (moved by D7c only); wasp_he23off under D7c; mol_metals under D7a, D7b, D7d; mol_diffusion under D6 and D4b; hydrostatic_column under D1a; the three hp_* data rows under D3b. lower_profile was identical under D5b-2 and D5b-3 and moved under D2b. The matrix cases wasp_full_newton, mol_lyman_werner, mol_ir_bands, mol_sec_ion and oxygen_chemistry were not run by any item of this series, so their movement is not known; the molecular ones carry a molecular base and are expected to move with D5b-2 and D2b, which is an expectation and not a measurement.

The refresh is one refresh, after the reference states are re-solved under boundary model v2 (D9 step 3), with the movement against the 2026-09-17 set measured and reported at that time.

D9 step 3 and the golden refresh of 2026-09-19

Approved by the user on 2026-09-19 ("남은 것도 전부 진행"). Before the refresh the series closed four further code defects that the catalog re-evaluation exposed (docs/lhs1140b_stationary_D9fix_20260919.md): the ghost H2 closure of D5b-2 contracted linearly (slope 0.74 to 0.90 per pass) and now takes a secant step on its scalar equation and tests the residual at the returned state; transfer_h2 of the atomic-to-molecular seed overwrote the H2 that the molecular reservoir's ghost already carries and now partitions the neutral pool, conserving H nuclei to 4e-16; bg_ready, which item L36e had narrowed to transported_rows_exist(), again holds for every molecular run, since the molecular seed's local chemical root reads the background (src/tests/molecular_seed 25/0); the parse-dump mode now stops before EXHALE_setup.out is opened, so the parse corpus no longer truncates the setup report stored beside a case's results. The rounding-dependent carrier_bound_default_halves row became a DIAGNOSTIC. The catalog

runner gained `run_case.sh --evaluate`, writable `_IC` copies, `pick_seed.py` tier 0 on `latest_certified`, and the demotion of a parent that the current binary refuses (`docs/lhs1140b_stationary_D8b_20260918.md`, addendum).

The three $x0.01$ -XUV atomic catalog states are refused on the final binary and the refusal is physical: they were solved on the pre-D2b blended base face, and under the accepted contact law their base face flux is 38 to 178 times the window mean. They are demoted in their indexes and re-solved as a catalog item.

A metal-bearing transported-ionization reference now exists: WASP-121b with $\text{He } 2^3\text{S}$ and solar trace metals, Ionization transport: True, certified at outer pass 29 from the certified `wasp_full_newton` state (`docs/iontrans_metals_reference_20260919.md`); the transported stage sources equal the sweep's rows at every cell exactly. It is the regression case `iontrans_metals`, in the default list, with its outer-pass cap pinned in the new one-line file `outer_passes` (read by `run_check.sh` as `EXHALE_OUTER_PASSES`). `run_check.sh` golden snapshots the two `*_adv.txt` products only where the case's route writes them.

The goldens were refreshed once, on EXHALE.x md5 2c3b0acc9983aec03bb4f844294fed18 (make -q clean), after a strict check of all eighteen cases on that binary. The previous set is kept as `backup/regression/golden_20260917/`. Movement against it, largest relative change of any value in the two state files, MEASURED:

case	Hydro_ioniz.txt	Ion_species.txt	nature of the case
<code>wasp_full_newton</code>	5.5e-11	1.4e-10	converged, certified
<code>wasp_full</code>	3.9e-4	2.0e-6	fixed-step snapshot
<code>wasp_he23off</code>	7.4e-5	5.0e-6	fixed-step snapshot
<code>mol_base_handoff</code>	1.2e-5	8.0e-5	fixed-step snapshot
<code>mol_carrier</code>	5.2e-6	8.0e-6	fixed-step snapshot
<code>mol_diffusion</code>	1.3e-5	9.0e-6	fixed-step snapshot
<code>mol_ir_bands</code>	1.9e-5	2.5e-6	fixed-step snapshot
<code>mol_lyman_werner</code>	4.2e-5	2.9e-6	fixed-step snapshot
<code>mol_metals</code>	1.3e-5	2.5e-6	fixed-step snapshot
<code>mol_sec_ion</code>	4.9e-2	1.8e-5	fixed-step snapshot
<code>lower_profile</code>	3.3e-4	2.6e-4	fixed-step snapshot
<code>roundtrip</code>	4.1e-3	1.8e-3	fixed-step snapshot
<code>hp_front</code>	1.6e-2	1.6e-1	fixed-step snapshot, transported stages
<code>hp_trace_seed</code>	9.0e-1	1.0e0	fixed-step snapshot, transported stages
<code>hp_zero_seed</code>	9.0e-1	1.0e0	fixed-step snapshot, transported stages
<code>hydrostatic_column</code>	1.0e-7	8.3e-1 (on values of order 1e-180)	static column
<code>oxygen_chemistry</code>	1.9e0	1.0e0	fixed-step snapshot, oxygen network
<code>iontrans_metals</code>	new	new	converged, certified

The one converged state that existed before moved by $1e-10$. The large movements are all fixed-step snapshots of runs that are not stationary, whose trajectory any change of the physics moves; the transported-stage cases carry D6, D7a to D7d and D3 corrections to their sources, and the oxygen case was already known to move by order unity (item L30c). `parse_golden/` was refreshed the same day on 882 live inputs, its previous copy having been out of date since the Jupiter-radius constant became the IAU equatorial value.

14. PLAN_20260919_rev1: the review folded in, the boundary composition closed (2026-09-19 to 2026-09-20)

Plan of record: `docs/PLAN_20260919_rev1.md`, which folds `docs/PLAN_20260919_review.md` into the first text `docs/PLAN_20260919.md` after checking each of the review's ten findings against the source and, where a finding was about behavior, against an execution on this tree. Nine were confirmed and one (R2, the solved ghost electron count) was confirmed with a narrower reading, as a defect of a diagnostic and not of the equation of state. The plan's section 1b re-ran the review's central case twice here rather than quoting it.

Every number below is READ from the item memo named beside it, from the plan, or from the file named; where a memo labels a number MEASURED, the measurement is that item's. Three things were measured for this section and are said to be: the md5 of the tree's EXHALE.x, that make `-q` finds it current, and the movement table of the golden refresh, read from the strict check that preceded it.

The order of work changed from the first text: measurement and record correctness came before the boundary investigation, because the investigation's evidence is read through them.

P0: the case list the counts are of

What was wrong. The catalog counts quoted in the plans were enumerations nobody could repeat: 88 indexes with 74 certified in the first text, 137 with 83 in the review, 136 with 83 in the advisor's check of the same day. None of the three stated which directories it counted or what "certified" meant.

What changed (`docs/lhs1140b_p0_p4a_20260919.md`). `models/status.py inventory [--write]` publishes `LHS1140b/models/CASE_INVENTORY.{md,json}`: the selection rule in words and the name of the function that IS the rule, the instant it was taken, the binary of record with its md5 and its source list, the criterion for certified (the index names a `latest_certified` whose entry carries no `stale_under`), the totals by kind of directory, and one row per state index. The rule is one row per `state_index.json` under `models/`, walked recursively, hidden directories included, with a `kind` column; a narrower count is taken by reading that column and never by walking the tree again with another rule.

How it was verified. MEASURED by running it, 2026-09-19: 137 state indexes, 83 certified, of which 84 ladder cases (74), 46 closure rungs (9), 4 diffusion_check (0), 2 archives and 1 study directory. The advisor's 136 differs by the tree and not by the rule: `.L22/i3_alt` published its first generation that day, and it is the only index of the catalog whose oldest generation was published then. The review's 93 with 83 is the same table read narrowly, the 84 ladder cases plus the last iterate of each of the nine closure ladders.

What it left open. Nothing of the rule. The count itself moved twice more in this series, to 85 certified, by the

measurements of the fourth catalog pass.

P4a: the four facts of how a run ended

What was wrong. `classify_ending` of `models/run_case.sh` returned solved on `INFO = 0` before it tested that a complete finite state existed, and it took `INFO` from the last inner (`JFNK|PTC`) `done info=` line when the outer route printed no verdict. `solver_from_log` of `publish_state.py` repeated the same fallback in its own code, so a manifest could carry a verdict no route gave.

What changed. The route is determined first, from the run's own `Restart intent: echo`, and only that route's terminal event is its verdict; an inner solver line is carried as evidence and never promoted. Four facts are kept apart and all recorded: the process ended, the route stated its outcome, a complete finite state of that pass is on disk, and the state as written certified. `state_missing`, `nonfinite_state` and the new `stale_state` are reachable from `info = 0` as well, and conflicting evidence has its own class (`conflicting_evidence`, which is `info = 0` with a refusing certificate). `run_case.sh` and `publish_state.py` now read ONE rule (`RUN_CASE_POLICY_ONLY=1`) and `status.py` delegates to it.

One correction the advisor made to the item: the marching route prints six different endings on one `-> stopped:` line, and calling all six a relaxation snapshot is wrong in two, since a run that requested no time integration never advanced the state it wrote and a stop on a NaN is not a state to carry forward. The rule now reads the binary's own reason and gives `no_integration` and `nonfinite_state` their own classes.

How it was verified. New fixture `tests/termination_classification.sh`, 14 checks on synthetic logs: RED 1 of 14 against the previous text, GREEN 14 of 14 after. Every existing catalog fixture passes (27, 16, 12, 11, 10, 18, 18). Read only, over the 398 stored generation records: 311 name a log on disk, 306 are still the record's own log, and 44 of the 306 would be classified differently, all of them `no_verdict` becoming `no_integration` (37 closure rungs of `atomic_photochem_gj1132_kzzprofile`, the 4 `diffusion_check` cases, 3 generations of `molecular_scalar_gj1132_kzz1e9/HeH0.55`). No record stored as solved moves.

What it left open. Nothing was re-classified in place: the 44 are evidence for P4c. A record binds its log by `PATH`, and that path is reused, which P4d then closed by binding the bytes.

P5a: the capacity contract of a certification report

What was wrong (`docs/lhs1140b_p5a_20260919.md`). `add_entry` in `certification.f90` returned silently when the report already held `cert_max_entries` entries, and every entry writer then assigned into `rep%(rep%n)`, which in that state is the PREVIOUS entry. MEASURED through the public entry writer on the entry text: the hydrogen entry keeps its name and carries the helium equation's measure, `silent_overwrite=T`.

What changed. `add_entry` returns the index of the entry it created and every writer assigns through that index, so no `rep%(rep%n)` remains in the module (MEASURED). An entry that does not fit stops the run before any field is written, naming the equation, the capacity and the entry that stands last. `cert_max_entries` is no longer a literal: it is the sum of the inventory the builder writes (3 hydrodynamic rows, 7 carrier balances, 2 stage nucleus sums, 1 He/H partition, 10 trace element balances, 2 level balances, 7 eliminated-species closures) and stays 32.

Why it mattered more than it looked. The capacity was exactly consumed: READ from the run logs of the binary of record, a transported-stage configuration prints active equations 11 of 32 in the inventory, so ONE added equation of any kind would have overwritten the last entry silently.

How it was verified. RED on the entry text for the three new rows, GREEN after; certification 123 of 123 (109 before the item), ionization_stage_flux 45, element_operator 40, adv_static_limit 53, krylov_and_dogleg 334, steady_species_rows 197, all passing. wasp_full at 400 steps and mol_carrier at its pinned 12000 are byte-identical in their data rows between a control build of the entry text and the measured build, as they must be, since no present configuration reaches the capacity.

What it left open. Nothing. Raising 32 to another constant was refused as a repair: the number is the size of the inventory and is now derived from it.

P4b: the seed of a new solve, resolved once

What was wrong (docs/lhs1140b_p4b_20260919.md). The manifest of molecular_scalar_gj1132_kzz1e9/HeH9.7 g0004_20260919T103629Z_9b8394a3 carried a null parent with "not recorded by the run that wrote this state", while the run record it names carries the seed g0003_...ffa78b36 of its own case. The runner knew the seed and the manifest did not, because run_case.sh passed the publisher the generation an internal continuation continued, which is empty for a first solve however it was seeded.

What changed. The runner resolves the seed ONCE into runs/<run id>/seed_identity.json (schema exhale_seed_identity/1) and hands that one file to the publisher and to write_reproduce.py, so the manifest and REPRODUCE.md cannot name different seeds. The record carries whether the seed was established, its kind, the case and generation it came from or the md5 of both halves where no generation exists, who chose it (the pick_seed.py tier, an explicit SEED=, a listed candidate, NOSEED=1, or the runner continuing its own generation), and what was done to it: used as it stands, interpolated onto this case's cell centers, or converted from an atomic state with what the conversion changed. A seed of the case being solved becomes the parent; a seed of another case does not, and says in words where it came from. The contract is LHS1140b/MODELS.md section 9.12.

How it was verified. tests/seed_provenance.sh 33 of 33, against 5 of 33 on the reconstructed entry text; the eight existing catalog fixtures pass; status.py inventory still reads 137 indexes with 83 certified. A bounded solve was run end to end through run_case.sh on a scratch copy of carrier_model_a_newton and published two generations, the first carrying the SEED= record and the second the continuation with the first as parent.

What it left open. Read only, of the 398 published manifests 226 carry a null parent: one has its own run record and is R9's case, five more can be read from a REPRODUCE.md and are each of ANOTHER case, and the remaining 220 predate the publisher storing a run id. Nothing was rewritten: that is P4c. The closure runner was believed to publish without a seed, which P4b-2 then found to be wrong in a different way.

Fixed in passing. tests/termination_classification.sh set its pass mark in the same second in which it wrote the state, so check T7 read stale_state about once in two runs; the mark is now set a minute into the past. The rule itself is not at fault: a solve never writes its state within a second of the mark.

P5b: the executable a test selects, and the one it says it ran

What was wrong (docs/lhs1140b_p5b_20260919.md). No suite under `src/tests/` printed or checked the identity of the binary it ran (READ: the string `md5` occurs in no pre-change suite script except where an OUTPUT file is hashed), and the selection was made by four different names. MEASURED: a run that set `EXHALE_EXE` to a missing file and reached `residual_determinism` was answered by the tree's own build, with no refusal and no statement.

What changed. One policy, stated once in `src/tests/exhale_exe.sh` and sourced by every suite that runs the binary: `EXHALE_EXE` selects it; `EXHALE_RESID_EXE`, `EXHALE_SPECIES_EXE` and `EXHALE_COUPLED_EXE` are accepted aliases; several names resolving to one file are one request, and names resolving to different files are refused rather than silently chosen. Every such suite prints the path, the `md5` and which name selected it before anything runs, asserts that the file about to run carries the requested `md5`, and, where it copies the binary into a working directory, asserts the `md5` of the copy. `EXHALE_EXE_IDENTITY_ONLY=1` states the identity and runs nothing.

How it was verified. The new sentinel suite `src/tests/executable_identity` runs 21 rows, none failing, in under a second, by pointing `EXHALE_EXE` at a copy of the binary whose `md5` differs and requiring each suite to report the COPY. RED and GREEN are both executed on the same request: the pre-change scripts do not refuse a missing file named in `EXHALE_EXE` and never name the file that ran; the delivered ones refuse it by name and exit 1. `run_mode` 32, `boundary_state` 38, `charge_exchange_rows` 251, `spectrum_type` 171, `molecular_seed` 26 and `fuv_band_ledger` 25 all pass.

What it left open. A standing failure the item exposed and did not touch: `residual_determinism's` `closure_spread_within_the_row_tolerance_atomic_elem_newton` measures $1.715\text{E}+04$ against a bound of 1, identically before and after the item on the binary of record. It belongs to the residual contract and joins the open list with the two `coupled_block_jacobian` rows of R5.

R2 and P1 step 1, experiments 1 and 2: the ghost's seed decides the certificate

What was wrong (docs/lhs1140b_p1_step1_20260919.md). R2: the solved ghost electron count summed H II, He II, He III and the metal stages and declared the molecular ions negligible, while `calc_ne`, which the production electron density comes from, counts H2+, H3+ and HeH+ with their charges. The defect is a diagnostic's, not the equation of state's, and 1 per cent is not a rounding argument for leaving the charges out.

What changed for R2. The count is now `calc_ne's` own count of the composition the sweep returned; the second sum is gone, and the comment at the assignment states what the count is of, its charge carriers, its units and its validity range. MEASURED: the reported count moves by $1.022\text{e}-2$ relative, which is the molecular share the review measured, and nothing numerical moves with it (`mol_carrier` at its pinned 12000 steps byte-identical in both files between a control build and the measured one).

The instrumentation, all default off. `EXHALE_GHOST_COMPOSITION_SEED` (the ghost enters the sweep from the state file's ghost rows, the previous sweep's ghost, or a named file, instead of the reservoir row whose partition is the first physical cell's), `EXHALE_GHOST_SEED_WRITE` (the installed ghost rows written in that format), and `EXHALE_GHOST_RECORD` (one record per evaluation: the ghost's thermodynamic state, counts, fractions, charge and elemental budgets, the residual it was accepted at, the partition residual, the closure passes, and the base row with both

face fluxes, their signed difference, the row scale and its rounding floor).

The decisive measurement. At one frozen physical column, reservoir, spectrum, options, executable and thread count, on the refused generation molecular_scalar_gj1132_kzz1e9/HeH9.7 g0004_...9b8394a3:

ghost seed	cell-1 mass row	in floors of that row	verdict
the first physical cell's partition (the rule then in force)	8.3353e-08	111.29	REFUSED at cell 1 against 7.5e-9
the ghost rows the state file carries	2.9229e-09	3.903	CERTIFIED, the stored verdict reproduced at cell 210
the same rows with 1e-3 of the H2 moved into H I	2.9248e-09	3.905	CERTIFIED, same verdict cell and measure

The two ghosts differ by $4.7\text{e-}3$ (H II, He II) to $5.5\text{e-}2$ (H2+) in the trace ions while p, rho, T and the particle count agree to $1\text{e-}14$ to $1\text{e-}10$, and BOTH are accepted at reaction residuals of about $3\text{e-}26$ with their imposed partition closed to 0. The exact replay is byte-identical in both state files and in the ghost record. Between the first two seeds the base face mass flux moves by $7.8\text{e-}8$ relative and the signed difference of the two face fluxes by $2.1\text{e-}4$, while the row moves by a factor 28.5: the row is not the difference of the two fluxes the record prints, the face areas and the cell volume stand between them, and both rows are above their own rounding floor.

How it was verified. New fixture boundary_state/ghost_seed_and_ghost_record.sh 9 of 9 (3 of 9 failing on the control build), boundary_state 38 of 38, carrier_retry 137 of 137, grid_and_gates 295 of 295.

What it left open. Whether the two accepted states are two roots of the ghost's system or one root the cell solver stops short of, which experiments 3 to 6 then answered.

P1 step 1, experiments 3 to 6: one root, and the sweep does not reach it

What the experiments found (docs/lhs1140b_p1_step1b_20260919.md). There is ONE root, and the composition sweep does not reach it: the ghost it returns keeps a measured fraction of the composition it was seeded with. Three measurements decide it.

1. It is not unresolved iteration. Over six decades of EXHALE_IEQ_TOL (the default $1.49\text{e-}8$ down to $1\text{e-}14$) the two states do not approach each other at all: each stays put to $1\text{e-}15$ while they remain $4.7\text{e-}3$ to $5.5\text{e-}2$ apart, and the cell-1 row reads $8.33528\text{e-}08$ and $2.92293\text{e-}09$ at EVERY rung. The ladder does reach the solver: at $1\text{e-}14$ hybrid1 answers info = 3 on 225 of 508 cells, so the returned iterate is already pinned far better than the difference.
2. Neither state is a fixed point. Seeded with the refused ghost, in the refused ghost's own column, the solve LEAVES it, returns a ghost $5.5\text{e-}2$ away in H2+ and certifies.
3. The seed-to-ghost map has a gain: seeds $9.7\text{e-}1$ apart return ghosts $4.7\text{e-}3$ apart, and a seed perturbed by $9.8\text{e-}1$ in H I returns a ghost $5.3\text{e-}5$ away. A converged root does not remember its start.

No present control names the shortfall: at both states the network residual is $3e-26$, `hybrd1` returns `info = 1`, the imposed partition closes to 0.0, and the cell is accepted as a converged root. Both also satisfy $x_{H2} = x_2 (1 - x_{ion})$ exactly, so the handoff closure is not what separates them.

The other three experiments: the temperature closure removes most of the difference (the two branches' $H2+$ fractions go from $5.5e-2$ apart to $4.3e-6$, the rows from a factor 28.5 to 1.34, and both certify); holding the interior's eliminated species changes the base row in no printed digit, so the interior refresh is not a term in the base row; and the reload ladder settles, the ordinary reload giving the refused state three times from either start and the file-seeded reload giving the certified state in at most two reloads. Nothing drifts and nothing cycles.

How it was verified. New fixture `interior_composition_held` 4 of 4 (1 of 4 failing on the control build), `ghost_seed_and_ghost_record` 9 of 9, `boundary_state` 38 of 38, `carrier_retry` 156 of 156.

What it left open, and what it decided. P1 step 2 is therefore an accuracy contract, not a branch statement (there is one root) and not a reformulated base row first (a row that hid this would hide a ghost 5 per cent wrong in its ionization). Two standing failures were reported and not changed: `coupled_source_step` fails 2 of its 44 rows on the entry text as well, which reads as a suite that no longer reaches the events it measures; and a metals-off run writes its metal columns at a floor of order $1e-301$ that does not survive a load and a write exactly.

P1 step 2: the boundary composition is a fixed point of the sweep

What changed (`docs/lhs1140b_p1_step2_20260920.md`). The sweep is a map from the composition it is entered at to the one it returns, and `ioniz_eq` now applies that map again at the ghost it returned, with the interior restored at each application, until the ghost's composition and its heavy-particle plus electron count have both stopped moving; at twenty applications it stops the run by name with both moves. What the routine returns for the physical cells is still one application of one map, so the steady residual is still one application of one operator. The seed is now the boundary model's own statement, "the reservoir row at the first sweep, the ghost the previous sweep returned after it", the model identity moves to `v3`, and `load_IC` reports a `v2` state instead of refusing it.

Why the accuracy is what it is. From step 1's measurements, $4.7e-3$ of ghost composition moves the base continuity row from 3.90 to 111.3 of its own rounding floors, so one floor of that row costs $4.4e-5$ of ghost composition; the inner composition solve is at the resolution of the arithmetic by $1e-13$, and the map reaches $1.5e-5$ after one application and $4e-13$ after two. The temperature leg is inside the contract: the ghost's particle plus electron count is what turns its temperature into the pressure the hydrostatic continuation builds, so it is held to the same accuracy (MEASURED: it closes to $3.7e-16$, $2.4e-16$ and $4.8e-15$ for the three seeds).

How it was verified. The refused generation CERTIFIES with the cell-1 row $2.923E-09$, reproducing the stored verdict at cell 210 from all three seeds; `boundary_state` 38 of 38 and the new fixture `ghost_composition_fixed_point` 11 of 11 (4 of 11 failing on the control build). Against the goldens, strict: `wasp_full`, `wasp_he23off` and `lower_profile IDENTICAL`, `hp_front` moving by $9.7e-12$. Movement carried to the golden refresh, nothing refreshed at the time: `mol_base_handoff` $9.9e-12$, `mol_diffusion` $4.4e-9$ and $7.4e-9$, `mol_carrier` $1.6e-11$ at 400 steps, `mol_metals` $4.2e-4$ in one trace column at 400 steps, with `log10 Mdot` unchanged in each; cost 1.18 times the wall clock on `mol_carrier` at 200 steps.

A rule the item found. Nothing that separates one sweep from the next may turn over between the applications. Letting

bg_ready turn over broke mol_carrier outright (185 cells clamped at step 0); it is held across them.

What it left open. The base row still rests on a near cancellation; the reservoir's prescribed particle count and the ghost's own are still two quantities (P6c then measured the distance); six molecular catalog states still refuse for reasons the ghost composition does not reach; the atomic ghost is outside the contract; and no full stationary solve had yet run under it.

P1 step 2b: the accuracy belonged in the composition's own units

What was wrong (docs/lhs1140b_p1_step2b_20260920.md). The contract introduced a regression: grid_and_gates coupled_carrier_h2 failed 2 of its 11 rows, both runs stopping with ERROR STOP ioniz_eq: the ghost composition is not a fixed point after twenty applications. The cause is none of the four candidates the item listed, each of which was tested and refuted. The contract asked an accuracy in a norm the sweep cannot deliver in and no quantity the boundary builds reads: each ghost species against its OWN value. MEASURED, the ghost settles into an exact two-cycle in which H I moves by $6.5\text{e-}12$ of the cell's composition and H3+ by $2.2\text{e-}12$, while against their own values those same moves read $4.1\text{e-}9$ and $5.1\text{e-}3$; the same cycle moves the ghost's particle plus electron count by $4.9\text{e-}12$ and the base row by 0.06 of that row's own rounding floor. A deterministic map on a revisited point cycles for ever, so the twenty-application bound was reached with certainty.

What changed. The move is now the largest absolute change of any species of the two lower ghost cells in the composition's own units, held to $1\text{e-}11$, and the count leg keeps $1\text{e-}6$ against its own value. The compare floor is gone, since in those units nothing needs keeping out. $1\text{e-}11$ is a quarter of the base row's rounding floor (one floor is bought by $3.8\text{e-}11$ of composition) and above every plateau measured ($6.8\text{e-}12$ on the failing case, $3.5\text{e-}12$ on the hot-Uranus carrier state, $1.3\text{e-}14$ to $3.3\text{e-}13$ on the LHS 1140 b generation). The reason the earlier states closed is measured: their trace ions sit three decades higher.

How it was verified. RED and GREEN on the fixture that carries the regression, exit status 1 then 0; grid_and_gates 305 of 305 including the two rows, boundary_state and ghost_composition_fixed_point passing, the refused generation still CERTIFYING with the cell-1 row $2.923\text{E-}09$ and the contract reaching $6.84\text{E-}14$ in three applications, and wasp_full (400) and lower_profile (pinned) byte-identical to the control. Movement of the molecular cases against the control, with log10 Mdot unchanged in all four: mol_carrier $8.5\text{e-}12$ and $4.1\text{e-}11$ at 400 steps, mol_metals $3.6\text{e-}11$ and $1.9\text{e-}05$ in one trace column, mol_base_handoff $5.0\text{e-}09$ and $8.4\text{e-}09$, mol_diffusion $3.3\text{e-}09$ and $6.1\text{e-}09$.

What it left open. The accuracy sits a factor 1.5 above the lowest plateau measured, so a configuration whose composition solve plateaus higher will stop the run; the bracket is stated in the code and the memo, and the answer if another case meets it is to measure that plateau and not to widen the number. The atomic ghost is still outside the contract, the base row still rests on a cancellation, and no full stationary solve had run under the contract when the item closed.

P6d: the charge-exchange turnover says which cell it answers for

What was wrong (docs/lhs1140b_p6d_p6e_20260919.md). cx_add_to_turnover read the thread-local rate coefficients and the metal base without the cell guard its neighbors carry, and the turnover sets a normalization that chemical acceptance reads.

What changed. The entry point takes the temperature of the cell whose turnover it is bounding and goes through `cx_require_cell`. The cell is optional in the interface, because two of the three call sites sit in a file another item owned this session; a call that states no cell is refused when the thread holds none at all.

How it was verified. Whether a production call ever ran on another cell's rates was MEASURED before the guard was added, with an instrumented build that counted and logged instead of stopping: on `wasp_full` (400), `mol_metals` and `mol_carrier` (12000), `hp_front` (100) and `iontrans_metals` (90 outer passes) the entry point was not reached at all, and a positive control shows the instrument would have seen it. READ from the source, the reason is structural: the branches that reach the turnover are entered only when a cell solve is rechecked, and the entry points that evaluate the measure on a finished state install the cell first. So the defect was a missing contract and not a wrong number in these runs. RED on the entry text is a compile refusal (the entry point has no cell to state); GREEN gives the four new rows, and `charge_exchange_rows` runs 255 assertions, all passing.

What it left open. The three call sites, one line each: two in `ionization_equilibrium.f90`, and the one in `constrained_chemical_equilibrium.f90` which sits on the molecular rung path that none of the five runs entered, so making it a stated cell without that measurement would turn an unexercised path into an abort.

P6e: the elemental base flux record belongs to an evaluation

What was wrong. `element_base_diffusive_flux` was assigned from $J_f(0)$ of the last operator call, so printing it after a different evaluation could describe a different state. The item found more than that: the old record was IDENTICALLY ZERO, not small, because the gradient and drift coefficients are built on faces 1 to N-1 and faces 0 and N are left at zero, so $J_f(0)$ was zero whatever the state. The flux the Dirichlet reservoir composition drives sits at face 1, and the plan's earlier figure of $-5.57\text{e-}13$ on `lower_profile` was that face's number, not the base face's.

What changed. The record is now `element_base_flux`: the evaluation that wrote it, the first solved face, and the advective and diffusive halves at the base face and at that face, in $\text{g cm}^{-2} \text{ s}^{-1}$ positive outward, with a reporter that says no evaluation has written one rather than printing an uninitialized number.

How it was verified. MEASURED at the last evaluation of two runs, the diffusive half at face 1 reading $-5.5826627\text{E-}13$ on `lower_profile` and $-3.0298600\text{E-}13$ on `mol_diffusion` while the base face's is a signed zero; and the second reason the record needs its evaluation, the advective half at the base is not steady on `lower_profile`, reading $+4.47\text{e-}8$ at evaluation 1079, $-3.35\text{e-}10$ at 11220 and $+6.29\text{e-}10$ at 12000 as the base sound wave turns the face flux over in sign. Report only, and it shows: `wasp_full` (400), `mol_metals`, `lower_profile` and `mol_diffusion` (12000 each) are byte-identical in both state files between a control build of the entry text and the measured one. `element_operator` 50 of 50 (RED is a link failure on the entry text), `stage_row_balance` 10 of 10.

What it left open. `element_flux_profile.txt` keeps its columns: adding the base face would add a row that `reduce_element_flux_windows` feeds into a reported product, so it waits for a golden-refresh decision. `element_base_flux` still has no reader; the stationary route's budget is where it belongs.

Corrected by the advisor while the item landed. `backup/regression/iontrans_metals/NOTE.txt` said "No golden yet", which stopped being true when the matrix was snapshot on 2026-09-19.

P6f: the transit spectrum of a solved state

What was wrong. The first text said the transit tool reads only the advection-corrected profiles. It does not: EXHALE_TRANSIT_STATE=solution selects the solved state pair (READ). What the item found instead were three defects of the tool (docs/lhs1140b_p6f_20260919.md): the default selection on a run with no _adv products died inside numpy.loadtxt with a bare FileNotFoundError; a saved curve named its inputs by path only, so a number could not be traced to the state that produced it; and the O I ground-term populations were read from OI_levels_adv.txt whatever the selection, so a solution-state O I line would have used post-processed level populations, and on a stationary state that file is absent and the line was skipped silently.

What changed. The refusal now names every missing file, says that a stationary solve writes no such products and names the other selection; every saved curve carries the path, md5 and size of each file read and the state's own provenance, coupling, boundary model and reservoir lines; the O I level file follows the selection; and the tool states what the two selections mean, that the _adv pair is a post-process of a marching state (a steady advective correction, first-order upwind at the bulk velocity, no eddy term, refused where the steady assumption fails) and the solved pair is the state the solver produced and the certification judged. No _adv file is manufactured for a stationary route.

What the choice costs, MEASURED on wasp_full, which has both states: He I 10830 peak depth -0.64 per cent of itself, H-alpha -1.28, H-beta -3.48, Ly-alpha unchanged at its saturated ceiling, and the optically thin metal lines +0.1 to +0.8. The cause is stated rather than guessed: the metastable helium column is 12 per cent LARGER in the solved state, and its line is nonetheless shallower because the advection correction moves that column outward (column-weighted mean radius 1.2941 against 1.3112 R_p), and a saturated line's disk-averaged depth follows how far the absorbing shell reaches. H-beta moves most because it is the line still on the linear part of the curve of growth.

How it was verified. src/tests/transit_state 12 of 12 (5 of 12 failing against the entry text), src/tests/transit_census 21 of 21, and the default path's ten products on wasp_full identical outside their comment headers. The certified stationary state iontrans_metals, which has no _adv products, now produces a full spectrum including the O I 1302 line.

What it left open. Nothing of the item. A run directory is written again by the next run, so the digest and not the path is what ties a curve to its numbers, and reading a resolved generation is preferred over a moving output/.

P6b: the time-step restriction of the cells that are evolved

What was wrong (docs/lhs1140b_p6b_p6c_20260920.md). eval_dt took $CFL \min_j dr_j / (|v_j| + c_j)$ over $1-N_g:N+N_g$, and two things are wrong with that for the scheme in use. The cell-centered width dr_j is not the width the spherical finite-volume divergence gives a cell at its outer face, $V_j/A_{\{j+1/2\}} < dr_j$, so the restriction was too LONG by dr/r (1.96e-4 on the production base grid, MEASURED). And the minimum ran over cells that are never evolved: MEASURED, an outermost ghost given a sound speed a thousand times the column's shortened the entry text's step by 89.7 per cent, although Apply_BC overwrites that row after every stage.

What changed. Each evolved cell now gets $CFL \ V_j / \max(A_{\{j-1/2\}} S_{\{j-1/2\}}, A_{\{j+1/2\}} S_{\{j+1/2\}})$ with S_f the fastest signal bounding that face, minimized over cells 1 to N, derived at the routine for the divergence RK_rhs assembles, the Riemann fluxes it uses and the SSP-RK3 that integrates it. R6 is kept: the base and outer faces are faces

of cells 1 and N and their Riemann problems do restrict them, through the ghost states that bound those faces. In the Cartesian uniform limit the expression returns $CFL = \Delta r / (|v| + c)$, so the CFL: key keeps its meaning and nothing is recalibrated.

How it was verified. Three assertions were added to `grid_and_gates/hydrostatic_residual`, reading 1.96e-04, 8.97e-01 and 1.96e-04 under the entry text and exactly 0 now. MEASURED movement: the first step shortens by 2.8e-4 to 3.2e-4 on every regression case. The case that reaches a solved steady state, `wasp_full_newton`, gives the same state to 5.7e-12 over ρ , v , p , T and the same $\text{Log}_{10} \dot{M} = 13.33$, from 4709 and 4712 marching steps. `wasp_full` and `wasp_he23off` reach their du stop in 12251/12261 and 12236/12246 steps with the same $\text{Log}_{10} \dot{M} = 13.36$, and every column is inside 1e-3 except the velocity of rows 226 to 230, $r = 1.074$ to 1.077 , where it alternates in sign from cell to cell and moves by 18 to 26 cm/s (the median velocity row moves 1.3e-3, and cell 226 is the cell whose mass row the certification already refuses). The fixtures pinned at a step count on a state still moving move with their trajectories, as a fixed-step snapshot of a run that is not stationary must. The stationary restart route is untouched: `carrier_model_a_newton` is byte-identical over three outer passes, because that route sets $\text{dtau0} = 1$ and the JFNK floor does not bind.

What it left open. Nothing of the derivation. `eval_dt` also supplies the direct steady route's initial pseudo-time, which prints $\text{PTC dtau0} = 1.47\text{E-}04$ under both binaries, unchanged at the precision that line carries.

P6c: there is no 4.7 per cent inconsistency

What was claimed, and what is measured. The first text carried a "4.7 per cent inconsistency" between the ghost's temperature and the level's. The item evaluated the four relations separately, each quantity at its own radius, on `carrier_model_a_newton` and on the LHS 1140 b He/H 9.7 generation P1 closed. The equation of state closes at 6.7e-16 and 2.2e-16 against a census summed from `Ion_species.txt` independently of the code, and the census agrees with the code's own counts bitwise; the two ghost cells lie on one isentrope whose index is the same from the pressure and from the temperature to 2.9e-07 and 1.2e-06; the ghost's own particles per unit mass and the reservoir's prescribed value stand at 2.7e-08 and 1.3e-07; and the composition and its count are fixed points of their own solve to 1.2e-07 and 4.0e-12.

Where the number came from. It is a distance between a particle count PRESCRIBED AT THE BASE LEVEL and a count SOLVED IN THE LOWER GHOST CELL one cell below it, which is a difference of location together with a difference of quantity. MEASURED, that ratio is 4.87e-02 on the LHS 1140 b state, whose base cells are 7.8e-02 pressure scale heights, and 5.50e-03 on the hot-Uranus grid of `carrier_model_a_newton`, the state D2b attributed it to: it follows the GRID. The ghost offsets are the cell average of a convex profile against the point value at its centre, and the two pressure offsets stand in the ratio 97.3 against the 104.2 of $(\Delta r/H)^2$. The ghost's temperature and the level's own stand at 9.1e-07 at the radius where both are defined, which is the size of the movement D2b itself measured when the conduction operator changed from one to the other (2.6e-05 in one velocity).

What changed. The code was left alone and three wordings were corrected, in `base_boundary.f90`'s comment, in the D2b memo and in the P1 step 2 memo; `hydrostatic_column` rerun with the rebuilt binary reproduces the data rows of both state files byte for byte. The comment at `conduction_base_level_T` of `viscous_conduction.f90`, whose file belonged to another item that day, was corrected afterwards by the advisor and now states the measured distances and

why the old number compared two locations.

What it left open. The real seam is smaller and is named: the ghost's p and ρ are built with the reservoir's particles per unit mass while the caloric map uses the composition's own count, $2.7e-08$ and $1.3e-07$ apart on these two states, and both counts are printed every run so that a state where it opens is visible.

P4b-2: a flux-closure rung publishes, with its seed

What was wrong (docs/lhs1140b_p4cd_20260920.md). The item as written said `run_closure.sh` publishes the rungs through the same publisher and hands it no seed. READ, the delivered text of `run_closure.sh` and of `src/utls/element_flux_closure.py`: neither names `publish_state.py` at all, so the rungs were publishing nothing, and their 46 indexes came from the legacy import. A rung's state was whatever `k<NN>/output/` happened to hold.

What changed. The closure runner publishes each rung through the one publisher and resolves its seed first: `k00`'s seed is the state the driver mapped onto the current cell centers, with the directory it was read from and the md5 of both halves; `k<NN>`'s is the rung before it; and the transformation is read from what the driver left, a rung carrying a seed. `log` having been mapped again and one with none copied as it stands. The publication is idempotent, and the ending rule is sourced from `run_case.sh` and never restated. The contract is `MODELS.md` section 9.13.

How it was verified. `tests/closure_seed_provenance.sh` 22 of 22, against 4 of 22 on the reconstructed entry text, with no binary and no closure driver run.

What it left open. The rung before is a state directory of its own, so its generation is not in the chain this rung's index resolves: parent stays null and says why, and the identity is in seed.

P4c: the provenance recovered without rewriting anything

What changed. `models/recover_provenance.py` writes `<state directory>/provenance/<generation id>/provenance_recovered.json` beside each generation and never inside it: what was recovered, the file it came from with that file's md5 and size, the rule, the date and one of three confidences, with every candidate kept where several match. A later recovery writes the next version and names what it supersedes; nothing is rewritten or deleted. A reader gets a recovered fact ONLY when it is labeled established, and says "recovered" rather than "recorded": `status.py`'s seed cell, and `pick_seed.py --seed-provenance`, which is off by default so every existing caller sees the same lines. The contract is `MODELS.md` section 9.14.

How it was verified. MEASURED over 398 published generations in 137 state directories, all of which gained an attachment: 176 established (172 the manifest's own parent, 4 a run record naming the generation), 10 inferred from a single `REPRODUCE.md` window, 212 not established, and $4 + 10 + 212$ is exactly the 226 null-parent population P4b measured. That recovers more than P4b's 1 and 5, because a run record names the generations it published whether or not the manifest names the run, and because a case keeps the records it superseded. MEASURED, not one of the 398 manifests has a modification time inside that day's work. `tests/provenance_recovery.sh` 34 of 34, against 6 of 34 on the entry text.

What it left open. The ten inferred facts stay inferred and no reader uses them: a run window is not a name.

P4d: the records bound to their logs, and the four that were wrong

What was wrong. A record named its log by a path the next run of the directory reuses, and six REPRODUCE.md files carried a superseded seed sentence.

What changed. At publication a record is bound to the log BYTES (certification.source_md5 and _bytes, ending.source_md5), and for what is already published each attachment states whether the file at that path still carries the generation's block.

How it was verified. MEASURED over the 398 records: 1 names no log, 86 name a log that is gone, and of the 311 that name one on disk, 226 still carry the block and 85 do not. The five P4d named are in those 85, and so are 80 more that name a pp.log a later evaluate pass overwrote, which the earlier count did not look at. The four .L35/ study records were rewritten with their old records kept whole beside them, and the rewrite exposed a second defect in the same four: their pass table and verdict belonged to the run that was STOPPED and not to the state in output/ (g1000_hllc stated 11 passes and info = 2 where its run.log ends info = 1 after 8).

What it left open. The two pre_D9s3/ copies were deliberately not rewritten, since a rewritten copy of a record is a record of nothing; each carries a NOTE.md stating, measured there, that its input.inp resolves the spectrum one directory short and that its index names generations it does not hold.

Two stale things this item found. MODELS.md sections 7 and 8 had been stale since P4a (two rows move to not solved, MEASURED to be P4a's classification and not this item's), and status.py's seed cell had been an unlabeled inference, printing a seed read from a run window as though it were recorded; its wall-clock and outer-pass cells still are.

The certification rule of 2026-09-20: a state that passes the certificate is certified

What was wrong. The three molecular states that PASS certification under the binary of record carried no certified generation, because run_case.sh --evaluate resolved latest_certified and refused where the index named none, and an evaluation made outside the runner publishes nothing. On the user's instruction this is backwards: a certificate is a measurement of the state as written, so a state that passes it is certified whatever the run that produced it ended as, and the procedure must not decide otherwise. The old refusals were the boundary and not the states: they were made under boundary model v2, whose ghost composition was seed dependent.

What changed. The rule is stated in one place in code (publish_state.apply_evaluation_verdict, with its halves demote_refused_parent and promote_certified_evaluation) and one place in prose (MODELS.md section 9.15): a certificate is a property OF THE STATE, measured, and the generations, the index, the parentage and the ending classes record the measurement and never confer it. --evaluate measures a named generation, or latest_certified where the index names one, or latest_complete where it does not, and says which it took and why; an evaluate child whose certificate says CERTIFIED makes that child latest_certified, with the index recording that the certificate came from an evaluation and under which binary. The guard rails, each tested: the certificate must be of the bytes the generation carries; the state must be complete and finite; an evaluate generation must be an evaluation of its parent's own two halves bitwise; and a refused child still demotes.

Four bookkeeping defects the instruction exposed, each fixed with the rule. (1) --evaluate refused to run where the

index named `no_latest_certified`, which is the defect the instruction names. (2) `certified_here` did not require `storage_complete`, so an incomplete or nonfinite state carrying a CERTIFIED block could have taken the reference (MEASURED: none of the 83 certified generations is affected). (3) A refusal was recorded as unknown: the evaluate route adds a further (certification) heading at the end of a refusing run saying how the pair was written, which measures nothing, and `certification_from_log` took the LAST heading, so `demote_refused_parent`, which fires on NOT CERTIFIED, would not have taken a certified reference away; the reader now takes the last heading whose own block states a verdict. (4) Wording that read as a verdict on the state: `status.py` and `pick_seed.py` said the index does not certify a case, where what is true is that no measurement has been recorded, which says nothing about whether its state solves the equations.

How it was verified. `models/tests/certified_evaluation.sh`, 18 checks: RED 3 of 18 on the entry text, GREEN 18 of 18 after. The three that pass on the entry text are the parent's manifest not being rewritten, a refused evaluation demoting, and the solve path under a wall ceiling, which is the measurement behind the statement that the solve path already did this. Every existing catalog fixture passes, eleven of them. `publish_state.py` `verify` passes on 135 of the 137 indexes, the two refusals being the `pre_D9s3` copies, which refuse identically on the entry text.

What it left open. `molecular_scalar_gj1132_kzz1e9/HeH9.7 g0004` now carries both `stale_under` from the 2026-09-19 refusal by the previous binary and `certified_under` from this pass: two measurements by two binaries, both kept, with the reference sitting on this pass's child.

The catalog under the contract: the third and fourth passes

The third pass (`docs/lhs1140b_catalog_refresh_20260919.md`, third pass), on the binary of record EXHALE_75d55d9d.x, md5 75d55d9d4fd0e748cd01d6e35713e34c, which differs from the previous catalog binary in nine of its 169 production sources, each named with the item that changed it.

- **83 of 83 re-evaluations CERTIFY.** Every case whose index named a `latest_certified` generation was handed to `run_case.sh --evaluate`, none refused, the wind-window mean of $\rho \propto r^2$ is character for character the parent's in all 83, the two-decimal log10 Mdot line is unchanged in all 83, and the largest change of the He 10830 red-pair equivalent width is 2.6e-14 per cent Å. The boundary composition contract applies to none of them and prints no line in their logs, its condition being a molecular network with helium: that is applicability and not a missing measurement.
- **The nine molecular states, evaluated read only: 3 CERTIFY, 6 refused.** All nine reach the ghost fixed point in two or three applications at 6e-15 to 5e-12 against 1e-11, so the contract is not what separates them: the six are refused by rows the ghost does not reach, a carrier balance H2 in the wind in all six, an interior energy row in three, and in the photochemical HeH9 the base rows at order one.
- **Five re-solves were run and none certified.** `kzz1e9/HeH9.7` converges and does not arrive: its gated carrier row falls monotonically to 5.13e-4 against 1e-5 when the 40 passes are spent, a factor 1.2 from where the previous pass left it. `kzz1e9/HeH2.13` does not converge, its carrier row banding between 5e-2 and 2e-1 with no trend. The three x0.01-XUV atomic cases stop as before, with the composition relaxing and the hydrodynamic rows not moving between passes. Nothing was deleted and every replaced case-level file is kept.

The fourth pass measured the nine molecular states again, this time through the runner and the publisher, under the certification rule above. The same three certify, and the catalog goes from 83 certified cases of 137 state indexes to 85: `molecular_scalar_gj1132_kzz1e9/HeH2.13` and `.../HeH9.7` are the first certified molecular cases of the catalog. The two `pre_D9s3` archival copies were not run and not repaired, their states being by md5 the generations of the two cases one level up, which is where they were measured. The verdicts, the refusing entries and the ghost numbers reproduce the third pass case for case.

Corrected by the advisor. `status.py` carried the binary of record as a literal, so the inventory it wrote named the previous binary although every run used the new one; it now takes `EXHALE_BIN` where a caller states one and otherwise the newest installed copy.

The typeset documents (`docs/EXHALE_physics_and_algorithms.pdf` 62 to 67 pages, `docs/EXHALE_user_manual.pdf` 91 to 94, `docs/EXHALE_BC_and_IC.pdf` 29 to 31, all rebuilt clean) carry what the code now does: the time step restriction with its derivation, the contact law of boundary model v2 and the composition contract of v3 with the accuracy and why it is that number, the certification inventory including the stage-sum entries and the report's capacity contract, the elemental base flux record, the transit state selection and what each selection means, the grid default and the pinning of older inputs, the diagnostics this series added, and the state file's new header lines. Two statements were wrong and are corrected in place: the branch handover was described as a smoothstep of half width $1e-6$, where the code makes a discrete choice with no width and the only declared width is the supersonic-outflow transition at $1e-8$; and the transit tool was described as reading only the advection-corrected profiles.

What a state that satisfies the flux criterion and not the certification costs

`docs/du_stop_vs_stationary_20260920.md`, with `docs/lhs1140b_catalog_state_20260920.tex` as the document it answers a question of. On one certified case, `atomic_scalar_gj1132_kzz1e9/HeH2.13`, a state that meets the flux-spread criterion the literature converges on ($du < 1e-3$, ATES and CETIMB, the latter asking that $\rho v r^2$ be constant with altitude) but is refused by this code's row-by-row certification differs from the certified stationary solution by at most 1.9 per cent in any profile, and that difference DOES reach a measured quantity: the He I 10830 red-pair equivalent width moves by 1.3 per cent and its depth by 1.1 per cent, while $\log_{10} \dot{M}$ does not move at the two decimals the code reports.

Where it sits is the point. Both states meet the flux criterion by four to eight decades, and $\rho v r^2$ is the one quantity that agrees between them to 0.055 per cent everywhere; what differs is the composition and the temperature of the emitting layer between 1.7 and 17 R_p , driven by the one row the certification refuses, the He/H elemental transport in the wind ($5.209E-05$ against $1.0E-05$ at cell 288, $r = 1.6951 R_p$). The criterion is blind by construction to the physics that sets the line. Against the observational side the shift is small; against the differences the catalog resolves between neighboring models it is not, the He/H rungs of this group standing 0.178 per cent A apart in the equivalent width where the shift is 0.018 per cent A, a tenth of one rung.

What was not measured, and why. The marching route does not reach $du < 1e-3$ on this case at all: MEASURED, a cold march is still three and a half decades above the threshold after 2106 steps and descending, and a seeded march starts one step from the threshold, turns around and climbs to 218 before oscillating between 45 and 190.

The compared state was therefore produced by the route the catalog uses, the partitioned stationary route stopped at one outer pass. A state reached by a different route, in particular one whose hydrodynamic rows are far from stationary while its flux happens to be flat, is not represented by this comparison and would be further away. Figures: docs/figures/du_stop_vs_stationary_profiles.pdf, _differences.pdf and _he10830.pdf.

What the user decided

- **The D2b direction rule**, carried over from the previous series, was accepted with the agreement check: the certification report states whether the base contact's direction, where the wind window decided it, agrees with the base face mass flux.
- **D1b was executed** on 2026-09-19: the Mixed-grid default base cell width is now exactly $2.0d-4 R_p$, and every live input with stored results carries Base grid [dr,cells]: 1.9999999494757503e-4 50, 770 files under LHS1140b/models, 51 under backup/regression and the rest across the planet, benchmark and example directories, with the frozen trees carrying a GRID_DEFAULT_NOTE.md. An input without the key and one stating the pinned width build one grid bit for bit.
- **Retention is kept**: every generation stays, a state whose certificate is now invalid or whose solve failed being a control for P1 and P2.
- **The instruction of 2026-09-20**: a state that satisfies the equations is certified, whatever the run that produced it ended as. Its four bookkeeping defects are the four above (the evaluate entry, the completeness guard, the refusal recorded as unknown, and the wording that read as a verdict on the state), and its result is the two first certified molecular cases of the catalog.
- **P2 stays closed unless the user opens it**. No solver control was changed by anything in this series.

The golden refresh of 2026-09-20

The binary of record is the tree's EXHALE.x, md5 f89569405648f9ef8ff8dc78b1ab6810 (MEASURED here), with make -q clean (MEASURED here). It carries one file beyond the catalog copy EXHALE_75d55d9d.x: the comment at conduction_base_level_T of viscous_conduction.f90, corrected on 2026-09-20 after P6c (MEASURED, the md5 of every other file of the 75d55d9d source list is the file's md5 in the tree now).

The goldens were refreshed once on that binary, over eighteen cases, the same list as the 2026-09-19 refresh, after a strict check of all eighteen on it. Movement against the 2026-09-19 set, READ from that strict check (REGRESSION_REL_TOL=0, OMP_NUM_THREADS=1), as the largest relative difference of any value in each file:

case	Hydro_ioniz.txt	Ion_species.txt	nature of the case
wasp_full_newton	1.4e-11	1.1e-10	converged, certified
wasp_full	1.9e-1 (velocity, row 226)	9.0e-4	du stop, 12261 steps
wasp_he23off	1.9e-1 (velocity, row 228)	6.6e-4	du stop, 12246 steps
hp_front	9.0e-5	6.2e-5	fixed-step snapshot, transported stages

case	Hydro_ioniz.txt	Ion_species.txt	nature of the case
hp_trace_seed	1.8e-4	2.3e-4	fixed-step snapshot, transported stages
hp_zero_seed	1.8e-4	2.3e-4	fixed-step snapshot, transported stages
hydrostatic_column	2.7e-2 (on values of order 1e-146)	9.998e-1 (on values of order 1e-181)	static column
lower_profile	1.3e-1	3.3e-2	fixed-step snapshot
mol_base_handoff	2.2e-2	1.6e-2	fixed-step snapshot
mol_carrier	2.2e-2	1.5e-2	fixed-step snapshot
mol_diffusion	2.2e-2	1.6e-2	fixed-step snapshot
mol_ir_bands	2.3e-2	1.5e-2	fixed-step snapshot
mol_lyman_werner	3.8e-2	2.6e-2	fixed-step snapshot
mol_metals	2.3e-2	1.5e-2	fixed-step snapshot
mol_sec_ion	1.9e0 (a velocity changing sign, row 192)	1.7e-2	fixed-step snapshot
oxygen_chemistry	1.9e0	1.0e0	fixed-step snapshot, oxygen network
roundtrip	7.6e-2	1.4e-3	fixed-step snapshot, 40 steps
iontrans_metals	identical	identical	converged, certified

The one converged marching case moved by $1.4\text{e-}11$ and the one certified stationary case did not move at all. Everything above $1\text{e-}3$ is a fixed-step snapshot of a run that is not stationary, whose trajectory the shorter time step of P6b moves by construction, or the velocity of the base region of the two du-stop cases, where the velocity alternates in sign from cell to cell and the absolute movement is 18 to 26 cm/s. The molecular cases carry the boundary composition contract as well.

After the refresh a strict re-check was run on the same binary and it finished: REGRESSION PASS (all cases byte-identical) over all eighteen cases (MEASURED, REGRESSION_REL_TOL=0, OMP_NUM_THREADS=1), in both state files and in the two _adv products where the case writes them (iontrans_metals writes none, so its two _adv files have no reference and are not compared). The refreshed set therefore reproduces on the binary of record.

Code-size summary: EXHALE vs. the original ATES (2026-06-07; updated 2026-09-11)

A line-by-line comparison of the Fortran source against the pristine upstream code (ATES/ATES-Code-main) quantifies the extent of the in-house extension. *Pure code lines (SLOC)* exclude blank lines and comments (the comment stripper is string-aware, so a ! inside a quoted string is not treated as a comment); the line-by-line diff additionally normalizes whitespace and case, so reformatting and comment-only edits are *not* counted as changes. The measurement script is src/utls/codesize.py, which reprints every number of this appendix and the byte-identical list of Table 34.

Measured 2026-09-11, i.e. after the whole of stage 1 and the stage-2 series through item P19 of section 8: the acceptance and certification contracts, the attempted-step controller, the partitioned stationary solver, the oxygen and carbon chemistry, the molecular infrared bands and the test suites those series brought with them.

What the script counts as *core*. Every .f90 file under src/ except those in src/modules/wind_ae/, which is the ported Wind-AE IC generator and is listed separately. That rule puts the standalone drivers under src/tests/ in the core figure, and they are now a large part of it, so the table splits the core row into the production build and those drivers. Only the production files are compiled into EXHALE.x; each test suite builds its own driver from src/tests/<suite>/run.sh.

	Fortran files	code lines (SLOC)
Original ATES (ATES-Code-main)	47	3 514
EXHALE core (excl. Wind-AE port)	169	71 503
of which the production build	95	50 762 ($\approx 14.4\times$)
of which the src/tests/ drivers	74	20 741
net change (core)	+122	+67 989 ($\approx 20.3\times$)
Wind-AE IC-generator port (wind_ae/)	+36	+4 756
EXHALE total	205	76 259 ($\approx 21.7\times$)

Line-by-line diff, core (comments + blanks excluded; whitespace/case normalized): **69 179** code lines added in EXHALE (new modules + the new side of modified lines) and **1 190** removed from the original (deleted + the old side of modified lines), for **70 369** total changed lines. At the file level, of the 47 original files **38 were modified**, **8 are unchanged** (all eight byte-for-byte; the list is Table 34) and **one was removed**: flux/speed_estimate_HLLC.f90, Toro's adaptive p_* estimate, was used by Num_Fluxes.f90 and never called, because the HLLC branch builds its own Davis/Einfeldt speeds inline. **123 new core modules** were added. Of those, 49 are part of the simulation build:

New module	Purpose	SLOC
time_step/steady_newton.f90	stationary solve of the hydrodynamic rows: residual $F(Y) = 0$ on the physical cells, Newton/Krylov with pseudo-transient continuation, line search, best-iterate ledger	9435
lower_atmosphere/diffusive_photochemistry.f90	vertical transport of the molecular carriers (H_2 , OH, H_2O , CO) solved together with the chemistry that makes and destroys them	2485
lower_atmosphere/molecular_infrared_data.f90	band and line data of the molecular infrared coolant (generated by cooling_data/molecular_infrared_bands.py)	2182
functions/binary_element_diffusion.f90	conservative binary H/He element transport (stage-resolved pair coefficients, donor-cell drift, metals slaved to H)	1898
time_step/attempted_step.f90	the attempted-step controller: one checkpoint, one trial, one adoption boundary, and the rollback contract	1608
time_step/certification.f90	one evaluator of the equations a run actually solves, and the contexts that read its verdict	1273
nonlinear_system_solver/constrained_chemical_equilibrium.f90	chemical equilibrium of the molecular H/He network in logarithmic number densities, element conservation as explicit residual rows	1173

New module	Purpose	SLOC
lower_atmosphere/h2_self_shielding_table.f90	H ₂ Lyman–Werner dissociation cross section with overlapping lines, and its two branching ratios	952
time_step/energy_semi_implicit.f90	implicit source step for the thermal energy of a cell, at fixed composition and heating	545
time_step/steady_residual.f90	finite-volume stationary residual $R = du/dt$ and its norms, shared by the diagnostic, the convergence monitor and the stationary solver	509
time_step/viscous_conduction.f90	Navier–Stokes viscous momentum force, its dissipation, and thermal conduction (Crank–Nicolson)	504
radiation/excited_hydrogen.f90	non-LTE H($n=2$) 2s/2p equilibrium (Christie, Arras & Li 2013) and the Balmer feedback	474
lower_atmosphere/element_inventory.f90	one stoichiometric map of the H, He, O and C nuclei of a cell, and the three constraints a composition state is judged by	448
radiation/charge_exchange.f90	charge-exchange network (Huang et al. 2023, Table 4)	446
files_IO/lower_atmosphere_profile.f90	lower-atmosphere profile reader: log- p interpolation, elemental reservoirs, $K_{zz}(p)$	367
states/caloric_eos.f90	caloric equation of state: internal energy density, pressure and temperature, with the H ₂ level ladder	340
lower_atmosphere/molecular_reaction_heat.f90	net chemical heat of the collisional reactions of the H ₂ /He network	339
lower_atmosphere/oxygen_rates.f90	rate coefficients and thermodynamic data of the oxygen-hydrogen chemistry, with their validity ranges	316
functions/element_census.f90	one definition of the elemental and charge content of a composition state, and the invariant checks built on it	307
nonlinear_system_solver/System_HeH_mol.f90	coupled H/He + molecular (H ₂ /H ₂ ⁺ /H ₃ ⁺ /HeH ⁺) system, turnover-scaled rows	303
radiation/electron_energy_degradation.f90	degradation of a fast photoelectron in a partly ionized H/H ₂ /He gas: the heat fraction and the ionizations of each target	287
radiation/lya_rt.f90	Ly α transfer by the escape-probability closure, evaluated in line each step	257
lower_atmosphere/molecular_infrared_cooling.f90	infrared exchange of the molecular layer: H ₂ O and CO vibration-rotation bands, H ₂ quadrupole and magnetic-dipole lines	231
states/base_boundary.f90	the lower boundary as a characteristic condition at the face, the number of conditions set by the Mach number there	199
functions/h2_photo_channels.f90	H ₂ photoabsorption resolved into mutually exclusive final-state channels, with one stoichiometric table	199
lower_atmosphere/lyman_werner.f90	H ₂ photodissociation in the Lyman and Werner bands (the Solomon process)	195
radiation/opacity_models.f90	opacity-model dispatcher for the photoionization cross sections (A/C/P/T)	180
files_IO/metals_input_read.f90	parse metals.inp: trace-metal abundances at runtime	177
functions/composition.f90	single point turning (ρ, f_{sp}) into every species density, n_e and n_{tot} , and converting pressure and temperature	162
flux/species_face_flux.f90	advection of the species mass fractions on the same faces, with the same mass flux, as the density	162
time_step/hydrodynamic_rows.f90	the hydrodynamic rows of the stationary residual in double precision, instantiated from the kind-generic text	161
nonlinear_system_solver/System_HeH_metals.f90	coupled H/He+metals ionization (+ analytic Jacobian)	150

New module	Purpose	SLOC
lower_atmosphere/mol_rates.f90	molecular reaction rates (Koskinen et al. 2022, Table 1)	150
lower_atmosphere/h3p_cooling.f90	optically thin H_3^+ infrared cooling (Miller et al. 2013)	148
nonlinear_system_solver/System_HeH_mol_metals.f90	molecular + metals merged ionization system	141
nonlinear_system_solver/newton_solver.f90	damped Newton with an analytic Jacobian for the ionization systems, with the MINPACK hybrd1 retry	128
init/species_table.f90	metadata table of the metal stages and of the base, molecular and oxygen species	128
lower_atmosphere/co_photodissociation.f90	CO photodissociation on the 912–1201 Å band	127
nonlinear_system_solver/ion_residual_core.f90	shared ionization-residual rows and analytic-Jacobian pieces of the <code>System_*</code> family	127
lower_atmosphere/water_photolysis.f90	H ₂ O and OH photodissociation in the four far-ultraviolet bands of the oxygen option	111
lower_atmosphere/co_self_shielding_table.f90	CO self-shielding function $\Theta[N(CO), N(H_2)]$ (Visser, van Dishoeck & Black 2009)	110
nonlinear_system_solver/ion_cell_state.f90	named-field cell state of the ionization residuals, in place of the position-packed params	93
lower_atmosphere/lower_column.f90	analytic lower-atmosphere column from the 1-bar level to the base pressure (Koskinen et al. 2022)	90
flux/low_mach_dissipation.f90	gated fourth-difference stress damping the two-cell velocity mode a contact-resolving flux leaves undamped	87
files_IO/opacity_input_read.f90	parse <code>opacity.inp</code>	85
radiation/hydrogen_n2_rates.f90	atomic data and rate coefficients of the H $n=2$ manifold and of the Ly α line, in one place	76
nonlinear_system_solver/System_HeH_TR_metals.f90	merged He 2 ³ S+metals system	68
lower_atmosphere/h2_vibrational_relaxation.f90	the fate and the size of the vibrational energy left in an H ₂ molecule, for both of its drivers	55
nonlinear_system_solver/constrained_equilibrium_probe.f90	standalone re-evaluation of one saved constrained-equilibrium state, without the hydrodynamics	15
production new-module total		30 003

The other 74 are the standalone test drivers under `src/tests/`, which did not exist in ATES at all. They state what a routine is allowed to return and are run by suite, not by the production binary:

Test driver	What it asserts	SLOC
tests/krylov_and_dogleg/krylov_and_dogleg_tests.f90	what the linear solve and the step geometry of the stationary solver are allowed to return	2675
tests/steady_species_rows/steady_species_rows_tests.f90	the unknown space of the stationary system, one row per transported balance	1806
tests/diffusion_tests.f90	the binary H/He element-diffusion operator: analytic limits and conservation	1051
tests/grid_and_gates/hydrostatic_residual.f90	the momentum row returns zero on a hydrostatic state	967

Test driver	What it asserts	SLOC
tests/carrier_retry/carrier_retry.f90	the carrier-local checkpoint and the rejection-and-retry controller, on a real molecular column	842
tests/certification/certification_contexts.f90	the certification evaluator and its contexts, against the predicates they replace	790
tests/species_face_flux/species_face_flux_tests.f90	species advection on the face mass fluxes, and the element rows of the diffusion operator	557
tests/physics_probe/weno3_reconstruction_order.f90	convergence order of the face reconstruction against the exact face value	482
tests/element_operator/element_operator_tests.f90	the boundary the element transport operator is measured under	471
tests/spectrum_type/loaded_sed_table_semantics.f90	what a row of a loaded SED table means, and what the production consumers make of it	417
tests/attempted_step/attempted_step_tests.f90	the checkpoint and the step-size policy of the attempted-step controller	415
tests/a2_m1/a2_m1_rate_check.f90	the oxygen rate coefficients against their published sources	406
tests/spectrum_type/balmer_band_quadrature.f90	quadrature of the Balmer-continuum band integrals of excited_hydrogen	367
tests/energy_update/conduction_floor_tests.f90	what the Crank–Nicolson transport stage returns, refuses and conserves	360
tests/physics_probe/metal_photoionization_fits.f90	every metal photoionization cross section against the published fit, and only where it is valid	343
tests/element_census_tests.f90	the elemental and charge invariants of a composition state	340
tests/energy_update/energy_update_tests.f90	the residual-controlled temperature solve of the implicit energy source step	334
tests/spectrum_type/loaded_sed_threshold_edges.f90	the photon grid built from a loaded SED, and the rates formed on it	304
tests/physics_probe/species_formation_energy_table.f90	the formation-energy table of the energy ledger, and the reaction energies that are its differences	275
tests/physics_probe/lya_band_transmission.f90	the stellar Ly α band reaching a cell through the atomic hydrogen column above it	272
tests/spectrum_type/wasp121b_sed_sensitivity.f90	the production band quantities of the three WASP-121 b spectrum candidates	262
tests/coupled_source_step/coupled_source_tests.f90	the coupled temperature-composition source step	260
tests/physics_probe/lyman_werner_cell_mean.f90	the Lyman–Werner rate a cell carries is the mean of the local rate over that cell	234
tests/physics_probe/lya_beam_cell_mean.f90	the stellar Ly α field of a cell is the beam mean over the cell’s own optical depth	229
tests/spectrum_type/balmer_continuum_field.f90	the two scalar Balmer-continuum rates of H($n=2$)	227
tests/physics_probe/positivity_limiter_scaling.f90	the positivity scaling of the reconstructed face states (Zhang & Shu 2010)	223
tests/carrier_returned_state_acceptance/carrier_returned_state_acceptance.f90	what makes a returned carrier state a solution, and what does not	222
tests/grid_and_gates/photon_grid_threshold_edges.f90	every active absorber’s threshold against the bin partition of the photon grid	213

Test driver	What it asserts	SLOC
tests/physics_probe/photoionization_field_substitution.f90	the order in which the field and the composition that attenuates it are brought into agreement	209
tests/adv_static_limit/stationarity_by_the_mass_row.f90	which test decides that the state the post-process was handed is stationary in a cell	199
tests/physics_probe/co_destruction.f90	the CO destruction rates, the shielding table, the deposited energies and the cell mean	195
tests/physics_probe/recombination_coefficient_fits.f90	every recombination coefficient against the published fit	189
tests/spectrum_type/photon_grid_n2_floor.f90	the photon grid built when the excited-hydrogen coupling is armed	184
tests/carrier_reference_scales/carrier_reference_scales.f90	the element a transported carrier is made of, and whether its chemistry rows can be differentiated	181
tests/carrier_constraint_attribution/carrier_constraint_attribution.f90	which residuals belong to the solver and which to a constraint	179
tests/adv_static_limit/enthalpy_flux_of_mass_divergence.f90	the third term of the advected energy equation	178
tests/grid_and_gates/free_outflow_boundary.f90	the outer free-outflow boundary on a supersonic wind, and its ghost state	174
tests/spectrum_type/planck_photon_field.f90	the incident flux put on the photon grid for the photospheric blackbody type	173
tests/a2_m2/a2_m2_kinetics_check.f90	the kinetics quantities the oxygen option adds on top of the rate audit	163
tests/physics_probe/lya_escape_probability.f90	the escape probability of the two-level closure against the published slab solution	163
tests/e1_h2/e1_h2_channel_check.f90	the mutually exclusive H ₂ photoabsorption channels	161
tests/adv_static_limit/upstream_caloric_energy.f90	the upstream energy of the advected energy equation	154
tests/physics_probe/photoionization_field_self_consistency.f90	the field one cell is solved in, against the composition that cell returns	148
tests/grid_and_gates/photon_grid_quadrature.f90	the photoionization rates of one unattenuated cell against the band quadrature	146
tests/physics_probe/co_helium_ion_sink.f90	He ⁺ + CO as a species sink, not only as a heat source	145
tests/physics_probe/riemann_wave_speeds.f90	signal speeds at a face, and the state one conservative update with that flux leaves behind	143
tests/physics_probe/helium_level_cooling_ledger.f90	which helium level each cooling channel of eval_cool is charged to	139
tests/physics_probe/heating_channel_closure.f90	one assembly of the volumetric heating rate, and the closure of its channels	136
tests/acceptance_classes/acceptance_class_contract.f90	one meaning of the ionization acceptance classes for every consumer	132
tests/physics_probe/opacity_model_p_ledger.f90	the radiative ledger of a cell whose cross sections are broadened by opacity model P	124
tests/physics_probe/photoevent_energy_ledger.f90	the energy an isolated photoabsorption event has to account for	117
tests/physics_probe/xuv_cell_mean_attenuation.f90	the beam attenuation the photoionization rate of a cell is evaluated with	115

Test driver	What it asserts	SLOC
tests/grid_and_gates/grid_window_indices.f90	the two index statements that decide which cells a quantity is read from	101
tests/physics_probe/h2_rovibrational_identity.f90	the identities tying the H ₂ partition function of the chemistry to the internal energy of the caloric EOS	96
tests/adv_static_limit/stationary_flux_refusal.f90	the sensitivity of the advected energy correction to a non-stationary mass flux	91
tests/physics_probe/ionization_threshold_turn_on.f90	one definition of each ionization threshold, and every cross section turning on exactly there	90
tests/grid_and_gates/grid_width_identity.f90	the cell-width and total-width identities of the radial grid	83
tests/physics_probe/h2_channel_detector_ratio.f90	the measured proton-to-H ₂ ⁺ ratio of Chung et al. (1993) against the four channels	83
tests/physics_probe/fine_structure_escape_probability.f90	the escape probability of the ground-term fine-structure lines, and the depth it is fed	80
tests/physics_probe/atomic_mass_and_radius_constants.f90	the helium mass and the astronomical constants of the input readers, one definition each	79
tests/physics_probe/h3p_cooling_limits.f90	the limits, the continuities and the published values of the H ₃ ⁺ cooling rate	79
tests/physics_probe/assertion_report.f90	one printed verdict line per assertion, and the failure count the driver exits on	74
tests/adv_static_limit/thermal_damkohler_limit.f90	the validity condition of the advection-corrected energy solve, as a function of the flow speed	72
tests/a2_m2/a2_m2_g3_run_check.f90	with every photolysis rate at zero, the solved O/OH/H ₂ O partition is the chemical equilibrium	71
tests/physics_probe/absorbed_fraction_switch.f90	$(1 - e^{-d})/d$, the fraction of the photons entering a cell that the cell absorbs	71
tests/constrained_network_layout/transported_proton_constraint_layout.f90	which species are unknowns of the continuation, and whether the system built is square	71
tests/test_columns.f90	synthetic columns whose species carry the mass density they are measured against	67
tests/physics_probe/h2_infrared_line_populations.f90	the LTE populations weighting the H ₂ quadrupole lines are those of the caloric EOS ladder	65
tests/physics_probe/wind_ae_bridge_composition.f90	the composition an EXHALE input.inp hands the Wind-AE solver	54
tests/physics_probe/voronov_carbon_ionization.f90	the electron-impact ionization of C ⁰ and C ⁺ against the published Voronov (1997) rows	51
tests/physics_probe/water_photolysis_lyman_alpha_yields.f90	the H ₂ O branching on the band that contains Ly α , against the published yields	49
tests/physics_probe/h3p_recombination_branching.f90	the two product channels of H ₃ ⁺ dissociative recombination	44
tests/energy_update/cooling_models.f90	analytic cooling laws for the tests of the implicit energy update	43
tests/species_masses/carrier_background_masses.f90	the mass the carrier transport gives each of its background collision partners	36
test-driver total		20 741
new-module total		50 744

So the 123 new core modules contribute 50 744 of the 69 179 added code lines; the remaining 18 435 added (and 1 121 of the 1 190 removed, the other 69 being the deleted HLLC estimator) are edits spread across the 38 modified files. The ten heaviest, by added code lines, are the main driver (EXHALE_main.f90, 3 498 added, 246 → 3 625 SLOC: the marching loop, the attempted-step adoption boundary, the run-mode contracts and the written-state assertions), the ionization equilibrium and its diagnostics (ionization_equilibrium.f90 2 274 and util_ion_eq.f90 1 803: the metal, molecular and oxygen networks, the acceptance classes, the photoionization field and the heating channels), the cooling module (Cool_coeff.f90, 2 250 added, 442 → 3 966 raw lines: CHIANTI metal-line fits, Fe tables, fine-structure statistical equilibrium, Penning and molecular rate coefficients), input parsing (input_read.f90, 1 753), the state reader and writer (load_IC.f90 1 063, write_output.f90 928, write_setup_report.f90 751), the advection-corrected post-processor (post_process_adv.f90, 647) and the cross sections (cross_sec.f90, 438). Those ten carry 15 405 of the 18 435. The rest of the edits are in the reconstruction, the fluxes, the boundary conditions, the grid, the time step, the energy equation and the System_* ionization systems, which wire the metals, the molecules, the opacity dispatcher, the Roche potential, the transonic/auto/Wind-AE IC, the excited-H and Ly α physics, the element diffusion and the Brent/Newton solver upgrades into the original solver.

In total the EXHALE core is now $\approx 20.3\times$ the original code base (3 514 → 71 503 SLOC), or $\approx 14.4\times$ counting only the production build (50 762 SLOC), touching almost all of the original files (38 of 47 modified, one deleted, plus 123 new); with the ported Wind-AE IC generator the full Fortran tree is 205 files / 76 259 SLOC ($\approx 21.7\times$). Beyond Fortran, the tree also carries the Python tooling (GUI, the EXHALE_transit.py transmission post-processor, plotting/analysis, the lower-atmosphere adapters and the elemental-flux closure driver) and 181 816 lines of documentation (187 md/tex files: the memos, the manual and this log) under docs/.

Fortran source inherited unchanged from ATES

Of the 47 upstream Fortran files (ATES/ATES-Code-main), **8 are byte-for-byte identical** in EXHALE (re-measured 2026-09-11): the numerics EXHALE reuses without touching, which are now the linear-algebra part of the included MINPACK nonlinear solver plus the gravity pre-evaluation. Table 34 lists them with their role. No file is identical after whitespace and case normalization but not byte-identical, so the normalized-unchanged set and the byte-identical set are the same eight.

Five files left the byte-identical set since the 2026-08-30 edition of this table, each for a documented reason rather than drift.

- nonlinear_system_solver/hybrd.f90, hybrd1.f90 and fdjac1.f90 carry a new unit_step_floor argument threaded from hybrd to fdjac1, which selects the forward-difference step of the Jacobian: `.false.` is MINPACK's own $h = \varepsilon|x_j|$ with the exact-zero fallback, `.true.` floors it at unit magnitude, $h = \varepsilon \max(1, |x_j|)$, which is the rule for unknowns already scaled so that unity is their natural size (the logarithmic unknowns of the constrained chemical equilibrium). The params dummy also became assumed-size, so one transport array serves systems of different length. hybrd1 passes `.false.` and is therefore unchanged in behavior.
- states/Source.f90 gained the well-balanced option of 2026-09-10 (item N37 of the stage-2 solver program,

docs/ISSUES_20260909.md): under Well balanced: True the gravitational source and the geometric pressure term are not evaluated here at all, because they cancel the equilibrium part of the momentum flux difference identically and RK_rhs assembles the row from what is left (Käppeli & Mishra 2014, J. Comput. Phys. 259, 199, eq. 2.26). The file also reads its pressure through caloric_eos instead of a constant γ .

- radiation/J_inc.f90 carries the single-spectrum-type decision: Spectrum type: now states the spectrum of the whole photon grid, the XUV and the part below 13.6 eV alike, and no band is filled from a type the input did not select. The file also became the one definition of the eV-to-Hz conversion and of the Balmer-continuum head $e_{\text{th,HI}}/4$.

The two wave-speed estimators are no longer in the “effectively untouched” category either. flux/speed_estimate_HLLC.f90 was deleted as dead code, as recorded above. flux/speed_estimate_R0E.f90 was rewritten (99 code lines added, 43 removed) into a star-state estimate that returns a status: the primitive-variable, two-rarefaction and two-shock branches of Toro (2009) chapter 9 are selected by the pressure ratio, the vacuum condition of his eq. 4.76 is tested first, and a face with no admissible star state is reported rather than given a square root of a negative ratio.

Table 34: The 8 Fortran files byte-identical between the original ATES (ATES-Code-main) and EXHALE (measured 2026-09-11), grouped by role.

File	Purpose
<i>MINPACK linear algebra and constants</i> (Argonne; dir nonlinear_system_solver/)	
qrfac.f90	QR factorization via Householder transformations
qform.f90	Accumulate the orthogonal factor Q from the QR factorization
r1updt.f90	Rank-one update of the QR factorization
r1mpyq.f90	Apply the Givens-rotation sequence of the rank-one update
dogleg.f90	Dogleg (trust-region) step combining Gauss–Newton and steepest descent
enorm.f90	Euclidean norm of a vector (overflow/underflow safe)
dpmpar.f90	Machine floating-point precision / range constants
<i>Grid setup</i>	
init/set_gravity_grid.f90	Pre-evaluate gravity at cell centers and interfaces