

EXHALE: physical equations and numerical algorithms

Contents

1. Introduction and conventions	3
1.1 What the code solves	3
1.2 Units	3
1.3 Notation	4
1.4 How to read this document	5
2. Hydrodynamics	5
2.1 Conserved variables and the conservation laws	5
2.2 The discrete divergence	5
2.3 The spherical pressure term and gravity	6
2.4 The gravitational potential	7
2.5 Equation of state	7
2.6 The well-balanced option	8
2.7 Low-Mach dissipation of the contact mode	9
2.8 Viscosity and heat conduction	9
2.9 The two-dimensional approximation	10
3. The stellar radiation field and its absorption	10
3.1 The incident spectrum	10
3.2 Photoionization cross sections	12
3.3 Transfer along the radial ray	13
3.4 Photoionization and photoheating rates	14
3.5 Secondary ionization by photoelectrons	15
3.6 Reprocessed and non-stellar fields	16
4. Ionization and chemical state	17
4.1 The species carried	17
4.2 The cell system	17
4.3 The rate coefficients	19
4.4 Solving one cell	20
4.5 Ionization carried by the flow	21
4.6 Excited hydrogen and the $\text{Ly}\alpha$ field	21
5. The thermal source: heating and cooling	23
5.1 The heating channels	23
5.2 The cooling channels	24

5.3	Advancing the temperature	26
6.	The molecular layer, the trace metals and element transport	27
6.1	The H ₂ /He reaction network	27
6.2	H ₂ photoabsorption resolved into channels	28
6.3	Lyman-Werner photodissociation	28
6.4	The caloric equation of state	29
6.5	Carrier transport	29
6.6	The oxygen option	30
6.7	The lower-atmosphere handoff	31
6.8	Trace metals	32
6.9	Binary element transport	32
7.	Numerical scheme	33
7.1	The radial grid	33
7.2	Reconstruction	34
7.3	Approximate Riemann solvers	36
7.4	Positivity repair of a stage	37
7.5	Time integration and the time step	37
7.6	Operator splitting of one step	38
7.7	The marching convergence measure and the stop conditions	39
8.	Boundary and initial conditions	39
8.1	The lower boundary	39
8.2	The upper boundary	41
8.3	Initial conditions	42
8.4	Restart	43
9.	The stationary state	43
9.1	The stationary residual	43
9.2	The acceptance test	44
9.3	Certification	45
9.4	The Jacobian-free Newton-Krylov solve	46
9.5	Pseudo-transient continuation	47
9.6	The partitioned outer iteration	48
10.	Output and post-processing	48
10.1	The advection-corrected profiles	48
10.2	The mass-loss rate	49
10.3	Output files	50

10.4 The transmission spectrum	50
--	----

11. The input keys that select physics	50
--	----

1. Introduction and conventions

1.1 What the code solves

EXHALE integrates the one-dimensional, spherically symmetric equations of radiation hydrodynamics for the escaping upper atmosphere of an irradiated planet. The radial domain runs from a base near the lower atmosphere, which a run may place at the planet radius or at a stated pressure level (the $1\ \mu\text{bar}$ level of the Koskinen et al. (2022) handoff is the one the molecular configurations use), out to the Roche-lobe (L1) radius of the planet–star system or, in the spherical domain mode, to a radius the user states. The unknowns advanced in time are the conserved hydrodynamic state $u = (\rho, \rho v, \mathcal{E})$ on a finite-volume radial grid, together with the ionization and chemical state of the gas: hydrogen and helium always, the metastable He 2^3S level, a set of trace metals, and a molecular network carrying H_2 , H_2^+ , H_3^+ , HeH^+ and, optionally, the oxygen carriers OH , H_2O and CO . The radiative source term couples the two: the stellar spectrum is attenuated along the radial column, photoionizes and heats the gas, and the gas cools through recombination, collisional and line processes whose rates depend on the composition the ionization solve returns.

The code is a heavily extended fork of ATES v2.0 (Caldirola et al., 2021; Biassoni et al., 2024). The hydrodynamic core, the finite-volume discretization, the three-stage strong-stability-preserving Runge-Kutta integration and the choice of approximate Riemann solvers are recognizably those of ATES; the equation of state, the lower boundary condition, the reconstruction, the steady-state solvers, the transported species and essentially all of the chemistry are not. Every physics extension is selected at run time by the presence of a key in `input.inp` or of a companion file (`metals.inp`, `opacity.inp`, `base.inp`), and every one of them defaults to off, so that a configuration that states none of them reproduces the atomic H/He problem.

1.2 Units

The equations are solved in dimensionless form. The normalization is fixed once, in `input_read.f90`, from the planet radius R_0 , the base temperature T_0 , and the base number density n_0 :

$$v_0 = \sqrt{\frac{k_B T_0}{m_H}}, \quad t_s = \frac{R_0}{v_0}, \quad p_0 = n_0 m_H v_0^2, \quad q_0 = \frac{n_0 m_H v_0^3}{R_0}, \quad (1)$$

where $m_H = 1.67353284 \times 10^{-24}$ g is the mass of the hydrogen *atom*, not the atomic mass unit. Radii are in units of R_0 , so the base level sits at $r = 1$; velocities in v_0 ; mass densities in $n_0 m_H$; pressures in p_0 ; temperatures in T_0 ; and volumetric energy rates in q_0 . The mass unit being the hydrogen atom, the density is the number of hydrogen-atom masses per unit volume, and a species contributes m_{species}/m_H to it; the species table therefore carries $m(\text{HeI}) = 3.9715259$ and not 4.

The gravitational potential is measured in v_0^2 , so it is fixed by the single dimensionless group

$$b_0 = \frac{GM_p m_H}{k_B T_0 R_0}, \quad (2)$$

the Jeans escape parameter at the base. Only three further dimensionless numbers enter the potential and the domain: $\tilde{a} = a_{\text{orb}}/R_0$, the mass ratio $M_\star/M_p$, and the outer radius.

The base level itself may be stated in three ways, and a run that states two that disagree is stopped at startup: Log10 lower boundary number density gives n_0 directly; Base BC: pressure states the base pressure in microbar and n_0 follows from $n_0 = p_{\text{base}}/(k_B T_0 \hat{n}_{\text{bc}})$; and a lower-atmosphere handoff (base.inp or a profile file) states p_{base} in bar and n_0 follows the same way. Here $\hat{n}_{\text{bc}}$ is the base particle count per H+He nucleus, which the composition module derives from the element ratios and, with a molecular base, from the H nuclei bound into H_2 (Sec. 2.5).

1.3 Notation

One symbol stands for one quantity throughout this document. The table below fixes the symbols that appear in more than one section. Symbols introduced inside a published fit and defined where they appear (the x of a Verner-type cross section, the λ of a recombination-cooling fit, the α of a spectral index) are local to that fit and are not in the table.

symbol	quantity	note
u	conserved state $(\rho, \rho v, \mathcal{E})$	the unknown of the hydrodynamics, Sec. 2.1
W	primitive state (ρ, v, p)	what the reconstruction works on
ρ, v, p, T	mass density, radial velocity, pressure, temperature	
$\mathcal{E}, \rho e, e, \varepsilon$	total energy density, internal energy density, internal energy per unit mass, internal energy per particle	$\mathcal{E} = \frac{1}{2} \rho v^2 + \rho e$
Φ	gravitational potential	Φ_i, Φ_c at interfaces and cell centers
γ_{eff}, c_s	effective adiabatic index, sound speed	from the caloric equation of state (14)
m_{H}	mass of the hydrogen atom	the code's mass unit, (1)
μ, κ	dynamic viscosity, heat conductivity	(21) only
n_X, n_{tot}, n_e	number density of species X , of all heavy particles, of free electrons	
$x_{X^{q+}}, f_i$	ionization fraction of a stage, mixing ratio n_i/n_{tot}	fractions of the element's own nuclei
$E, h\nu$	photon energy	always subscripted where a particular energy is meant (E_k, E_{th})
$F_E, \mathcal{F}_k$	incident and attenuated flux per unit photon energy	(31)
σ, τ	photoionization cross section, optical depth	
Γ, α, β	photoionization rate, recombination coefficient, electron-impact ionization coefficient	Sec. 4.2
$\mathcal{H}, \Lambda$	volumetric heating rate, volumetric cooling rate	$\text{erg cm}^{-3} \text{ s}^{-1}$; $Q = \mathcal{H} - \Lambda$
ξ_{day}	dayside dilution of every stellar beam	Table 1
J_i, J	diffusive number flux of a carrier, diffusive mass flux of an element	Secs. 6.5, 6.9
dF, S, R	discrete flux difference, source, stationary residual	(6)–(11), (108)
$A_{\pm}, \Delta V_j, \Delta r_j$	face areas, cell volume, cell width	(5)

1.4 How to read this document

Every equation below is the discrete operator the code applies, taken from the source and not from a derivation. Where the code's expression differs from the textbook form (the spherical pressure term under PLM, the gravitational work term folded into the energy flux, the mean molecular weight that is never formed), the code's form is given and the difference is stated. Routines are named with the file they live in, for instance `RK_rhs.f90`, so that a statement here can be checked against one place in the tree.

Section 2. states the conservation laws, the equation of state and the source terms. The sections that follow it state the radiation field, the ionization and chemical network, the thermal source and the molecular and metal chemistry. Section 7. then gives the grid, the reconstruction, the Riemann solvers, the time integration and the operator splitting; Sec. 8. the boundary and initial conditions; and Sec. 9. the stationary residual, its acceptance test and the solvers that drive it to zero. A section on the post-processing tools closes the physics account, and Sec. 11. tabulates the input keys that select physics, with the default each of them takes when the key is absent.

2. Hydrodynamics

2.1 Conserved variables and the conservation laws

The conserved state of a cell is the triple

$$u = (\rho, \rho v, \mathcal{E})^\top, \quad \mathcal{E} = \frac{1}{2}\rho v^2 + \rho e, \quad (3)$$

with ρe the internal energy density. The map between u and the primitive triple $W = (\rho, v, p)$ is `UW_conversions.f90`: the density and the momentum are trivial, and the pressure follows from the caloric equation of state of Sec. 2.5, $p = p(\rho, \rho e)$, evaluated with the composition of the cell the state belongs to. For a reconstructed face state that cell is the cell the state was extrapolated from, because the composition ratios the equation of state needs are ratios of number densities and are therefore independent of the density.

The equations integrated are the Euler equations in spherical symmetry with gravity and a radiative source,

$$\begin{aligned} \frac{\partial \rho}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v) &= 0, \\ \frac{\partial(\rho v)}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v^2) + \frac{\partial p}{\partial r} &= -\rho \frac{\partial \Phi}{\partial r} + S_\mu, \\ \frac{\partial \mathcal{E}}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 (\mathcal{E} + p) v) &= -\rho v \frac{\partial \Phi}{\partial r} + Q + S_\mathcal{E}, \end{aligned} \quad (4)$$

where $Q = \mathcal{H} - \Lambda$ is the net radiative heating minus cooling per unit volume (Sec. 5.) and $S_\mu, S_\mathcal{E}$ are the viscous and conductive sources of Sec. 2.8, both off unless a run asks for them. The factor 4π that would appear in the face areas and the cell volume cancels between them and is simply not carried.

2.2 The discrete divergence

`RK_rhs.f90` assembles the right-hand side of (4) as a flux difference dF and a source S , so that one forward Euler stage of the marching loop is $u \leftarrow u - \Delta t (dF - S)$ and the stationary residual is $R = dF - S$ with the radiative source subtracted

from its third row. With

$$A_{\pm} = r_{j\pm 1/2}^2, \quad \Delta V_j = \frac{r_{j+1/2}^3 - r_{j-1/2}^3}{3}, \quad \Delta r_j = r_{j+1/2} - r_{j-1/2}, \quad (5)$$

and $F_{\pm}$ the numerical flux at the two faces of cell j (Sec. 7.3), the three rows are

$$dF_1(j) = \frac{A_+ F_+^{\rho} - A_- F_-^{\rho}}{\Delta V_j}, \quad (6)$$

$$dF_2(j) = \frac{A_+ F_+^{\rho v} - A_- F_-^{\rho v}}{\Delta V_j} + w_p \frac{p_R - p_L}{\Delta r_j}, \quad (7)$$

$$dF_3(j) = \frac{A_+ F_+^{\mathcal{E}} - A_- F_-^{\mathcal{E}} + \delta F_3}{\Delta V_j}, \quad (8)$$

with $F^{\rho} = \rho v$, $F^{\mathcal{E}} = v(\mathcal{E} + p)$, and p_L, p_R the face pressures the Riemann solver returns at the lower and upper face. The weight w_p and the content of $F^{\rho v}$ depend on the reconstruction, because the spherical pressure term is split differently in the two schemes (Sec. 2.3).

The term δF_3 in (8) is the gravitational work,

$$\delta F_3 = A_+ F_+^{\rho} [\Phi_i(j) - \Phi_c(j)] - A_- F_-^{\rho} [\Phi_i(j-1) - \Phi_c(j)], \quad (9)$$

with Φ_i the potential at the cell interfaces and Φ_c at the cell centers (`set_gravity_grid.f90`). Expanding the bracket separates (8) into the energy flux of Caldiroli et al. (2021), their Eq. (14), $F^{\mathcal{E}} + F^{\rho} \Phi$ taken at the interface potential, minus $\Phi_c(j)$ times the mass-flux divergence, which is their energy source $\Phi_i L_j^{\rho}$. The two forms are algebraically the same operator; the code carries it as one term so that the mass flux the work term rides on is the very flux the continuity row differenced.

2.3 The spherical pressure term and gravity

The pressure gradient in spherical symmetry is

$$\frac{\partial p}{\partial r} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 p) - \frac{2p}{r}, \quad (10)$$

and the two reconstructions use the two sides of this identity. Under PLM, `Phys_flux` adds p to the momentum flux, so the first term of (10) is inside dF_2 and the geometric correction is carried by the source; $w_p = 0$. Under WENO3 the pressure is left out of the flux and the whole gradient enters as the single face difference $(p_R - p_L)/\Delta r_j$; $w_p = 1$. `Source.f90` accordingly returns

$$S_1 = 0, \quad S_2 = -\frac{\rho_L + \rho_R}{2} \frac{\Phi_i(j) - \Phi_i(j-1)}{\Delta r_j} + \underbrace{\frac{A_+ - A_-}{\Delta V_j} p_c}_{\text{PLM only}}, \quad S_3 = 0, \quad (11)$$

where ρ_L and ρ_R are the two reconstructed face densities that bound the cell and p_c is the cell's own pressure from the caloric equation of state. Because $(A_+ - A_-)/\Delta V_j \rightarrow 2/r$ as the cell narrows, the PLM branch of (11) is the second term of (10).

The three physical terms of the momentum equation, the ram divergence, the spherical pressure gradient and the weight $\rho \partial \Phi / \partial r$, are gathered out of these pieces by `momentum_row_terms_of_cell` in the same file and stored. They are not dF_2 and S_2 : under PLM each of those holds a part of order $2p/r$ that cancels against the other, so a magnitude built from them alone reads $2p/r$ on a uniform pressure at rest, where the physical force is zero. The gathered terms are what the stationary residual scales the momentum row by (Sec. 9.1).

2.4 The gravitational potential

`grav_field.f90` carries two potentials, selected by `Domain` mode. The default is the full Roche potential of a synchronously rotating planet,

$$\Phi(r) = -\frac{b_0}{r} - \frac{b_0 \mathcal{M}}{\tilde{a} - r} - \frac{b_0(1 + \mathcal{M})}{2\tilde{a}^3} \left(\frac{\tilde{a} \mathcal{M}}{1 + \mathcal{M}} - r \right)^2, \quad \mathcal{M} = \frac{M_\star}{M_p}, \quad \tilde{a} = \frac{a_{\text{orb}}}{R_0}, \quad (12)$$

in code units, which is Eq. (2) of Caldiroli et al. (2021): the planetary term, the stellar tidal term and the centrifugal term about the center of mass. With `Domain` mode: `Spherical` the last two terms are dropped and $\Phi = -b_0/r$, which is what allows the domain to be carried past the L1 radius. The analytic derivative $d\Phi/dr$ is provided beside it and is used by the initial-condition sonic-point search; the flux assembly itself never differentiates Φ , it differences the tabulated Φ_i and Φ_c , which are evaluated once at startup at the true face and center radii.

The default outer radius in Roche mode is the L1 (Hill) radius $r_{\text{max}} = (3\mathcal{M})^{-1/3}\tilde{a}$; `Outer` radius overrides it.

2.5 Equation of state

The thermal equation of state is that of an ideal gas of all particles including free electrons,

$$p = (n_{\text{tot}} + n_e) k_B T \quad \Longleftrightarrow \quad T = \frac{p}{n_{\text{tot}} + n_e} \text{ in code units,} \quad (13)$$

which is `comp_T_from_p` and `comp_p_from_T` of `composition.f90`. The particle and electron counts are built in one place, `get_species_densities` in the same file, from the density and the species mass fractions: `calc_ntot` adds one particle for every species the run carries (H I, H II, He I, He II, He III, every metal ion stage, and each of H_2 , H_2^+ , H_3^+ , HeH^+ , OH, H_2O , CO), and `calc_ne` adds the free electrons weighted by the net charge of each species. The $\text{He } 2^3S$ column is not added by either: it is an excited level of He I and its particle and its nucleus are already counted there. Trace metals enter both sums only under the `eos_metals` policy of Sec. 6.8, on by default, so that the legacy H/He-only budget can be reproduced exactly.

The code never forms a mean molecular weight for the equation of state. The mean molecular weight of Caldiroli et al. (2021), their footnote 1, is a derived diagnostic here; (13) with the explicit particle count is what the hydrodynamics uses, so the temperature of a cell is a function of its composition and never of a fitted composition ratio.

The *caloric* half of the equation of state is `caloric_eos.f90`. For an atomic gas it is the monatomic relation $\rho e = p/(\gamma_{\text{ad}} - 1)$ with $\gamma_{\text{ad}} = 5/3$, exactly as in ATES. This is not valid in a molecular layer: at the 400 to 1300 K of the H_2 base the rotational ladder of H_2 is populated (the $J = 1$ level lies at 170 K), so an H_2 molecule stores about $\frac{5}{2}k_B T$ and the vibrational ladder adds to that above ~ 800 K. Where the gas is 80% H_2 the heat capacity is then wrong by a factor 1.67. The mixture relation used instead is

$$\rho e = \frac{3}{2}(n_{\text{tot}} + n_e)k_B T + n(\text{H}_2)k_B u_{\text{rv}}(T), \quad \frac{C_V}{k_B} = \frac{3}{2}(n_{\text{tot}} + n_e) + n(\text{H}_2)c_{\text{rv}}(T), \quad (14)$$

so that the exponent making $\rho e = p/(\gamma_{\text{eff}} - 1)$ exact is $\gamma_{\text{eff}} = 1 + (n_{\text{tot}} + n_e)k_B/C_V$, a function of both the composition and the temperature. The mean rovibrational energy u_{rv} and the heat capacity c_{rv} are evaluated from the complete bound rovibrational ladder of $\text{H}_2 X^1\Sigma_g^+$, 302 levels to 51966 K, that is to the 4.478 eV dissociation limit (Roueff et al., 2019, Table 2 through VizieR); the same partition function is the internal partition function of the $\text{H} + \text{H} \leftrightarrow \text{H}_2$ equilibrium

constant (73), so the energy and the chemistry cannot disagree. H_2^+ , H_3^+ , HeH^+ and the oxygen carriers are counted as monatomic, a stated approximation valid while they stay trace species (their largest ratio to $n(\text{H}_2)$ in the molecular layer of the reference configuration is 1.7×10^{-7}). Dissociation is not a heat capacity and is not in (14); it is a chemical source term. The branch between the two forms is structural, not a limit: a run with no molecules and a cell with $n(\text{H}_2)$ exactly zero both take the constant- γ_{ad} expression verbatim. The key Caloric EOS: monatomic forces the monatomic form everywhere.

The mapping $\rho e \rightarrow p$ in the molecular branch is the inverse of (14) and costs one safeguarded Newton iteration per cell; the mapping $p \rightarrow \rho e$ is direct. The sound speed used by the Riemann solvers and by the time step is $c_s = \sqrt{\gamma_{\text{eff}} p / \rho}$ evaluated with the cell's own composition.

Two base composition scalars follow from the same module. The gas mass per hydrogen nucleus is $m_{\text{H-nuc}} = m(\text{H}) + m(\text{He})y + \sum_e A_e X_e$ with $y = n_{\text{He}}/n_{\text{H}}$; the base particle count per H+He nucleus is

$$\hat{n}_{\text{bc}} = \frac{1+y+\sum_e X_e}{1+y} - \frac{1}{2} \frac{x_{\text{H}_2}}{1+y}, \quad x_{\text{H}_2} = \frac{2q_{\text{H}_2}(1+y)}{1+q_{\text{H}_2}}, \quad (15)$$

where x_{H_2} is the fraction of base hydrogen nuclei bound into H_2 and the last term is present only with Molecular base: True. The mixing ratio q_{H_2} is the one a lower-atmosphere handoff states, or otherwise the chemical-equilibrium fit (78) at (p_{base}, T_0) (Sec. 6.7). The same x_{H_2} is imposed on the inflowing ghost composition, so the equation of state and the species state describe one gas.

2.6 The well-balanced option

Well balanced: True (default off) changes the coordinate the pressure is carried in. The local hydrostatic equilibrium of cell j is the constant-density one through its own (ρ_j, p_j) ,

$$p_{\text{eq},j}(r) = p_j - \rho_j [\Phi(r) - \Phi(r_j)], \quad (16)$$

(Eq. 16 of Käppeli & Mishra 2016), whose two face values are $P_{\text{up}}(j) = p_j - \rho_j [\Phi_i(j) - \Phi_c(j)]$ and $P_{\text{dn}}(j) = p_j + \rho_j [\Phi_c(j) - \Phi_i(j-1)]$. The reconstruction then works on the departure from (16) rather than on the pressure itself, the Riemann pressure jump is formed as $\Delta p = \Delta p_{\text{eq}} + q_R - q_L$ from quantities that are all of the size of the departure, and the momentum row is assembled from the face pressure measured against each side's own equilibrium. The two identities that make the substitution exact are

$$\begin{aligned} A_+ P_{\text{up}} - A_- P_{\text{dn}} - (A_+ - A_-) p_j &= -\rho_j \{A_+ [\Phi_i(j) - \Phi_c(j)] + A_- [\Phi_c(j) - \Phi_i(j-1)]\}, \\ P_{\text{up}} - P_{\text{dn}} &= -\rho_j [\Phi_i(j) - \Phi_i(j-1)], \end{aligned} \quad (17)$$

the first for the PLM assembly and the second for WENO3. Their left-hand sides are the equilibrium part of the pressure terms the row carries and their right-hand sides are the discrete gravitational source, so the two cancel in the algebra, Source.f90 returns $S_2 = 0$ and neither side is ever evaluated. This is the momentum source of Käppeli & Mishra (2014), their Eq. (2.26), in spherical geometry. What cancels is still the physics the row balances, so its magnitude,

$$\mathcal{W}_j = \frac{|\rho_j \{A_+ [\Phi_i(j) - \Phi_c(j)] + A_- [\Phi_c(j) - \Phi_i(j-1)]\}|}{\Delta V_j} \quad (\text{PLM}), \quad \mathcal{W}_j = \frac{|\rho_j [\Phi_i(j) - \Phi_i(j-1)]|}{\Delta r_j} \quad (\text{WENO3}), \quad (18)$$

is formed separately and is the term the momentum residual is read against under this option.

2.7 Low-Mach dissipation of the contact mode

The HLLC flux resolves the contact wave exactly, so the dissipation it applies to the entropy family is proportional to the contact speed $S_* \simeq v$ and vanishes as the flow stalls; the acoustic families keep theirs at $|v| \pm c_s$. A stagnant layer can then carry a $2\Delta r$ pattern in v and T that nothing in the scheme removes. Low-Mach damping: (default off, `low_mach_dissipation.f90`) adds a gated fourth-difference stress (Jameson et al., 1981) to the numerical momentum flux,

$$D_p^{j+1/2} = \epsilon_4 g(M) \rho_f \lambda_f (v_{j+2} - 3v_{j+1} + 3v_j - v_{j-1}), \quad D_{\mathcal{E}}^{j+1/2} = v_f D_p^{j+1/2}, \quad (19)$$

with ρ_f, v_f the face averages and $\lambda_f = \frac{1}{2}[(|v| + c_s)_j + (|v| + c_s)_{j+1}]$ the face spectral radius. The gate $g = [\max(0, 1 - M_f^2/M_{\text{th}}^2)]^2$, written in M^2 so that it is smooth where the velocity changes sign, is unity at rest and identically zero above M_{th} (default 10^{-3}). The pair $(D_p, D_{\mathcal{E}})$ conserves total energy exactly and converts kinetic energy into heat and never the reverse; the mass flux is untouched. The term is added inside `RK_rhs.f90`, so the marching right-hand side and the stationary residual solve the same equation, which is what distinguishes it from the Shapiro filter of Sec. 7.6.

2.8 Viscosity and heat conduction

Viscosity: True and Conduction: True (both default off, `viscous_conduction.f90`) add the Navier-Stokes level of the same equations, the level the CETIMB code (the time-dependent upper-atmosphere model of Koskinen and collaborators, Koskinen et al. 2013, 2022) carries. For the Newtonian stress in spherical symmetry the only nonzero components are $\tau_{rr} = \frac{4}{3}\mu(v' - v/r)$ and $\tau_{\theta\theta} = \tau_{\varphi\varphi} = -\frac{1}{2}\tau_{rr}$, so exactly

$$S_\mu = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \tau_{rr}) + \frac{\tau_{rr}}{r}, \quad q_\mu = \frac{4}{3}\mu \left(\frac{\partial v}{\partial r} - \frac{v}{r} \right)^2, \quad S_{\mathcal{E}} = v S_\mu + q_\mu + \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \kappa \frac{\partial T}{\partial r} \right). \quad (20)$$

These differ from the expressions printed as Eqs. (B5) and (B6) of Koskinen et al. (2022): those carry a different gradient term ($\mu'v'$ against $\frac{4}{3}\mu'v/r$), a geometric coefficient $16/3$ against $8/3$, and a dissipation that is not sign definite, and the printed pair does not satisfy $v S_\mu + q_\mu = \nabla \cdot (\tau \cdot v)$. Physical correctness being the criterion, (20) is what is implemented and the discrepancy is recorded in the source.

Neither Koskinen paper prints the transport coefficients, so the atomic hydrogen conductivity of Watson et al. (1981) is used,

$$\kappa(T) = 4.45 \times 10^4 \left(\frac{T}{1000\text{K}} \right)^{0.7} \text{ erg cm}^{-1} \text{ s}^{-1} \text{ K}^{-1}, \quad \mu(T) = \frac{4}{15} \frac{m_{\text{H}}}{k_B} \kappa(T), \quad (21)$$

the second being the Chapman-Enskog/Eucken relation for a monatomic gas, Prandtl number $2/3$. Both are the *neutral* values: above the ionization front the electron conductivity is far larger and that regime is not covered, which is stated at the routine. In code units $\mu_{\text{code}} = \mu_{\text{cgs}}/(n_0 m_{\text{H}} v_0 R_0)$, the inverse Reynolds number, and $\kappa_{\text{code}} = \frac{15}{4} \mu_{\text{code}}$.

Both operators are assembled once as tridiagonal coefficient triplets on the same faces and volumes as (5), and the same triplets serve as the source entering the stationary residual and as the matrix of a Crank-Nicolson update in the marching loop, so the marched fixed point and the zero of the residual are the same state. The inner boundary takes the flux through the base face from the anchored ghost, held fixed during the implicit step; the outer boundary carries zero diffusive flux, which is the condition an extrapolated outflow ghost admits. A step whose solve leaves a cell below the temperature floor of the equation of state fails with a named status rather than returning the floor, because returning it would add energy with no source behind it.

2.9 The two-dimensional approximation

A one-dimensional radial model of an irradiated sphere has to state how the stellar flux and the mass loss are shared over the planet. The mandatory key `2D approximate` method carries that statement, in the four forms of Table 1. The dilution factor ξ is applied to every stellar beam at the point where the beam is used, and in one place only (`dayside_dilution` in `parameters.f90`): the XUV photon grid, the stellar Ly- α beam and the FUV bands all read it. The stated fluxes therefore keep their meaning as the flux at the planet's orbit, and the correction to the mass-loss rate is applied once, at the end of the run (114).

Table 1: The `2D approximate` method options, as `input_read.f90` normalizes them, `dayside_dilution` applies ξ and `EXHALE_main.f90` applies the $\dot{M}$ factor.

Key word	Stored as	ξ	$\dot{M}$ factor
<code>Mdot/4</code>	<code>Mdot/4</code>	1	1/4
<code>Rate/2</code>	<code>Rate/2 + Mdot/2</code>	1/2	1/2
<code>Rate/4</code>	<code>Rate/4 + Mdot</code>	1/4	1
<code>alpha</code>	<code>alpha</code>	1	1

The first row is the recipe of Salz et al. (2016) that the ATES paper adopts: the undiluted substellar rates with the resulting mass-loss rate divided by four. The second is the tidally locked reading, dayside averaged rates with the outflow leaving one hemisphere; the third the rapidly rotating one, rates reduced by four with the outflow leaving the whole surface. These are the three recipes of Caldiroli et al. (2021), their Sec. 2.4.

The fourth is different in kind. With `alpha` the run states a coefficient α and the attenuated field carries the further factor $1/(1 + \alpha\tau)$ inside the column integral (Sekiya et al., 1980; Odert et al., 2020), so that the cell-mean attenuation becomes

$$\langle e^{-\tau} \rangle_{\text{cell}} = \int_0^1 \frac{e^{-\tau_{\text{out}} - s \delta\tau}}{1 + \alpha(\tau_{\text{out}} + s \delta\tau)} ds, \quad (22)$$

which is the cell-mean attenuation (30) with the further factor inside the integral; it has no closed form and is integrated by composite three-point Gauss-Legendre on segments of optical depth at most 0.25. With $\alpha = 0$, the default and the only value for which the field is a pure exponential, it reduces to (30) itself.

The range of validity is stated at the routine: $\xi = 1/2$ is the illuminated hemisphere seen by an *optically thin* absorber. Where a band is shielded the true shell average is lower, the slant columns away from the substellar point being longer; it is 0.20 to 0.33 for the Lyman-Werner band over a molecular layer. No key asks for that refinement, so a shielded band is diluted by the optically thin factor.

3. The stellar radiation field and its absorption

3.1 The incident spectrum

The star is represented by a plane beam that falls radially onto the substellar column. Its spectral shape is chosen once, by the `Spectrum` type: key of `input.inp`, and that single choice fills *every* band of the photon grid: the

extreme ultraviolet and X-ray bands, and also the part below the hydrogen Lyman edge where the He 2^3S metastable, the low-ionization-potential metals and hydrogen in $n = 2$ absorb. The three types are implemented in `J_inc.f90` (`stellar_flux_ev`) and `sed_read.f90`:

Power law (`is_PL_sed`). Two power laws of one common index $\alpha \equiv \text{PLind}$, normalized separately on the EUV interval $[E_{\text{low}}, E_{\text{mid}}]$ and on the X-ray interval $[E_{\text{mid}}, E_{\text{top}}]$. With L_X and L_{EUV} the two band luminosities, a the orbital distance and $\mathcal{R} = 10^{L_X - L_{\text{EUV}}}$,

$$J_{\text{XUV}} = \frac{10^{L_X} + 10^{L_{\text{EUV}}}}{4\pi a^2}, \quad J_X = \frac{\mathcal{R}}{1 + \mathcal{R}} J_{\text{XUV}}, \quad J_{\text{EUV}} = \frac{1}{1 + \mathcal{R}} J_{\text{XUV}}, \quad (23)$$

and, for $\alpha \neq -1$ and $P_1 = \alpha + 1$,

$$J_{\text{inc}}(E) = \begin{cases} \frac{P_1 J_{\text{EUV}}}{E_{\text{mid}}^{P_1} - E_{\text{low}}^{P_1}} E^\alpha, & E < E_{\text{mid}}, \\ \frac{P_1 J_X}{E_{\text{top}}^{P_1} - E_{\text{mid}}^{P_1}} E^\alpha, & E \geq E_{\text{mid}}, \end{cases} \quad (24)$$

the logarithmic form replacing the ratio at $\alpha = -1$, and the X-ray normalization set to zero when `thereis_Xray` is false. Below E_{low} the same law is extrapolated, which is what the type means rather than an accident.

Planck (`spectrum_is_planck`). The photospheric blackbody seen at the orbit,

$$F_\nu = \pi B_\nu(T_{\text{eff}}) \left(\frac{R_\star}{a} \right)^2, \quad B_\nu = \frac{2h\nu^3/c^2}{e^{h\nu/kT_{\text{eff}}} - 1}, \quad (25)$$

converted to a flux per unit photon energy by $F_E = F_\nu d\nu/dE$ with $d\nu/dE = 2.417989242 \times 10^{14} \text{ Hz eV}^{-1}$. For $h\nu/kT_{\text{eff}} > 700$ the Wien form replaces $1/(e^x - 1)$ so that no exponential overflows. The absolute scale comes from T_{eff} and $R_\star$, not from the two luminosities, so the grid-integrated flux generally differs from J_{XUV} ; `write_setup_report.f90` states both.

Load (`do_read_sed`). A measured spectral energy distribution read by `read_sed` (`sed_read.f90`): two columns, bin central wavelength in $\text{\AA}$ and flux at the planet in $\text{erg cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$, ascending in wavelength. Between two rows the field is interpolated log-linearly in $\log E$, exact for the power-law segments a tabulated continuum is made of; outside the band it returns zero, and `spectrum_covers_ev` must be asked first so that the run stops rather than integrating a zero the input never stated.

A monochromatic option (`is_monochr`) puts the whole $10^{L_{\text{EUV}}}/4\pi a^2$ at a single energy E_{low} .

Bands of the photon grid. `set_energy_vectors.f90` lays the grid out over at most six bands,

$$[E_{\text{floor}}, E_{\text{abs}}], [E_{\text{abs}}, 13.598 \text{ eV}], [13.598, 24.587], [24.587, 54.418], [54.418, E_{\text{mid}}], [E_{\text{mid}}, E_{\text{top}}],$$

with fixed bin budgets (20, 89, 50, 50, 50, 50). E_{low} , E_{mid} and E_{top} are stated by the `[E_low, E_mid, E_high]` line of `input.inp` (the tutorial run uses 13.60, 123.98 and $1.24 \times 10^3 \text{ eV}$; with `Use only EUV? True` the top is held at $1.24 \times 10^3 \text{ eV}$ and the X-ray normalization is zero). E_{abs} is `sub_lyman_absorber_threshold_ev()`, the lowest ionization threshold of an absorber of the bulk gas below the hydrogen edge: the He 2^3S metastable at 4.768 eV when it is tracked, or the neutral stage of an active low-IP metal (K I 4.341, Na I 5.139, Ca I 6.113, Mg I 7.646, Fe I 7.902, Si I

8.152, S I 10.36 eV) when one is lower. E_{floor} is `photon_grid_floor_eV()`, which lowers that further to $E_{\text{th}}(\text{H}, n=2) = E_{\text{th}}(\text{H})/4 = 3.3996 \text{ eV}$, the head of the Balmer continuum, whenever the excited-hydrogen coupling is armed. The interval $[E_{\text{floor}}, E_{\text{abs}}]$ carries no bulk-gas absorber: it exists so that the $\text{H}(n=2)$ band is stated, that rate being a closed-form integral over it (Sec. 4.6) and not a sum over these bins.

Every ionization threshold is a bin edge. A photoionization rate is the rectangle sum $\sum_k F_k \sigma_k \Delta E_k / E_k$, so a bin that contains a threshold would charge the absorber its cross section, read at the bin center, over the part of the bin below the threshold where it cannot absorb at all. `ionization_thresholds_active` therefore collects every threshold the run carries (H I, He I, He II, He 2³S, H₂ and its two channel thresholds, and every photoionizable stage of an element with nonzero abundance together with its subshell turn-ons); each band is cut at those falling strictly inside it, and its budget is shared over the subintervals in proportion to their logarithmic widths (largest remainder, at least one bin each). The bins are geometric and E_k is the *geometric* mean of the edges, the more accurate abscissa of a rectangle rule for a power-law integrand. A grid whose points sat *on* the thresholds instead overstates the exact P_{HI} , P_{HeI} and P_{HeII} by factors 1.095, 1.016 and 1.030.

Dayside dilution. Every stellar beam of the code is multiplied by the dilution ξ_{day} of Table 1: the XUV grid in `set_energy_vectors`, $F_{\text{XUV}} \rightarrow \xi_{\text{day}} F_{\text{XUV}}$; the stellar Ly α beam in `lya_rt.f90` and `excited_hydrogen.f90`; and the four far-ultraviolet bands through `fuv_band_flux`.

Bands stated by their integrated flux. Four far-ultraviolet bands lie almost entirely below the photon grid and are given by scalar keys instead, in the band order of `water_photolysis.f90`: LW (912–1201 Å, Stellar LW flux), B2 (the H I Ly α line at 1215.67 Å, Stellar Ly α flux), B3 (1231–1450 Å) and B4 (1451–2304 Å). `fuv_band_flux` returns $\xi_{\text{day}} F_{\text{band}}$, the dilution being applied where the beam is used so that the keys keep their stated meaning, the band flux at the planet's orbit.

3.2 Photoionization cross sections

All cross sections are carried in units of 10^{-18} cm^2 (`cross_sec.f90`) and are evaluated once, at the grid centers, into the vectors `s_hi`, `s_hei`, `s_heii`, `s_heiTR`, `s_h2` and the table `sigma_tab`.

Hydrogenic ions (H I, He II), function `sigma`. With $\epsilon = \sqrt{E/E_{\text{th}} - 1}$,

$$\sigma(E) = \frac{6.3}{Z^2} \left(\frac{E_{\text{th}}}{E} \right)^4 \frac{\exp[4 - 4 \arctan(\epsilon)/\epsilon]}{1 - \exp(-2\pi/\epsilon)}, \quad E > E_{\text{th}}, \quad (26)$$

and $6.3/Z^2$ exactly at threshold, zero below it. E_{th} is the *measured* potential of the ion, the same energy the grid uses as a band edge and the photoelectron is charged against.

He I ground singlet The Verner et al. (1996) single-shell fit,

$$x = \frac{E}{E_0} - y_0, \quad z = \sqrt{x^2 + y_1^2}, \quad Q = 5.5 - P/2, \quad \sigma = \sigma_0 [(x-1)^2 + y_w^2] z^{-Q} \left(1 + \sqrt{z/y_a} \right)^{-P}, \quad (27)$$

with the published parameters $(E_0, \sigma_0, y_a, P, y_w, y_0, y_1) = (13.61, 949.2, 1.469, 3.188, 2.039, 0.4434, 2.136)$ and E_{th} taken as the global He I threshold (it only gates the fit). The legacy ATES two-term fit $0.6935/[(E/100)^{1.82} + (E/100)^{3.23}]$ is restored by `ates_photoion_rate` (default off).

He 2³S metastable Two forms of (27) joined by a log-linear bridge: wing A from the 4.768 eV threshold to the Cooper minimum at 34.70 eV, wing B from the resonance-averaged bump at 45.59 eV upward, and a power law joining them continuously in between. Each wing is a fit at the few-percent level to the Norcross (1971) and TOPbase background.

Metal ions `metal_photoion_sigma` returns the *shell sum* over the seven subshells of Verner et al. (1996) and Verner & Yakovlev (1995); the 1995 form is (27) with $y_0 = y_1 = 0$ and $Q = 5.5 + \ell - P/2$. Above the E_{max} of the 1996 outer-shell table the handover to the 1995 subshell fit is made, as in Verner's own `phfit2`; extrapolating the outer-shell fit is not conservative (the Fe I row rises as E^{2-Q} and reaches 24.9 Mb at 1240 eV against a true 0.51 Mb). An inner-shell absorption is charged to one ionization of the absorbing stage with a photoelectron of $h\nu - E_{\text{th}}(\text{outer})$, which approximates the Auger cascade and undercounts the charge state by at least one stage; this is the approximation the published exoplanet photochemistry models make (Cecchi-Pestellini et al., 2009; Locci et al., 2022). Fluorescence is neglected: with $E_{\text{top}} = 1240$ eV only C, N, O and Na I can have a K hole, and their K yields (Krause, 1979) are 0.28, 0.52, 0.83 and 2.3 percent.

H₂ `sigma_H2` is a piecewise *measured* total photoabsorption: $k_{\text{bx}} \times$ Backx et al. (1976) Table 1 over 15.4–18 eV with $k_{\text{bx}} = 1.016415$, Samson & Haddad (1994) Table 1 over 18–300 eV, and $k_{\text{hi}} \times$ the Yan et al. (1998) eq. (19) $E^{-7/2}$ sum-rule tail above 300 eV with $k_{\text{hi}} = 0.885805$, all interpolated in $\log \sigma$ against $\log E$. No piece of the Yan fit survives below 300 eV: its three published pieces do not meet at their own join energies, stepping by a factor 1.719 at 18 eV and 0.619 at 85 eV.

Opacity models `opacity_models.f90` dispatches the cross sections through a model selected in the optional `opacity.inp`: A analytic (the curves above, the default), C constant above threshold, P constant with the Robinson & Catling (2012) pressure broadening $f(p) = 1 + a(p/p_{\text{piv}})^n$, and T a two-column table. In model P the multiplier $f(p)$ of a cell is stored in `opa_pf(j)` and carried identically in the optical depth and in the local absorption, so that the photons a cell removes from the beam are the photons it absorbs.

3.3 Transfer along the radial ray

The transfer is frequency-integrated over the grid of Sec. 3.1 and is pure absorption: there is no scattering term and no diffuse field in the XUV. For cell j the column of each absorber *outside* the cell is accumulated from the top down by `advance_starward_columns` (`util_ion_eq.f90`) on the same rectangle rule the grid uses,

$$N_a^{\text{out}}(j) = \sum_{j' > j} n_a(j') \Delta r_{j'} R_0 f(p_{j'}), \quad (28)$$

and the cell's *own* depth is formed from its own density rather than as a difference of two columns, which is the same number in exact arithmetic and loses its digits to cancellation for a thin cell. The two optical depths at photon energy

E_k are

$$\tau_k^{\text{out}} = \sum_a \sigma_{a,k} N_a^{\text{out}}, \quad \Delta\tau_k = \sum_a \sigma_{a,k} n_a(j) \Delta r_j R_0 f(p_j), \quad (29)$$

the sum running over H I, the He I ground singlet, He II, He 2³S when tracked, H₂ when the molecular chemistry is on, and every photoionizable metal stage.

The attenuated flux the cell's *rate* sees is the *mean* of the beam over the cell, not its value at either face (cell_mean_attenuation in utilities.f90). Since the densities are uniform inside a cell, $\tau(s) = \tau^{\text{out}} + s \Delta\tau$ and

$$\langle e^{-\tau} \rangle_{\text{cell}} = e^{-\tau^{\text{out}}} \frac{1 - e^{-\Delta\tau}}{\Delta\tau}, \quad (30)$$

the second factor being absorbed_fraction_per_unit_depth, which switches to its series $1 - \Delta\tau/2 + \Delta\tau^2/6$ below $\Delta\tau = 10^{-8}$. Only this mean makes the photons the cell absorbs equal the photons the beam loses across it; a one-point rule at the inner face is low by 42 percent at $\Delta\tau = 1$, the depth a cell carries at the ionization front. The optional alpha form of the 2-D approximation replaces (30) by (22), which has no closed form.

The working flux of the cell is therefore

$$\mathcal{F}_k(j) = F_{\text{XUV},k} \langle e^{-\tau} \rangle_{k,\text{cell}} f(p_j), \quad (31)$$

the model-P factor being folded once into the shared flux weight rather than into each cross section, which is identical because every integrand below is linear in exactly one cross section.

3.4 Photoionization and photoheating rates

Every integral is the rectangle sum over the grid, $\sum_k (\cdot) \Delta E_k$, with ΔE_k the exact bin width, so $\sum_k \Delta E_k = E_{\text{top}} - E_{\text{floor}}$ exactly. For absorber a the photoionization rate per particle and the photoheating rate of *one* particle of that absorber are

$$P_a = \sum_k \mathcal{F}_k \frac{\sigma_{a,k}}{E_k} \Delta E_k \times 10^{-18} \text{ erg}^{-1} \text{ eV}, \quad (32)$$

$$h_a^{(1)} = \sum_k \mathcal{F}_k \left(1 - \frac{E_{\text{th},a}}{E_k} \right) f_{\text{heat}}(E_k) \sigma_{a,k} \Delta E_k \times 10^{-18} \text{ [erg s}^{-1}\text{]}, \quad (33)$$

where the factor $1 - E_{\text{th}}/E$ is the share of the photon left as photoelectron kinetic energy (photoelectron_share; the key photoheat_photon_fraction, default off, replaces it by a constant fraction), and f_{heat} is the secondary-ionization heat fraction of Sec. 3.5. The heating of the cell is the contraction of these one-particle rates with the composition,

$$\mathcal{H} = \sum_a n_a h_a^{(1)}, \quad (34)$$

performed in exactly one place (photoheating_of_cell) and split by absorber into the six columns H I, He I, He II, He 2³S, H₂, metals. The heating efficiency written to the output is

$$\eta_{\text{heat}} = \frac{\mathcal{H}}{\sum_k \mathcal{F}_k \left(\sum_a \sigma_{a,k} n_a \right) \Delta E_k \times 10^{-18}}, \quad (35)$$

the ratio of the energy deposited to the energy absorbed, set to zero in a transparent or unilluminated cell.

Molecular channels. H_2 absorption is resolved into four mutually exclusive final states whose cross sections are *shares* of σ_{H_2} , so the opacity and the total H_2 destruction rate are unchanged. Each channel is charged its own threshold in (33), and the two deposits that are not a photoelectron, the kinetic energy release of the double channel and the internal excitation the neutral channel radiates away, are stated with the channels themselves in Sec. 6.2.

Self-consistency of the field with the composition. Nothing in (29) depends on any cell inside j , which is what lets the equilibrium sweep march outside in and re-form the field of one cell against its own composition (Sec. 4.4); the cell mean (30) is formed once per pass so that no second attenuation is stacked on the first. The rates are constants of the cell's nonlinear solve, carrying the field dependence inside the residual meaning a re-integration of the photon grid at every residual evaluation.

3.5 Secondary ionization by photoelectrons

A photoelectron faster than $E_{\text{sec}} = 30 \text{ eV}$ does not thermalize completely: part of its energy ionizes further neutrals. `electron_energy_degradation.f90` divides the labor between two published determinations. Shull & van Steenberg (1985) supply the energy budget of the ionization channels as a function of the ionized fraction x ,

$$\phi_{\text{HI}}(x) = 0.3908 (1 - x^{0.4092})^{1.7592}, \quad \phi_{\text{HeI}}(x) = 0.0554 (1 - x^{0.4614})^{1.6660}, \quad (36)$$

whose form vanishes at $x = 1$ where no neutral is left. Dalgarno et al. (1999) supply the heating fraction and everything that depends on molecular hydrogen; their Table 4 mean energies per ion pair, $W(x, E_0) = a(1 + cx^b)$, give the energy dependence,

$$f_{\text{HI}}(x, E_0) = \phi_{\text{HI}}(x) \frac{W_{\text{H}^+}(x, 1 \text{ keV})}{W_{\text{H}^+}(x, E_0)}, \quad f_{\text{HeI}}(x, E_0) = \phi_{\text{HeI}}(x) \frac{W_{\text{He}^+}(x, 1 \text{ keV})}{W_{\text{He}^+}(x, E_0)}, \quad (37)$$

so that the high-energy limit is exactly Shull & van Steenberg's. The division of that energy among the targets is recomputed on the cell's own densities, using the collision weight 1.89 that Dalgarno et al.'s eqs. (9) and (10) imply (their coefficients 1.89 and 0.53 are reciprocal to 0.17 percent), with $r_{\text{ref}} = n(\text{HeI})/n(\text{HI}) = 0.1$ the reference composition of the fits:

$$w = (n_{\text{HI}} + 1.89 n_{\text{H}_2}) r_{\text{ref}} f_{\text{HI}} + n_{\text{HeI}} f_{\text{HeI}}, \quad c_{\text{HI}} = \frac{f_{\text{ion}} r_{\text{ref}} f_{\text{HI}}}{w}, \quad c_{\text{H}_2} = 1.89 c_{\text{HI}}, \quad c_{\text{HeI}} = \frac{f_{\text{ion}} f_{\text{HeI}}}{w}, \quad (38)$$

with $f_{\text{ion}} = f_{\text{HI}} + f_{\text{HeI}}$. These coefficients are *per target particle* (units cm^3), so $n_{\text{HI}} c_{\text{HI}} + n_{\text{H}_2} c_{\text{H}_2} + n_{\text{HeI}} c_{\text{HeI}} = f_{\text{ion}}$ identically and nothing is divided by a vanishing neutral density. The heat fraction is

$$f_{\text{heat}}(E_0) = 1 - [1 - \eta_{\text{mix}}(x, E_0)](1 - x) + f_{\text{diss}} + f_{\text{vib}} + f_{\text{fluor}}, \quad (39)$$

with $\eta_{\text{D}} = 1 + (\eta_{\text{D},0} - 1)/(1 + cx^b)$ their eq. (14) and η_{mix} their eq. (12) mixing of the H/He and H_2 /He efficiencies. The factor $(1 - x)$ is this code's closure and not either source's: the channels that compete with heat need a neutral to act on, and Dalgarno et al.'s form leaves 10 percent of the primary energy unthermalized even at $x = 1$. The three molecular terms (fragment kinetic energy, direct vibrational excitation of $v = 1, 2$, and the vibrational energy the B and C states return on fluorescing) vanish identically with $n(H_2)$; the last two carry the collisional quench fraction of `h2_vibrational_relaxation.f90`, since a radiated excitation is not heat.

The secondary rates are added to the primary ones of (32): for an absorber a whose photoelectron energy exceeds E_{sec} ,

$$P_{\text{HI}}^{\text{sec}} = \sum_k \mathcal{F}_k \frac{\sigma_{a,k} n_a}{E_k} c_{\text{HI}}(E_k) \frac{E_k - E_{\text{th},a}}{E_{\text{th}}(\text{HI})} \Delta E_k \times 10^{-18} \text{ erg}^{-1} \text{ eV}, \quad (40)$$

and likewise for He I and H₂; the H₂ secondaries additionally make one proton per 22 H₂⁺ ions (Dalgarno et al., 1999, after their eq. 10), which is an added yield and not a share.

Staged activation. The key `Secondary_ionization` is *on* by default (`use_sec_ion = .true.`) but *staged*: the runtime flag `sec_ion_active` stays false until the wind has first converged without the coupling, because from a cold initial condition the base feedback amplifies the startup transient into a runaway (`EXHALE_main.f90`, the block that sets `sec_ion_active = .true.` once any of the mass-flux, step-change, plateau or steady gates is met, and again before the JFNK handover so that Newton is never asked to solve a system the marched state does not approximately satisfy). `Secondary_ionization: Immediate` applies it from step 0 and `False` disables it entirely. Photoelectrons between 13.6 and 30 eV are still treated as depositing all their energy as heat, which is a limit of the published tables and not a physical threshold.

3.6 Reprocessed and non-stellar fields

Three fields in the code are not the direct stellar beam.

Helium recombination radiation. With `He_rec_coupling` (default on), the photons emitted when He II recombines are coupled on the spot to the ionization of H I, H₂ and the metals (`he_rec_coupling` in `util_ion.eq.f90`). Each cascade exit is represented at one photon energy: the ground-capture continuum starting at $E_{\text{th}}(\text{He I})$, the $2^1P \rightarrow 1^1S$ resonance line at 21.2 eV, the $2^3S \rightarrow 1^1S$ line at 19.8 eV, and the part of the 2^1S two-photon continuum above the hydrogen edge. The last is not a line: integrating the Drake et al. (1969) shape over 13.598–20.62 eV gives 0.5564 ionizing photons per decay carrying 8.9646 eV, a mean in-band photon energy of 16.110 eV and hence 2.512 eV of photoelectron energy per photon, against 3.511 eV for a flat distribution over the same window.

Lyman-Werner band. The 912–1201 Å beam is solved once for all of its absorbers together (`fuv_lw_photon_field`), because an H₂O molecule cannot absorb a photon an H₂ line has already taken. Its transmission down to a face is $T = (1 - A)e^{-\tau_c}$ with $A = \int \sigma_{\text{pump}} f_{\text{shield}}(N_{\text{H}_2}) n_{\text{H}_2} dr$, and across one cell the identity

$$T^{\text{out}} - T^{\text{in}} = e^{-\tau^{\text{out}}} [(1 - A^{\text{out}}) (1 - e^{-\Delta\tau}) + \Delta A e^{-\Delta\tau}] \quad (41)$$

is exact, so the photons the cell removes split into a continuum share and a line share with nothing left over. The continuum factor $e^{-\tau_c}$ is the Draine & Bertoldi (1996) eq. (40) term, identically 1 where the oxygen option is off. Line self-shielding is the tabulated function of (T, n, N_{H_2}) of Sec. 6.3. CO is shielded on the same beam by the Visser et al. (2009) function $\Theta(N_{\text{CO}}, N_{\text{H}_2})$ (`co_self_shielding_table.f90`).

Ly α and the Balmer continuum are described with the excited-hydrogen model in Sec. 4.6.

4. Ionization and chemical state

4.1 The species carried

The composition of a cell is the vector $f_sp(j, \cdot)$ of species fractions, laid out once in `species_table.f90`:

column(s)	species	note
1–2	H I, H II	fractions of the hydrogen nuclei
3–5	He I, He II, He III	fractions of the helium nuclei; column 3 <i>contains</i> the metastable
6	He 2 ³ S	the metastable triplet, tracked when Include He23S? is true (default on)
7–33	the 27 metal ion stages	C, O, N, Mg, Si, Ca, Fe in three stages; Na, K, S in two
34–37	H ₂ , H ₂ ⁺ , H ₃ ⁺ , HeH ⁺	molecular option (<code>thereis_mol</code> , default off)
38–40	OH, H ₂ O, CO	oxygen option (<code>thereis_oxychem</code> , default off)

Ten elements carry a nuclear abundance n_X/n_H , set by `metals.inp` (`metals_input_read.f90`) or by a lower-atmosphere handoff. Masses are in units of the hydrogen *atom*: helium weighs $m_{He}/m_H = 3.9715259$ and the metal weights are the CIAAW values converted from the unified mass unit by $1\text{ u} = 0.99223573 m_H$.

4.2 The cell system

For each cell the code solves a system of algebraic balance equations at fixed temperature, density and radiation field. The unknowns are stage *fractions*, and one stage of each element is closed as the remainder rather than being an unknown: $n(\text{HI}) = (1 - x_{H^+})n_H$, $n(\text{HeI}) = (1 - x_{He^+} - x_{He^{2+}})n_{He}$, and for a metal element $n(X^0) = (1 - x_{X^+} - x_{X^{2+}})n_X$. The layout is

rows	contents
1–3	$x_{H^+}, x_{He^+}, x_{He^{2+}}$
4	He 2 ³ S (triplet systems) <i>or</i> H ₂ (molecular systems)
5–8	H ₂ ⁺ , H ₃ ⁺ , HeH ⁺ , He 2 ³ S (molecular systems)
$m_{\text{base}} \dots$	two rows per metal element (<code>metal_row_base()</code>)
$i_{\text{ox}} \dots$	OH and H ₂ O (oxygen option)

with $m_{\text{base}} = 4$, or 5 with the triplet. In the molecular layout $i_{\text{ox}} = 8$, or 9 with the triplet, and $m_{\text{base}} = i_{\text{ox}}$, raised by two when the oxygen chemistry is on (`metal_row_base`, `oxygen_row_base`: one definition, read by the residual and by the driver alike). Seven residual routines cover the configurations: `System_H`, `System_HeH`, `System_HeH_metals`, `System_HeH_TR`, `System_HeH_TR_metals`, `System_HeH_mol` and `System_HeH_mol_metals`. The shared rows live once, in `ion_residual_core.f90`.

The H/He rows. With photoionization rates Γ , case-B recombination coefficients α and electron-impact ionization coefficients β , `heh_rows` writes

$$0 = n_{\text{HI}}\Gamma_{\text{HI}} + (n_{\text{HI}}\beta_{\text{HI}} - \alpha_{\text{HII}}n_{\text{HII}})n_e, \quad (42)$$

$$0 = n_{\text{HeI}}\Gamma_{\text{HeI}} + (n_{\text{HeI}}\beta_{\text{HeI}} - \alpha_{\text{HeII}}n_{\text{HeII}})n_e, \quad (43)$$

$$0 = n_{\text{HeII}}\Gamma_{\text{HeII}} + (n_{\text{HeII}}\beta_{\text{HeII}} - \alpha_{\text{HeIII}}n_{\text{HeIII}})n_e, \quad (44)$$

with n_e the charge sum of every tracked ion, $n_e = n_{\text{HII}} + n_{\text{HeII}} + 2n_{\text{HeIII}} + \sum_i Z_i n_{\text{m},i} + n_{\text{H}_2^+} + n_{\text{H}_3^+} + n_{\text{HeH}^+}$ (`calc_ne`), so charge neutrality is imposed identically rather than solved for. Each metal element contributes

$$0 = n_{X^0}\Gamma_{X^0} + (n_{X^0}\beta_{X^0} - \alpha_{X^+}n_{X^+})n_e, \quad 0 = n_{X^+}\Gamma_{X^+} + (n_{X^+}\beta_{X^+} - \alpha_{X^{2+}}n_{X^{2+}})n_e, \quad (45)$$

the second row being replaced by $x_{X^{2+}} = 0$ for a two-stage element, and the whole pair by $x_{X^+} = x_{X^{2+}} = 0$ for an element with zero abundance.

The triplet row. With the metastable tracked, the He rows are rewritten in the form of Oklopčić & Hirata (2018) (rows 2 and 3 summed, the triplet and the singlet ground state distinguished) and the fourth row is the 2^3S balance,

$$0 = -n_{2^3\text{S}}\Gamma_{2^3\text{S}} + n_e [n_{\text{HeII}}\alpha_{2^3\text{S}} + n_{1^1\text{S}}q_{13} - n_{2^3\text{S}}(q_{31a} + q_{31b}) - n_{2^3\text{S}}\beta_{2^3\text{S}}] - n_{2^3\text{S}}(A_{31} + n_{\text{HI}}Q_{31}), \quad (46)$$

where q_{13} is the collisional excitation out of the singlet ground state, q_{31a} and q_{31b} the collisional conversions to 2^1S and 2^1P , $A_{31} = 1.272 \times 10^{-4} \text{ s}^{-1}$ the forbidden radiative decay, $\beta_{2^3\text{S}}$ the electron-impact ionization of the metastable at its own 4.8 eV threshold, and Q_{31} the *total* $\text{He}(2^3\text{S})+\text{H}$ ionization rate coefficient. Both channels of Q_{31} quench the metastable, so the sink takes the sum; only the Penning branch $f_P = 0.9$ leaves a lasting proton, and it appears as the source $f_P Q_{31} n_{2^3\text{S}} n_{\text{HI}}$ in the H^+ row. The remaining 10 percent is associative ionization to HeH^+ , which the atomic systems drop (in an atomic gas HeH^+ dissociatively recombines straight back) and the molecular system carries explicitly.

The molecular rows. `mol_heh_rows` adds four species balances on the Koskinen et al. (2022) Table 1 network (Sec. 6.1). Schematically, writing g for photorates and k_n for the Table-1 coefficients,

$$\begin{aligned} \text{H}_2 : \quad 0 &= k_6 n_e n_{\text{H}_3^+} + k_9 n_{\text{H}_2^+} n_{\text{HI}} + k_{11} n_{\text{H}_3^+} n_{\text{HI}} + k_{15} n_{\text{HI}}^2 \\ &\quad - [g_{\text{H}_2} + g_{\text{LW}} + (k_{10} + k_{13})n_{\text{HII}} + k_{12}n_{\text{tot}} + k_{14}n_e + k_8 n_{\text{H}_2^+} + (k_{17} + k_{20} + k_{23})n_{\text{HeII}} + k_{18}n_{\text{HeH}^+} + k_{\text{H}_2}^{\text{ion}} n_{2^3\text{S}}] n_{\text{H}_2}, \end{aligned} \quad (47)$$

$$\begin{aligned} \text{H}_2^+ : \quad 0 &= (g_{\text{H}_2} - g^{\text{di}} - g^{\text{dd}} - g^{\text{nd}}) n_{\text{H}_2} + k_{10} n_{\text{HII}} n_{\text{H}_2} + k_{11} n_{\text{H}_3^+} n_{\text{HI}} + k_{19} n_{\text{HeH}^+} n_{\text{HI}} + k_{23} n_{\text{HeII}} n_{\text{H}_2} \\ &\quad + f_P k_{\text{H}_2}^{\text{ion}} n_{2^3\text{S}} n_{\text{H}_2} - (k_5 n_e + k_8 n_{\text{H}_2} + k_9 n_{\text{HI}}) n_{\text{H}_2^+}, \end{aligned} \quad (48)$$

$$\text{H}_3^+ : \quad 0 = k_8 n_{\text{H}_2^+} n_{\text{H}_2} + k_{13} n_{\text{HII}} n_{\text{H}_2} + k_{18} n_{\text{HeH}^+} n_{\text{H}_2} - [(k_6 + k_7)n_e + k_{11} n_{\text{HI}}] n_{\text{H}_3^+}, \quad (49)$$

$$\text{HeH}^+ : \quad 0 = k_{20} n_{\text{HeII}} n_{\text{H}_2} + (1 - f_P) n_{2^3\text{S}} (Q_{31} n_{\text{HI}} + k_{\text{H}_2}^{\text{ion}} n_{\text{H}_2}) - (k_{16} n_e + k_{18} n_{\text{H}_2} + k_{19} n_{\text{HI}}) n_{\text{HeH}^+}, \quad (50)$$

and the H^+ row gains $(g^{\text{di}} + 2g^{\text{dd}})n_{\text{H}_2} + k_9 n_{\text{H}_2^+} n_{\text{HI}} + k_{17} n_{\text{HeII}} n_{\text{H}_2} - (k_{10} + k_{13})n_{\text{HII}} n_{\text{H}_2}$. The factor 2 on the double photoionization branch is the stoichiometry, two protons out of one event, while the H_2 destroyed is still one per event, which is why (47) is not doubled with it.

4.3 The rate coefficients

All of the following live in `Cool_coeff.f90` unless stated. The default set is selected by `Legacy_HHe_rates`: `False` (default) and `Atomic_rate_set`: `default`; the two alternatives are the Hui & Gnedin (1997) fits and the Koskinen et al. (2022) Table 1 fits, and the three meet at the accessor functions at the end of that module.

Recombination. Radiative recombination follows the Badnell (2006, 2023 update) total ground-level fit

$$\alpha_{\text{RR}}(T) = \frac{A}{\sqrt{T/T_0} \left(1 + \sqrt{T/T_0}\right)^{1-B'} \left(1 + \sqrt{T/T_1}\right)^{1+B'}}, \quad B' = B + Ce^{-T_2/T}, \quad (51)$$

implemented as `rr_badnell` (its dummy `T0` keeps the published notation and is not the code's temperature scale). Dielectronic recombination of He II adds the adf09 three-term sum $T^{-3/2} \sum_i c_i e^{-E_i/T}$. Case B is formed by subtracting the $n = 1$ direct-capture level fit of Mao & Kaastra (2016),

$$\alpha_1(T) = \frac{a_0 \theta^{-b_0 - c_0 \ln \theta} (1 + a_2 \theta^{-b_2})}{1 + a_1 \theta^{-b_1}}, \quad \theta = k_B T \text{ [eV]}, \quad (52)$$

so that $\alpha_B = \alpha_A - \alpha_1$ for H II, He II and He III. The metal stages take Badnell RR + adf09 DR through the same two forms, with the one gap (Ca II, K-like) handled separately and noted at the code site.

Collisional ionization. Voronov (1997) Table 1,

$$\beta(T) = A \frac{(1 + P\sqrt{U}) U^K e^{-U}}{X + U}, \quad U = \frac{\Delta E}{k_B T}, \quad (53)$$

with $(\Delta E, P, A, X, K)$ equal to $(13.6, 0, 2.91 \times 10^{-8}, 0.232, 0.39)$ for H I, $(24.6, 0, 1.75 \times 10^{-8}, 0.180, 0.35)$ for He I and $(54.4, 1, 2.05 \times 10^{-9}, 0.265, 0.25)$ for He II. The ΔE here is Voronov's own fit coefficient, not the measured potential; the energy *removed* per ionization is the measured one, E_{th} in erg.

Charge exchange. `charge_exchange.f90` carries the 63 reactions of Huang et al. (2023) Table 4 plus one row that is not in it. The default active set is Group A, the 23 metal+H/H⁺ reactions; `cx_full` adds the metal+He (C) and metal+metal (D) groups. The He $\leftrightarrow$ H pair (Group B), $\text{He}^0 + \text{H}^+ \rightleftharpoons \text{He}^+ + \text{H}^0$, is applied separately by `he_h_cx_fvec` so that it is available in *every* system containing helium and is never counted twice; it is gated by `He_H_charge_exchange` (default on). Group E is the one reaction outside Table 4, $\text{O}^{2+} + \text{H}^0 \rightarrow \text{O}^+ + \text{H}^+$, on by default at the published rate because its absence leaves the O III profile wrong by decades in the region a transit probes; `cx_O2p_H 0` in `metals.inp` restores the Table-4-only set exactly. Each reaction enters as $R = k n_{\text{donor}} n_{\text{acceptor}}$, so the zero-abundance guard is automatic.

Penning ionization of the metastable. The total He(2³S)+H ionization rate coefficient is the Maxwell-Boltzmann average of the Cohen & Lane (1971) and Morgner & Niehaus (1979) cross sections as collected by García Muñoz (2025),

$$Q_{31}(T) = 10^{-9} \exp \left[-\frac{86.4804}{T} - 0.286766 \ln T + 0.0868445 (\ln T)^2 - 5.73001 \times 10^{-3} (\ln T)^3 \right] \text{ cm}^3 \text{ s}^{-1}, \quad (54)$$

smooth, positive and single-peaked at 3.8×10^3 K. The code notes at that site why the two-branch power law of Taylor et al. (2025) Table 2 is *not* used: it jumps by a factor 1.5724 at 4000 K, which a rate coefficient cannot do, and it disagrees

with that paper's own Figure 19; the expression above agrees with Figure 19 to 10–20 percent. The corresponding $\text{He}(2^3\text{S})+\text{H}_2$ coefficient is $k = aT^b e^{c/T}$ with the Cohen & Lane (1977) amplitudes of the García Muñoz (2025) network file, reproducing its Table A.5 to 0.13 percent. Both split 0.9:0.1 between the Penning and the associative branch.

4.4 Solving one cell

The solver. `solve_ieq` (`newton_solver.f90`) tries a damped Newton iteration with an *analytic* Jacobian first (`newton_dense`: backtracking Armijo line search on $\|f\|_2$, at most 60 iterations and 20 line-search steps), and on any failure restores the entry guess and calls the in-tree MINPACK `hybrd1` from the same guess, so the result can never be worse than the forward-difference solve. Each residual routine ships its own exact Jacobian; the photoionization rates are constants of the solve, so those Jacobians are exact for the system actually solved (Sec. 3.4). Run-wide counters record how many cells each route handled, and `EXHALE_FORCE_HYBRD1` forces the MINPACK path for an A/B comparison. `use_newton_ieq` is on by default.

The sweep. `ioniz_eq` (`ionization_equilibrium.f90`) takes the temperature and the total density as *inputs* and overwrites the species fractions in place; chemistry rearranges nucleons and electrons and creates no mass, so the mass density is never corrected by this step and the mass sum of the returned composition is a check against it (`rho_recon`), never a correction. The grid is traversed from the top down in blocks; the columns of (28) are advanced with the composition already solved, so the field each cell is solved at is self-consistent with the states above it in one traversal. Inside a cell the pair (field, composition) is iterated at most `xuv_self_field_passes` times, stopping when the accepted stage fractions move by less than `xuv_self_field_tol`. Cells within a block are independent and the sweep is OpenMP-parallel with the per-cell scratch declared `threadprivate`; blocks are of fixed even length so that the vectorized transcendental calls pair the same lanes at any thread count.

Acceptance classes. Every cell's returned state carries a class, and the contract is that classes 1, 2, 3 and 5 are *roots* of the requested equations (each passed the same two tests, a state inside the element simplex and a normalized reaction residual at or below `ieq_res_tol`) and differ only in how the root was found:

class	meaning
1	root of the last attempt made
2	root of the best earlier attempt
3	root after projecting the closest attempt onto the element budget
4	not a root; accepted under the relaxation amnesty and reported
5	root of the constrained element-conserving continuation solve
6	not a root, residual above <code>ieq_nonroot_res_cap</code> : the cell <i>keeps the composition it entered the sweep with</i>

A residual that is not a finite number is a class-4 non-root by construction, since NaN passes every tolerance test. A consumer counting states its chemistry does not describe counts classes 4 and 6 and nothing else.

The constrained continuation. `constrained_chemical_equilibrium.f90` is the second formulation of the molecular network, and it removes the possibility of an unphysical iterate instead of repairing one. The unknowns are $u_k = \ln n_k$, so $n_k > 0$ for every iterate, converged or not (the logarithmic composition variables of the CEA method, Gordon & McBride 1994); element conservation is an explicit residual row per element rather than a closure, so a species carrying nuclei of two elements (HeH^+ , OH , H_2O) satisfies both rows at the accepted root; and n_e is the charge-neutrality sum, never an unknown. Because the network is bistable in its starting point (a molecular basin in the shielded base, an atomic basin above the $\text{H}_2 \rightarrow \text{H}$ front), the solve is a natural-parameter continuation (Allgower & Georg, 2003) in the radiation field: λ scales the field from six decades below the cell's own, where the composition is known in closed form from the H_2 dissociation equilibrium of the local p, T , up to $\lambda = 1$, each rung starting from the previous rung's root and a failed rung being bisected geometrically. The rungs use the general MINPACK `hybrd` and not `hybrd1`, whose fixed initial trust radius $100\|x\|$ would open a first step of e^{10^4} in every density when $\|u\| \sim 130$. A transported fraction, when one is imposed, enters as a further constraint row.

4.5 Ionization carried by the flow

Where the recombination time is not short compared with the flow time, the local balance of Sec. 4.2 is not the answer. Two places in the code carry the ionization state along the flow instead.

The implicit advection systems. `System_implicit_adv_H`, `_HeH` and `_HeH_TR` solve, per hydrogen nucleus and with $c_1 = \Delta r/v$ the residence time of the cell,

$$0 = x_{\text{HI}}^{\text{up}} - x_{\text{HI}} + c_1 [-(\Gamma_{\text{HI}} + \beta_{\text{HI}}x_e n_{\text{H}})x_{\text{HI}} + \alpha_{\text{HII}}x_{\text{HII}}x_e n_{\text{H}}], \quad (55)$$

$$0 = x_{\text{HeI}}^{\text{up}} - x_{\text{HeI}} + c_1 [-(\Gamma_{\text{HeI}} + \beta_{\text{HeI}}x_e n_{\text{H}})x_{\text{HeI}} + \alpha_{\text{HeII}}x_{\text{HeII}}x_e n_{\text{H}}], \quad (56)$$

$$0 = x_{\text{HeIII}}^{\text{up}} - x_{\text{HeIII}} + c_1 [(\Gamma_{\text{HeII}} + \beta_{\text{HeII}}x_e n_{\text{H}})x_{\text{HeII}} - \alpha_{\text{HeIII}}x_{\text{HeIII}}x_e n_{\text{H}}], \quad (57)$$

with $x_e = x_{\text{HII}} + (\text{He}/\text{H})_{\text{local}}(x_{\text{HeII}} + 2x_{\text{HeIII}}) + x_e^{\text{metal}}$, and the same $\text{He} \leftrightarrow \text{H}$ charge-exchange terms on rows 1 and 2. The helium ratio is the global one on the legacy path and the *local*, radius-dependent one when the element diffusion is on, since the global value would misstate n_e by the separation factor. These are the systems of the `_adv` post-process (Sec. 10.1).

Transported protons. With `Ionization transport` (default off) H^+ is carried as a fifth transported species on the hydrodynamic face mass flux inside the Runge-Kutta stages (98), in the same way the molecular carriers are (Sec. 6.5), and its local balance is then a constraint rather than a closure.

4.6 Excited hydrogen and the $\text{Ly}\alpha$ field

`excited_hydrogen.f90` solves the $2s/2p$ statistical equilibrium of Christie et al. (2013, their eqs. (12)–(13)) as a 2×2 linear system,

$$\begin{pmatrix} L_{2p} & -M_{12} \\ -M_{21} & L_{2s} \end{pmatrix} \begin{pmatrix} n_{2p} \\ n_{2s} \end{pmatrix} = \begin{pmatrix} S_{2p} \\ S_{2s} \end{pmatrix}, \quad n_{2p} = \frac{S_{2p}L_{2s} + M_{12}S_{2s}}{L_{2p}L_{2s} - M_{12}M_{21}}, \quad (58)$$

whose sources are the recombination cascade, collisional excitation from $1s$ and radiative pumping by the $\text{Ly}\alpha$ mean intensity, and whose sinks are radiative decay, the two-photon continuum, collisional de-excitation, ℓ -mixing and photoionization by the stellar Balmer continuum. The atomic data are in one place (`hydrogen_n2_rates.f90`): $A_{2p1s} = 6.2649 \times 10^8 \text{ s}^{-1}$ (NIST ASD, the nonrelativistic dipole value carrying the reduced-mass factor), $A_{2s1s} = 8.26 \text{ s}^{-1}$, $g_{1s} = g_{2s} = 2$, $g_{2p} = 6$, $f_{\text{Ly}\alpha} = 0.4162$, the Christie et al. Table 2 collisional rates such as $q_{1s2p} = 1.71 \times 10^{-8} t_4^{-0.077} e^{-118400/T}$, their detailed-balance inverses, and the Draine (2011) case-B recombination and its $2s/2p$ split,

$$\alpha_B = 2.54 \times 10^{-13} t_4^{-0.8163-0.0208 \ln t_4}, \quad \alpha_{2s} = (0.282 + 0.047 t_4 - 0.006 t_4^2) \alpha_B, \quad (59)$$

with $t_4 = T/10^4 \text{ K}$. The removal of a $2p$ atom without emission of a line photon is

$$D_{2p} = n_e q_{2p1s} + \Gamma_{2p} + n_e C_{2p2s} \frac{A_{2s1s}}{A_{2s1s} + n_e (q_{2s1s} + C_{2s2p}) + \Gamma_{2s}}. \quad (60)$$

Balmer continuum. The $n = 2$ photoionization rate and the photoelectric heating it delivers are integrals over $[E_{\text{th}}(\text{H}, n=2), E_{\text{th}}(\text{H})]$ of the run's own spectrum with the hydrogenic cross section $\sigma_2(E) = \sigma_2^{\text{thr}} (E_2/E)^3$, $\sigma_2^{\text{thr}} = 1.4 \times 10^{-17} \text{ cm}^2$:

$$\Gamma_2 = \frac{\sigma_2^{\text{thr}} E_2^3}{h \nu_{\text{eV}}} \int F_E E^{-4} dE, \quad h_2^{(1)} = \sigma_2^{\text{thr}} E_2^3 \int F_E (E^{-3} - E_2 E^{-4}) dE. \quad (61)$$

The quadrature resolves the field it integrates, which a fixed rule cannot: for a loaded table the nodes are the table's *own* rows, and each segment is integrated in closed form because the interpolation is a power law there; for the power law the whole band is closed form; only for the smooth Planck field is a composite five-point Gauss-Legendre rule on a geometric subdivision used. The feedback is lagged-explicit: it is computed once per timestep before the ionization and energy solve, and the frozen arrays are read inside `ioniz_eq` and never inside its Newton iteration. The whole module is gated by `use_excited_H`, which is armed by supplying `Stellar Teff [K]`: and `Stellar radius [R_sun]`; it is *off* by default.

The $\text{Ly}\alpha$ field. `lya_rt.f90` closes the line with an escape probability rather than a Monte Carlo, because the columns involved reach $\tau_0 \sim 10^8$ at line center. With $\Delta\nu_D = \nu_{\text{Ly}\alpha} \sqrt{2kT/m_{\text{H}}}/c$, the Voigt parameter $a = (A_{2p1s}/4\pi)/\Delta\nu_D$ and the line-center depth accumulated cell by cell from $n_{\text{HI}} C_{\text{Ly}\alpha}/\Delta\nu_D$, the escape probability of a photon emitted at the cell center is

$$\beta = \frac{1}{1 + \langle N \rangle}, \quad \langle N \rangle = \frac{4\sqrt{6}}{\pi^2} \frac{\tau_{\text{up}} + \tau_{\text{dn}}}{2} \Psi(\xi), \quad \xi = \frac{\tau_{\text{dn}}}{\tau_{\text{up}} + \tau_{\text{dn}}}, \quad \Psi(\xi) = \sum_{n \text{ odd}} \frac{\sin(n\pi\xi)}{n^2}, \quad (62)$$

which is Neufeld (1990) eq. (3.27) at zero continuum destruction, reducing for a mid-plane source to Harrington (1973) eq. (40), $\langle N \rangle = 0.909316B$. The form $1/(1 + \langle N \rangle)$ and not $1/\langle N \rangle$ is used because $\langle N \rangle$ counts absorptions per photon generated, so a photon is emitted $1 + \langle N \rangle$ times before it leaves; this is exact at both ends and needs no join between the thin and thick limits. Under $R_{\text{II}}\text{-A}$ redistribution $\langle N \rangle$ is independent of a , so the Voigt parameter survives only in the one-flight penetration of the stellar beam. The trapped internal field and the stellar term are

$$J_{\text{int}} = \frac{2h\nu^3}{c^2} \frac{g_{1s}}{g_{2p}} \frac{P(1-\beta)}{(A_{2p1s}\beta + D_{2p}) n_{\text{HI}}}, \quad J_{\star} = \frac{\xi_{\text{day}} F_{\text{Ly}\alpha}}{4\pi \Delta\nu_{\star}} \langle T_s E \rangle, \quad J_{\text{Ly}\alpha} = J_{\text{int}} + J_{\star}, \quad (63)$$

with $P = \alpha_B n_e n_{\text{HII}} + C_{1s2p} n_e n_{\text{HI}}$ the internal production, $T_s = \text{erfc}(x_1/\sqrt{2}X_s)$ the fraction of the broad stellar profile whose wings reach depth τ , and $\langle \cdot \rangle$ the mean of the beam over the cell's own optical depth. The published solution

requires $(a\tau)^{1/3} > 10$ (Neufeld section Va); cells outside that are counted by `lya_wing_domain_record` and are carried there by the thin limit $\beta \rightarrow 1$, which (62) reaches smoothly. The alternative `jlya_mode = 1` reads an externally computed $J_{\text{Ly}\alpha}(r)$, for instance from the LaRT Monte Carlo code, and interpolates it onto the grid.



5. The thermal source: heating and cooling

5.1 The heating channels

The volumetric heating rate of a cell is a sum of physically distinct deposits, and there is exactly one list of them (`util_ion_eq.f90`): `heating_of_composition` fills every column of `heat_chan`, the total it returns is the sum of those columns in that order, and the header of `output/Heating_breakdown.txt` is built from the same names, so a deposit cannot exist in the energy equation and be missing from the breakdown.

#	name	what it is
1–6	<code>heat_HI . . . heat_metals</code>	photoionization of each absorber, $n_a h_a^{(1)}$ of (33)–(34)
7	<code>heat_Hpe[excitedH]</code>	photoelectric heating of $\text{H}(n=2)$ by the Balmer continuum, (61)
8	<code>heat_Hdx[Lyα-deexc]</code>	collisional de-excitation of the Ly α -pumped $n=2$ atoms (<code>incl_deexc_heat</code> , default on)
9	<code>heat_He_recomb</code>	photoelectrons of the H I , H_2 and metal ionizations driven by helium recombination radiation
10–12	<code>heat_He23S_*</code>	$\text{He}(2^3\text{S})+\text{H}$ Penning and associative branches, and $\text{He}(2^3\text{S})+\text{H}_2$ Penning
13–14	<code>heat_H2_LW_*</code>	kinetic energy of the Lyman-Werner dissociation fragments, and collisional de-excitation of the H_2 fluorescence
15	<code>heat_mol_chem</code>	net chemical heat of the collisional reactions of the H_2/He network
16–19	<code>heat_FUV_photolysis,</code> <code>heat_oxygen_collisional,</code> <code>heat_CO_Hep_transfer,</code> <code>heat_CO_photodissoc</code>	the oxygen option's deposits

Channels 1–6 are the photoheating integrals of Sec. 3.4, already carrying the heating efficiency f_{heat} of (39). Channel 15 exists because the return path that makes ionization energy a non-channel in an atomic gas is different in a molecular one (`molecular_reaction_heat.f90`). Follow one closed cycle:



The cycle returns to H_2 having turned one H_2 into $H+H$ and having left $I(H_2) - D_0(H_2) = 15.43 - 4.48 = 10.95$ eV in the gas, with no photon anywhere in it to carry that away. Measured on the converged $He/H = 0.0793$ hot Uranus, that is 81 percent of the total heating rate at $1.02 r_{\text{base}}$. Molecular reaction heat is therefore *on* by default; it is unreachable in an atomic run, so no non-molecular result moves.

5.2 The cooling channels

`eval_cool` (`util_ion_eq.f90`) assembles the cooling rate of every cell as

$$\Lambda = n_e (\Lambda_{\text{ff}} + \Lambda_{\text{cx}} + \Lambda_{\text{rec}} + \Lambda_{\text{ci}}) + \Lambda_{\text{metal}} + \Lambda_{H_3^+} + \Lambda_{H_2} + \Lambda_{H_2O} + \Lambda_{CO}, \quad (64)$$

in $\text{erg cm}^{-3} \text{ s}^{-1}$, with n_e the same charge sum the ionization balance uses (`calc_ne`), molecular donors included, since inside a deep molecular base H_3^+ is the dominant ion. The breakdown written to `output/Cooling_breakdown.txt` is read straight off the arrays (64) is built from, so the columns sum to Λ exactly.

Recombination. $\Lambda_{\text{rec}} = \lambda_{\text{HII}} n_{\text{HII}} + \lambda_{\text{HeII}} n_{\text{HeII}} + \lambda_{\text{HeIII}} n_{\text{HeIII}}$ with, for hydrogen (Hui & Gnedin, 1997),

$$\lambda_{\text{HII}}(T) = 3.435 \times 10^{-30} T \frac{x^{1.970}}{[1 + (x/2.250)^{0.376}]^{3.720}}, \quad x = \frac{2 \times 157807}{T}. \quad (65)$$

The helium terms are kT times the *same* recombination coefficient the ionization balance runs on, so electrons are not removed at one rate and the gas charged at another.

Collisional ionization. Λ_{ci} charges each ionization its measured potential in erg,

$$\Lambda_{\text{ci}} = E_{\text{th}}^{\text{HI}} \beta_{\text{HI}} n_{\text{HI}} + E_{\text{th}}^{\text{HeI}} \beta_{\text{HeI}} n_{1^1\text{S}} + E_{\text{th}}^{\text{HeII}} \beta_{\text{HeII}} n_{\text{HeII}} + E_{\text{th}}^{2^3\text{S}} \beta_{2^3\text{S}} n_{2^3\text{S}}, \quad (66)$$

the He I term acting on the ground *singlet* $n(1^1\text{S}) = n(\text{HeI}) - n(2^3\text{S})$, because the metastable sits 19.8 eV up and carries its own 4.8 eV threshold in the last term. The same subtraction is made in the photoheating and in the collisional excitation below, so the metastable is never charged the ground-state coefficients as well.

Collisional excitation. Λ_{cx} is the sum of three fits,

$$\lambda_{\text{HI}} = \frac{7.5 \times 10^{-19}}{1 + \sqrt{T/10^5}} e^{-118348/T}, \quad \lambda_{\text{HeI}} = 1.1 \times 10^{-19} T^{0.082} e^{-2.3 \times 10^5/T}, \quad \lambda_{\text{HeII}} = \frac{5.54 \times 10^{-17} T^{-0.397}}{1 + \sqrt{T/10^5}} e^{-473638/T}, \quad (67)$$

contracted with n_{HI} , $n(1^1\text{S})$ and n_{HeII} . The first of these is the $\text{Ly}\alpha$ -dominated hydrogen line cooling, and it is the optically thin fit as written: *no escape probability multiplies the $\text{Ly}\alpha$ cooling channel*. λ_{HI} is `lambda_coex_HI` of `Cool_coeff.f90` and `eval_cool` contracts it with n_{HI} and n_e with no such factor anywhere. The two escape probabilities the code does compute enter elsewhere: β of `lya_rt.f90` (62) enters the $H(n=2)$ populations and the pumping field, and β_{ul} of `fine_structure_line_transfer` enters the metal ground-term statistical equilibrium (68); neither is read by any cooling coefficient. The asymmetry is deliberate: multiplying λ_{HI} by an escape probability would leave the trapped fraction with no owner while keeping the de-excitation heating channel 8 as an independent source, that is it would double count. A ledger carrying both is the stated target and is not built.

With the metastable tracked, two further terms are added to Λ_{cx} , both scaling with the explicitly computed $n(2^3\text{S})$: the $2^3\text{S} \rightarrow 2^3\text{P}$ excitation followed by 10830 Å decay and escape, and the $2^3\text{S} \rightarrow 2^1\text{S}$ (0.80 eV) and $2^3\text{S} \rightarrow 2^1\text{P}$ (1.40 eV) collisional conversions, each removing its threshold energy from the electron gas. The ground-to-triplet excitation q_{13} is deliberately excluded here, being already inside λ_{HeI} .

Free-free. $\Lambda_{\text{ff}} = 1.426 \times 10^{-27} \sqrt{T} \sum_{\text{ions}} Z_{\text{ion}}^2 \bar{g}(T, Z_{\text{ion}}) n_{\text{ion}}$, with $\bar{g}$ the thermally averaged Gaunt factor of van Hoof et al. (2014), interpolated linearly in $x = \log_{10}(Z^2 157807/T)$ and clamped to the tabulated range. Only $Z_{\text{ion}} = 1$ and 2 occur. The molecular ions H_2^+ , H_3^+ and HeH^+ do donate to n_e but are deliberately left out of this charge sum: they exist only in the $T \sim 10^3$ K molecular base, where this hot-plasma expression is an extrapolation whose emission comes out at frequencies the atmosphere is not thin to, and where H_3^+ and metal lines dominate by far.

Metal lines. $\Lambda_{\text{metal}} = n_e \sum_i n_{\text{m},i} c_i(T)$ with c_i a closed-form analytic fit to the CHIANTI v11.0.2 optically thin line-cooling curve of that ion. The C/N/O fits are the default (`cno_chianti = .true.;` `cno_cool 0` in `metals.inp` restores the legacy set ported from AIOLOS), and Mg I/II, Ca II, Na I and Fe I/II follow the same route. Two entries are not in CHIANTI v11 and are built otherwise: the Na I D doublet with a $\bar{g}$ factor 0.2 calibrated against the CHIANTI Mg II and Ca II values, and the Fe I permitted multiplets.

Two density-dependent replacements are made on top of those coronal curves, because a coronal fit assumes $\Lambda \propto n_e$ and no level saturation while the base of an irradiated atmosphere runs at $n_e \sim 10^9$, $n_{\text{HI}} \sim 10^{14} \text{ cm}^{-3}$, decades above the critical densities of the lines involved.

- *Fe II* takes a two-dimensional statistical-equilibrium table $\Lambda_{\text{eff}}(T, n_e) = (\sum_u n_u A_{ul} \Delta E_{ul}) / n_e$, whose $1/n_e$ cancels the n_e of the assembly and so leaves the correct saturated emission; at low n_e it reduces to the coronal rate. Without it the forbidden $a^6\text{D}$ lines overestimate the base cooling by about four decades.
- *Ground-term fine structure* of C I ([C I] 609.1, 370.4 μm), C II (157.7 μm), N II (205.3, 121.8 μm) and O I (63.2, 145.5, 44.1 μm) is replaced by the exact statistical equilibrium of the ground term,

$$\sum_j f_j R_{ji} = f_i \sum_j R_{ij}, \quad \sum_i f_i = 1, \quad R_{ul} = C_{ul} + \beta_{ul} A_{ul}, \quad R_{lu} = C_{ul} \frac{g_u}{g_l} e^{-E_{ul}/kT}, \quad (68)$$

with $C_{ul} = n_e k_{e,ul}(T) + n_{\text{HI}} k_{\text{H},ul}(T)$ and $k_{e,ul} = 8.629 \times 10^{-6} \Upsilon / (g_u \sqrt{T})$, the effective collision strengths Υ being quartic fits in $x = \log_{10}(T/10^4)$ to the CHIANTI .scups values over 10^3 – 10^5 K (largest error 4.8 percent) and the H-atom rates taken from Yan & Babb (2023) for C I and N II, Abrahamsson et al. (2007) for O I and Barinovs et al. (2005) for C II. The emitted power per ion is $W_{\text{FS}} = \sum_{u>l} f_u \beta_{ul} A_{ul} k_B E_{ul}$ and the coefficient returned is $\Lambda_{\text{eff}} = W_{\text{FS}}/n_e + \Lambda_{\text{rem}}$, Λ_{rem} being a refit of the coronal curve keeping only the channels that leave the ground term.

The line trapping β_{ul} of these fine-structure lines is the line-center escape probability formed from the columns above and below the cell (`fine_structure_line_transfer`), and it enters *inside* the statistical equilibrium as $A_{ul} \rightarrow \beta_{ul} A_{ul}$ rather than as a factor on the result: in the subcritical limit the cooling is set by the collisional excitation rate and is independent of β , and in the saturated limit the escaping flux is proportional to it. $\beta = 1$ reproduces the optically thin result exactly, and that is the value every other metal ion keeps. With `Base IR field: True` (default off) the same

lines also *absorb* the thermal infrared of the lower atmosphere, a blackbody at T_0 covering the sky fraction the base subtends, through the photon occupation number $\bar{n}_{ul}$; this gives the layer a radiative equilibrium floor instead of letting it cool into vacuum.

Molecular infrared. $\Lambda_{H_3^+}$ is the Miller et al. (2013) emission of one H_3^+ molecule in LTE,

$$\Lambda_{H_3^+} = 4\pi n_{H_3^+} E(T) s(T, n_{H_2}), \quad \ln E(T) = \sum_n C_n T^n \text{ [W molecule}^{-1} \text{ sr}^{-1}], \quad (69)$$

their Table 5 (the “using $z(t)$ ” column) with the non-LTE departure factor s of their Table 6, whose only collider is H_2 through the proton-hopping reaction with the Oka & Epp (2004) coefficient $2 \times 10^{-15} \text{ m}^3 \text{ s}^{-1}$. Table 6 covers $n(H_2) = 10^6\text{--}10^{14} \text{ cm}^{-3}$; below that the code adopts the linear low-collider limit $s(T, n) = s(T, 10^6)n/10^6$ and records the cell as evaluated outside the tabulated range. This is the dominant coolant of a molecular base: the atomic channels above are all exponentially suppressed at $T \sim 10^3 \text{ K}$, so leaving it out leaves that gas with no radiative loss at all.

With Molecular IR bands: True (default off) three further net exchanges are added (molecular_infrared_cooling.f90): the H_2 quadrupole and magnetic-dipole line spectrum, and the H_2O and CO vibration-rotation bands. Each is a *net* rate,

$$\Lambda_X^{\text{net}} = n_X [E_X(T) - W_{\text{dil}} E_X^{\text{abs}}(T; T_{\text{rad}})], \quad E_X(T) = 4\pi \sum_b \sigma_b(T) \int_b B_\nu(T) d\nu, \quad (70)$$

so each vanishes at its own radiative equilibrium temperature instead of running the layer down to nothing; a column is negative wherever that channel heats. The dilution W_{dil} is the same $0.5 f_{\text{sky}}$ the H_3^+ closure uses and is zero when Base IR field is off, which reduces these channels to pure emitters. Without them a converged molecular layer radiates itself down to 190–400 K against an equilibrium temperature of 1100–1400 K.

5.3 Advancing the temperature

In the marching loop. The temperature of a cell is advanced by `solve_energy_semi_implicit` (energy_semi_implicit.f90), the inner step of the coupled temperature-composition source step: the marching loop alternates it with the composition sweep until the pair stops moving, so the composition is fixed for one pass and not for the whole step. With the composition and the photoheating frozen over the step,

$$\rho e(T) - \rho e_{\text{old}} = \Delta t [\mathcal{H} - \Lambda(T)], \quad \text{i.e.} \quad R(T) = \varepsilon(T) - \varepsilon_{\text{old}} - a[\mathcal{H} - \Lambda(T)] = 0, \quad a = \frac{\Delta t}{n_{\text{tot}} + n_e}, \quad (71)$$

with $\varepsilon = \rho e / (n_{\text{tot}} + n_e)$ the internal energy per particle, which is what `internal_energy_per_particle` returns. The caloric internal energy ρe is that of (14). The root is found by a safeguarded Newton-bisection hybrid on a physical bracket whose ends are the floor of the equation-of-state validity and a ceiling above the heating-only update: the search starts from the explicit-Euler point, brackets the root as it goes, and once both ends are known takes a Newton step whenever that step stays strictly inside the bracket and the bracket keeps halving, bisection otherwise. The Newton derivative is exact in the equation-of-state term (`heat_capacity_per_particle`) and a secant estimate in $d\Lambda/dT$, which affects only the path, the safeguard owning the convergence.

What is tested before the update is returned is the residual at the *returned* temperature against $|R|/s$ with $s = |\rho e_{\text{old}}| + \Delta t(|\mathcal{H}| + |\Lambda|)$, and admissibility; the cooling written back is $\Lambda(T_{\text{returned}})$, evaluated there and not at a previous iterate.

There is no clamp: a cell that does not meet the tolerance, that has no bracket, or that reaches the equation-of-state floor is a named failure, nothing is assembled from it, and the step controller retakes the step at half Δt as in Sec. 7.5. The floor is not a reservoir: reaching it means the balance asks for a temperature below the range in which the equation of state and the rate fits are defined, and a floor representing real external heating would have to be a specified reservoir with its own budget. `use_semi_implicit_energy` is on by default.

In the advection post-process. `T_equation.f90` is *not* the marching loop's temperature solve. Its module `equation_T` is used by `post_process_adv.f90` (Sec. 10.1) and by test drivers, and by nothing else; what it solves is the *steady* internal-energy balance of the profile, upwind differenced,

$$\nabla \cdot (u\mathbf{v}) + p \nabla \cdot \mathbf{v} = \rho v \frac{de}{dr} - p v \frac{d \ln \rho}{dr} + h \nabla \cdot (\rho \mathbf{v}) = \mathcal{H} - \Lambda, \quad (72)$$

with e the internal energy per unit mass and $h = e + p/\rho$ the enthalpy per unit mass. The upstream term carries the *upstream* cell's specific internal energy formed at the upstream composition and temperature, not this cell's caloric state evaluated at T_{up} : the rovibrational heat capacity belongs to the gas that holds the molecules, so across a dissociation front two cells at the same temperature store different energy. The root is bracketed rather than guessed (`solve_T_brent`, on by default through `use_brent_tsolve`): 80 logarithmically spaced probes from $\max(0.05T_{\text{guess}}, 1 \text{ K})$ to $4T_{\text{guess}}$ locate the *first* sign change, i.e. the lowest root, and Brent's method closes the bracket to $2\epsilon|b| + \text{tol}/2$ with $\text{tol} = 10^{-10}$ in T/T_0 , which is what keeps the solve off the hot spurious root; `hybrd1` on the same residual is the fallback. The metal densities and the fine-structure escape probabilities are held fixed while the root finder varies T , the latter because the optical depth is a column over the whole atmosphere and not a function of this cell's trial temperature.

The heating efficiency. The diagnostic η_{heat} of (35), written as column `eta` by `write_output.f90`, is a property of the composition the sweep returned and is not a free parameter anywhere in the code; the closed-form parameterizations of the same quantity that the literature uses are reproduced offline by `eta_approx.py` for comparison only.

6. The molecular layer, the trace metals and element transport

Everything in this section is *off* by default. A run acquires the molecular layer by asking for it (`Molecular chemistry: True`, which also switches `Molecular` base on so that the base particle count and the chemistry describe one gas), the metals by the presence of `metals.inp` or of elemental abundances in a handoff file, and the element transport by `He_diffusion`. An atomic, metal-free run never enters any of the code described here.

6.1 The H₂/He reaction network

`mol_rates.f90` carries the reaction rate coefficients of Koskinen et al. (2022) Table 1 as published, in cgs ($\text{cm}^3 \text{s}^{-1}$, and $\text{cm}^6 \text{s}^{-1}$ for the two three-body entries R13 and R15, which the code multiplies by the total heavy-particle density). EXHALE carries one temperature, so the heavy-particle and the electron temperature of that table are the same variable. The photorates P1–P5 of the paper are marked “SC” there, computed from cross sections, stellar flux and columns; in EXHALE they are the integrals of Sec. 3.4 and are not part of this module.

One entry is deliberately *not* transcribed. R12, the thermal dissociation of H_2 , is built by detailed balance from the R15 recombination,

$$k_{12}(T) = \frac{k_{3b}(T)}{K_{\text{eq}}(T)}, \quad (73)$$

with K_{eq} the $\text{H} + \text{H} \rightleftharpoons \text{H}_2$ equilibrium constant built from the same internal partition function $Q_{\text{H}_2}(T)$ the caloric equation of state uses (Sec. 6.4), because the two Table-1 entries are the same channel in opposite directions and their source measurements do not overlap in temperature. The third body of R12, R13 and R15 is the total gas-particle density excluding electrons, so every heavy particle is an equally efficient third body; a monatomic third body has no internal modes to take up the released energy, so these three rates with He as M are upper bounds. Two further limits in a helium-dominated gas, the missing $\text{H}_2^+ + \text{He} \rightarrow \text{HeH}^+ + \text{H}$ channel and the R16–R20 disagreements with García Muñoz (2025), are recorded at the code site rather than patched, since fixing either would replace a Table-1 entry with a rate from another compilation.

The eight balance rows the network makes are (47)–(50) together with the H^+ , He^+ , He^{2+} and $\text{He } 2^3\text{S}$ rows of Sec. 4.2.

6.2 H_2 photoabsorption resolved into channels

`h2_photo_channels.f90` splits the one measured cross section $\sigma_{\text{H}_2}(E)$ into four mutually exclusive final states whose sum is identically σ_{H_2} , so no consumer of the opacity changes and the total H_2 destruction rate is what it was; what changes is which fragments the source terms are told to make.

$$\sigma_N = f_n(E) \sigma_{\text{H}_2}(E) \quad \text{H}_2 + h\nu \rightarrow \text{H} + \text{H} \text{ (33–41 eV window)}, \quad (74)$$

$$\sigma_M + \sigma_S + \sigma_D = [1 - f_n(E)] \sigma_{\text{H}_2}(E) \quad \text{the three ionizing channels}, \quad (75)$$

with f_n the neutral fraction of Chung et al. (1993) Table 1, carried as a *fraction* so that it rides on this code's σ_{H_2} normalization rather than importing theirs. Inside the ionizing part the split follows the Chung et al. Table II ratio column $r(E) = \sigma(\text{H}^+)/\sigma(\text{H}_2^+)$, a ratio of time-of-flight detector signals and therefore a ratio of *events*, one count per event and not one per proton; `frac_H2_dissociative_ionization` is that column converted to $\sigma(\text{H}^+)/[\sigma(\text{H}^+) + \sigma(\text{H}_2^+)]$ and tabulated at 69 energies from 18.076 to 124 eV, linearly interpolated, zero at and below threshold and held flat above 124 eV. The thresholds are 15.4 eV (M, leaves H_2^+), 18.076 eV (S, leaves $\text{H} + \text{H}^+$), 51.4 eV (D, two protons), and none of its own for N. With the default `h2_double_ionization = 'off'` only the S channel is resolved separately.

The energy of one absorbed photon is accounted channel by channel by `h2_channel_energy_recipients`, the single statement of what each channel charges, and the recipients of each sum to $h\nu$ once. Two entries are fixed with no free parameter: the double channel's vertical threshold 51.4 eV (Yan et al., 1998) exceeds the asymptotic product energy $D_0 + 2I(\text{H}) = 31.675$ eV by exactly the kinetic energy release of the two receding protons, 19.725 eV; and the neutral window leaves both fragments in $n = 2$ (the $\text{Q2 } ^1\Pi_u$ state whose fluorescence curve overlaps this cross section), so $2E(n=1 \rightarrow 2)$ departs as prompt $\text{Ly}\alpha$ and two-photon continuum and is not heat.

6.3 Lyman-Werner photodissociation

`lyman_werner.f90` adds the Solomon process, $\text{H}_2(X) + h\nu(11.2\text{--}13.6\text{ eV}) \rightarrow \text{H}_2(B, C) \rightarrow \text{H} + \text{H}$, which the Koskinen et al. Table-1 network leaves out (their photorates start at the 15.4 eV edge). The band is 912–1201 Å: 912 Å is the

H Lyman edge, and 1201 Å is both the blue edge of band B2 and the red end of the Lyman and Werner line list the shielding table is built from. Draine & Bertoldi's own 912–1110 Å is not used, because a planetary base runs at 700–3200 K where the vibrationally excited levels pump in lines longward of 1110 Å that carry about a third of the dissociations at saturation. One band, one line list, one normalization.

The optically thin rate is formed from the band-integrated *energy* flux at the planet and the mean photon energy of a flat F_λ band, $\langle h\nu \rangle = 2hc/(912 + 1201 \text{ Å}) = 11.7354 \text{ eV}$:

$$k_{\text{LW}}^{\text{thin}} = \sigma_{\text{LW}} \frac{\xi_{\text{day}} F_{\text{LW}}}{\langle h\nu \rangle}. \quad (76)$$

The band-mean σ_{LW} is not the effective cross section $2.263 \times 10^{-18} \text{ cm}^2$ that Draine & Bertoldi (1996) Table 1 and 2 imply for a flat- F_λ field restated per 912–1201 Å photon: the rate is the level-resolved table of `h2_self_shielding_table.f90`, normalized per photon of the same band, and that number is only the limit it approaches in optically thin interstellar conditions.

That table is a line-by-line calculation with *overlapping* lines, and it stores three different counts because three are needed: σ_{diss} , dissociations per unit incident band photon fluence, which carries the self-shielding, the line overlap and the trapping of the fluorescent photons and sets the rate and the fragment heating; $p_{\text{single}} = D/(D + \sum A_{ul})$, dissociations per pump with every fluorescent photon free to escape, which sets the vibrational heat return because a trapped decay plus its re-absorption moves one molecule down and another up and deposits nothing; and $p_{\text{eff}} = D/(D + \sum A_{ul} P_{\text{loss}})$, dissociations per photon removed from the beam, which is the band photon ledger. They are related by $\sigma_{\text{diss}} = p_{\text{eff}} \sigma_{\text{pump}}$, and the beam's own loss is served from the same table, so the photons the beam loses and the dissociations the rate spends are one absorption with one normalization. Line overlap is not a correction at these columns: at $n_{\text{H}} = 10^{13} \text{ cm}^{-3}$ the surviving pumping relative to a one-line-at-a-time calculation is 0.45 at $N_{\text{H}_2} = 10^{20} \text{ cm}^{-2}$ and 900 K, and 4.8×10^{-5} at $4.4 \times 10^{21} \text{ cm}^{-2}$ and 2700 K, which is where the base of a molecular layer sits. The rate enters (47) as g_{LW} and is zero when no band flux is supplied.

6.4 The caloric equation of state

The caloric equation of state that gives the bound H_2 its rovibrational heat capacity is (14) of Sec. 2.5, where it is stated with its validity and its Caloric EOS: `monatomic_switch.caloric_eos.f90` supplies the four maps the hydrodynamics needs, $(\rho, \rho e) \rightarrow T$, $(\rho, \rho e) \rightarrow p$, $(\rho, p) \rightarrow \rho e$ and $(\rho, p) \rightarrow \gamma_{\text{eff}}$, and γ_{eff} itself for the wave speeds.

6.5 Carrier transport

Everywhere else in the code the composition of a cell is the root of a local algebraic system. `diffusive_photochemistry.f90` solves, instead, the continuity equation of the neutral carriers H_2 , OH , H_2O and CO ,

$$\frac{\partial n_i}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} [r^2 (n_i v + J_i)] = P_i - L_i, \quad J_i = -n_{\text{tot}} (D_i + K_{zz}) \frac{\partial f_i}{\partial r} - n_i D_i \left(\frac{1}{H_i} - \frac{1}{H_{\text{atm}}} \right), \quad (77)$$

with $f_i = n_i/n_{\text{tot}}$. Between them these four carry every H nucleus that is not atomic or in a molecular ion and every O and C nucleus that is not a free atom. H_2^+ , H_3^+ , HeH^+ and $\text{O}(^1\text{D})$ are left local; on the HD 209458 b molecular example the three molecular ions together carry at most 1.2×10^{-9} of the H nuclei, and the H-nucleus budget is closed against

the element total rather than against a sum of transported species, so whatever they hold is simply not available to the carriers. All four transported species are neutral, so the ambipolar field term of the general drift-diffusion equation is absent rather than neglected.

D_i is Blanc's law over the background carriers (H I, H II, H₂, H₂⁺, H₃⁺, He I, He II, He III, HeH⁺), each pair from the rigid-sphere coefficient of Banks & Kockarts (1973). The ion-neutral polarization enhancement is not added: the transported species only exist where the gas is neutral (electron fraction per H nucleus at most 10⁻⁷ over the cells that hold H₂), so the ion pairs carry that share of the Blanc friction sum, and for H₂O the induced-dipole formula would be the wrong one anyway, a permanent dipole of 1.85 D dominating the capture rate. K_{zz} is read from `kzz_cell` and nowhere else; that array is filled once, from the scalar `He_Kzz` or from the lower-atmosphere profile's own K_{zz} column. Settling is $G_i = (m_i - \bar{m})g/(kT)$ with $\bar{m} = \rho/n_{\text{tot}}$ and g from the same potential the hydrodynamics uses; thermal diffusion is not carried ($\alpha_T = 0$), there being no evaluated α_T for H₂O, OH or CO in an H₂/He bath.


$P_i - L_i$ is not written in this module: it is obtained by calling the very rows the local equilibrium solve uses, `mol_heh_rows` row 4 for H₂ and `oxygen_carrier_rows` for OH and H₂O, at the trial densities with the background frozen, and the Jacobian is a forward difference of those same calls, so a rate that changes there changes here. CO has no chemical source and only its transport moves it, with a ceiling from `limit_to_element_budget` that keeps it from carrying the oxygen and the carbon out of the wind, where the molecule cannot survive and where the O I and C II lines are the coolant.

The discretization is cell-centered f_i , face-centered fluxes on `r_edg`, one backward-Euler step per call, Newton with a backtracking line search, block tridiagonal with 4×4 blocks. Faces 0 and N carry zero diffusive flux; the drift term switches from central to donor cell on a Péclet test. The element reservoirs at the base are Dirichlet, imposed through `melem_ab`, the base density and the equation of state, but the *partition* of each element among its carriers is not imposed anywhere, which is what makes the comparison against a photochemical handoff a test rather than a tautology.

The material advection of a carrier is *not* in these rows. It is the divergence of the hydrodynamic face mass flux, $F_\rho(j)Y_c(j)/m_c$, taken with the mass row itself inside the Runge-Kutta stages (98) on the same faces, areas, volumes and time step (`species_face_flux.f90`), so a cell can only lose the carrier the mass row says it loses and a uniform partition is preserved to round-off. That this half must ride on the *face* mass flux is a numerical statement about the front: a first-order donor-cell difference on the cell velocity carries a numerical diffusivity $|v|\Delta r/2$, which on the 500-cell hot Uranus grid is 8.5, 20 and 22 times the molecular D of H₂ at 1.15, 1.30 and 1.50 R_p . The key is Molecular carrier transport, default off.

6.6 The oxygen option

With `thereis_oxychem` (default off) three neutral carriers OH, H₂O and CO join the network. `oxygen_rates.f90` carries the rate coefficients, each transcribed from a named publication and each checked against two independent ports of the same literature that are on this machine (the `Photochem_zahnle_earth.yaml` network and the `VULCAN_NCHO_photo_network.txt`), every disagreement being recorded at the coefficient. `water_photolysis.f90` supplies the photolysis,



in the four far-ultraviolet bands of Sec. 3.1, whose edges are fixed by the branching ratios rather than chosen: the three-body branch opens only in the interval containing $\text{Ly}\alpha$ and is 12 percent there, so $\text{Ly}\alpha$ needs its own band on branching grounds alone. The first branch alone carries 54.5 percent of the H_2O loss at the HD 189733 b base, which makes the OH/ H_2O cycle catalytic rather than a one-way sink. `co_photodissociation.f90` destroys CO on the same Lyman-Werner beam, using a band-mean dissociation cross section times the band photon flux and the Visser et al. (2009) shielding function $\Theta(N_{\text{CO}}, N_{\text{H}_2})$ rather than their $k_0\Theta$ form, because EXHALE has a band flux and not an interstellar field and k_0 is a statement about a particular field. The other CO destruction channel is the He^+ charge transfer $\text{He}^+ + \text{CO} \rightarrow \text{C}^+ + \text{O} + \text{He}$ at the Langevin value $1.6 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$, which appears in the He^+ row of (48)’s system and whose +2.2117 eV appears in heating channel 18, one event counted once.

6.7 The lower-atmosphere handoff

The base of the escape model sits at a stated pressure level, and three routes supply the state there.

The analytic column. `lower_column.f90` integrates the hypsometric relation from the 1-bar level up to the base pressure (default 1 μbar), isothermal at T_{eq} with spherical gravity, using the mean molecular weight of the chemical-equilibrium $\text{H}_2/\text{H}/\text{He}$ partition of Visscher et al. (2006) as fitted by Koskinen et al. (2022) eqs. (11)–(13),

$$q_{\text{H}_2} = \frac{1.9845 + 10^u - \sqrt{10^u (3.9690 + 10^u)}}{2.3670}, \quad u = -\frac{23672}{T} - \log_{10} p [\text{bar}] + 6.2645, \quad (78)$$

with q_{He} from the elemental He/H and $q_{\text{H}} = 1 - q_{\text{H}_2} - q_{\text{He}}$. It returns the radius of the base level and the base mixing ratios, and also a fully atomic bracket, because in chemical equilibrium $q_{\text{H}_2}(1 \mu\text{bar})$ stays large up to $T \sim 2000 \text{ K}$ while photochemistry dissociates H_2 there: a large q_{H_2} on a hot planet is the signal that a photochemical handoff is needed. The validation case is the Koskinen et al. (2022) Model A Uranus-mass planet at 0.05 au, for which the column gives $r_0(1 \mu\text{bar}) = 1.34 R_{1\text{bar}}$ with $q_{\text{H}_2} \simeq 0.84$.

base.inp, the scalar handoff. An optional file of “key value” lines, written by `src/utls/run_lower.py` or `src/utls/vulcan_to_base.py`. Every key belongs to one of five categories and the category states what the value may do to the wind: *provenance* (comments only); *EOS boundary* ($T_{\text{base}} \rightarrow T_0$, $r_{\text{base}} \rightarrow R_0$, p_{base} , and $q_{\text{H}_2\text{base}}$, the H_2 volume mixing ratio of the base gas); *elemental reservoir* (HeH_{base} and $\langle \text{El} \rangle_{\text{H}_{\text{base}}}$ for the ten elements, overriding `metals.inp` for that element); *initial guess* (no key today); and *boundary constraint* ($K_{\text{zz}_{\text{base}}} \rightarrow \text{he}_{\text{kzz}}$). $q_{\text{H}_2\text{base}}$ is the q_{H_2} of (15) and is imposed on the H_2 partition of the inflowing ghost species as well; it is refused above $0.5/(0.5 + \text{He}/\text{H})$. Species mixing ratios other than $q_{\text{H}_2\text{base}}$ stay comments: no part of the code consumes them, so presenting them as physics would be false. A `base.inp` that states p_{base} fixes the base level and n_0 is derived from it as in Sec. 1.2, instead of being read from `input.inp`.

Lower atmosphere profile, the tabulated handoff. `lower_atmosphere_profile.f90` reads the lower model’s solution over an *interval* of pressure that reaches above the level where EXHALE places its base. It supplies the state at the matching level p_{match} (T , r , q_{H_2} , and the elemental ratios He/H and El/H), which is what the base equation of state and the elemental reservoirs are built from, and K_{zz} as a *profile*, because the homopause, the only consequential

thing K_{zz} does, is where $K_{zz} = D_{12}$, a property of two profiles, and the molecular coefficient rises by nearly five decades between the base and $1.2R_p$. Every intensive quantity is interpolated linearly in $\log p$, the variable both models solve on; extrapolation is never performed. The values go in through exactly the doors the scalar keys use, so nothing in the base equation of state changes; n and ρ carried by the file are *not* imposed, the base density remaining n_0 .

6.8 Trace metals

`metals.inp` (`metals_input_read.f90`) sets the elemental abundances n_X/n_H by number, one ion per line with the neutral label naming the element; a second statement of the same element stops the run, since two abundances for one element are two answers to one question. The same file also carries `cno_cool`, `eos_metals`, `cx_full` and `cx_O2p_H`. The ionization of each element is solved in the same coupled system as hydrogen and helium (`System_HeH_metals.f90` and its triplet and molecular counterparts), through the rows (45); the code carries no parallel decoupled metal solver.

`eos_metals` (default `on`) puts the metals into the bulk gas budget consistently: their mass in ρ and hence in μ , v_0 , p_0 ; their electrons in n_e ; and their nuclei in n_{tot} and in the ghost pressure, matching the charge balance the ionization systems already use. The legacy trace approximation (metals in the ionization and cooling but not in the bulk gas) is `eos_metals 0`. A metals-off run is identical either way, every metal sum vanishing.

6.9 Binary element transport

`binary_element_diffusion.f90` treats the gas as two components moving through each other with the single mass-averaged velocity of the hydrodynamics: component 1 is the hydrogen carriers together with the trace metals and the heavy nuclei bound in the molecular carriers, and component He is helium in all stages. Since $\rho_1 + \rho_{\text{He}} = \rho$ exactly, the two diffusive mass fluxes close, $J_1 = -J_{\text{He}}$, and the composition update creates no mass. The transported variable is the helium *mass* fraction $X = \rho_{\text{He}}/\rho \in [0, 1]$, which is bounded at both ends unlike the ratio n_{He}/n_H that the earlier trace kernel carried and that diverges exactly in the helium-rich limit this operator exists for. The equation is

$$\frac{\partial(\rho X)}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} [r^2 (\rho X v + J)] = 0, \quad J = -\rho(D_{12} + K_{zz}) \frac{\partial X}{\partial r} - \rho D_{12} G X(1-X), \quad (79)$$

$$G = \frac{(m_{\text{He}} - m_{c1})m_{\text{H}}g - (\bar{Z}_{\text{He}} - \bar{Z}_{c1})eE}{kT} + \alpha_T \frac{d \ln T}{dr} - \frac{d \ln \psi}{dr} \quad [\text{cm}^{-1}]. \quad (80)$$

The masses and charges in G are *carrier* quantities: $\psi = n_{1c}/n_H$ is the number of collision partners each hydrogen nucleus is spread over (1 atomic, 1/2 fully molecular) and $m_{c1} = m_1/\psi$. The gradient coefficient collapses to ρD_{12} for any ψ , so the Fickian term is the same in the molecular and the atomic region; the piece of the mole-fraction gradient the chemistry carries does not cancel and rides in G as $-d \ln \psi / dr$, which vanishes identically wherever ψ is constant. The ambipolar field is computed and not assumed, following Koskinen et al. (2013) section 2.1,

$$eE = -\frac{1}{n_e} \frac{d(n_e kT)}{dr} = -kT \frac{d \ln(n_e T)}{dr}, \quad (81)$$

in the logarithmic form, which is exact for the barometric stratification of the base and the same second-order central difference elsewhere. It is gated by `he_ambipolar` (default `on`); α_T is 0 by default (`He_alphaT`).

The equation is split. The advection of the element mass fraction is a divergence of the same face mass flux the density rides on and is carried inside the Runge-Kutta stages (98), so each element's nucleus total over the domain changes only by its boundary fluxes to round-off. What is left for the implicit step here is $\rho(X^{\text{new}} - X^{\text{adv}})/\Delta t = -r^{-2}\partial_r(r^2 J)$, taken at the composition the stages advected; the operator advects only when it is *given* the face mass flux, which happens in the fixed-wind relaxation that has no stages to ride on.

The discrete operator is one backward-Euler step with frozen coefficients, faces carrying r^2 areas, Newton on a tridiagonal Jacobian (the step is nonlinear in X through the product $X(1-X)$). The gradient term is central; the drift term keeps a Péclet hybrid, central where $|B|\Delta r \leq 2(A+E)$ and donor cell otherwise. The two factors of $X(1-X)$ come from *opposite* sides of the face, because the drift is a counter-flow and a donor-cell rule must take each element's mass fraction from the cell that element leaves:

$$J_{\text{drift}}(f) = \begin{cases} -B_f X(j+1) [1-X(j)], & B_f \geq 0 \text{ (helium inward)}, \\ -B_f X(j) [1-X(j+1)], & B_f < 0. \end{cases} \quad (82)$$

A donor at $X = 0$ then sends nothing and an acceptor at $X = 1$ receives nothing, so a cell on either end of the axis cannot be pushed past it; with the Péclet switch the same holds in the central branch, and the row argument (time term, gradient flux, upwind advection and drift all of one sign at a cell that first touches an end) gives $0 \leq X \leq 1$ on the residual itself rather than through a comparison principle. Measured, the worst excursion outside $[0, 1]$ before the assertion clip is -1.05×10^{-1} .

`project_elements` then rebuilds the species vector on the new element totals: helium-bearing species are scaled by $r_{\text{He}} = n_{\text{He}}^{\text{new}}/n_{\text{He}}^{\text{old}}$, component-1 species by r_1 , and a species carrying a nucleus of each (HeH^+) by $\min(r_1, r_{\text{He}})$, the nuclei this undercounts for the element with the larger factor being deposited into that element's neutral ground species. Every operation is a multiplication by a nonnegative factor or an addition, so no negative intermediate can arise and both element totals are met exactly. The invariant is that the species handed back carry the mass the species handed in carried, to round-off, which requires r_1 to be the ratio of the two component-1 *masses* read from the cell. The result is the initial guess for `ioniz_eq`, which owns the split *within* an element while this step owns the element totals.

The whole operator is gated by `He_diffusion`, default off. With `he_metal_diffusion` (default off) each trace element additionally carries its own mass fraction and its own binary coefficient; otherwise the metals are frozen to hydrogen, i.e. they ride in component 1 at the metal/H ratio the cell holds, and what `project_elements` preserves for them is that ratio and not the density.

7. Numerical scheme

7.1 The radial grid

The domain $[1, r_{\text{max}}]$ is discretized into N finite-volume cells (`Grid cells:`, default 500) with $N_g = 2$ ghost cells at each end; all grid arrays run over $1 - N_g \leq j \leq N + N_g$. `define_grid.f90` builds the cell centers r_j , the interfaces $r_{j+1/2}$ as their midpoints, and the widths $\Delta r_j = r_{j+1/2} - r_{j-1/2}$. The last is an exact identity of the discretization and not an approximation: it is the same face pair that forms ΔV_j in (5), the same pair the gravitational source differences, and the same width the column integrals $n_j \Delta r_j$ and the cell optical depths use.

Three grid types are available, as in ATES. Uniform takes $\Delta r = (r_{\max} - 1)/N$. Stretched places the centers at $r_j = r_{\max}^{(j-1+N_g)/(N+2N_g-1)}$. Mixed, the type every shipped configuration uses, lays N_{low} uniform cells of width Δr_{base} on the base and then stretches the remaining $N_{\text{up}} = N - N_{\text{low}}$ cells by a factor k that solves

$$\frac{1 - k^{N_{\text{up}}}}{1 - k} = \frac{r_{\max} - r_{N_{\text{low}}}}{\Delta r_{\text{base}}} \quad (83)$$

by Newton-Raphson to 10^{-8} , which is Eq. (10) of Caldiroli et al. (2021). The two numbers are the key Base grid [dr, cells]: and default to $\Delta r_{\text{base}} = 2 \times 10^{-4} R_p$ and $N_{\text{low}} = 50$, the values ATES hard-codes. The widths are then smoothed once by a 1-2-1 pass running downward from the junction to the base, the centers are rebuilt from the smoothed widths, and the whole grid is rescaled back onto $[1, r_{\max}]$. The requirement behind Δr_{base} is that the pressure scale height $H = k_B T / (\bar{m} g)$, with $\bar{m} = \rho / n_{\text{tot}}$ the mean particle mass, be resolved by several cells at the base; a configuration that violates it is the one that produces a base-layer momentum imbalance no solver removes.

Two windows are located once, at grid construction: $j_{\min}$, the first cell with $r \geq r_{\text{esc}}$ (Escape radius [R_p]:), which the marching convergence measure of Sec. 7.7 is taken over, and j_{flux} , the first cell with $r \geq r_{\text{flux}}$ (default $1.2 R_p$), which the stationary flux gate uses. Both are clamped to mid-domain, with a message, where the requested radius lies outside the domain: a deep Roche-lobe-filling case can have $r_{\max}$ smaller than the default escape radius, and an empty window would make the measure $-\infty$ and the state converged at step zero.

7.2 Reconstruction

Both schemes reconstruct the *primitive* variables $W = (\rho, v, p)$ to the two sides of every face, and both write the left state of face j from cell j and the right state of face j from cell $j + 1$.

PLM. PLM_rec.f90 builds a limited slope from the three cell averages $j - 1, j, j + 1$ and evaluates it at the two faces,

$$W_{L,j+1/2} = W_j + \frac{1}{2} \sigma (r_{j+1} - r_j), \quad W_{R,j-1/2} = W_j - \frac{1}{2} \sigma (r_j - r_{j-1}), \quad (84)$$

with σ the generalized minmod (monotonized central) slope of the right, left and central difference quotients at $\theta = 2$,

$$\sigma = \text{minmod} \left(\theta \frac{\Delta W_{j+1/2}}{r_{j+1} - r_j}, \theta \frac{\Delta W_{j-1/2}}{r_j - r_{j-1}}, \frac{W_{j+1} - W_{j-1}}{r_{j+1} - r_{j-1}} \right), \quad (85)$$

the slope being zero whenever the three arguments do not share a sign. This is the generalized minmod limiter as Käppeli & Mishra (2016) state it, at the least diffusive value for which the scheme stays total-variation diminishing; ATES uses the same limiter and attributes it to Kurganov & Tadmor (2000). The difference from ATES is that the quotients here are formed on the actual center spacing of a stretched grid. The same scalar limiter is exported as PLM_rec_scalar, so that a transported mass fraction rides on the same discretization as the density whose face flux carries it.

WENO3. Reconstruction.f90 uses the third-order WENO reconstruction with the nonlinear weights of Yamaleev & Carpenter (2009). With $\Delta W_{j\pm 1/2}$ the two interface jumps bounding cell j ,

$$\tau = \Delta W_{j+1/2} - \Delta W_{j-1/2}, \quad S_0 = 1 + \frac{\tau^2}{\Delta W_{j+1/2}^2 + \Delta r_j^2}, \quad S_1 = 1 + \frac{\tau^2}{\Delta W_{j-1/2}^2 + \Delta r_j^2}, \quad (86)$$

which is their $1 + \tau/(\epsilon + \beta_r)$ with their already-squared τ (their Eq. 22) and the floor $\epsilon = \Delta r_j^2$ folded in. The face states are

$$\begin{aligned} W_{L,j+1/2} &= W_j + \frac{S_0 C_2(j) \Delta W_{j+1/2} + D_1(j) S_1 C_1(j-1) \Delta W_{j-1/2}}{S_0 + D_1(j) S_1}, \\ W_{R,j-1/2} &= W_j - \frac{D_2(j) S_0 C_2(j) \Delta W_{j+1/2} + S_1 C_1(j-1) \Delta W_{j-1/2}}{D_2(j) S_0 + S_1}, \end{aligned} \quad (87)$$

with the purely geometric coefficients $C_1(j) = \Delta V_{j+1}/(\Delta V_j + \Delta V_{j+1})$, $C_2 = 1 - C_1$, $D_1(j) = \Delta V_{j+1}/(\Delta V_j + \Delta V_{j-1})$ and $D_2(j) = \Delta V_{j-1}/(\Delta V_j + \Delta V_{j+1})$. Each candidate is the linear interpolant of two cell averages *in the volume coordinate*, so the jump to a neighbor is weighted by the cell's own volume share; these shares, and not the nonlinear weights, are what set the order on a stretched grid. The ideal weights $d_0 = 2/3$, $d_1 = 1/3$ of the uniform-grid scheme enter only through their ratio, which is what D_1 and D_2 carry: exactly $1/2$ on a grid uniform in the volume coordinate and $1/2 + O(h)$ otherwise.

Two limits of the scheme are stated at the routine. What runs is a third-order WENO reconstruction *with* those weights and not the energy-stable ESWENO scheme of the same paper, whose extra dissipation flux (their Eq. 48) is not computed here. And the floor $\epsilon = \Delta r_j^2$ has the required h^2 scaling but drops the solution scale of the published prescription, so it is not invariant under a rescaling of the solution; what that costs is the stencil biasing in the tenuous outer domain and not the order, which is flat over four decades of the coefficient.

Positivity of the face states. Neither scheme has a positivity property: a face value of ρ or p can cross zero while every cell average on the stencil is positive, and at Mach 60 the thermal pressure is 2×10^{-4} of the total energy density, so a reconstruction across a jump tips it negative and the sound speed takes the square root of it. `positivity_limited_faces` therefore scales each offending face state *continuously* toward its own cell average,

$$W_{\text{face}} \leftarrow W_{\text{avg}} + \theta (W_{\text{rec}} - W_{\text{avg}}), \quad \theta = \min_{q \in \{\rho, p\}} \frac{q_{\text{avg}} - \varepsilon}{q_{\text{avg}} - q_{\text{rec}}}, \quad \varepsilon = \epsilon_{\text{mach}} q_{\text{avg}}, \quad (88)$$

which is the linear scaling idea of Zhang & Shu (2010a) applied to the primitive vector, and not their limiter: they scale the conserved vector in two steps with θ the root of a quadratic, one θ per cell against one per face state here, and their floor is absolute against one unit in the last place of the cell average here. What carries over is their Lemma 2.5, that a convex combination of an admissible average with the reconstruction is admissible; their Theorem 2.1 does not, its hypotheses being a Gauss-Lobatto quadrature set and a CFL condition nothing here imposes. Continuity is the point: a hard switch at the crossing is a step discontinuity of the residual and no stationary solve can descend across one. A face whose reconstruction is admissible has $\theta = 1$ and is not rewritten.

The PLM to WENO3 continuation. The two-stage recipe Reconstruction scheme: PLM+WENO3 marches with PLM until the convergence measure falls below the first `du_th` value (or its plateau is detected) and then continues with WENO3. Replacing one discretization by another between two time steps is a jump of the right-hand side, so the key Reconstruction continuation: interpolates instead. `reconstruction_continuation_rhs` (`steady_residual.f90`) evaluates the homotopy

$$R_\lambda(u) = (1 - \lambda) R_{\text{PLM}}(u) + \lambda R_{\text{WENO3}}(u), \quad (89)$$

returning its flux-difference and source parts separately, and blends the stored interface fluxes, the face pressures and the three momentum terms on the same λ . Everything the two schemes disagree about is inside the two evaluations:

the face reconstruction, the extrapolated reconstructed-face boundary states, and the pressure split of Sec. 2.3. At $\lambda \leq 0$ and $\lambda \geq 1$ a single evaluation is taken and the two pure operators are reproduced exactly; only the open interval costs two right-hand sides. Because the admissible set $\rho > 0$, $\rho e > 0$ is convex and each stage is then a convex combination of two forward Euler stages, the continuation cannot manufacture a positivity failure that neither scheme has. The ramp is adaptive: λ advances by $\Delta\lambda$ where the relative state change of the step stays below a tolerance and $\Delta\lambda$ is halved where it does not.

7.3 Approximate Riemann solvers

`Num_Fluxes.f90` carries three fluxes, selected by the mandatory key `Numerical_flux:`. Each side's total energy and sound speed are built with its *own* heat capacity, which is what the caloric equation of state makes different across a face inside the H_2 front; the physical flux is

$$F(W) = (\rho v, \rho v^2 + p, v(\mathcal{E} + p)), \quad (90)$$

the bracketed pressure being present under PLM without the well-balanced option, as Sec. 2.3 requires.

HLLC. The three-wave solver of Toro et al. (1994), with the Davis/Einfeldt speed bounds of Batten et al. (1997),

$$S_L = \min(0, v_L - a_L, v_R - a_R), \quad S_R = \max(0, v_L + a_L, v_R + a_R), \quad (91)$$

$$S_* = \frac{p_R - p_L + \varphi_L v_L - \varphi_R v_R}{\varphi_L - \varphi_R}, \quad \varphi_K = \rho_K(S_K - v_K), \quad (92)$$

and the star states

$$u_K^* = \frac{\rho_K(S_K - v_K)}{S_K - S_*} \left(1, S_*, \frac{\mathcal{E}_K}{\rho_K} + (S_* - v_K) \left[S_* + \frac{p_K}{\varphi_K} \right] \right)^\top, \quad (93)$$

the flux being F_K , $F_K + S_K(u_K^* - u_K)$ or the mirror of those according to the signs of S_L , S_* , S_R . Because S_L and S_R are clamped at zero, a face with every physical wave moving outward returns $F(W_L)$ and never forms an expression in the right state. The face pressure p_{out} that the momentum assembly of (7) needs is the pressure of whichever side the branch took. HLLC is the flux every shipped configuration under `examples/` and `backup/regression/` states; the key is mandatory and has no default.

Roe. The linearized solver of Roe (1981), with the averages

$$\tilde{\rho} = \sqrt{\rho_L \rho_R}, \quad \tilde{v} = s v_L + (1-s) v_R, \quad \tilde{H} = s H_L + (1-s) H_R, \quad s = \frac{\sqrt{\rho_L}}{\sqrt{\rho_L} + \sqrt{\rho_R}}, \quad (94)$$

$\tilde{a} = \sqrt{(\bar{\gamma} - 1)(\tilde{H} - \tilde{v}^2/2)}$, and the flux

$$F_{j+1/2} = \frac{1}{2} (F_L + F_R) - \frac{1}{2} \sum_{i=1}^3 \tilde{\alpha}_i |\tilde{\lambda}_i| \tilde{K}^{(i)}, \quad (95)$$

with the wave strengths $\tilde{\alpha}_{1,3} = (\Delta p \mp \tilde{\rho} \tilde{a} \Delta v) / 2 \tilde{a}^2$ and $\tilde{\alpha}_2 = \Delta \rho - \Delta p / \tilde{a}^2$. The single index $\bar{\gamma}$ is the arithmetic mean of the two face gammas, exact wherever the two sides agree; a run whose caloric equation of state makes the index vary across the front, that is molecular chemistry with the ladder heat capacity, is refused this flux at startup rather than being given a wave speed with no bound property. A Harten-Hyman entropy correction is applied to the acoustic

eigenvalues from the star state that `speed_estimate_ROE.f90` builds (the primitive-variable, two-rarefaction and two-shock estimates of Toro (2009) chapter 9, selected by the pressure ratio). Where that estimate has no admissible star state, two rarefactions separating fast enough leave vacuum, or no branch returns a positive star pressure and positive star densities, the face is given the HLLE flux instead,

$$F_{j+1/2} = \frac{S_R F_L - S_L F_R + S_L S_R (u_R - u_L)}{S_R - S_L}, \quad \begin{aligned} S_L &= \min(v_L - a_L, \tilde{v} - \tilde{a}), \\ S_R &= \max(v_R + a_R, \tilde{v} + \tilde{a}), \end{aligned} \quad (96)$$

which is positively conservative under those bounds (Einfeldt et al., 1991), where the Roe flux is not. The number of such faces is counted and reported.

Local Lax-Friedrichs.

$$F_{j+1/2} = \frac{1}{2} [F_L + F_R - \alpha (u_R - u_L)], \quad \alpha = \max(|v_L| + a_L, |v_R| + a_R), \quad (97)$$

the coefficient being the spectral radius of the flux Jacobian (Rusanov, 1961). It is the most diffusive of the three and resolves no contact wave; it is also the flux the positivity repair below substitutes at a single face, because with cell-average input states it maps positive density and positive internal energy to positive density and positive internal energy under $\Delta t(|v| + c)/\Delta r \leq 1$ (Perthame & Shu, 1996; Zhang & Shu, 2010b).

7.4 Positivity repair of a stage

A Runge-Kutta stage can leave the admissible set $\rho > 0$, $\rho e > 0$ even with admissible face states, because the source terms are bounded by no flux choice. `positivity_limited_fluxes(RK_rhs.f90)` repairs such a stage by replacing, at the offending cells only, the high-order interface fluxes with the first-order Lax-Friedrichs flux of the neighboring cell averages and rebuilding those cells' update from the mixture; the replacement is swept until no cell is inadmissible or until both faces of a cell are already first order, in which case no flux choice repairs it and the caller halves the time step. Replacing one face rather than all of them is the flux correction of Hu et al. (2013). The stored face fluxes are written back after the repair, because the transported species rows ride on F^ρ and a species flux built on a discarded high-order flux would not sum to the mass update the density took. Every count is reported at the end of a run, separately for repaired faces belonging to accepted steps, so that a run in which nothing fired is identifiable as the run an unguarded build gives.

7.5 Time integration and the time step

The marching loop of `EXHALE_main.f90` advances the hydrodynamics with the three-stage strong-stability-preserving Runge-Kutta method (Gottlieb & Shu, 1998),

$$\begin{aligned} u^{(1)} &= u^n - \Delta t (dF - S)|_{u^n}, \\ u^{(2)} &= \frac{1}{4} \left[3u^n + u^{(1)} - \Delta t (dF - S)|_{u^{(1)}} \right], \\ u^{n+1} &= \frac{1}{3} \left[u^n + 2u^{(2)} - 2\Delta t (dF - S)|_{u^{(2)}} \right], \end{aligned} \quad (98)$$

each stage followed by the boundary condition, by the admissibility test of Sec. 7.4, and by the species advection sub-step that rides on the face mass fluxes that stage built. A stage that cannot be repaired causes the whole step to be discarded and retaken from u^n at half Δt , up to a bounded number of bisections.

The time step is the CFL condition `eval_dt.f90`,

$$\Delta t = C_{\text{CFL}} \min_j \frac{\Delta r_j}{|v_j| + c_{s,j}}, \quad C_{\text{CFL}} = 0.6 \text{ by default}, \quad (99)$$

with $c_s = \sqrt{\gamma_{\text{eff}} p / \rho}$ from the cell's own composition. With Time stepping: Local each cell instead takes its own $\Delta t_j = C_{\text{CFL}} \Delta r_j / (|v_j| + c_{s,j})$; since the fixed point $dF = S$, $\mathcal{H} = \Lambda$ is independent of the step, this is a legitimate acceleration toward a steady state, and it makes the trajectory meaningless as a time history. In the global mode the local array is uniformly the global step, which keeps the update arithmetic identical to the scalar form.

7.6 Operator splitting of one step

One accepted marching step applies the following operators, in this order, with the state after each one carried into the next (`EXHALE_main.f90`):

1. the three Runge-Kutta stages (98), each with its reconstruction, flux assembly, boundary fill, positivity test and species advection sub-step;
2. the conversion to primitive variables, the composition-dependent particle and electron counts, and the temperature (13);
3. the advective half of the element transport written into the species vector, followed by the implicit diffusive half (`binary_element_diffusion.f90`, Sec. 6.9), active only with `He_diffusion`;
4. the vertical transport of the molecular carriers with their chemistry (`diffusive_photochemistry.f90`, Sec. 6.5), active only with the carrier transport option;
5. the refresh of the lagged $H(n=2)$ populations and their Balmer source (`excited_hydrogen.f90`, Sec. 4.6);
6. the coupled source step: a bounded fixed-point iteration alternating the composition equilibrium at the current temperature (`ioniz_eq`, Sec. 4.4) with the semi-implicit energy update (71) at the returned composition, exiting when the temperature and the composition both stop moving;
7. the boundary condition;
8. the Crank-Nicolson viscous and conductive stage (20), where those options are on, and the boundary condition again;
9. the periodic Shapiro filter, where it is on, and the boundary condition again.

Steps 3, 4, 8 and 9 are no-ops unless their key is present. The Shapiro filter is the 1-2-1 low-pass pass $u_j \leftarrow u_j + \frac{1}{4}\epsilon(u_{j-1} - 2u_j + u_{j+1})$ applied every `shapiro_every` steps, the way CETIMB damps the base sound wave (Koskinen et al., 2013). It is off by default and is recorded as a validity state of every step it runs on, because unlike the low-Mach term of Sec. 2.7 it alters the marched state with no source term behind it and therefore has no counterpart in the stationary residual.

7.7 The marching convergence measure and the stop conditions

The measure the marching stops on is the radial spread of the mass flux over the escape window,

$$du = \frac{\max_{j \geq j_{\min}} F_j - \min_{j \geq j_{\min}} F_j}{|\bar{F}|}, \quad F_j = \rho_j v_j r_j^2, \quad (100)$$

with $\bar{F}$ the mean over the same window. A steady spherical wind carries one mass flux at every radius, which is the stop ATES uses ($\Delta \dot{M} / \dot{M} < 10^{-3}$) and the constancy of $F_c = \rho v r^2$ CETIMB asks for. The sign is kept and the normalization is the mean: taking $|F|$ first would read a window whose flux reverses, an inflow at one radius and an outflow at another, as flat. It is evaluated once per step and read by every consumer, so the stop, the trigger arming, the staged secondary-ionization activation of Sec. 3.5 and the handoff to the steady solver all speak about one number. This is emphatically *not* a velocity change, and a single-stage PLM run stalls at a few percent spread.

A second measure, the largest relative change of the conserved state over a step,

$$dtu = \max_{k \in \{1,2,3\}} \max_{j_{\min} \leq j \leq N} \left| 1 - \frac{u_{k,j}^{n+1}}{u_{k,j}^n} \right|, \quad (101)$$

says that the state has stopped changing; a row that was exactly zero at the start of the step has no relative change defined and the measure is taken as unbounded rather than dropped.

A run stops when one of the following holds, and reports which: $du < du_th$ (default 10^{-3}) with the trigger armed and the flux level stable; $dtu < dtu_th = 10^{-8}$; du settled on a plateau over N_stall steps (default 2000) within $stall_tol$ (default 10^{-6}); the steady gates of Sec. 9.2 where `Resid tol:` was stated; the Newton finish returning a state; or the hard cap `count_max` (10^6). The triggers are *armed* the first time the measure is seen at or above its threshold, so only a descending crossing can fire them and a run whose initial condition already carries a flat mass flux, which a transonic one does, cannot stop at step zero. With `Solver: Newton` the marching is a warm-up only: the du and dtu stops are disabled and the loop hands over once du falls below `newton_du_switch` (default 10^{-2}) or plateaus near it. Both measures can be small at a premature balance of the operator split while the energy residual is of order 30, so for a quantitative mass-loss rate the stationary finish is not optional.

8. Boundary and initial conditions

8.1 The lower boundary

The lower boundary is a characteristic condition imposed *at the base face* $r_{-1/2}$, with the number of conditions set by the sign and magnitude of the Mach number there (`base_boundary.f90`). At that face the eigenvalues are $v - c$, v and $v + c$, and a wave enters the domain when its eigenvalue is positive, the radial coordinate increasing outward:

$0 < v < c$	subsonic inflow	2 reservoir conditions, 1 interior
$-c < v < 0$	flow reversal	1 reservoir condition, 2 interior
$v > c$	supersonic inflow	3 reservoir conditions, 0 interior
$v < -c$	supersonic outflow	0 reservoir conditions, 3 interior.

(102)

The interior relation is the linearized $v-c$ compatibility condition in its LODI form (Thompson, 1987; Poinso & Lele, 1992),

$$p_b - \rho_i c_i v_b = p_i - \rho_i c_i v_i, \quad (103)$$

with the subscript i denoting the interior state continued to the face. The closed-form Riemann invariant $v - 2c/(\gamma - 1)$ is deliberately not used: it assumes a single constant γ , and the caloric equation of state gives every cell its own $\gamma_{\text{eff}}(x_{\text{H}_2}, T)$, so that invariant is wrong wherever the H_2 fraction or the temperature varies across the boundary stencil, which is the whole molecular base. The sound speed c_i is the interior one, because the outgoing wave is an interior wave.

The reservoir is (p, s) . The two reservoir conditions are the pressure and the specific entropy of the lower atmosphere at the base *level*, not the pressure and the density. The distinction matters as soon as the base composition changes: with (p, s) the temperature follows the composition, with (p, T) the entropy does. The entropy is carried implicitly, as the isentrope through $(p_{\text{base}}, T_{\text{base}})$ at the base composition, which is what `base.inp` and the lower-atmosphere profile reader state. The level is $r = 1$ and not the face, so the level the user states does not move when the grid does; the reservoir state is carried from the level to the face along its own hydrostatic isentrope.

Continuation along a hydrostatic isentrope. Both the interior state and the reservoir state are carried to the face by `continue_hydrostatic_isentrope`, which uses the fact that at rest and at constant entropy the Bernoulli function is uniform,

$$u(T) + T - \frac{\Phi(r)}{\hat{n}} = \text{const} \implies u(T_b) + T_b = u(T_a) + T_a - \frac{\Phi(r_b) - \Phi(r_a)}{\hat{n}}, \quad (104)$$

with u the internal energy per particle of the caloric equation of state and $\hat{n}$ the particles per unit mass, solved by Newton with $dh/dT = c_v + 1$; the density then follows from $d \ln \rho = c_v d \ln T$ as $\rho_b = \rho_a (T_b/T_a)^{c_v(T_{\text{mid}})}$. The midpoint heat capacity makes this exact to third order in $\ln(T_b/T_a)$, and the routine is only ever used over half a cell and over the two ghost cells, where that logarithm is below 0.1 and the density error is below 10^{-7} relative. It is not a general isentrope integrator over a scale height and is not used as one.

Continuing along the isentrope rather than extrapolating linearly is what makes the condition well balanced: in a hydrostatic atmosphere at rest the interior face pressure equals the reservoir's and (103) returns $v_b = 0$ exactly, at any resolution. The interior state at the face is built from cell 1 alone, never from the reconstruction stencil, because that stencil reaches into the ghost and a boundary condition that reads its own output is circular. The interior velocity is carried to the face at constant $\rho v r^2$ rather than copied.

What is held, and the branches. The pressure is a reservoir condition in every subsonic branch, $p_b = p_{\text{res}}$, and the velocity then follows from (103). The entropy is a reservoir condition only while gas enters; on reversal the entropy at the face is the interior's, advected out. The handover between the two is a C^1 cubic smoothstep in the face Mach number over a window of half width 10^{-6} , which is the one designed-in constant of the module: the branch is a genuine change in the number of conditions and therefore a kink in the residual, and the stationary solver needs a differentiable one. The window sits far below any intended operating point, the base face Mach number of the hot-Uranus configuration being about 6×10^{-6} . A supersonic outflow branch hands the whole face state to the interior through the same smoothstep.

A supersonic *inflow* would need a third reservoir condition which a (p, s) reservoir does not have, so the state is not a solution of a well-posed problem there; the run reports the crossing and does not invent a velocity. A compatibility relation is linear and far from a solution can ask for any velocity, so the face state is capped at Mach 5 and the events are counted.

Holding the incoming acoustic invariant $p_b + \rho c v_b = p_{\text{res}}$ instead of the static pressure is present in the source as a weight that is zero. Both forms are exactly well balanced at rest; on a moving solution the invariant makes the static face pressure fall below the reservoir's by a fraction M of itself where a real reservoir's Bernoulli drop is a fraction M^2 , which is why the weight is zero.

The ghost cells. `base_ghost_averages` fills the two lower ghosts with the *volume averages* of the hydrostatic isentrope through the face state, continued below it, evaluated by 8-point Gauss-Legendre quadrature weighted by r^2 , with the velocity carrying a constant $\rho v r^2$ so the reconstruction sees no spurious velocity gradient. Writing the face value into a ghost cell average instead is a half-cell error: the two differ by $\Delta r/2H$, which is first order in Δr in the state and *zeroth* order in the momentum residual it produces, an imbalance that no resolution removes. `Rec_BC` then puts the face state itself into the left slot of the base face; the right slot is the interior's reconstruction of cell 1 down to that face and is deliberately left alone, the Riemann solver being what mediates the two.

The base Mach number is recorded every step and a first crossing into supersonic inflow is warned about, as a validity check on the boundary that is independent of the chemistry (`Apply_BC.f90`).

8.2 The upper boundary

No characteristic enters the domain at an outflowing outer face, so the boundary imposes no condition and the ghost is a continuation of the interior. The rule is stated once, for every reconstruction: which characteristics cross a face is a property of the flow and not of the scheme that reads the ghost. The continuation (`free_outflow_ghost`) is the isothermal hydrostatic one,

$$\rho_g = \rho_N \exp \left[-\frac{\Phi_c(N+k) - \Phi_c(N)}{p_N / \rho_N} \right], \quad p_g = p_N \exp[\dots], \quad v_g = v_N, \quad (105)$$

the state of rest the last physical cell carries outward under the code's own potential with the isothermal sound speed squared p/ρ held at cell N 's value. Since ρ and p are scaled by the same factor, the ghost is at the *same* temperature as cell N , so the ionization sweep, the column integral and the sound speed of the time step all see the temperature they saw before, at the density the stratification gives.

A zero-gradient copy would assert $dp/dr = 0$ at the outer face, a hydrostatic imbalance of the order of the local weight that no resolution removes, and under PLM it is worse: the forward difference entering the limiter is then exactly zero, so the limited slope of cell N vanishes and that cell carries no pressure gradient at all. On an analytic hydrostatic column the momentum residual of cell N in units of its own weight converges at order 1.80 for (105), 1.42 for a linear extrapolation and not at all (0.006) for the zero-gradient copy. The velocity stays a copy: the outgoing characteristics carry it out of the domain and the boundary has nothing of its own to say about it; a gradient-carrying ghost velocity either drives the run into a base-breathing limit cycle or divides by a ghost density the stratification can make arbitrarily small. A ghost that leaves the admissible set falls back on the zero-gradient copy and is counted.

For the reconstructed face states, Rec_BC extrapolates outward with a zero-gradient copy under PLM and a linear extrapolation under WENO3.

8.3 Initial conditions

set_IC.f90 builds four families, selected by IC mode: (or by the older Transonic IC and Hot Parker IC keys, which take precedence).

Cold hydrostatic (cold, the default). An isothermal hydrostatic atmosphere at the base temperature,

$$\rho(r) = \rho_{bc} \exp[-b_{\text{eff}}(\Phi_c(r) - \Phi_c(0))], \quad p = \frac{\hat{n}_{bc}}{\rho_{bc}} \rho, \quad T = 1, \quad (106)$$

with a small linear velocity seed $v = \frac{1}{2}(r - r_0)$ and a nearly neutral composition. The factor b_{eff} starts at 1 and is raised in steps of 0.2 until the mid-domain density falls below 10^{-8} , so that a low-gravity planet does not start with a domain at uniform density; the value used is reported. This is the ATES initial condition.

Transonic (transonic). The steady isothermal-wind profile obtained from the algebraic Bernoulli integral of the isothermal wind equation in the code's own potential,

$$\frac{1}{2}v^2 - c^2 \ln v - 2c^2 \ln r + \Phi(r) = B, \quad (107)$$

with B fixed at the sonic point, where $v = c$ and $d\Phi/dr = 2c^2/r$. The subsonic root is taken inside the sonic radius and the supersonic root outside it, and the density follows from $\rho v r^2 = \dot{M}$ with $\dot{M}$ pinned to the base density. It exists only where the potential admits an interior sonic point; where it does not, the cold hydrostatic profile is kept. It is needed for deep Roche-lobe-filling cases whose sonic point sits near L1 and which launch no wind from a hydrostatic start.

Hot Parker warm seed (hot_parker). Seeding the full isothermal Parker wind, its density as well as its velocity, destabilizes the run: the hot, low-density Parker column is globally inconsistent with the code's non-isothermal equilibrium and drains onto the base. The warm seed therefore keeps the stable cold hydrostatic *density* and overlays only a warm temperature and ionization ramp and the Parker *velocity* as a head start, both anchored to the cold static base through a cubic smoothstep window that is zero at $r = 1$ and unity above a stated transition radius. This removes the inverted base pressure gradient that otherwise drives an inflow and a thermal runaway.

Automatic selection (auto). One exact probe decides between the cold and the transonic family with no tunable threshold: does the *cold* base sound speed admit an interior sonic point in the actual potential? If it does the transonic profile is taken, which catches both the deep Roche-lobe cases, where the Roche $d\Phi/dr \rightarrow 0$ toward L1 always crosses the critical condition, and the low-gravity cases whose sonic radius falls inside the domain; if it does not, the cold hydrostatic profile is taken. The hot Parker seed is deliberately not auto-selected.

Wind-AE (windae). The ported Wind-AE solver in src/modules/wind_ae/ is run in process to build a warm-start profile on the code's own grid, which is then loaded through the restart path.

Every family fills the whole species vector, and a molecular run seeds the H_2 column from the same two functions the inflowing ghost uses: the handoff value where a lower atmosphere states one (Sec. 6.7), and the chemical-equilibrium mixing ratio (78) at (p_{base}, T_0) otherwise, clipped at the element-ratio ceiling $q_{\text{max}} = 0.5/(0.5+y)$. The reason is that the optically thick base equilibrium is strongly molecular, and a zero- H_2 start makes the base composition solve land on the spurious atomic root.

Two of the families set `force_start`, which suppresses the convergence test for the first iterations: a transonic profile already carries $\rho v r^2 = \text{const}$, so the marching measure of (100) would be near zero at step 0 and the run would exit before the cold wind had heated to its steady state.

8.4 Restart

`Load IC?` `True` restores the state from `Hydro_ioniz_IC.txt` and `Ion_species_IC.txt` (`load_IC.f90`). Files carrying the `#` columns schema header are read by label, so every species the file holds, metal ionization stages included, is restored; a headerless legacy file takes the original fixed seven-column read and the metals are rebuilt from the abundances, which is reported. An element the file does not carry, or carries as identically zero, is initialized from its abundance exactly as a cold start does. An element whose reservoir a lower-atmosphere handoff states has its loaded column renormalized onto that reservoir by one factor common to all its ionization stages. The file's own He/H ratio is compared against the input's, and a state written at a different element ratio is rescaled or refused rather than silently adopted, the input file being the authority on composition.

`Restart intent`: states what the loaded state is continued as, and is refused when no state is being loaded. `trajectory` continues the history, including its elapsed time; `relaxation` treats the state as a numerical iterate; `stationary` asks for a stationary solve from it, and takes a second word, `evaluate` to only measure the residual of the loaded state, or `equilibrate` to bring the composition to equilibrium before the solve. `Restart option change`: names the entries of the state file's option line that are allowed to differ from this run's, and a token that decides how many unknowns the state has is refused: naming it cannot make the file's rows this run's rows.

9. The stationary state

Marching to a steady state is only one of the two ways EXHALE reaches one. The other is to solve for it directly, as the root of the stationary residual of the very same discrete operator. This section states that residual, the measure by which a state is judged steady, and the two solvers that drive it to zero.

9.1 The stationary residual

`assemble_residual` (`steady_residual.f90`) evaluates the right-hand side of one forward Euler step of the marching operator and subtracts the radiative source from its third row,

$$\begin{aligned} R_1 &= dF_1 - S_1, \\ R_2 &= dF_2 - S_2 - S_\mu, \\ R_3 &= dF_3 - S_3 - (\mathcal{H} - \Lambda) - S_\mathcal{E}, \end{aligned} \tag{108}$$

with dF and S from (6) to (11), evaluated through the same `reconstruction_continuation_rhs` the marching stages call, so that the fixed point of the marching and the zero of the residual are the same state by construction. The lowest ghost cell has no lower face and therefore no balance to state; its row is explicitly zero, so that no reader of R finds a heating rate standing alone in a row that has no fluxes.

A residual row is a dimensional quantity and has to be divided by something before it can be compared with a tolerance. Every row is divided by the largest term *that row itself contains* (`residual_row_scale`):

$$s_1(j) = \frac{\max(|F_{j-1/2}|A_-, |F_{j+1/2}|A_+)}{\Delta V_j}, \quad s_2(j) = \max(|\text{ram}|, |\partial_r p|, \rho|\partial_r \Phi|, |S_\mu|), \quad s_3(j) = \max(|dF_3|, |S_3|, \mathcal{H}, \Lambda). \quad (109)$$

Since ΔV_j is exactly the volume the flux difference was divided by, R_1/s_1 is the fractional change of the mass flux across the cell, and the mass row having no source, that fraction is the whole of what the row says. The momentum scale uses the three gathered physical terms of Sec. 2.3 and not dF_2 and S_2 , for the reason given there.

A signal-speed bound $D_k(|v| + c_s)/\Delta r$ is not usable as the scale of these rows: nothing in any of them moves at c_s , so in a quasi-hydrostatic layer that bound stands far above the terms the row holds and R_1 divided by it is the fractional flux error per cell *times* the local Mach number, which at Mach 5×10^{-5} is a blindness of four decades.

These are the scales of the *measure* only. The Newton system and its merit keep the state scales D_k of `state_scales_of_cell`, so the merit and the acceptance test are deliberately not one expression apart.

9.2 The acceptance test

`steady_gates_met` is the only definition of an accepted steady state in the code: every route that may stop on “the equations are satisfied”, the two stationary solvers and the marching loop under `Resid tol:`, calls it. Four gates:

1. the *residual gate*, $\|R\| < \text{resid_tol}$, with $\|R\|$ the maximum over physical cells and rows of $|R_k(j)|/s_k(j)$;
2. the *flux gate*, the spread of the Riemann face mass flux $F_{f+1/2}r_{f+1/2}^2$ over $r \geq r_{\text{flux}}$ below `flux_spread_th` (default 2×10^{-5} at $r_{\text{flux}} = 1.2R_p$). A residual norm does not say this: a slowly varying flux has a small local derivative, and two states that differ by a factor 25 in the spread differ by only 1.9 to 5.7 in every residual norm tried. It is taken on the *face* flux and not on the cell-centered $\rho v r^2$, because the finite-volume update moves the face flux and the centered product is a reconstruction of the state;
3. the *chemistry gate*, no cell carrying a composition that is not a root of the chemical network, that is no cell in acceptance class 4 or 6 of Sec. 4.4. Heat and cool in the energy row are evaluated on the accepted composition, so on such a cell the row is a number computed on a state the chemistry does not describe;
4. the *carrier gate*, where $n(\text{H}_2)$ is among the unknowns: the carrier row balanced to `carrier_resid_th` of its own largest terms.

A caller that cannot supply the third makes no statement about the chemistry and the test rests on the others; the marching stop is such a caller.

9.3 Certification

The four gates above judge the hydrodynamic rows and at most one species row. A configuration that also carries an elemental transport balance, a level population or a set of eliminated species would reach its answer with nothing said about those equations. `certification.f90` closes that: it lists every equation the configuration makes independent and returns, for each, one of three statuses, *not applicable*, *evaluated* with its residual measured on *this* state, or *unavailable*. The third is the point of the construction: a routine that returns zeros when its frozen background is not ready is indistinguishable at the call site from a balanced equation, and an active equation with no evaluator returns nothing at all. Both now say so, and a state carrying either is written and not certified.

The inventory covers the three hydrodynamic rows, each transported carrier balance, the He/H partition and the trace elemental transport balances, the He 2^3S and H($n=2$) level balances, and the eliminated-species closure of the coupled cell system. Every row measure is the *maximum* over physical cells of $|\text{res}|/\max(\text{scale}, \text{floor})$, never a volume average: a front or a thin layer must not be averaged away by the wind that carries most of the volume. The volume average is reported beside it and is never decisive. The measurement has no side effects; every module array the carrier evaluator writes is held aside around the call, allocation status included, because a certification is a measurement of a state and not an operation on it.

The tolerances are one per kind of row, so that the report says which number refused a state: 3×10^{-12} for mass, 10^{-8} for momentum, 10^{-6} for energy, 10^{-5} for an elemental or carrier balance in the wind, and the composition solve's own root tolerance for the closure and the level balances. One number for the three hydrodynamic rows is not possible: on one converged state the three reach values seven decades apart and their one-ulp arithmetic floors two decades apart.

Two of these are functions and not constants. A species row is gated only in the wind, $r \geq 1.20R_p$, and reported without gating below it: the balance is a cancellation between terms whose ratio is set by the flow, the advective divergence carrying the row in the wind and the eddy and settling terms carrying it below the homopause, each of the latter orders of magnitude larger than their difference, so the precision the same equation can be resolved to is not one number. And the mass row's tolerance is a function of the cell,

$$\text{tol}_1(j) = \max \left[3 \times 10^{-12}, \min(1, c_{\text{round}} f(j)) \right], \quad c_{\text{round}} = 10, \quad (110)$$

anchored on the rounding floor of that cell's own continuity row,

$$f(j) = \epsilon_{\text{mach}} \frac{\max_{\text{faces}} [\rho(|v| + c_s)A]}{\max_{\text{faces}} |AF|}. \quad (111)$$

The reasoning is that each interface flux is assembled from two reconstructed states and from their jumps, and where the column is nearly hydrostatic those jumps stand many decades below the states, so a correctly rounded interface flux carries the last bit of the $O(\rho c_s)$ quantities the Riemann solve is built from and not the last bit of its own value ρv . The smallest flux difference the arithmetic can resolve is then $\epsilon \rho(|v| + c_s)A/\Delta V$, which is one ulp in the wind and about ϵ/Mach in a layer whose flux is a tiny residue of the momentum. Measured at the base of three reference configurations that floor spans four decades, 5.7×10^{-14} to 3.2×10^{-11} , so a single fixed number would hold one state to eight ulps of its own base layer and another to a tenth of one ulp. The margin $c_{\text{round}} = 10$ stands a factor 3.7 above the largest measured ratio between the step the row takes when one ulp is added to every density and the estimate (111). The

ceiling of unity is not an operative branch on any state measured: the row measure is the fractional change of the mass flux across a cell, so a state whose estimated floor reaches 1 is one whose mass row cannot be judged at all.

9.4 The Jacobian-free Newton-Krylov solve

The unknowns of the direct solve are the conserved state on the physical cells,

$$Y = (\rho_j, (\rho v)_j, \mathcal{E}_j, \{y_{s,j}\})_{j=1}^N, \quad n_{\text{var}} = 3 + n_{\text{spec}}, \quad (112)$$

where the optional species rows $y_{s,j}$ are the elemental mass fractions of the diffusing elements and the transported molecular carriers, registered by `set_transported_species_rows` from the configuration. The ghost cells are not unknowns: the boundary condition fills them from the interior at every residual evaluation. The ionization state and the temperature are eliminated *locally* inside the residual, that is, given Y the equilibrium sweep of Sec. 4.4 re-solves the composition and returns the consistent heating and cooling of Sec. 5., so that $F(Y)$ is a residual of size $n_{\text{var}}N$ and not of the full chemical system.

The solve is a Jacobian-free Newton-Krylov iteration in the sense of Knoll & Keyes (2004): the Krylov method needs only the action of the Jacobian on a vector, which is taken as a forward difference of the residual, so the nonlocal radiative coupling never has to be assembled.

The outer iteration of `solve_steady_jfnk` is, per step: evaluate the full residual F at the iterate; build the banded finite-difference Jacobian of the *frozen-radiation* residual as a preconditioner; solve $(I/\Delta\tau + J)\delta Y = -F$ by right-preconditioned GMRES with the matrix-vector product taken matrix free, so that the nonlocal radiative coupling is captured; take a step along a controlled direction; and ramp $\Delta\tau$.

The banded preconditioner. WENO3 reaches two cells on either side and within a cell every variable couples, so in the flat ordering $Y(n_{\text{var}}(j-1)+k)$ a column couples to rows within $|\text{row} - \text{col}| \leq 3n_{\text{var}} - 1$; the half-bandwidths are that number and a graph coloring with stride $n_{\text{color}} = k_l + k_u + 1 = 6n_{\text{var}} - 1$ makes the row supports of same-color columns disjoint, so that the frozen-radiation residual, which is strictly local, is captured exactly by that many finite-difference probes. For $n_{\text{var}} = 3$ this is 8 and 17. The band is factored by LAPACK `dggbtrf` and applied as the right preconditioner of the Krylov cycle. The finite-difference and Krylov probe steps are taken in the diagonally scaled variables of `cell_state_scales`, the characteristic magnitude of each unknown in its own cell; a probe point that leaves the set of states the code can describe, a chemical balance row going to NaN or overflowing, is answered by halving the step and sampling closer to the iterate, at most six times, and never by accepting a non-finite residual or inventing a derivative.

The step control. Where the system carries no species row the step is the Krylov direction with a backtracking line search on the merit $\|F/D\|_2$, with a Grippo non-monotone reference, the worst true merit of the last five iterates, so that sideways steps are admissible and the solve can leave a bad neighborhood. Where the system carries a species row the step control is a scaled trust region with a Powell dogleg: the path runs from the origin to the Cauchy point s_U , the minimizer of the linear model along the steepest descent direction, and from there to the Krylov point s_N , and the step

is where that path first meets the ball of radius Δ ,

$$s = a_U s_U + a_N s_N, \quad \begin{cases} a_U = 0, a_N = 1 & \|s_N\| \leq \Delta, \\ a_U = \Delta / \|s_U\|, a_N = 0 & \|s_U\| \geq \Delta, \\ a_U = 1 - \tau, a_N = \tau & \text{otherwise,} \end{cases} \quad (113)$$

with τ the root of $\|s_U + \tau(s_N - s_U)\| = \Delta$ in $[0, 1]$. Returning the two coefficients rather than the vector is deliberate: the image As follows from As_U and As_N by linearity and costs no further matrix-free product. The Krylov leg is solved without the pseudo-transient shift, $As_N = -r_0$ rather than $(I/\Delta\tau + A)s_N = -r_0$, because the dogleg falls monotonically along its path only when that leg is the model's own minimizer. A species unknown also carries a box, the fraction-to-the-boundary rule holding the step inside the physical range of a mass fraction, and an element with no reservoir is excluded from the registry altogether: its unknown would sit exactly on its own lower bound with no room below, and the boundary rule then returns zero for any direction with a component pointing there, so the whole Krylov direction would be refused.

The ledger and the stop. The best iterate reached so far is kept, with its composition, its flux spread and whether it met the gates. The non-monotone search accepts sideways steps, which is what lets the solve leave a bad neighborhood and also what lets it circle, so when the best iterate has not improved for 20 outer iterations the solve returns to it and switches to a monotone Armijo acceptance, and if that buys nothing in another 20 it stops and reports that it stagnated, a different statement from “no descent direction exists”. The loop-top stop requires the certified rows of Sec. 9.3 and not only the merit: the two are different functionals, and a window on the merit alone bounds nothing about the quantity the state is judged by. The return flag is 0 when the returned state met the acceptance gate and every judged row, 1 at the iteration limit, 2 when the solve ended without a state meeting the gate, and 3 at an iterate where no feasible descent direction exists. A flag of 2 is not a solver failure: the solve converged and the certification refused the state it hands back, naming the row, and that state is written with the refusal recorded rather than marched on.

9.5 Pseudo-transient continuation

`solve_steady_ptc` is the same residual solved with the banded Jacobian used as the *operator* and not only as a preconditioner: per iteration $(I/\Delta\tau + J)\delta Y = -F$ is solved by LAPACK banded LU, the step is taken with a backtracking line search and a positivity test, and $\Delta\tau$ follows the switched-evolution-relaxation ramp $\Delta\tau \leftarrow \Delta\tau \|R\|_{\text{old}}/\|R\|_{\text{new}}$, cut on a failed step. As $\Delta\tau \rightarrow \infty$ this is Newton's method; at small $\Delta\tau$ it behaves like explicit relaxation, which is the robust startup the continuation exists for. The route is not available with a carrier unknown: it builds its Jacobian from the frozen residual, which by construction performs no equilibrium sweep and therefore has no carrier row, and giving it one would mean assembling that row from a background nothing in the routine refreshes. The caller is told so rather than being handed a silently three-row answer.

The starting $\Delta\tau$ matters: the marching handoff starts the continuation from $\Delta\tau_0 = 1$ and the two direct routes from the CFL step.

9.6 The partitioned outer iteration

A species the run moves by an operator-split transport step instead of by a Newton row, which is the diffusing elements always and the molecular carriers whenever `Coupled carrier solve` is off, has a stationary balance no hydrodynamic solve touches; a single steady solve would freeze it at whatever state it was handed. `steady_wind_with_element_diffusion` in `EXHALE_main.f90` alternates the two: solve the wind at fixed composition, relax the composition to its own steady state in that wind, solve again. The composition relaxations are the fixed-wind element relaxation of `binary_element_diffusion.f90` (Sec. 6.9), under-relaxed with a factor ω starting at 0.5, and the fixed-wind carrier relaxation of `diffusive_photochemistry.f90` (Sec. 6.5), bounded by a movement trust rather than damped. The advecting flux handed to the element operator is the face mass flux of the state just solved, read from the row terms that state assembled, so the operator is given its wind and does not reach into the residual for one.

What is accepted is only a state on which the certification of the *full* set of active equations passes, evaluated on the refreshed state the routine would hand back. A hydrodynamic solve returning 0 while an alternated species row refuses is not an accepted state: that flag is a statement about the rows in the Newton registry, and the alternated rows stand outside it by construction. Conversely a nonzero hydrodynamic flag does not end the loop by itself, because the composition update that follows it still makes progress.

The quantity the alternation waits on is the worst gated species row of the certification, judged in the wind. Where it fails to fall over a pass, both updates are shortened, ω halved down to a floor of 0.125 and the carrier movement bound halved down to 10^{-3} , and three consecutive passes without a fall end the loop rather than spend the budget on a fixed point the alternation is not approaching. The endings are named: the joint certification passes; a row of the state is not finite; the element composition update is refused; the joint measure stops falling; the pass budget ends uncertified.

Two rules keep the state self-consistent. The progress control stands *ahead* of the composition update, so the pass that ends the iteration takes no update and what is handed back is the very state the certification of that pass was evaluated on; and the last pass of the budget takes no update either, an update existing only to be consumed by the next hydrodynamic solve. Taking it would replace a stationary wind by a state whose hydrodynamic rows the update itself spoiled, by eleven decades in the mass row on a molecular configuration.

With `Coupled carrier solve: True` the alternation does not arise for the carriers at all: $n(\text{H}_2)$ becomes a fourth Newton unknown per cell, one solve returns a wind and a carrier partition that are steady states of each other, and the outer loop runs once. With no alternated species at all, which is what an atomic run and a fully coupled run both give, the body is one steady solve followed by the state refresh and the flag is the solve's own.

10. Output and post-processing

10.1 The advection-corrected profiles

For the reason given in Sec. 4.5, `post_process_adv.f90` re-solves the profile once, at the end of the run, with the ionization advected: the systems (55)–(57) are solved cell by cell from the base outward, each cell taking the upstream cell's solved fractions as its x^{up} , and the temperature is then re-solved on the steady internal-energy equation (72) with

the Brent bracket of Sec. 5.3.

Three properties of that pass are worth stating.

- The third term of (72), the enthalpy flux carried by the divergence of the mass flux, $hr^{-2}d(\rho vr^2)/dr$, vanishes for a stationary mass flux and for nothing else, so a correction built without it would describe a converged wind and no other state. The divergence is formed with the same operator the mass row of the state uses, $(A_p F_p - A_m F_m)/dV$ over the control volume whose two bounding points are the two points the upwind energy difference is taken between, so the term is an exact zero wherever those two carry the same ρvr^2 .
- A cell is corrected only where the correction means something. Above the Damköhler number 10^2 , formed with the *slowest* relaxing species of the solved system, the gas reaches local equilibrium many times over while it crosses the cell and the equilibrium solution already solves the advection equation, so the cell is pinned to it. Below an equilibrium ion fraction of 10^{-6} the residual, which carries the neutral fraction, cannot represent the ion density to better than about a percent. Where the mass row of the supplied state exceeds `adv_conditional_tol` as a fraction of its own largest term, the cell keeps the run's own value; each cell's verdict travels in the output file as `adv_T_status`, `adv_comp_status` and `adv_mass_row`.
- The reconstruction treats the gas as H/He plus trace metals and *excludes* the molecular species: H_2 , H_2^+ , H_3^+ and HeH^+ are not passed in and are not re-solved, so inside a molecular layer the `_adv` particle count omits the neutral H_2 and the molecular-ion electrons. The output file says so in its own header rather than only in the source.

The first of those gates has a fourth companion: wherever $v \leq 0$ on either face, the cell keeps the converged equilibrium ionization and temperature, because the upwind discretization takes the upstream state from the cell below, which is not the upstream cell when the gas moves inward, and the residence time $\Delta r/v$ is then negative as well. The relaxation rate entering the Damköhler number is the total rate at which each solved population is destroyed and re-formed, and the *minimum* over the species the system solves is taken, because $He(2^3S)$ relaxes orders of magnitude more slowly than hydrogen ($A_{31} = 1.272 \times 10^{-4} \text{ s}^{-1}$ sets the floor) and the vector is only as local as its slowest member.

10.2 The mass-loss rate

The quantity the run reports as `Log10` of `steady-state Mdot` is measured at one radius, in `EXHALE_main.f90`:

$$\dot{M} = 4\pi \rho(j) v(j) r(j)^2 n_0 v_0 m_H R_0^2 \quad \text{at} \quad j = \max(N-20, 1), \quad (114)$$

the offset of 20 cells keeping the evaluation far enough from the outer boundary, and the floor at $j = 1$ keeping it out of the base ghosts where the boundary closure and not the wind sets ρ and v . A logarithm is taken only where the flow there is directed outward: $\rho v \leq 0$ is an inflow, which is not a mass-loss rate, and the run then reports the signed flux instead of a logarithm. The dayside convention of Table 1 is applied here and only here: $\log_{10} \dot{M} \rightarrow \log_{10} \dot{M} - \log_{10} 2$ for `Rate/2 + Mdot/2`, and $-\log_{10} 4$ for `Mdot/4`.

A *quantitative* $\dot{M}$ must come from a state the steady solver has finished. A run stopped on the `du` threshold (100) alone carries a path-dependent spread of several percent in (114); the two-stage marching of Sec. 7.2 followed by the stationary solve of Sec. 9.4 is the working recipe, and it is what the regression matrix uses.

10.3 Output files

Every file written to output/ carries a schema header, so a reader adapts instead of counting columns. Hydro_ioniz.txt opens with # EXHALE schema 2 and then

```
# columns r[Rp] rho[mH/cm3] v[cm/s] p[cgs] T[K] heat[erg/cm3/s] cool[erg/cm3/s]
```

(with three further status columns on the _adv file). Column 2 is ρn_0 , the *mass* density in units of m_H per cm^3 with the metals included under the eos_metals policy, not a number density. The first and last Ng rows of every file are ghost cells filled by the boundary conditions, and their $\rho v r^2$ is a boundary extrapolation. Ion_species.txt carries the species fractions in the species_table order of Sec. 4.1; Heating_breakdown.txt and Cooling_breakdown.txt carry the channel decompositions of Sec. 5., each built from the array the energy equation itself used, so the channel sum and the heat or cool column of Hydro_ioniz.txt are the same number to round-off; Excited_H.txt, Lyman_Werner.txt, FUV_bands.txt, Oxygen_chemistry.txt and OI_levels.txt carry the optional physics. The Python loader examples/exhale_io.py reads all of them through those headers (exhale_io.Run('output/') gives .r, .T, .Mdot and the rest), and it should be used in place of any new column parser.

10.4 The transmission spectrum

EXHALE_transit.py, with the library exhale_transit_lib.py, builds transit transmission spectra from the _adv profiles: the He I 10830 Å triplet, Ly α , H α and H β , and the metal doublets Mg II h&k, Ca II H&K and Na I D, the metal lines being skipped automatically for a metals-off run. It integrates the Voigt line opacity over impact parameter, averages over the stellar disk, and convolves with the instrument profile and the planet's rotation; the Balmer branch builds the non-LTE $2s/2p$ populations following Christie et al. (2013), the same model Sec. 4.6 solves inside the Fortran code. An optional triaxial Roche geometry comes from roche_recon.py. Overrides are environment variables (EXHALE_TRANSIT_*: resolving power, He window, rotation period, azimuthal samples) and the output keeps the historical tpm_*.txt names, each with its own # transit_schema 1 metadata block. The physics of that tool is documented separately in docs/transmission_spectrum.pdf and is not repeated here.

11. The input keys that select physics

Runtime configuration is by the presence of a key, not by a compile flag (Sec. 1.1). Three companion files are read at startup and their mere presence is itself a switch: metals.inp turns the trace metals on, opacity.inp the tabulated continuum opacities, and base.inp the lower-atmosphere handoff.

Table 2 lists the keys that select physics or the numerical scheme, with the default the code takes when the key is absent. The defaults were read from parameters.f90 and the parsing from input_read.f90; keys that only name a file, a path or an output option are omitted. Four lines of the core block are mandatory and have no default: they are marked as such.

Table 2: Physics and algorithm keys of `input.inp`, with the value the code takes when the key is absent.

Key	Default	What it selects, and where
<i>Geometry, grid and the one-dimensional approximation</i>		
2D approximate method:	mandatory	Mdot/4, Rate/2, Rate/4 or alpha: the dayside dilution of every stellar beam and the factor on $\dot{M}$ (Table 1; <code>parameters.f90</code> , <code>utilities.f90</code>).
Grid type:	mandatory	Uniform, Mixed or Stretched (<code>define_grid.f90</code>).
Grid cells:	500	Number of physical cells; ghosts are added on top (<code>define_grid.f90</code>).
Base grid [dr,cells]:	2×10^{-4} , 50	Width and count of the uniform base region of the Mixed grid (<code>define_grid.f90</code>).
Domain mode:	Roche	Spherical drops the tidal and centrifugal terms of (12) (<code>grav_field.f90</code>).
Outer radius [R_p]:	L1 radius	Outer boundary; required in spherical mode (<code>input_read.f90</code>).
<i>Discretization</i>		
Numerical flux:	mandatory	HLLC, ROE or LLF (<code>Num_Fluxes.f90</code>); ROE is refused with the molecular ladder equation of state.
Reconstruction scheme:	mandatory	PLM, WENO3 or the two-stage PLM+WENO3 (<code>Reconstruction.f90</code>).
Reconstruction continuation:	off	$\Delta\lambda$ of the homotopy (89) between the two operators (<code>steady_residual.f90</code>).
CFL:	0.6	CFL number of (99) (<code>eval_dt.f90</code>).
Time stepping:	global	Local gives every cell its own pseudo-time step (<code>eval_dt.f90</code>).
Well balanced:	off	Carries the departure from the cell's own hydrostatic equilibrium instead of the state (<code>Reconstruction.f90</code> , <code>RK_rhs.f90</code>).
Low-Mach damping:	off, $M_{th} = 10^{-3}$	Gated fourth-difference dissipation (19) of the contact mode (<code>low_mach_dissipation.f90</code>).
Shapiro filter:	off, every 4 steps	1-2-1 low-pass filter on the marched state (<code>Apply_BC.f90</code>).
Caloric EOS:	ladder	monatomic forces $\gamma = 5/3$ everywhere (<code>caloric_eos.f90</code>).
Energy solver:	semi-implicit	Explicit reverts the source update to forward Euler (<code>energy_semi_implicit.f90</code>).
<i>Boundary and initial conditions</i>		

Table 2 continued

Key	Default	What it selects, and where
Base BC:	density	pressure <microbar> sets the base level from a pressure instead of a density (input_read.f90, base_boundary.f90).
IC mode:	cold	cold, transonic, hot_parker, auto or windae (set_IC.f90).
Transonic IC:	off	Older synonym of IC mode: transonic.
Hot Parker IC:	off, $T_{\text{wind}} = 10^4$ K	Older synonym of IC mode: hot_parker.
Restart intent:	relaxation	What a loaded state is continued as: trajectory, relaxation or stationary (load_IC.f90).
<i>Convergence and the stationary solvers</i>		
du_th [PLM,WENO3]:	10^{-3}	Threshold on the mass-flux spread (100); a second value makes the run two-stage (EXHALE_main.f90).
Escape radius [R_p]:	mandatory	Inner edge of the window (100) is taken over (define_grid.f90).
Resid tol:	off	Residual convergence instead of the flux spread; the solvers use 10^{-5} when it is absent (steady_residual.f90).
Flux spread tol:	2×10^{-5} , $r_{\text{flux}} = 1.2$	The flux gate of the acceptance test (steady_residual.f90).
Solver:	marching only	Newton hands over to the stationary solve once the spread falls below 10^{-2} (steady_newton.f90).
Stall:	10^{-6} , 2000 steps	Plateau detector of the marching stop (EXHALE_main.f90).
Level tol:	off	Mass-flux level stability required beside every marching stop.
Max steps:	10^6	Hard cap on marching iterations.
Newton solver:	on	False reverts the composition solve to MINPACK hybrd1 (newton_solver.f90).
Brent solver:	on	False reverts the post-process temperature solve (T_equation.f90).
<i>Atomic hydrogen and helium</i>		
Include He23S?	on	The He 2^3S metastable as a solved level (System_HeH_TR*.f90).
ATES_photoionization_rate	off	Reverts He I photoionization to the legacy ATES fit (cross_sec.f90).
Legacy_HHe_rates	off	Reverts H/He recombination and collisional ionization to the ATES fits (util_ion_eq.f90).

Table 2 continued

Key	Default	What it selects, and where
Atomic rate set:	default	Koskinen2022 swaps four atomic H/He rate coefficients (util_ion_eq.f90).
Secondary_ionization	on, staged	Secondary ionization by fast photoelectrons; Immediate applies it from step 0 (electron_energy_degradation.f90).
He_rec_coupling	on	He II recombination radiation ionizing H I (util_ion_eq.f90).
He_H_charge_exchange	on	The He/H charge-exchange pair (charge_exchange.f90).
Photoelectron heating:	excess	Whether a photoionization deposits $h\nu - E_{\text{th}}$ or a stated fraction of $h\nu$ (util_ion_eq.f90).
Deexc heat	on	Collisional de-excitation heating of H($n=2$) (excited_hydrogen.f90).
<i>Radiation field</i>		
Spectrum type:	mandatory	Load, Power-law, Planck or Monochromatic: the spectrum of the whole photon grid (sed_read.f90, J_inc.f90).
Stellar Teff: + Stellar radius:	0 (off)	Enable the diluted-blackbody Balmer continuum and the H($n=2$) model (excited_hydrogen.f90).
Stellar Ly α flux:	0	Stellar Ly- α at the orbit; also band B2 of the oxygen photolysis (lya_rt.f90).
Jlya escape-prob	parameterized	The Neufeld escape-probability Ly- α transfer (lya_rt.f90).
Jlya RT file:	none	Import a $J_{\text{Ly}\alpha}(r)$ profile computed elsewhere.
Ly α stellar halfwidth:	70 km/s	Half-width of the stellar Ly- α line.
Ly α stellar boost:	5.0	Bounded buildup of the trapped stellar beam.
Ly α absorbing bottom	reflecting	Closes the Ly- α domain from below with an absorber.
Stellar LW flux:	0	Lyman-Werner band flux driving H ₂ photodissociation (lyman_werner.f90).
Stellar FUV B3/B4 flux:	0	The two longer FUV bands of the oxygen photolysis (water_photolysis.f90).
<i>Molecules, metals and the lower atmosphere</i>		
metals.inp present	metals off	Trace metals solved in the same coupled system as H and He (System_HeH_metals.f90).
eos_metals	1	Metals in the bulk mass, electron and particle budget (composition.f90).
cno_cool	1 (CHIANTI)	0 reverts C/N/O line cooling to the legacy analytic fits (Cool_coeff.f90).

Table 2 continued

Key	Default	What it selects, and where
Molecular chemistry:	off	The $\text{H}_2/\text{H}_2^+/\text{H}_3^+/\text{HeH}^+$ network (System_HeH_mol.f90, mol_rates.f90).
Oxygen chemistry:	off	OH, H_2O and CO with their photolysis (oxygen_rates.f90, water_photolysis.f90).
Molecular carrier transport:	follows oxygen chemistry	Vertical transport of the molecular carriers (diffusive_photochemistry.f90).
Coupled carrier solve:	off	$n(\text{H}_2)$ as a fourth Newton unknown per cell instead of an alternation (steady_newton.f90).
Ionization transport:	off	H^+ carried with the flow as a fifth carrier.
Molecular base:	off	Removes the H nuclei bound into H_2 from the base particle count (15) (composition.f90).
Molecular reaction heat:	on	Deposits the energy the collisional molecular reactions release (molecular_reaction_heat.f90).
Molecular IR bands:	off	H_2 , H_2O and CO infrared line cooling (molecular_infrared_cooling.f90).
Base IR field:	off	The infrared coolants see the thermal radiation of the atmosphere below the base instead of vacuum.
Lower atmosphere:	none	analytic or vulcan lower-atmosphere pre-step (lower_column.f90).
Lower atmosphere profile:	none	The lower atmosphere handed over as a table over pressure (lower_atmosphere_profile.f90).
Lower column:	off	1-bar radius of the analytic column.

Diffusion and molecular transport

He_diffusion	off	He/H diffusive separation as a transported element (binary_element_diffusion.f90).
He_metal_diffusion	off	Extends the same operator to the trace metals.
He_Kzz	0	Constant eddy diffusion coefficient where no profile states one.
He_alphaT	0	Thermal-diffusion factor.
He_ambipolar	on	Ambipolar term of the element diffusion.
Viscosity:	off	Radial viscous force and its dissipation (20) (viscous_conduction.f90).
Conduction:	off	Heat conduction with $\kappa(T)$ of (21).

References

Abrahamsson, E., Krems, R. V., & Dalgarno, A. 2007, ApJ, 654, 1171

- Allgower, E. L., & Georg, K. 2003, *Introduction to Numerical Continuation Methods*, SIAM Classics in Applied Mathematics 45 (Philadelphia: SIAM)
- Backx, C., Wight, G. R., & Van der Wiel, M. J. 1976, *J. Phys. B*, 9, 315
- Badnell, N. R. 2006, *ApJS*, 167, 334
- Banks, P. M., & Kockarts, G. 1973, *Aeronomy* (New York: Academic Press)
- Barinovs, Ģ., van Hemert, M. C., Krems, R., & Dalgarno, A. 2005, *ApJ*, 620, 537
- Batten, P., Clarke, N., Lambert, C., & Causon, D. M. 1997, *SIAM J. Sci. Comput.*, 18, 1553
- Biassoni, F., Caldiroli, A., Gallo, E., et al. 2024, *A&A*, 682, A115
- Caldiroli, A., Haardt, F., Gallo, E., et al. 2021, *A&A*, 655, A30
- Cecchi-Pestellini, C., Ciaravella, A., Micela, G., & Penz, T. 2009, *A&A*, 496, 863
- Christie, D., Arras, P., & Li, Z.-Y. 2013, *ApJ*, 772, 144
- Chung, Y. M., Lee, E.-M., Masuoka, T., & Samson, J. A. R. 1993, *J. Chem. Phys.*, 99, 885
- Cohen, J. S., & Lane, N. F. 1971, *Chem. Phys. Lett.*, 10, 623
- Cohen, J. S., & Lane, N. F. 1977, *J. Chem. Phys.*, 66, 586
- Dalgarno, A., Yan, M., & Liu, W. 1999, *ApJS*, 125, 237
- Draine, B. T. 2011, *Physics of the Interstellar and Intergalactic Medium* (Princeton: Princeton Univ. Press)
- Draine, B. T., & Bertoldi, F. 1996, *ApJ*, 468, 269
- Drake, G. W., Victor, G. A., & Dalgarno, A. 1969, *Phys. Rev.*, 180, 25
- Einfeldt, B., Munz, C.-D., Roe, P. L., & Sjögreen, B. 1991, *J. Comput. Phys.*, 92, 273
- García Muñoz, A. 2025, *A&A*, 698, A199
- Gordon, S., & McBride, B. J. 1994, *Computer Program for Calculation of Complex Chemical Equilibrium Compositions and Applications*, NASA RP-1311
- Gottlieb, S., & Shu, C.-W. 1998, *Math. Comput.*, 67, 73
- Harrington, J. P. 1973, *MNRAS*, 162, 43
- Hu, X. Y., Adams, N. A., & Shu, C.-W. 2013, *J. Comput. Phys.*, 242, 169
- Huang, C., Koskinen, T., Lavvas, P., & Fossati, L. 2023, *ApJ*, 951, 123
- Hui, L., & Gnedin, N. Y. 1997, *MNRAS*, 292, 27

- Jameson, A., Schmidt, W., & Turkel, E. 1981, AIAA Paper 81-1259
- Käppeli, R., & Mishra, S. 2014, J. Comput. Phys., 259, 199
- Käppeli, R., & Mishra, S. 2016, A&A, 587, A94
- Knoll, D. A., & Keyes, D. E. 2004, J. Comput. Phys., 193, 357
- Koskinen, T. T., Harris, M. J., Yelle, R. V., & Lavvas, P. 2013, Icarus, 226, 1678
- Koskinen, T. T., Lavvas, P., Huang, C., et al. 2022, ApJ, 929, 52
- Krause, M. O. 1979, J. Phys. Chem. Ref. Data, 8, 307
- Kurganov, A., & Tadmor, E. 2000, J. Comput. Phys., 160, 241
- Locci, D., Cecchi-Pestellini, C., & Micela, G. 2022, PSJ, 3, 1
- Mao, J., & Kaastra, J. 2016, A&A, 587, A84
- Miller, S., Stallard, T., Tennyson, J., & Melin, H. 2013, J. Phys. Chem. A, 117, 9770
- Morgner, H., & Niehaus, A. 1979, J. Phys. B, 12, 1805
- Neufeld, D. A. 1990, ApJ, 350, 216
- Norcross, D. W. 1971, J. Phys. B, 4, 652
- Odert, P., Erkaev, N. V., Lammer, H., et al. 2020, A&A, 638, A49
- Oka, T., & Epp, E. 2004, ApJ, 613, 349
- Oklopčić, A., & Hirata, C. M. 2018, ApJ, 855, L11
- Perthame, B., & Shu, C.-W. 1996, Numer. Math., 73, 119
- Poinsot, T. J., & Lele, S. K. 1992, J. Comput. Phys., 101, 104
- Robinson, T. D., & Catling, D. C. 2012, ApJ, 757, 104
- Roe, P. L. 1981, J. Comput. Phys., 43, 357
- Roueff, E., Abgrall, H., Czachorowski, P., et al. 2019, A&A, 630, A58
- Rusanov, V. V. 1961, J. Comput. Math. Phys. USSR, 1, 267
- Salz, M., Czesla, S., Schneider, P. C., & Schmitt, J. H. M. M. 2016, A&A, 586, A75
- Samson, J. A. R., & Haddad, G. N. 1994, J. Opt. Soc. Am. B, 11, 277
- Sekiya, M., Nakazawa, K., & Hayashi, C. 1980, Prog. Theor. Phys., 64, 1968
- Shull, J. M., & van Steenberg, M. E. 1985, ApJ, 298, 268

- Taylor, J., Oklopčić, A., Rustamkulov, Z., et al. 2025, *ApJ*, 989, 68
- Thompson, K. W. 1987, *J. Comput. Phys.*, 68, 1
- Toro, E. F., Spruce, M., & Speares, W. 1994, *Shock Waves*, 4, 25
- Toro, E. F. 2009, *Riemann Solvers and Numerical Methods for Fluid Dynamics*, 3rd edn. (Berlin: Springer)
- van Hoof, P. A. M., Williams, R. J. R., Volk, K., et al. 2014, *MNRAS*, 444, 420
- Verner, D. A., & Yakovlev, D. G. 1995, *A&AS*, 109, 125
- Verner, D. A., Ferland, G. J., Korista, K. T., & Yakovlev, D. G. 1996, *ApJ*, 465, 487
- Visscher, C., Lidders, K., & Fegley, B., Jr. 2006, *ApJ*, 648, 1181
- Visser, R., van Dishoeck, E. F., & Black, J. H. 2009, *A&A*, 503, 323
- Voronov, G. S. 1997, *At. Data Nucl. Data Tables*, 65, 1
- Watson, A. J., Donahue, T. M., & Walker, J. C. G. 1981, *Icarus*, 48, 150
- Yamaleev, N. K., & Carpenter, M. H. 2009, *J. Comput. Phys.*, 228, 3025
- Yan, M., Sadeghpour, H. R., & Dalgarno, A. 1998, *ApJ*, 496, 1044
- Yan, P.-G., & Babb, J. F. 2023, *MNRAS*, 518, 6004
- Zhang, X., & Shu, C.-W. 2010a, *J. Comput. Phys.*, 229, 8918
- Zhang, X., & Shu, C.-W. 2010b, *J. Comput. Phys.*, 229, 3091