Quasi-Random Photon Launching in MoCHII: Implementation, Validation, and the Diffuse-Bin Fix

Abstract

MoCHII has a randomized quasi-Monte Carlo (RQMC) photon launch, the default since 2026-09-25 (`par%launch_sequence = 'sobol'`): Owen-scrambled Sobol coordinates replace the Mersenne Twister draws used to select a packet's source component, photon energy, direction, and entry surface, while every post-launch transport event (forced interaction, Henyey–Greenstein scattering, Russian roulette) stays pseudo-random. The Sobol point is addressed by the *global* photon number, so the launch set is independent of the MPI task count. On the fixed-state analytic attenuation gate the cell-wise $\Gamma_{\rm HI}$ error scales as $\sim 1/N$ rather than the Monte Carlo $1/\sqrt{N}$, giving effective photon gains of $29\times$ at 2^{17} packets and $222\times$ at 2^{20}, at unchanged wall time and with no significant bias (measured 2026-07-18, when the launch was added and the frequency was still sampled and transported as a discrete bin). The stage-2 source gates (multiple point sources, external field) improve in step, and a diffuse-field variant reduces the replicate scatter of the ionized volume by $1.73\times$. This work also uncovered and fixed an unrelated defect in the diffuse-field energy-to-bin mapping that had become active once threshold-aligned bins (`par%ion_align_edges`) became the default: a closed-form log-grid formula mis-binned recombination photons emitted at an ionization threshold, sending He I ground-recombination photons into the bin below the He I edge; the corrected binary-search mapping lowers the mean neutral-helium fraction on a diffuse Strömgren sphere by 50.5%. The design rationale is recorded in `md/QUASI_RANDOM_LAUNCH.md`.

Contents

1. Motivation and design **2**

2. Method **2**

2.1 Sobol point by global index . . . 2
2.2 Owen scrambling by hashing . . . 3
2.3 Fixed dimension layouts . . . 3
2.4 Common random numbers and replicates . . . 5

3. Implementation **5**

4. The diffuse bin-mapping fix **5**

5. Validation **6**

5.1 Generator gates (Q0/Q2) . . . 6
5.2 Regression of the Mersenne Twister path . . . 6
5.3 MPI-count independence . . . 7
5.4 Variance gate: analytic attenuation . . . 7
5.5 Full-physics Strömgren smoke . . . 7
5.6 Stage-2 source gates . . . 7

5.7 Diffuse variance 8

6. Usage guidance and limitations **8**

References **8**

1. Motivation and design

Unless dust scattering is switched on (`par%ion_dust_scatter`, default `.false.`), MoCHII's ionizing band has no scattering: given a packet's launch energy, position, and direction, the attenuation along its ray is integrated *analytically* (the zero-variance edge walk). Consequently the residual noise in the stellar radiation field comes almost entirely from the sampling of the launch variables — which source component, which photon energy, and which direction each packet takes — and *not* from the transport estimator. This is exactly the situation an RQMC launch is built for: a low-dimensional integral with a deterministic post-launch response, so a low-discrepancy launch set directly reduces the noise in global photoionization rates, ionized volume, and the iteration-to-iteration convergence measures without touching the transport.

The design is deliberately hybrid. Only the fixed-dimensional launch step uses scrambled Sobol coordinates; every conditional, variable-length draw after launch stays on the Mersenne Twister. Dust-scattering histories consume a variable number of uniforms whose meaning depends on the packet's previous events, so mapping them onto sequential Sobol dimensions would be both unstable and high-dimensional. Keeping them pseudo-random preserves the unbiased scattered field and confines the low-discrepancy structure to the launch cube, where it helps most.

Two properties of MoCHII make the launch cube favorable. First, the photon energy is the inverse of the source's continuous spectral CDF (`energy_sampler_mod`) at one stratified Sobol coordinate, so every energy interval — in particular every tally bin — of probability p receives pN packets to order one for $N = 2^m$, removing the binomial bin-count noise that plain Monte Carlo inherits. (When the launch was added the energy was a discrete bin index drawn from the band luminosity CDF; the continuous sampling replaced it on 2026-07-28, and the stratification argument is the same.) Second, the launch is at most seven-dimensional (nine for the diffuse field), well within the range where scrambled Sobol nets retain useful variance reduction even though AMR cell boundaries and ionization fronts make the integrand only piecewise smooth. No ideal $N^{-1}(\log N)^d$ rate is assumed; the gain is measured (Section 5.).

2. Method

2.1 Sobol point by global index

The generator (`src/qmc_mod.f90`) is self-contained; no external library is used. The unscrambled Sobol integer for the 0-based global photon index i and dimension d is the random-access XOR of direction numbers over the set bits of i,

$$x_d(i) = \bigoplus_{k\,:\,\text{bit } k \text{ of } i \text{ set}} V_d(k), \tag{1}$$

with 32-bit direction numbers $V_d(k)$. The cost is $O(\text{popcount}(i))$ integer XORs, with no stored state and no sequential dependence, so any packet's coordinates are computed directly from its global number. Dimension 1 is the van der Corput

sequence in base 2 ($V_1(k) = 2^{32-k}$); dimensions 2–12 use the Joe & Kuo (2008) `new-joe-kuo-6` primitive polynomials and initial direction integers, expanded by the Joe–Kuo recurrence. `QMC_MAXDIM` = 12 covers the seven-dimensional stellar superset and the nine-dimensional diffuse launch. The unscrambled coordinates reproduce `scipy.stats.qmc.Sobol` (which uses the same table) exactly.

2.2 Owen scrambling by hashing

A bare Sobol sequence has no ordinary statistical error estimate and can form coherent patterns with a Cartesian or octree grid, so the coordinates are Owen-scrambled. Full tree-based scrambling is unnecessary; MoCHII uses hash-based nested-uniform scrambling (Laine & Karras 2011; Burley 2020): reverse the 32 bits of x_d, apply a keyed Laine–Karras avalanche permutation (each bit is allowed to affect only less-significant bits), and reverse back. This is $O(1)$ per coordinate and preserves the net's equidistribution while producing statistically independent randomized replicates. The scrambled integer k maps to the open unit interval as

$$u = (k+\tfrac{1}{2})2^{-32} \in (0,1), \tag{2}$$

so every coordinate is strictly interior, removing a failure class for any later mapping that takes a logarithm.

Three decorrelated scramble streams share the one Sobol net: the stellar launch (`QMC_STREAM_STELLAR`), the diffuse launch (`QMC_STREAM_DIFFUSE`), and the FUV stratum of the stellar launch (`QMC_STREAM_FUV`), used when `par%add_fuv` runs with `par%fuv_sampling = 'separate'`: the ionizing packets then keep the stellar-stream indices $0 \ldots N-1$ they have without the FUV extension, and the FUV packets are a prefix of their own stream, indexed from 0 and balanced at every power of 2 of `no_photons_fuv`. A distinct 32-bit key for each dimension and stream is derived deterministically from `par%qmc_seed` by an integer avalanche hash, so the streams behave as independent replicates while all remain balanced scrambles of the same net; the stellar and diffuse keys are the ones they had before the third stream was added.

2.3 Fixed dimension layouts

The meaning of every dimension is fixed and never reassigned conditionally: a component-dependent reassignment would shift all later coordinates and destroy the intended low-dimensional projections. A point component simply leaves its surface dimensions unused; unused-but-fixed dimensions are harmless because the projections onto the consumed subset remain those of the scrambled net.

A single internal point source (stage 1) uses the three-dimensional layout of Table 1; a single plane-parallel slab source uses six dimensions (energy, entry face, two in-face coordinates, μ, ϕ); every other source configuration uses the seven-dimension superset of Table 2; diffuse recombination packets use the nine-dimension layout of Table 3 on the separate diffuse stream.

The five diffuse channels are the ground captures onto H I, He I and He II, the He I excited cascade, and the He III cascade (`gen_diffuse_photon_qmc`). For the three ground channels dimension 8 inverts the energy distribution of `par%diffuse_energy_model` (default `'milne'`, the Milne free-bound spectrum; `'exponential'` is the $E_{\rm th}+kT\,\mathrm{Exp}(1)$ draw the launch used when it was added); for the He I cascade it picks the decay branch and dimension 9 draws the two-photon energy from the tabulated Drake shape; the He III cascade reuses dimensions 8 and 9 for its path pick (free-bound, Lyα, two-photon) and energy, so it adds no dimension.

Table 1: Single internal point source (`QMC_NDIM_USED` = 3).

dim	variable	mapping
1	photon energy	inverse of the continuous spectral CDF
2	polar direction	$\mu = 2u - 1$
3	azimuth	$\phi = 2\pi u$

Table 2: Superset layout for every configuration other than a single point or a single slab source (multiple point sources, external field, or a mixed set).

dim	variable	mapping
1	source component	component luminosity CDF
2	photon energy	that component's continuous spectral CDF
3	polar / incidence	point: $\mu = 2u - 1$; external: Lambert $\mu = \sqrt{u}$
4	azimuth	$\phi = 2\pi u$
5	entry surface	rectangular entry face / sphere entry $\cos\theta$
6	first surface coordinate	in-face coordinate / sphere entry azimuth
7	second surface coordinate	in-face coordinate (unused for the sphere)

Table 3: Diffuse recombination packets (`par%recombination_continuum = 'transported'`), on the diffuse stream.

dim	variable	mapping
1	emitting leaf	leaf luminosity CDF (binary search)
2–4	position in leaf	uniform, $2u - 1$ per axis
5	polar direction	$\mu = 2u - 1$
6	azimuth	$\phi = 2\pi u$
7	emission channel	channel CDF over the five channels (see text)
8	first energy variate	ground-continuum energy; He I or He III cascade path pick
9	second energy variate	He I two-photon or He III cascade energy (else unused)

2.4 Common random numbers and replicates

Using the same scrambled launch set at every ionization/thermal iteration is a form of common random numbers: it correlates the radiation estimates before and after an opacity update and reduces noise in their difference, which stabilizes the convergence measures and separates front motion from sampling noise. The scramble key is `par%qmc_seed`, held fixed through all nonlinear iterations of one solve. A fixed scramble can, however, imprint a repeatable quadrature pattern that the nonlinear solution adapts to, making iteration differences look quieter than the true sampling uncertainty; convergence under one scramble is therefore not a substitute for replicate agreement. For a production error estimate, run independent replicates with different `qmc_seed` values — at least four, more when cost permits — and take their scatter as the RQMC uncertainty.

3. Implementation

The generator lives in `src/qmc_mod.f90`; the launch mappings that consume its coordinates are `gen_ion_photon_qmc` in `src/ion_band_mod.f90` (stellar) and `gen_diffuse_photon_qmc` in `src/diffuse_mod.f90` (diffuse). Two parameters control the option:

- `par%launch_sequence = 'sobol'` (the default since 2026-09-25) or `'random'` (the Mersenne Twister MT19937-64 path, the default before then and pinned by the inputs written before then);
- `par%qmc_seed` (default `12345`) — the scramble/replicate key, fixed across the gas iterations.

The stellar loop in `src/ionizing_field_mod.f90` addresses the Sobol point by the global photon number $i-1$ (0-based), and the diffuse loop does the same on its own stream indexed by the global diffuse photon number. Because the coordinates depend only on the global index and not on which rank handles a packet, the launch *set* is identical for any MPI task count, and neither schedule offered by `par%use_master_slave` — the static cyclic split or the batches rank 0 serves on demand — changes the sample.

Everything after launch stays pseudo-random: forced first interaction, Henyey–Greenstein direction sampling, subsequent optical depths, and Russian roulette all draw from the rank-local Mersenne Twister exactly as before. Bitwise MPI-count independence of a *full* run is therefore obtained only in absorption-only transport, where the post-launch path is deterministic; with dust scattering enabled the launch set is still reproducible but the scattered histories are not. The multi-wavelength dust/SED launch (`src/sed_mod.f90`) is not exercised by the ionizing calculation and is left on its alias sampler; its inverse-CDF conversion is documented as unnecessary for now.

4. The diffuse bin-mapping fix

This work uncovered a defect independent of QMC, in the diffuse-field packet generator, and fixes it.

The bug. `gen_diffuse_photon` mapped a sampled photon energy to its band bin with a closed-form expression that assumed the legacy log-uniform bin grid. Once threshold-aligned bins (`par%ion_align_edges`) became the default, the

bin edges sit *on* the ionization thresholds, and the closed form mis-assigns an energy at a threshold: a 24.587 eV He I ground-recombination photon landed in the bin *below* the He I edge, where it could never ionize helium.

The fix. The band edge array is now stored once (`ion_eedge`, valid for both grids), and `ion_bin_of(eph)` finds the bin by binary search: `ion_eedge`$(b) \leq E_{\rm ph} <$ `ion_eedge`$(b{+}1)$, so an energy exactly on a threshold edge falls in the bin *above* it — the physically correct side for a recombination photon emitted at threshold$+\delta$. Both the pseudo-random and the QMC diffuse generators call `ion_bin_of`. Since the continuous-energy transport of 2026-07-28 a packet carries the energy it was drawn with (`photon%energy_eV`), and its cross sections are evaluated there (`ion_packet_mod`); the bin is only the tally index it lands in, so the mis-binning described here could no longer keep a photon from ionizing helium.

Measured impact (2026-07-18, binned transport). On a converged diffuse-field Strömgren sphere on the aligned (default) grid, buggy versus fixed: the mean $x_{\rm HeI}$ drops 50.5% (0.0147 $\rightarrow$ 0.0073), the mean $x_{\rm HeII}$ rises 0.75%, and $V_{\rm ion}/R_{\rm eff}$ are unchanged (hydrogen is stellar-dominated). Unit checks: on the legacy log grid the binary search reproduces the closed form over a two-million-point energy scan except at a single exact-edge floating-point tie; on the aligned grid the H I/He I/He II thresholds are bin edges and threshold$+\delta$ lands in the bin above. Runs without the transported continuum are arithmetically untouched, and the recorded diffuse validations in the project log predate the aligned-edge default, so the bug had contaminated no recorded gate.

5. Validation

Every number below was measured and independently verified on 2026-07-18, when the launch was added: `'random'` was then the default, the photon frequency was sampled and transported as a discrete bin (the continuous photon-energy sampling came on 2026-07-28), and the diffuse ground continua used the exponential energy draw. They have not been remeasured on the current code.

5.1 Generator gates (Q0/Q2)

The unscrambled Sobol coordinates for dimensions 1–12 match `scipy.stats.qmc.Sobol(d=12, scramble=False)` exactly (sorted integer lattices, maximum code difference 0). The Owen-scrambled sets are strictly interior; the one-dimensional dyadic counts are exact at every level; the dimension-$(1,2)$ $(0,m,2)$-net and the elementary-interval occupancy are preserved under scrambling, for seeds `12345/777/2024` and both streams (stellar and diffuse; the FUV stream came later). Different seeds give different but individually balanced nets (maximum coordinate shift $|\Delta u| = 0.84$–0.98), and the stellar and diffuse streams differ like independent replicates.

5.2 Regression of the Mersenne Twister path

With `launch_sequence='random'`, at one MPI task under `-fp-model precise`, the pre-change and post-change builds are *bit-identical* for single-source, multi-source, and external inputs (worst $|\Delta| = 0$). The diffuse log-grid input sits at the code-generation rounding floor (maximum relative difference 6.8×10^{-16}, with the bins proven identical). At more than one task the difference equals the MPI `ALLREDUCE` reordering floor ($\sim$ 1 ULP, the same as running one binary twice).

5.3 MPI-count independence

The Sobol launch set is task-count independent: `'sobol'` rates at 4 tasks versus 8 tasks agree to 6.0×10^{-15} (stage 1) and 2.5×10^{-14} (stage 2) relative, whereas the `'random'` contrast differs at order unity (each rank owns a different Mersenne Twister stream).

5.4 Variance gate: analytic attenuation

The cleanest test is the fixed-state G0 gate (`tests/g0_gamma`): $\Gamma_{\rm HI}$ against the exact analytic attenuation, comparing four ordinary Monte Carlo replicates with four RQMC replicates. Table 4 reports the cell-wise and radial-profile RMS error and the effective photon-number gain (the number of Monte Carlo packets needed to reach the RQMC RMS).

Table 4: Fixed-state $\Gamma_{\rm HI}$ error, Monte Carlo (MC) versus RQMC, four replicates each. "Gain" is the effective photon-number gain.

	MC RMS	RQMC RMS	gain
$N = 2^{17}$ *packets*			
cell-wise	16.9%	3.1%	28.8×
radial profile	0.58%	0.13%	20.4×
$N = 2^{20}$ *packets*			
cell-wise	6.0%	0.40%	222×
radial profile	0.24%	0.12%	4.1×

The RQMC cell-wise error scales as $\sim 1/N$ (Monte Carlo $\sim 1/\sqrt{N}$), so the gain grows with N. The RQMC bias is $\sim 0.06\%$, stable, and smaller than the Monte Carlo bias (0.12–0.41%): no significant bias. One caveat is reported honestly: at 2^{20} the RQMC radial-profile error saturates at a systematic floor (all four replicates give an identical 0.119%, with a fixed −0.06% bias) that is band-discretization systematics, not sampling noise, so the 4.1× profile gain there is a lower bound. Sequence generation is a few tens of integer operations per packet, negligible against ray traversal, so the wall time is unchanged.

5.5 Full-physics Strömgren smoke

On a full-physics Strömgren run, $R_{\rm eff}$ under `'sobol'` differs from `'random'` by –0.039% (within noise). The launch-noise reduction shows up directly in the convergence measure: the `'sobol'` run reached the volume-integrated criterion in 57 iterations where the `'random'` run hit the 60-iteration cap unconverged. The $x_{\rm HI}$ slices are clean — no angular spokes and no grid-aligned artifacts (inspected).

5.6 Stage-2 source gates

The existing stage-2 analytic checkers, rerun under `'sobol'` at their recorded criteria, improve or match the `'random'` numbers (Table 5).

Table 5: Stage-2 source gates under `'sobol'` versus the recorded `'random'` values.

gate	`'sobol'`	`'random'`
two-point superposition (median)	0.14%	0.76%
two-point superposition (mean ratio)	0.9989	—
external rec, interior $\langle J\rangle/J$	0.9984	0.9981
external sph, interior $\langle J\rangle/J$	0.9990	0.9991
mixed point+external (median)	0.13%	0.75%

5.7 Diffuse variance

On a diffuse-field Strömgren sphere from a neutral start, the replicate scatter (standard deviation over three seeds) of $V_{\rm ion}$ and $R_{\rm eff}$ is 1.73× smaller for RQMC than for Monte Carlo, and unbiased. This is a smaller gain than the stellar launch, as the design anticipated: the emitting-leaf CDF is strongly discontinuous and changes as the gas state iterates, and the diffuse cube is higher-dimensional.

6. Usage guidance and limitations

`par%launch_sequence = 'sobol'` (the default) selects the RQMC launch; prefer a power-of-two photon count, `par%no_photons` = 2^m, where the Sobol net has its cleanest balance. Keep `par%qmc_seed` fixed across the nonlinear iterations of one solve (common random numbers), and change it only to generate independent replicates; take the scatter of at least four replicates as the RQMC error estimate. The inputs recorded before 2026-09-25 pin `'random'`, so their gates reproduce.

The gains follow the design expectation. The largest reductions are for global, direct-field quantities: direct stellar photoionization rates, integrated ionized volume and effective Strömgren radius, volume-integrated convergence measures, smooth radial profiles, and early iterations run with a modest photon budget. Smaller or inconsistent gains are expected for the maximum error in a single I-front cell, narrow shadow boundaries and clump skins, highly localized line emissivities, high-resolution image pixels, scattering-dominated paths, and diffuse-field-dominated cells. Low discrepancy does not remove physical discretization errors — frequency binning, AMR resolution, atomic-data fits, and nonlinear stopping tolerances remain independent error sources — and no universal speedup factor should be assumed; a useful reduction is best reported as an effective photon-number gain. By design, variable-length scattering histories remain on the Mersenne Twister; extending RQMC into them would need a maximum-order dimension policy and a separate validation campaign.

References

- Sobol', I. M. 1967, *USSR Comput. Math. Math. Phys.*, 7, 86.
- Owen, A. B. 1997, *SIAM J. Numer. Anal.*, 34, 1884; 1998, *ACM Trans. Model. Comput. Simul.*, 8, 71 (nested-uniform scrambling).

- Joe, S., & Kuo, F. Y. 2008, *SIAM J. Sci. Comput.*, 30, 2635 (`new-joe-kuo-6` direction numbers).
- Laine, S., & Karras, T. 2011, in *Proc. High-Performance Graphics* (stratified sampling for stochastic transparency).
- Burley, B. 2020, "Practical Hash-based Owen Scrambling," *J. Comput. Graph. Tech.*, 9, 1.
- Seon, K.-I. 2026, `md/QUASI_RANDOM_LAUNCH.md` (MoCHII design document).