

MoCHII User Guide

Abstract

How to build and run MoCHII (MOnte Carlo for HII regions): requirements, the complete input-parameter reference, output-file formats, worked examples keyed to the regression tests, and the runbook for adding an ion. The physics and validation are documented separately in docs/MoCHII_physics.pdf; the atomic-data fitting pipeline in docs/MoCHII_fitting.pdf.

Contents

1. Requirements and build	1
2. Input parameters	2
2.1 Photons and RNG	2
2.2 Grid and dust density	3
2.3 Ionizing/FUV band and the source	3
2.4 Initial state and iteration	4
2.5 Thermal balance and metals	5
2.6 I-front re-refinement (G4)	6
2.7 Dust in the band and dust emission	6
2.8 Peel-off imaging	7
2.9 Output	7
3. Output files	7
3.1 <base>_rates.h5	7
3.2 Text outputs	8
3.3 Reading outputs in Python	8
4. Worked examples (the regression tests)	9
5. Adding an ion (the G5 runbook)	10
6. Practical notes	10

1. Requirements and build

- Intel oneAPI MPI Fortran (mpiifort/mpiifx; GNU mpif90 supported by the Makefile), HDF5 (default prefix /data/opt/hdf5_intel; override with make HDF5_PREFIX=...), CFITSIO.
- The SEDust dust-emission library: SEDust_lib/ in this tree holds libsedust.a and its .mod files (copied from the MoCafe v2.00 build). Only needed at link time; the SEDust *data* directory is referenced at run time (Section 2.7).

- Python 3 with numpy/scipy (h5py, astropy, matplotlib, PyNeb for the test and pipeline scripts).

Build:

```
cd MoCHII
make          # -> MoCHII.x   (always a full clean rebuild)
make DEBUG=1  # bounds/traceback build
```

The Makefile intentionally performs a full rebuild every time (the MoCafe policy): there is no inter-module dependency tracking, and stale `.mod` files after editing `define.f90` otherwise cause runtime namelist errors.

Run:

```
mpirun -np 8 ./MoCHII.x input.in
```

MoCHII is MPI-only; the octree and all leaf arrays live in MPI-3 shared memory (one copy on each node).

2. Input parameters

All input is one `¶meters` namelist assigning fields of `par`. Parameters not listed keep the defaults shown. Stage switches compose: `gas_niter = 0` gives a G0 single pass (rates only); adding `gas_niter > 0` iterates the ionization (G1); `solve_te` adds the thermal balance (G2); `use_metals` the trace-metal registry; `diffuse_field` the explicit diffuse field (G3, forces case A); `refine_front` the I-front re-refinement (G4); `ion_add_dust` the dust coupling.

2.1 Photons and RNG

parameter	default	meaning
no_photons	1e5	source packets in each iteration
no_print	1e7	progress-print stride
iseed	0	RNG seed (0 = from clock/pid)

2.2 Grid and dust density

parameter	default	meaning
grid_type	'car'	'amr' (octree) or 'uniform' (single-level leaf list stored in raster order; the shared walks step by integer index arithmetic instead of the neighbor table, no tree in memory; refine_front unavailable)
amr_file	"	generic AMR leaf file (FITS/HDF5/text): columns x, y, z, level, nH [, metallicity, xHI, ndust]
distance_unit	'kpc'	'kpc'/'pc'/'au'/'" (code units)
dust_model	'global_dgr'	leaf grey opacity: 'none', 'global_dgr' ($n_{\text{H}} c_{\text{ext}} \cdot \text{DGR}$), 'from_file' (ndust column), 'laursen09' (static xHI column), 'laursen09_live' (computed x_{HI} , refreshed each iteration)
cext_dust	1.6059e-21	reference extinction per H [cm^2] at λ_{ref}
DGR, Z_global, Z_ref	1e-2, 0.0134, 0.0134	dust-model scalings
f_ion_dust	0.01	laursen09 survival of dust in ionized gas
f_pah, f_ion_pah	0, 0	PAH share of the reference extinction and its own ionized-gas survival (laursen09_live)

2.3 Ionizing/FUV band and the source

parameter	default	meaning
use_ion_band	F	must be T (the MoCHII transport)
nnu_ion	16	log-spaced bins; gates use 32; raise when widening the band
eion_min, eion_max	13.598, 100	ionizing band edges [eV]
add_fuv	F	FUV extension: extra log bins on [efuv_min, eion_min] below the unchanged ionizing bins; transports FUV grain heating and FUV photoionization of low-threshold metals (Mg I, C I, S I, Fe I); luminosity stays the IONIZING luminosity (Q_{H} preserved, the FUV part rides on top)
efuv_min	6.0	FUV lower edge [eV]
nnu_fuv	8	FUV bins (16 recommended with metals)
tstar	-999	Planck source temperature [K]
ion_spectrum	"	alternative 2-column spectrum file (E [eV], L_E arb.)
luminosity	1.0	BAND luminosity [erg/s]
xs_point, ys_point, zs_point	0	point-source position (code units, recentered box)

2.4 Initial state and iteration

parameter	default	meaning
xHI_init	1.0	initial $n_{\text{HI}}/n_{\text{H}}$ (an xHI file column wins); an ionized start (1e-3) converges much faster than neutral
xHeI_init, xHeII_init	1, 0	initial He fractions
He_abund	0.1	$n_{\text{He}}/n_{\text{H}}$
gas_niter	0	iterations (0 = G0 rates-only pass)
gas_tol	1e-3	convergence tolerance on x_{HII} (measure set by conv_crit)
conv_crit	'cell'	'cell': cell-max tests ($\max \Delta x_{\text{HII}} $ — stalls at the front-cell Monte Carlo noise floor $\sim 2\text{--}4\text{e-}2$ at $4\text{e}7$ packets); 'vol': volume-integrated tests $\sum V \Delta x_{\text{HII}} / \sum V x_{\text{HII}} < \text{gas_tol}$ and $\sum V n_e \Delta T_e / \sum V n_e T_e < \text{gas_tol_te}$ (recommended; noise-robust, and the n_e weight removes vacuum and no-heating cells). The vol measures floor at their own Monte Carlo level ($\sim 3\text{e-}3$ and $7\text{e-}3$ at $8\text{e}6$ packets, scaling as $1/\sqrt{N}$) — set the tolerances just above it
ion_relax	1.0	under-relaxation on x (< 1 damps front oscillation)
case_ab	'B'	case A/B recombination ('A' forced by the diffuse field)

2.5 Thermal balance and metals

parameter	default	meaning
solve_te	F	bisection on T_e coupled with the ionization
te_fixed	1e4	fixed T_e [K] when solve_te is off
te_min, te_max	1e3, 5e4	bisection bracket [K]
gas_tol_te	1e-3	convergence on $\max \Delta T_e /T_e$
atomic_dir	'data/atomic'	coefficient files (relative to the run directory)
use_metals	F	registry cascade + Tier-1 metal cooling
abund_C/N/O/Ne/S	HII20 values	$n(X)/n(H)$; an element is active when > 0
abund_Ar/Mg/Fe	0	further registry elements (gas-phase values; Mg/Fe are heavily depleted)
abund_Si/Cl/Ca	0	five-stage registry elements (gas-phase; Si and especially Ca are heavily depleted, Cl nearly undepleted — Orion-like values $3e-6 / 3e-7 / 2e-8$); adds [Si II] 34.8 μm , [Si III] 1883/1892, [Si IV] 1394/1403, [Cl II], the [Cl III] 5518/5538 density pair, [Cl IV], [Ca II], [Ca V]
ion_metal_abs	T	metal photoionization absorption in the band opacity (active with use_metals); the only gas opacity in the FUV bins
emis_output	F	write _emis (HDF5/FITS): leaf-by-leaf line emissivities + the state needed for maps (Section 3.)
diffuse_field	F	explicit ground-recombination packets (G3)
hei_diffuse	F	fourth diffuse channel (needs diffuse_field): He I excited-level recombination photons (19.82 eV, 584 Å, two-photon) — correct energy bookkeeping; moves Te by $\sim 1\%$ at HII40, He split unchanged
gaunt_vh14	F	free-free Gaunt factors from the van Hoof et al. (2014) table (data/gauntff_vh14.dat) instead of Hummer (1988); the two agree to $\lesssim 0.5\%$ in T_e
metal_ne	F	PDR: electrons from the metal cascade join the n_e closure (the only electrons beyond the I-front)
metal_heat	F	PDR: metal photoheating in the thermal balance
grain_pe	F	PDR thermal package: grain photoelectric heating + grain recombination cooling (Bakes & Tielens 1994) from the local FUV field (needs add_fuv), plus H-impact [C II] 158 μm / [O I] 63 μm cooling; grain terms scaled by $pe_scale \times$ the leaf dust amount relative to MW (from the leaf opacity)
pe_scale	1.0	photoelectric-efficiency scale factor

2.6 I-front re-refinement (G4)

parameter	default	meaning
refine_front	F	rebuild the octree on the I-front once
refine_iter	20	at which iteration
refine_lbase, refine_lmax	5, 7	base/front levels
refine_eps	1e-2	front flag: $\epsilon < x_{\text{HI}} < 1-\epsilon$
refine_dx	0.2	or a face-neighbor x_{HI} jump above this

2.7 Dust in the band and dust emission

parameter	default	meaning
ion_add_dust	F	dust absorption in the band (dusty Strömgren)
ion_dust_kext	"	kext table with EUV coverage (e.g. data/kext_albedo_WD_MW_3.1_60_D03.all_2009); required
ion_dust_scatter	F	sample dust scattering (HG, albedo weights)
dust_sed	F	SEDust stochastic dust SED at output time
dust_model_sed	'astrodust'	'astrodust'/'dl07'/'zubko'
sed_workdir	"	a SEDust sed/ tree (dielectric tables are read relative to it), e.g. the MoCafe copy .../MoCafe_v2.00/SEDust/sed
sed_qtable, sed_sizedist	MoCafe defaults	optics/size-distribution tables (relative to sed_workdir)
sed_NT, sed_Tlo, sed_Thi	200, 2.7, 5e3	SEDust temperature grid
dust_emis_bands	unset	up to 8 wavelengths [μm] (e.g. 8, 24, 70, 160): SEDust band emissivities of each leaf $\rightarrow$ _dustemis (blocks emis_dust/wl_dust [$\text{erg s}^{-1} \text{cm}^{-3} \mu\text{m}^{-1}$]); the Python map maker projects them into dust-band images (the IR is optically thin)
sed_pah_live	F	x_{HII} -dependent PAH survival in the emission: the leaf spectrum is assembled from the model channels with the PAH channel weighted by $(x_{\text{HI}} + f_{\text{ion,pah}} x_{\text{HII}})$ (needs a model with a 'PAH' channel, i.e. astrodust); turns the 8- μm map into the classic PAH ring at the ionization front

2.8 Peel-off imaging

parameter	default	meaning
ion_peel	F	after convergence, one extra transport pass peels direct (stellar + diffuse) and dust-scattered contributions toward the observers $\rightarrow$ <code>_image</code> (blocks <code>direct_ion/scatt_ion</code> , plus <code>_fuv</code> with <code>add_fuv</code>) [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]
peel_bins	F	bin-resolved cubes <code>direct_cube/scatt_cube</code> (n_x, n_y, n_ν) in addition to the band-integrated channel images (hardness maps, synthetic EUV/FUV photometry)
obsx, obsy, obsz	unset	observer positions (or directions) in code units; or use <code>alpha/beta/gamma</code> angle lists [deg]; default: one observer on +z far away
distance	auto	observer distance (code units)
nxim, nyim	129, 129	image pixels
dxim, dyim	auto	pixel scale [deg]; auto-sized to cover the box

2.9 Output

`out_file` (base name; extension follows `file_format = 'hdf5' or 'fits'`).

3. Output files

3.1 <base>_rates.h5

One dataset for each extension, leaf-ordered:

extension	content
Gamma_HI/HeI/HeII	photoionization rates [s^{-1}]
Heat_HI/HeI/HeII	photoheating per particle [erg/s]
Heat_dust	grain heating [$\text{erg s}^{-1} \text{cm}^{-3}$] (when <code>ion_add_dust</code>)
T_dust	equilibrium mixture dust temperature [K]
x_HI, x_HeI, x_HeII, n_e, T_e	converged state
x_<el>_stages	metal ion fractions (stage, leaf)
J_nu	band mean intensity (bin, leaf) [$\text{erg/s/cm}^2/\text{Hz/sr}$]
E_bin, dE_bin	band grid [eV]
LeafXYZ	leaf centers (code units)

3.2 Text outputs

file	content
_lines.txt	line luminosities: SH95 H I and He II recombination lines (1640, 4686, ... when an He III zone exists) + Tier-2 collisional lines; columns elem, stage, $\lambda[\text{\AA}]$, L , $L/L(\text{H}\beta)$
_emis.h5	(with emis_output) leaf-by-leaf line emissivities: emis_h (SH95 lines) and emis_<el>_<stage> blocks [$\text{erg s}^{-1} \text{cm}^{-3}$; $\sum \text{emis} \cdot V = L$, rows match _lines.txt], wl_* wavelengths [$\text{\AA}$], plus LeafXYZ, LeafSize, n_H, n_e, T_e, x_HI/HeI/HeII, x_<el>_stages — self-contained for line maps
_nebcont.txt	nebular continuum L_ν (fb+ff tables, Milne, two-photon)
_dustir.txt	equilibrium-mixture dust IR spectrum (1–1000 μm) with the energy-closure header
_dustsed.txt	SEDust stochastic dust SED with PAH bands (when dust_sed)
_dustemis.h5	(with dust_emis_bands) SEDust band emissivities of each leaf, same block layout as _emis — readable by EmisData for dust-band maps
_image.h5	(with ion_peel) peel-off images for each observer: direct_ion/scatt_ion (+_fuv) [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$], suffix _obs<k> when several observers

3.3 Reading outputs in Python

tools/python/mochii_output.py is an importable module (works directly in a notebook) that reads every section file (HDF5 or FITS; read_sections), the line table (read_lines_table), and the text spectra (read_spectrum), and builds 2D maps from the emissivity file:

```
import sys; sys.path.insert(0, ".../MoCHII/tools/python")
import matplotlib.pyplot as plt
from mochii_output import EmisData

d = EmisData("run_emis.h5")
d.info()                # blocks, fields, line count
d.line_list()           # every stored line (block, wl)

# ready-made figures (notebook): returns (fig, ax)
fig, ax = d.plot_line("o_3", 5007.)          # log S map
fig, ax = d.plot_line("h", 6563., unit="intensity",
                      photons=True)         # photons/s/cm^2/sr
fig, ax = d.plot_field("T_e", weight="EM")
fig, axs = plt.subplots(1, 2); d.plot_line("s_2", 6717., ax=axs[0]) \
    ; d.plot_line("s_2", 6731., ax=axs[1])    # doublet side by side
```

```
# raw arrays, when the figure is yours to build
img, ext = d.line_map("o_3", 5007., axis="z", npix=512)
img, ext = d.line_map("h", 6563., unit="intensity", photons=True)
img, ext = d.field_map("T_e", weight="EM") # LOS-weighted average
Q = d.line_luminosity("h", 4861., photons=True) # photon rate [1/s]
```

Line maps are flux-conserving: each leaf's luminosity is deposited with exact area overlap onto the pixel grid (AMR leaf sizes handled), so $\sum S A_{\text{pix}} = L_{\text{line}}$ to machine precision. `unit='flux'` (default) gives surface brightness [$\text{erg s}^{-1} \text{cm}^{-2}$]; `unit='intensity'` gives specific intensity [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$] = $S/4\pi$ (isotropic emission, optically thin); `photons=True` converts either to photon units (divides by hc/λ of the line). A `slab=(lo,hi)` cut restricts the line of sight. Field maps average `T_e`, `n_e`, `x_HI`, ... along the line of sight with weight '`EM`' ($n_e^2 V$), '`ne`', '`nH`', or '`V`'. The same module doubles as a command-line tool for quick looks:

```
python3 mochii_output.py run_emis.h5 # list lines
python3 mochii_output.py run_emis.h5 --line o_3 5007 \
    --npix 512 --log --out o3.png
python3 mochii_output.py run_emis.h5 --line h 6563 \
    --unit intensity --photons # photons/s/cm^2/sr
python3 mochii_output.py run_emis.h5 --field T_e --weight EM
```

4. Worked examples (the regression tests)

Each `tests/` directory is self-contained: grid builder or a symlinked grid, the input file, and a `check_*.py` gate script.

directory	what it demonstrates / gate
<code>g0_gamma</code>	rate integrals vs. analytic attenuation (bias +0.004%)
<code>g1_stromgren</code>	Strömgren sphere, case B fixed T_e ; AMR $\equiv$ uniform
<code>g2a_thermal</code>	thermal H/He nebula vs. the 1D exact reference
<code>g2_hii</code>	MOCASSIN Lexington HII20/HII40 benchmarks + line fluxes
<code>g4_refine</code>	I-front re-refinement vs. native level 7
<code>g4_tng</code>	Illustris-TNG post-processing demo (shadows, maps)
<code>d_dusty</code>	dusty Strömgren; scattering bracket; T_d/IR

A minimal thermal + metals + dust input (from `d_dusty`):

¶meters

```
par%no_photons = 4.0e7      par%iseed = 1234
par%grid_type = 'amr'      par%amr_file = 'amr_L6.fits'
par%distance_unit = 'pc'   par%dust_model = 'global_dgr'
par%use_ion_band = .true.  par%nnu_ion = 32
par%eion_min = 13.598     par%eion_max = 100.0
```

```

par%tstar = 4.0e4                par%luminosity = 3.178e38
par%xHI_init = 1.0e-3           par%He_abund = 0.1
par%gas_niter = 60              par%gas_tol = 2.0e-3
par%solve_te = .true.          par%atomic_dir = '../data/atomic'
par%use_metals = .true.
par%ion_add_dust = .true.
par%ion_dust_kext = '../data/kext_albedo_WD_MW_3.1_60_D03.all_2009'
par%cext_dust = 4.868e-22       par%DGR = 1.0
par%out_file = 'run.h5'         par%file_format = 'hdf5'
/

```

5. Adding an ion (the G5 runbook)

Adding an ion touches only data:

1. `tools/fitting/fit_cooling_tier1.py`: add the ion with a T_i ladder guessed from its `.elvlc` level energies; run; check the reported max error and the overlay figure.
2. `tools/fitting/fit_nlevel_tier2.py`: add the ion with the level count needed for its diagnostics; run.
3. `tools/fitting/make_element_data.py`: add the element to `ELEMENTS` (Z , stage count) and its thresholds; charge exchange entries where sources exist; run. The CHIANTI `.rrparams/.drparams` types 1–3 are handled (RR/RR2/DR/DR2 lines).
4. Fortran: a `par%abund_<El>` field and one entry in the `species_setup` name/abundance lists (only for a NEW element; new stages of a known element need nothing).
5. Verify: extend `verify_nlevel_pyneb.py` with a diagnostic ratio where PyNeb has data; run a smoke test and check the new lines in `_lines.txt`.

Data caveats met so far: Ar II has no CHIANTI v11 level/collision data (stage tracked without lines); Fe III collision strengths differ from PyNeb's BB14 by up to ~16% in 4658/4702 (known spread for iron); charge-exchange stage labels differ between the Huang (2023) and MOCASSIN transcriptions for Mg/Fe (the Huang labels are adopted; see `make_element_data.py`).

6. Practical notes

- **Bins**: 32 bins give ~2% rate-integral discretization; use 64 for sub-percent work. The `add_fuv` extension carries its own bins below `eion_min`, so the ionizing resolution is unaffected.
- **FUV and the PDR**: without `add_fuv` there is no radiation beyond the I-front and the metals there recombine toward neutral; with it, FUV penetrates the neutral gas (dust extinction only) and produces the PDR-side C II

and Mg II. Metal photoheating and grain photoelectric heating are not in the thermal balance, so PDR-zone temperatures are indicative only.

- **Convergence:** use `conv_crit = 'vol'` for production — the cell-max criterion stalls at the front-cell noise floor (and, with `add_fuv`, at no-heating cells where T_e pins to `te_min`). Both measures are printed each iteration. The G3 diffuse field smooths the front and converges far better.
- **Starts:** prefer an ionized initial state; a neutral start advances the front $\sim$ one cell per iteration.
- **text tables:** must cover the band (the astrodust table stops at 912 Å; use the D03 table for EUV work). Tables are sorted on read (the D03 file has an out-of-order seam).
- **Re-refinement:** one event per run; the old shared-memory windows are reclaimed only at exit.
- **SEDust:** `sed_workdir` must point at a SEDust `sed/` tree; the exact stochastic solve costs ~ 0.05 – 0.1 s for each gas leaf (distributed over ranks).