

Modern Astronomy

Part 1. Interstellar Medium (ISM)

Lecture 1

September 09 (Monday), 2024

updated on 09/02, 10:08

선광일 (Kwang-il Seon)
UST / KASI

Professors & Classroom

Professors:

- 1- 4th weeks: Prof. Seon, Kwang-II (선광일, kiseon@kasi.re.kr)
- 5- 8th weeks: Prof. Kim, Sang Chul (김상철, sckim@kasi.re.kr)
- 9-12th weeks: Prof. Lee, Sang-Sung (이상성, sslee@kasi.re.kr)
- 13-16th weeks: Prof. Shin, Jihey (신지혜, jhshin@kasi.re.kr)

Day & Time:

Monday 2-5PM

Classroom:

LWC(이원철홀): 220 (K.I. Seon / J. Shin)

JYS(장영실홀) : 222 (S.C. Kim), 331-2 (S.S. Lee)

Syllabus

Week	Date	
1	09-09	Introduction / Interstellar Medium (ISM)
2	09-09	Atomic Gas / Multiphase Medium
3	09-16	Molecular Clouds and Dust
4	09-23	Formation of Stars and Planets
5	09-30	Stars: The Hertzsprung-Russell Diagram
6	10-07	The Evolution of Stars
7	10-14	Star Deaths
8	10-21	The Milky Way Galaxy
9	10-28	Galaxies beyond the Milky Way
10	11-04	Hubble's Law and Distance Scale
11	11-11	Active Galaxies 1
12	11-18	Active Galaxies 2
13	11-25	General Relativity and Friedmann Equation
14	12-02	Evolution of Universe and Inflation Cosmology
15	12-09	Density Perturbations and Nonlinear Structure Formation
16	12-16	Survey and Computer Simulation

References for Part 1

- Introduction to the Interstellar Medium (Jonathan P. Williams)
- Fundamentals of Astrophysics (Stan Owocki)
- Interstellar Medium
 - ▶ Introductory
 - Introduction to the Interstellar Medium - Jonathan P. Williams
 - The Physics of the Interstellar Medium - J. E. Dyson & D.A. Williams
 - ▶ Intermediate
 - Interstellar and Intergalactic Medium - Barbara Ryden

Unit of distance

- **AU (the Astronomical Unit):** the average distance of the Earth from the Sun

$$\text{AU} = 1.50 \times 10^{11} \text{ m}$$

- ***parallax:*** As the Earth goes around the Sun, the nearby stars change their positions very slightly with respect to the faraway stars. This phenomenon is known as parallax. The angle θ is half of the angle by which this star appears to shift with the annual motion of the Earth and is defined to be the parallax.



- ***parsec (pc):*** the distance where the star has to be so that its parallax turns out to be $1''$.

As seen in January As seen in July

$$\begin{aligned}
 \text{pc} &= 3.09 \times 10^{16} \text{ m} \\
 &= 3.26 \text{ light years} \\
 &= 206,265 \text{ AU}
 \end{aligned}$$

Introduction to the Interstellar Medium

What is the ISM?

What is the ISM?

- The ISM is anything not in stars. (D. E. Osterbrock)
- Just what it says: The stuff between the stars in and around galaxies, especially our own Milky Way.
- It is made up almost entirely of gas with tiny (solid) particles called dust grains.
 - In addition to these, the ISM includes radiation, cosmic rays, and magnetic fields.

Why do we study the ISM?


- The ISM is the most beautiful component of galaxies. (B. T. Draine)
- The ISM is beautiful, both in the literal sense, as in images of colorful nebulae, and in the physics that helps us understand our origins and the way the Universe works. (J. P. Williams)
- The ISM is everywhere and it affects all sorts of observations, but more often as an essential complement for understanding the Galaxy.
- The ISM is the most important component of galaxies, for it is the ISM that is responsible for forming the stars that are the dominant sources of energy.

History of ISM Studies

the Cat's Eye Nebula (planetary nebula/ HST image)



The first nebula spectrum: the Cat's Eye Nebula (NGC 6543; W. Huggins, 1864)



- William Herschel resolved some nebulae into stars. He thought that he had discovered star birth (it actually ejected by a dying star). In the 1860s, William Huggins demonstrated that some nebulae have **emission line spectra**, rather than the **absorption line spectra**.

- **Hypothetical elements:**


Nebula means cloud in Latin.

- ◆ Huggins attributed 4959Å, observed in the Cat's Eye Nebula, to “nebulum” (or “nebulium”), and 5007Å line to Nitrogen => Ira Bowen discovered that these lines were actually forbidden [O III] lines.
- ◆ aurorium : 5577Å in the spectrum of the aurora borealis => turned out to be [O I]
- ◆ coronium: 5303Å in the spectrum of the Sun’s corona => Fe XIV

Interstellar Dust

- The existence of dust had been hinted at by the presence of dark nebulae (Barnard 68).
 - ◆ The dark nebulae were originally thought to be due to a lack of stars, but later recognized as being clouds of obscuring material.
- Vesto Slipher (1912) discovered that the spectrum of the nebula surrounding the Pleiades shows a continuum with absorption lines superposed.
 - ◆ He conjectured that this is light from stars, reflected from “fragmentary and disintegrated matter”, or dust.

V. Slipher (/slaifer/ 1875-1969) is the first one who measured radial velocities for galaxies and discovered that distant galaxies are redshifted. He was also the first to relate these redshifts to velocity.



Barnard 68 (at $d \sim 150$ pc), in the constellation Ophiuchus.



The Pleiades cluster & surrounding reflection nebulae

Interstellar gas that is invisible to the eye

- Initially, bright nebulae were thought of as isolated clouds in (nearly) empty space.
- In 1901, Johannes Hartmann found:
 - ◆ the spectrum of binary Delta Orionis (a spectroscopy binary system) shows a narrow calcium absorption line (at $\lambda 3934$) that is in **stationary** in addition to the **time-varying**, broad absorption lines due to the orbital motion of the stars
 - ◆ the stationary Ca absorption line was caused by a gas cloud somewhere along the line of sight to Delta Orionis.
- Later, similar “stationary lines” were found along the sightlines to many other bright stars.
 - ◆ The lines were all narrow, and had strengths correlated with the distance to the background star.
 - ◆ Using higher resolution spectrographs, they had been revealed to have complex structures, consisting of many narrower lines with different radial velocities.
 - ◆ This led to the realization that the ISM has a complex structure, consisting neither of smooth uniform gas nor of isolated blobs drifting about in a near-vacuum.

Diversity of the ISM

Ionized nebulae


- H II regions
 - are regions of interstellar gas heated and photoionized by embedded O or B-type stars with $T_{\text{eff}} > 25,000 \text{ K}$.
 - In 1939, Bengt Stromgren developed the idea that bright nebulae with strong emission lines are regions of photoionized gas, surrounding hot star or other source of ionizing photons.
 - ex) Orion Nebula
- Planetary nebulae
 - are regions of ejected stellar gas heated and photoionized by the hot remnant stellar core, which is becoming a white dwarf.
 - ex) Ring Nebula, Cat's Eye Nebula
 - Ring Nebula:
 - ◆ central region: blue color, from He II 4686.
 - ◆ middle region: blue-green colors from [O III] 4959, 5007
 - ◆ outer reddish colors from H α 6563, [N II] 6548, 6583



Orion Nebula ($d \sim 410 \text{ pc}$)
HST image




Cat's Eye Nebula (HST image)



Ring Nebula (HST image)

H II regions





[Left] True-color optical image of the Rosetta Nebula. The reddish glow is from $\text{H}\alpha$ line emission from recombination of the ionized hydrogen. The central cavity has been evacuated by the strong, high-speed stellar winds from the central cluster of hot stars.


[Right] Composite false-color image showing the emission in $\text{H}\alpha$ (red), and lines of [O III] (green) and [S II] (blue). Credit: NASA/HST [Owocki]

- Supernova remnants

- are regions of gas heated by the blastwave from a supernova explosion.
- Crab Nebula
 - ◆ a young ($t \sim 1000$ yr) pulsar-containing supernova remnant
 - ◆ are filled in with luminous gas.
 - ◆ are photoionized by its central pulsar.
 - ◆ are sometimes called ‘plerions’ meaning “full.”
- Cygnus Loop (Veil Nebula)
 - ◆ most of the gas has been plowed up by the blast wave, leaving the center part empty.
 - ◆ The visible loop (or veil) is where the gas has cooled to $T \sim 10,000$ K.
 - ◆ is a middle-aged supernova remnant ($t \sim 10^4$ yr).



-
- Warm Ionized Medium (Diffuse Ionized Gas)
 - About 20-80% of the ionized hydrogen in our galaxy lies in the relatively low density WIM.
 - Balmer line emission from recombining hydrogen fills the entire sky.
 - Although many ionized nebula (Orion, Crab, Cat's eye, etc) can be seen as the bright red blotches, they are not the dominant repository of recombining hydrogen in our galaxy.



All-sky map of H α (6563Å) in a log scale from 0.03 Ry to 160 Ry.
Ry (rayleigh) = $10^6/4\pi$ photons cm $^{-2}$ s $^{-1}$ Hz $^{-1}$ sr $^{-1}$

-
- Neutral Hydrogen Gas
 - All-sky map of H I 21-cm line intensity from the LAB survey (Kalberla et al. 2005), with an angular resolution ~ 0.6 deg.
 - Scale gives $\log_{10} N(\text{HI}) [\text{cm}^{-2}]$. The LMC and SMC are visible, with a connecting H I “bridge”.




Plate 3 in [Draine]

ISM in external galaxies



[Left] HST optical image of M51 (Whirlpool galaxy). The reddish blotches are from $H\alpha$ line emission from giant H II regions, which arise when dense regions of hydrogen are photoionized by the UV radiation from numerous, recently formed, hot massive stars. Note their proximity to dark bands formed from absorption of background stellar light by cold interstellar dust, which outline the galactic spiral arms.

[Right] Composite image of M51 from 4 space missions. X-rays (purple) detected by the Chandra X-ray Observatory reveal point-like sources (black holes and neutron stars in binary systems) as well as diffuse gas. Optical data from HST (green) and infrared emission from the Spitzer Space Telescope (red) both highlight long lanes in the spiral arms that consist of stars and gas laced with dust. UV light (blue) from GALEX comes from hot, young stars, showing how well these track the H II giants and star-forming GMCs along the spiral arms.

Credit: NASA/HST/CXO/SST/GALEX [Owocki]


Baryonic Matter

Definitions:

- Baryons = protons, neutrons and matter composed of them (i.e. atomic nuclei)
- Leptons = electrons, neutrinos
- In astronomy, however, the term '**baryonic matter**' is used more loosely to refer to **matter that is made of protons, neutrons, and electrons**, since protons and neutrons are always accompanied by electrons. Neutrinos, on the other hand, are considered non-baryonic by astronomers. (Note that black holes are also included as baryonic matter.)



The baryonic mass density

- 7% : stars + compact objects (such as stellar remnants, brown dwarfs, and planets)
- 1% : interstellar medium (ISM), filling the volume between stars.
- 5%: circumgalactic medium (CGM), bound within the dark halo of a galaxy, but outside the main distribution of stars.
- 4% : intracluster medium (ICM) of clusters of galaxies, bound to the cluster as a whole, but not to any individual galaxy.
- 40%: diffuse intergalactic medium (DIM), made of low density, mostly photo-ionized gas ($T < 10^5$ K).
- 43% : warm-hot intergalactic medium (WHIM), made of shock-heated gas ($10^5 \text{ K} < T < 10^7 \text{ K}$).



Mass flow of the baryons in galaxies

- At early times, the baryonic mass in galaxies was primarily in the gas of the ISM. As galaxies evolve, the ISM is gradually converted to stars, and some part of the interstellar gas may be ejected from the galaxy in the form of galactic winds, or in some cases stripped from the galaxy by the IGM.
- About 10% of the baryons in the Milky Way are to be found in the ISM.





Flow of baryons in the Milky Way.



B. T. Draine

Credit: NASA, Night Sky Network

Structure of the Milky Way



2MASS survey
 $\sim 5 \times 10^8$ stars
 blue = 1.2 μm
 green = 1.65 μm
 red = 2.2 μm



IRAS+COBE
 100 μm
 dust emission

(Jansky, $\text{Jy} = 10^{-23} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}$)

This artist's conception shows the various parts of our galaxy, and the position of our Sun.

-
- Total mass of the Milky Way $\sim 10^{11} M_{\odot}$ ($M_{\odot} = 1.989 \times 10^{33}$ g)
 - stars $\sim 5 \times 10^{10} M_{\odot}$
 - dark matter $\sim 5 \times 10^{10} M_{\odot}$
 - interstellar gas $\sim 7 \times 10^9 M_{\odot}$ (mostly H + He)
 - ◆ Hydrogen mass: neutral H atoms $\sim 60\%$, H₂ molecules $\sim 20\%$, ionized H⁺ atoms $\sim 20\%$

Phase	$M(10^9 M_{\odot})$	fraction
Total H II (not including He)	1.12	23%
Total H I (not including He)	2.9	60%
Total H ₂ (not including He)	0.84	17%
Total H II, H I and H₂ (not including He)	4.9	
Total gas (including He)	6.7	

ISM = dust + gas

Dust

- dust = tiny grains of solid material
 - Historically, courses on the ISM have dealt with “non-stellar stuffs.”
 - The dust and gas strongly influence each other.
 - ◆ Dust reprocesses starlight, altering the radiation field passing through the gas.
 - ◆ Dust is made of **refractory** elements, so creating dust alters the chemical abundances of the surrounding gas.
 - ◆ Dust grains are a **leading source of free electrons** in the neutral interstellar gas.
 - ◆ **Gas molecules form on the surfaces of dust grains.**

Gas

- Interstellar gas occupies the same region as stars.
- Stars are made from interstellar gas, and emit stellar winds into the ISM over the course of their lives. When massive stars reach the end of their lifetimes, they inject enriched gas at high speeds into the surrounding interstellar gas.
- Stars emit photons that are capable of exciting the interstellar gas. The emission lines have strong diagnostic power, enabling us to determine densities, temperatures, and ionization states of interstellar gas.

Abundance of elements in the local ISM

Element	Abundance (ppm)	Atomic number	1 st ionization energy (eV)
hydrogen (H)	911,900	1	13.60
helium (He)	87,100	2	24.59
oxygen (O)	490	8	13.62
carbon (C)	270	6	11.26
neon (Ne)	85	10	21.56
nitrogen (N)	68	7	14.53
magnesium (Mg)	40	12	7.65
silicon (Si)	32	14	8.15
iron (Fe)	32	26	7.90
sulfur (S)	13	16	10.36

(ppm = parts per million)

H : 91.2% by number

He: 8.7%

others: 0.1%

The interstellar gas is primarily H and He resisting from the Big Bang.

A small amount of heavy elements was produced as the result of the return to the ISM of gas that has been processed in stars and stellar explosions.

Asplund (2009)

$$M(Z > 2)/M_{\text{H}} = 0.0199; M(\text{total})/M_{\text{H}} = 1.402$$

solar metallicity:

$$Z_{\odot} = M(Z > 2)/M_{\text{tot}} \approx 0.02$$

Metallicity = the abundance of the elements heavier than hydrogen and helium

Density of the ISM

By terrestrial standards, the ISM is an almost perfect vacuum.

- The typical distance between stars is about $2 \text{ pc} = 6 \times 10^{16} \text{ m}$.
 - This is ~ 100 million times greater than the solar radius and 4000 times greater than the size of its heliosphere.

- ISM density
 - Total ISM mass is $\sim 7 \times 10^9 M_\odot$.
 - Approximating the Galaxy as a cylinder with a radius $R \sim 10 \text{ kpc}$ and a scale height $H \sim 250 \text{ pc}$, this implies an average density

$$\rho = \frac{M}{\pi R^2 \times 2H} \approx 3 \times 10^{-21} \text{ kg m}^{-3}$$

$$\rho = n_H m_H + n_{\text{He}} m_{\text{He}} \approx \left(1 + \frac{1}{10} \times 4\right) n_H m_H \quad \longleftarrow$$

$$\begin{aligned} n_{\text{He}} &\sim 0.1 n_H \\ m_{\text{He}} &\sim 4 m_H \end{aligned}$$

$$n_H \approx \frac{\rho}{1.4m_H} \simeq 1.3 \text{ cm}^{-3}$$

-
- Density of Air
 - From the ideal gas law using the pressure at sea level, $P = 10^5 \text{ N m}^{-2}$ (1 bar), and temperature $T \approx 300 \text{ K}$, we get

$$n = P/k_B T \approx 2.4 \times 10^{19} \text{ cm}^{-3}$$

This density is **19 orders of magnitude higher than the average density in the ISM.**

- The extremely low density in the ISM mean that particle collisions are relatively rare, which allows us to observe some physical processes that we don't see on Earth (e.g., forbidden lines).

Typical pressure & Energy densities

Typical pressure of the ISM

- $P = nk_B T \sim 4 \times 10^{-13} \text{ dyn cm}^{-2} \sim 4 \times 10^{-19} \text{ atm}$ (atmospheric pressure; 1 bar = 0.987 atm)
- Here, Boltzmann constant, $k_B = 1.38 \times 10^{-16} \text{ cm}^2 \text{ g s}^{-2} \text{ K}^{-1}$
- This is extremely low pressure compared to the atmospheric pressure around us.

Energy density

$$\begin{aligned}\varepsilon &= \frac{3}{2} n k_B T \\ &\sim 6 \times 10^{-13} \text{ erg cm}^{-3} \\ &\sim 0.4 \text{ eV cm}^{-3}\end{aligned}$$


Type	Energy density (eV cm ⁻³)
Thermal energy	0.4
Turbulent kinetic energy	0.2
Cosmic microwave background	0.2606
Far-infrared from dust	0.3
Optical/near-IR from stars	0.6
Magnetic energy	0.9
Cosmic rays	1.4

- All of them are comparable in energy density.
- All energy densities in the ISM are roughly half an electron-volt per cubic centimeter.
- The near-equipartition is partly coincidental.
 - ◆ The fact that the energy density in the CMB is similar to the other energy densities is surely accidental.
 - ◆ But the other energy densities are in fact coupled, roughly regulated by feedback mechanisms between them.

Radiative Transfer

Radiative Transfer

- ***Radiative Transfer*** describes how radiation is affected as it propagates to the observer through intervening gas and dust media.




Definition: Energy Flux

- Definition
 - Consider a small area dA , exposed to radiation for a time dt .
 - Energy flux F is defined as ***the net energy dE passing through the element of area in all directions in the time interval dt*** so that

$$dE = F \times dA \times dt$$


- Note that F ***depends on the orientation of the area element dA .***
- Unit: erg cm⁻² s⁻¹

- F_ν is often called the “flux density.”
- Radio astronomers use a special unit to define the flux density:
1 Jansky (Jy) = 10^{-23} erg s⁻¹ cm⁻² Hz⁻¹



Inverse Square Law

- Flux from an isotropic radiation source, i.e., a source emitting equal amounts of energy in all directions.



- Because of energy conservation, flux through two shells around the source must be the same.


$$4\pi r_1^2 F(r_1) = 4\pi r_2^2 F(r_2)$$

- Therefore, we obtain the inverse square law.

$$F = \frac{\text{const.}}{r^2}$$

Definition: Specific Intensity or Surface Brightness

- Recall that ***flux is a measure of the energy carried by all rays passing through a given area***
- Specific intensity is the energy carried along by individual rays.***



- Let dE_ν be the amount of radiant energy which crosses the area dA in a direction \mathbf{k} within solid angle $d\Omega$ about in a time interval dt with photon frequency between ν and $\nu + d\nu$.
- The monochromatic specific intensity I_ν is then defined by the equation.

$$dE_\nu = I_\nu(\mathbf{k}, \mathbf{x}, t) dA_{\mathbf{k}} d\Omega d\nu dt$$

- Unit: $\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1} \text{Hz}^{-1}$
- From the view point of an observer, the specific intensity is called ***surface brightness***.

Relation between the flux and the specific intensity


- Let's consider a small area dA , with light rays passing through it at all angles to the normal vector \mathbf{n} of the surface.
- For a ray centered about \mathbf{k} , the area normal to \mathbf{k} is

$$dA_{\mathbf{k}} = dA \cos \theta$$

- By the definition,

$$F_{\nu} dAd\nu dt = \int I_{\nu}(\mathbf{k}, \mathbf{x}, t) dA_{\mathbf{k}} d\Omega d\nu dt$$

- Hence, net flux in the direction of \mathbf{n} is given by integrating over all solid angles:



$$F_{\nu} = \int I_{\nu} \cos \theta d\Omega = \int_0^{2\pi} \int_0^{\pi} I_{\nu} \cos \theta \sin \theta d\theta d\phi$$

[Note] **flux** = “sum of all ray vectors” which is then projected onto a normal vector
intensity = magnitude of a single ray vector

Note


- Intensity can be defined as per wavelength interval.

$$\begin{aligned} I_\nu |d\nu| &= I_\lambda |d\lambda| & \leftarrow \left| \frac{d\nu}{d\lambda} \right| &= \frac{c}{\lambda^2} = \frac{\nu}{\lambda} & \leftarrow & \nu = \frac{c}{\lambda} \\ \nu I_\nu &= \lambda I_\lambda \end{aligned}$$

- Integrated intensity is defined as the intensity over all frequencies.

$$I = \int_0^\infty I_\nu d\nu = \int_0^\infty I_\lambda d\lambda$$

- In astrophysics, we plot the ***spectral energy distribution (SED)*** as νI_ν versus ν or λI_λ versus λ .



Definition: Luminosity

- To determine the energy per unit time, we integrate flux over area.
 - **Monochromatic luminosity**: Considering a sphere centered on a source with radius R , the monochromatic luminosity is

$$\begin{aligned}L_\nu &= R^2 \int d\Omega F_\nu \\&= 4\pi R^2 F_\nu \quad \text{for an isotropic source}\end{aligned}$$

- The **bolometric luminosity** is

$$\begin{aligned}L_{\text{bol}} &= \int L_\nu d\nu = \int L_\lambda d\lambda \\&= 4\pi R^2 \int F_\nu d\nu\end{aligned}$$

Definition: The magnitude scale

- For historical reasons, fluxes in the optical and infrared are measured in magnitudes.
- On the basis of naked eye observations, the Greek astronomer Hipparchus (190-120 BC) classified all the stars into six classes according to their apparent brightness.
 - The brighter ones belong to the first magnitude class. The faintest ones belong to the sixth magnitude class.
- Pogson (1856) noted that the faintest stars visible to the naked eye are about 100 times fainter compared to the brightest stars.
 - The brightest and faintest stars differ by five magnitude classes.
 - Therefore, stars in two successive classes should differ in apparent brightness by a factor $100^{1/5}$.
- Note that the human eye is more sensitive to a geometric progression ($I_0, 2I_0, 4I_0, 8I_0, \dots$) of intensity rather than an arithmetic progression ($I_0, 2I_0, 3I_0, 4I_0, \dots$). In other words, ***the apparent magnitude as perceived by the human eye scales roughly logarithmically with the radiation flux.***

-
- Suppose two stars have apparent brightnesses F_1 and F_2 and their magnitude classes are m_1 and m_2 .

$$\frac{F_2}{F_1} = (100)^{\frac{1}{5}(m_1 - m_2)}.$$

- Then, on taking the logarithm of this, we find

$$m_1 - m_2 = 2.5 \log_{10} \left(\frac{F_2}{F_1} \right).$$

- This is the definition of ***apparent magnitude*** denoted by m , which is a measure of the apparent brightness of an object in the sky.
 - Note that the magnitude scale is defined in such a fashion that ***a fainter object has a higher value of magnitude.***

-
- We need a measure that quantifies the luminosity or intrinsic brightness of an object.
 - The ***absolute magnitude*** of a celestial object is defined as ***the magnitude it would have if it were placed at a distance of 10 pc.***
 - If the object is at a distance d pc, then $(10/d)^2$ is the ratio between its apparent brightness and the brightness it would have if it were at a distance of 10 pc

$$\frac{F(d)}{F(10)} = \left(\frac{10 \text{ pc}}{d}\right)^2$$


- Then, the relation between apparent magnitude m and absolute magnitude M is

$$m - M = 2.5 \log_{10} \left(\frac{d}{10 \text{ pc}}\right)^2 = 5 \log_{10} \left(\frac{d}{10 \text{ pc}}\right)$$

- The difference $m - M$ is called the ***distance modulus***.

Filters and Wavebands

- Common bandpasses



Johnson system (or Johnson-Morgan system)

U (ultraviolet)	365 nm
B (blue)	440 nm
V (visible)	550 nm
R (red)	641 nm
I (near-infrared)	0.896 μm
J	0.900 μm
H	1.22 μm
K	2.19 μm

- These are the central wavelengths of each band, which extend ~10% in wavelength to either side.
- Magnitude at each bandpass is denoted by m_U , m_B , m_V , m_R , m_K , etc.

- Zero-points in the Vega magnitude system
 - Note that the magnitude scale has been relatively defined.
 - ***The zero-points are defined such that the magnitude of a standard star (Vega) is zero in all wavebands.***

AB magnitude

- Oke & Gunn (1983) defined the AB magnitude system.
- The monochromatic AB magnitude is defined as follows:

$$m_{\text{AB}} = -2.5 \log_{10} f_\nu(\text{Jy}) + 8.90 \approx -2.5 \log_{10} \left(\frac{f_\nu}{3631 \text{ Jy}} \right) \quad \text{Here, Jy} = 10^{-23} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}$$

$$m_{\text{AB}} = -2.5 \log_{10} f_\nu(\text{erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}) - 48.60$$

- The bandpass AB magnitude is defined so that the zero point corresponds to a bandpass-averaged spectral flux density of about 3631 Jy.

$$m_{\text{AB}} \approx -2.5 \log_{10} \left(\frac{\int f_\nu (e_\nu/h\nu) d\nu}{\int 3631 \text{ Jy} (e_\nu/h\nu) d\nu} \right)$$

Here, e_ν is the “equal-energy” filter response function, expressed in terms of per unit energy. If the filter responses are expressed as quantum efficiencies, (i.e., in terms of the response per photon) rather than per unit energy. The $1/h\nu$ term is folded into the definition of e_ν and should not be included.


- Modern systems of passbands, such as the SDSS ugriz filter system are on the AB magnitude system.

Constancy of Intensity

- How does specific intensity changes along a ray in free space
 - Suppose a bundle of rays and any two points along the rays and construct areas dA_1 and dA_2 normal to the rays at these points.
 - What are the energies carried by the rays passing through both areas?


energy passing through 1

$$dE_1 = I_1 dA_1 d\Omega_1 d\nu dt$$




energy passing through 2

$$dE_2 = I_2 dA_2 d\Omega_2 d\nu dt$$




- Here, $d\Omega_1$ is the solid angle subtended by dA_2 at the location 1 and $d\Omega_2$ is the solid angle subtended by dA_1 at the location 2.

: Radiative Transfer Equation in free space



Conservation of energy:
Because energy is conserved,




$$dE_1 = dE_2 \rightarrow I_1 = I_2$$

- Conclusion (***the constancy of intensity***): $I_1 = I_2$
 - the specific intensity remains the same as radiation propagates through free space.
- We receive the same specific intensity at the telescope as is emitted at the source.
 - Imagine looking at a uniformly lit wall and walking toward it. As you get closer, a field-of-view with fixed angular size will see a progressively smaller region of the wall, but this is exactly balanced by the inverse square law describing the spreading of the light rays from the wall.


Radiative Transfer Equation in reality

- In reality, as a ray passes through matter, energy may be added, subtracted, or scattered from it by emission, absorption, or scattering.
- The intensity will not in general remain constant.
- These interactions are described by the ***radiative transfer equation***.



Emission

- If the radiation travels through a medium which is itself emitting radiation, that will add to the energy:



- Spontaneous “**emission coefficient**” or “**emissivity**” j_ν is the amount of energy emitted per unit time, per unit solid angle, per unit frequency, and per unit volume:

$$dE = j_\nu dV d\Omega dt d\nu \quad (j_\nu : \text{erg cm}^{-3} \text{ s}^{-1} \text{ sr}^{-1} \text{ Hz}^{-1})$$

- In going a distance ds , a beam of cross section dA travels through a volume $dV = dA ds$. Thus the intensity added to the beam is by ds is

$$dI_\nu = j_\nu ds \qquad \longleftrightarrow \qquad dE = (dI_\nu) dA d\Omega dt d\nu$$

-
- Therefore, the equation of radiative transfer for pure emission becomes:


$$\frac{dI_\nu}{ds} = j_\nu$$

- If we know what j_ν is, we can integrate this equation to find the change in specific intensity as radiation propagates through the medium:

$$I_\nu(s) = I_\nu(0) + \int_0^s j_\nu(s')ds'$$

Absorption

- If the radiation travels through a medium which absorbs radiation, the energy in the beam will be reduced:



- Let n denote the number density of absorbers (particles per unit volume).
- Assume that each absorber has a cross-sectional area of σ_ν (in units of cm^2).
- If a beam travels through ds , total area of absorbers is

$$\text{number of absorbers} \times \text{cross section} = (n \times dA \times ds) \times \sigma_\nu$$

Fraction of radiation absorbed = fraction of area blocked:

$$\frac{dI_\nu}{I_\nu} = - \frac{ndAds\sigma_\nu}{dA} = - n\sigma_\nu ds \quad \longrightarrow \quad \frac{dI_\nu}{ds} = - \alpha_\nu I_\nu$$

$$dI_\nu = - n\sigma_\nu I_\nu ds \equiv - \alpha_\nu I_\nu ds$$

- **Absorption coefficient** is defined as $\alpha_\nu \equiv n\sigma_\nu$ (units: cm^{-1}), meaning the ***total cross-sectional area per unit volume***.

$$\alpha_\nu = n\sigma_\nu \quad [\text{cm}^{-1}]$$

$$= \rho\kappa_\nu$$

where ρ is the mass density and κ_ν is called the **mass absorption coefficient** or the opacity coefficient.

-
- Rearranging the previous equation, we obtain the equation of radiative transfer for pure absorption:


$$\frac{dI_\nu}{ds} = -\alpha_\nu I_\nu$$

- The amount of reduced energy depends on how much radiation we already have.
- Integrate to find how radiation changes along path:

$$\int_0^s \frac{dI_\nu}{I_\nu} = - \int_0^s \alpha_\nu(s') ds'$$

$$[\ln I_\nu]_0^s = - \int_0^s \alpha_\nu(s') ds'$$

$$I_\nu(s) = I_\nu(0) \exp \left[- \int_0^s \alpha_\nu(s') ds' \right]$$




-
- If the absorption coefficient is a constant (example: a uniform density gas of ionized hydrogen), then we obtain

$$I_\nu(s) = I_\nu(0)e^{-\alpha_\nu s}$$

specific intensity
after distance s

initial intensity
at $s = 0$.

radiation exponentially
absorbed with distance



- ***Optical depth:***
 - Imagine radiation traveling into a cloud of absorbing gas, exponential defines a scale over which radiation is attenuated.
 - When $\int_0^s \alpha_\nu(s')ds' = 1$, the intensity will be reduced to $1/e$ of its original value.

- We define the optical depth τ_ν as:

$$\tau_\nu(s) = \int_0^s \alpha_\nu(s') ds' \quad \text{or} \quad d\tau_\nu = \alpha_\nu ds \quad \longrightarrow \quad I_\nu(\tau_\nu) = I_\nu(0)e^{-\tau_\nu}$$

- A medium is said to be **optically thick** at a frequency ν if the optical depth for a typical path through the medium satisfies:

$$\tau_\nu(s) > 1$$

- The medium is **optically thin** if, instead:

$$\tau_\nu(s) < 1$$

- An optically thin medium is one which a typical photon of frequency ν can pass through without being (significantly) absorbed.

Mean Free Path

- From the exponential absorption law, the **probability of a photon absorbed** between optical depths τ_ν and $\tau_\nu + d\tau_\nu$ is

$$|dI_\nu| = \left| \frac{dI_\nu}{d\tau_\nu} \right| d\tau_\nu \quad \& \quad |dI_\nu| \propto P(\tau_\nu) d\tau_\nu \quad \rightarrow \quad P(\tau_\nu) = e^{-\tau_\nu}$$

= probability density function for the absorption at an optical depth τ_ν .

- The mean optical depth traveled is thus equal to unity:

$$\langle \tau_\nu \rangle = \int_0^\infty \tau_\nu P(\tau_\nu) d\tau_\nu = \int_0^\infty \tau_\nu e^{-\tau_\nu} d\tau_\nu = 1$$

- The mean free path is defined as the average distance a photon can travel through an absorbing material until it is absorbed.** In a homogeneous medium, the mean free path is determined by

$$\langle \tau_\nu \rangle = \alpha_\nu \ell_\nu = 1 \quad \rightarrow \quad \ell_\nu = \frac{1}{\alpha_\nu} = \frac{1}{n\sigma_\nu}$$


- A local mean path at a point in an inhomogeneous material can be also defined.

Radiative Transfer Equation

- ***Radiative transfer equation*** with both absorption and emission is

$$\frac{dI_\nu}{ds} = -\alpha_\nu I_\nu + j_\nu$$

absorption emission



- We can rewrite the radiative transfer equation using the optical depth as a measure of 'distance' rather than s :

$$\frac{dI_\nu}{\alpha_\nu ds} = -I_\nu + \frac{j_\nu}{\alpha_\nu}$$

$$\frac{dI_\nu}{d\tau_\nu} = -I_\nu + S_\nu$$

- where $S_\nu \equiv j_\nu/\alpha_\nu$ **is called the *source function***. This is an alternative and sometimes more convenient way to write the equation.


Thermal equilibrium

- In general, equilibrium means a state of balance.
- Thermal Equilibrium
 - ***Thermal equilibrium refers to steady states of temperature, which defines the average energy of material or photons.***
(for ideal gas, $E_{\text{avg}} = \frac{3}{2}k_{\text{B}}T$)
 - In a state of (complete) ***thermodynamic equilibrium (TE)***, no net flows of matter or of energy, no phase changes, and no unbalanced potentials (or driving forces), within the system. ***In TE, matter and radiation are in equilibrium at the same temperature T.***
 - When the material is (locally) in thermodynamic equilibrium, and only the radiation field is allowed to depart from its TE, we refer to the state of the system as being in ***local thermodynamic equilibrium (LTE)***
 - In other words, if the **material is (locally) in thermodynamic equilibrium** at a well-defined temperature T , ***it is said to be in local thermodynamic equilibrium (LTE) even if it is not in equilibrium with the radiation field.***

Blackbody

- Imagine a container bounded by opaque walls with a very small hole.

- ***Photons will be scattered and absorbed many times, (and eventually trapped and completely absorbed in the box).*** Under such conditions, the particles and photons continually share their kinetic energies. In perfect thermal equilibrium, ***the average particle kinetic energy will equal to the average photon energy, and a unique temperature T can be defined.***




- A **blackbody** is an idealized physical body that absorbs all incident radiation regardless of frequency or angle of incidence (i.e., perfect absorber). The above cavity can be regarded to be a blackbody.
- Radiation from a blackbody in thermal equilibrium is called the **blackbody radiation**.

Blackbody radiation is the universal function.

- Now, consider another cavity (box 2), also at the same temperature, but made of different material or shape and connect two cavities with a filter transparent only in the narrow frequency range ν and $\nu + d\nu$.

- In equilibrium at T , radiation should transfer no net energy from one cavity to the other. Otherwise, one cavity will cool down and the other heats up; this violates the second law of thermodynamics.
- Therefore, the intensity or spectrum that passes through the holes should be a universal function of T and should be isotropic.
- The intensity and spectrum of the radiation emerging from the hole should be independent of the wall material (e.g., wood, copper, or concrete, etc) and any absorbing material that may be inside the cavity.



- The universal function is called the Planck function $B_\nu(T)$.
- This is the blackbody radiation.