

Interstellar Medium (ISM)

Week 8

2025 April 25 (Friday), 3PM

updated on 05/08, 21:55

선광일 (Kwangil Seon)
KASI / UST

- The UV, visible and IR spectra of H II regions are very rich in emission lines, primarily **(1) collisional excited lines of metal ions and (2) recombination lines of hydrogen and helium**. H II regions are also observed at radio wavelengths, emitting radio free-free emission from thermalized electrons and radio recombination lines from highly excited states of H, He, and some metals (e.g., H 109α and C lines).
- Three processes govern the physics of H II regions:
 - **Photoionization Equilibrium**: the balance between photoionization and recombination. This determines the spatial distribution of ionic states of the elements in the ionized zone.
 - **Thermal Balance** between heating and cooling. Heating is dominated by photoelectrons ejected from hydrogen and helium with thermal energies of a few eV. Cooling is mostly dominated by electron-ion impact excitation of metal ion followed by emission of “forbidden” lines from low-lying fine structure levels. It is these cooling lines that give H II regions their characteristic spectra.
 - **Hydrodynamics**, including shocks, ionization and photodissociation fronts, and outflows and winds from the embedded stars.

-
- Ionization fraction
 - In the CNM, the fractional ionization $x = n_e/n_{\text{H}} \sim 0.001$. In the WNM, $x \sim 0.1$.
 - In the WIM, $x \sim 0.7$. In the HIM, $x \sim 1.0$.
 - Although the number of free electrons are small in the neutral ISM, free electrons play a role in bringing the WNM to kinetic equilibrium. *Free electrons photo-ejected from dust grains are the major heat source in the neutral medium.*
 - H II regions, with $T \sim 10,000$ K and $n \sim 0.3 \text{ cm}^{-3}$, contributes only a few percent of the mass of the ISM, and no more than ten percent of its volume.
 - Classical H II regions and planetary nebulae have a similar temperature, but *planetary nebulae have a higher density.*

Ionization Energy

- First ionization energy = energy required to remove the most loosely bound electron in a neutral atom in its ground state.
 - The first ionization energy of hydrogen is $I_H = 13.59844 \text{ eV}$.
 - The highest first ionization energy is that of He, with $I_{\text{He}} = 24.6 \text{ eV} = 1.91I_H$.
 - The second ionization energy of helium is $I_{\text{HeII}} = 54.4 \text{ eV} = 4I_H$.
 - The lowest first ionization energy of astrophysically interesting elements is that of potassium (K, $Z=19$), with $I_K = 4.3 \text{ eV}$. (Rubidium, cesium, and francium have lower first ionization energies, but they are not much seen in the ISM.)
 - Thus, photoionization of neutral atoms will be done by UV photons in the wavelength range $\lambda = 500 - 3000 \text{\AA}$ (corresponding to $E = 4.1 - 24.8 \text{ eV}$).
- A hydrogenic (hydrogen-like) ion with atomic number Z has an ionization energy of $Z^2 I_H$.
 - 54.4 eV for He II, 122.4 eV for Li III, and so forth.

Photoionization

- The (nonrelativistic) quantum mechanics of hydrogen-like ions (with only one electron) give an analytic expression for the ground-state photoionization (photoelectric) cross section.

$$\sigma_{\text{pi}}(\nu) = \sigma_0 \left(\frac{Z^2 I_{\text{H}}}{h\nu} \right)^4 \frac{e^{4-4 \arctan(x)/x}}{1 - e^{-2\pi/x}}, \quad x \equiv \sqrt{\frac{h\nu}{Z^2 I_{\text{H}}} - 1} \quad \text{for } h\nu > Z^2 I_{\text{H}}$$

- The cross section at threshold is

$$\sigma_0 \equiv \frac{2^9 \pi}{3e^4} \alpha \pi a_0^2 Z^{-2} = 6.304 \times 10^{-18} Z^{-2} \text{ cm}^{-2}$$

fine-structure constant
($\alpha \equiv e^2/hc = 1/137.04$, $e = 2.71828\dots$)


- The photoionization cross section is reasonably approximated by a power-law:

$$\sigma_{\text{pi}}(\nu) \approx \sigma_0 \left(\frac{h\nu}{Z^2 I_{\text{H}}} \right)^{-3} \quad \text{for } Z^2 I_{\text{H}} \lesssim h\nu \lesssim 100 Z^2 I_{\text{H}}$$

- At high energies, the asymptotic behavior is:


$$\sigma_{\text{pi}}(\nu) \approx \frac{2^8}{3Z^2} \alpha (\pi a_0^2) \left(\frac{h\nu}{Z^2 I_{\text{H}}} \right)^{-3.5} \quad \text{for } h\nu \gg Z^2 I_{\text{H}}$$

The hydrogen photoionization cross section becomes equal to the Thomson (Compton) scattering cross section for $h\nu \approx 2.5 \text{ keV}$; above this energy, photoionization of H is dominated by Thomson scattering rather than photoelectric absorption.




Photoionization cross section for hydrogen(H^0), hydrogenic helium (He^+), and neutral helium (He^0).
[Fig. 4.1 in Ryden]

- For atoms with three or more electrons, the energy dependence of the photoionization cross section is considerably more complicated because there is more than one available channel.
 - Convenient analytic fits to the contribution of individual shells to photoionization cross section are given by Verner & Yakovlev (1995) and Verner et al. (1996). <https://www.pa.uky.edu/~verner/photo.html>



※ Ionization threshold for $H_2 = 15.4$ eV [Fig. 13.1 in Draine]



[Fig. 13.2 in Draine]

-
- Photoionization rate (the probability of photoionization per unit time, for a single atom that undergoes photoionization)

$$\zeta_{\text{pi}} = \int_{\nu_1}^{\infty} \sigma_{\text{pi}}(\nu) c \frac{u_{\nu}}{h\nu} d\nu \quad \nu_1 = Z^2 I_{\text{H}}/h = 3.29 \times 10^{15} \text{ Hz} \text{ (for hydrogenic ions)}$$

= (cross section) x (flux of ionizing photons by number)

$$\text{incident photon flux} = 4\pi \frac{J_{\nu}}{h\nu} = c \frac{u_{\nu}}{h\nu}$$

- Since the photoionization cross section decreases fairly steeply with increasing photon energy, **most photoionization occurs by photons with energies just above the ionization energy (13.6 eV for hydrogen)**, in a range of the spectrum where the background is produced mainly by hot stars.
- The volumetric photoionization rate, for instance, for hydrogen is


$$\frac{dn_p}{dt} = n_{\text{H}^0} \zeta_{\text{pi}} \quad [\text{cm}^{-3} \text{ s}^{-1}]$$

= (# of atoms/volume) x (ionization rate per atom)

where n_p and n_{H^0} are the number density of proton (ionized hydrogen) and the number of neutral hydrogen atom, respectively.

Radiative Recombination (RR)

- The cross section for the radiative recombination can be obtained using the photoionization cross section and the **Milne relation**, which is derived from the principle of detailed balance.
- Consider an ion with its electron in some configuration that we will refer to as the “core”. In a low-density plasma, free electrons can undergo transitions to bound states by emission of a photon. The electron is captured into some specific state $n\ell$ that will initially unoccupied.



- The RR rate coefficient for electron capture directly to level $n\ell$, with emission of a photon of energy $h\nu = I_{n\ell} + E$ (where $I_{n\ell}$ is the bounding energy required for ionization from level $n\ell$ and E is the captured electron energy), is

$$\alpha_{n\ell}(T) \equiv \langle \sigma_{\text{rr},n\ell} v \rangle = \left(\frac{8kT}{\pi m_e} \right)^{1/2} \int_0^\infty \sigma_{\text{rr},n\ell}(E) \frac{E}{kT} e^{-E/kT} \frac{dE}{kT}$$

The integral indicates an average over the Maxwell distribution for electrons.

- The volumetric rate of RR, for instance for hydrogen, can be written as

$$\frac{dn_p}{dt} = -n_e n_p \sum_{n\ell} \alpha_{n\ell}$$

Notice that an electron of any energy can trigger a collisional de-excitation as well as RR.

- Properties of radiative recombination

- In general, $\alpha_{n\ell}$ is a decreasing function of T , although it depends weakly on temperature. Therefore, **it's easier to recombine with a slow electron than with a fast electron.**
- In general, $\alpha_n = \sum_\ell \alpha_{n\ell}$ summed over all applicable values of ℓ , is a decreasing function of n , implying that **it's easier to recombine to a low energy level than to a high energy level.**

- Recombination rate coefficient α

- Maxwellian distribution $\bar{f}(\mathbf{v})d^3\mathbf{v} = \left(\frac{m}{2\pi kT}\right)^{3/2} \exp\left(-\frac{m\mathbf{v}^2}{2kT}\right) d^3\mathbf{v}$
 $f(E)dE = \frac{2}{\sqrt{\pi}} \left(\frac{E}{kT}\right)^{1/2} \exp\left(-\frac{E}{kT}\right) \frac{dE}{kT}$ Here, $E = \frac{1}{2}mv^2$ (the energy per particle)
- Then, the **radiative recombination rate coefficient**, $\alpha(T)$ is given by

$$\begin{aligned}\langle \sigma_{\text{rec}} v \rangle &= \left(\frac{2}{m_e}\right)^{1/2} \langle \sigma_{\text{rec}} E^{1/2} \rangle \\ &= \left(\frac{8kT}{\pi m_e}\right)^{1/2} \int_0^\infty \sigma_{\text{rec}} \frac{E}{kT} \exp\left(-\frac{E}{kT}\right) d(E/kT)\end{aligned}$$

Notice that Ryden calls α the “recombination rate in her book.”
But, the recombination rate is $n_e \langle \sigma_{\text{rec}} v \rangle$.

$$\therefore \alpha(T) = \langle \sigma_{\text{rec}} v \rangle \approx \left(\frac{8}{\pi m_e kT}\right)^{1/2} \sigma_{\text{rec},0} I_{\text{H}} \propto T^{-1/2} \quad \Leftarrow \quad \sigma_{\text{rec}} \approx \sigma_{\text{rec},0} \left(\frac{E}{I_{\text{H}}}\right)^{-1} \text{ for hydrogen}$$

- Photoionization Equilibrium

$$n(X^r)n_\gamma\sigma_{\text{pho}}c = n(X^{r+1})n_e\sigma_{\text{rec}}v$$

$$\frac{n(X^{r+1})n_e}{n(X^r)} = \frac{n_\gamma \langle \sigma_{\text{pho}} \rangle_{h\nu} c}{\langle \sigma_{\text{rec}} v \rangle_v}$$




σ_{pho} = cross-section for photoionization,
depending on the photon energy $h\nu$

σ_{rec} = cross-section for recombination,
depending on the electron velocity v

averaged over the photon energy spectrum
averaged over the Maxwellian velocity distribution

- Recombination to the ground state - On The Spot approximation**

- If the recombination is to the ground state of hydrogen ($n = 1$), the energy of the emitted photon is $E + I_H \geq I_H$. Thus, the emitted photon is guaranteed to have an energy of at least 13.6 eV, and will be capable of photoionizing any neutral hydrogen atom that it encounters. Thus, in regions that are optically thick to UV light at photon energies just above I_H , the emitted photon will be rapidly destroyed in photoionizing a nearby hydrogen atom.



II Case A and B (Radiative Recombination of Hydrogen)

- **On-the-spot (OTS) approximation:**
 - In optically thick regions, it is assumed that every photon produced by radiative recombination to the ground state of hydrogen is immediately, then and there, destroyed in photoionizing other hydrogen atom.
 - In the on-the-spot approximation, **recombination to the ground state has no net effect on the ionization state of the hydrogen gas.**
- Baker & Menzel (1938) proposed two limiting cases:
 - **Case A: Optically thin** to ionizing radiation, so that every ionizing photon emitted during the recombination process escapes. For this case, we sum the radiative capture rate coefficient α_{nl} over all levels nl .
 - **Case B: Optically thick** to radiation just above $I_H = 13.60 \text{ eV}$, so that ionizing photons emitted during recombination are immediately reabsorbed, creating another ion and free electron by photoionization. In this case, the recombinations directly to $n = 1$ do not reduce the ionization of the gas: **only recombinations to $n \geq 2$ act to reduce the ionization.**
- **Case B in photoionized gas:** Photoionized nebulae around OB stars (H II regions) usually have large enough densities of neutral H. For this situation, case B is an excellent approximation.
- **Case A in collisionally ionized gas:** Regions where the hydrogen is collisional ionized are typically very hot ($T > 10^6 \text{ K}$) and contain a very small density of neutral hydrogen. For these shock-heated regions, case A is an excellent approximation.

- ***Radiative recombination rate coefficients:***

- In Case A, the relevant radiative recombination rate coefficient is found by summing over all energy levels of the hydrogen atom:

$$\alpha_{A,H}(T) \equiv \sum_{n=1}^{\infty} \sum_{\ell=0}^{n-1} \alpha_{n\ell}(T)$$

$$\approx 4.18 \times 10^{-13} T_4^{-0.721 - 0.021 \ln T_4} [\text{cm}^3 \text{s}^{-1}] \quad \text{for } 0.3 \lesssim T_4 \lesssim 3 \quad (T_4 \equiv T/10^4 \text{ K})$$


- In Case B, the relevant radiative recombination rate coefficient is found by summing over all energy levels other than the ground state:

$$\alpha_{B,H}(T) \equiv \sum_{n=2}^{\infty} \sum_{\ell=0}^{n-1} \alpha_{n\ell}(T) = \alpha_{A,H}(T) - \alpha_{1s}(T)$$

$$\approx 2.59 \times 10^{-13} T_4^{-0.833 - 0.034 \ln T_4} [\text{cm}^3 \text{s}^{-1}] \quad \text{for } 0.3 \lesssim T_4 \lesssim 3$$

- The percentage of radiative recombinations that go directly to the ground state is 30% at $T = 3000 \text{ K}$ but increases to 46% at $T = 30,000 \text{ K}$. Thus, the distinction between Case A and Case B becomes increasingly important at higher temperatures.

$$\frac{\alpha_{1s,H}}{\alpha_{A,H}} = 1 - \frac{\alpha_{B,H}}{\alpha_{A,H}} = 1 - 0.619 T_4^{-0.112 - 0.013 \ln T_4}$$



H II Regions and Strömgren Spheres

- **Strömgren Sphere:**

- Following Strömgren (1939), we consider the simple idealized problem of a fully ionized, spherical region of uniform medium plus a central source of ionizing photons.
- The ionization is assumed to be maintained by absorption of the ionizing photons radiated by a central hot star. The central source produces ionizing photons, with energy $\nu > \nu_0 = I_{\text{H}}/h$ at a constant rate Q_0 [photons s⁻¹].
- At a distance r from the central star, the balance equation between ionization and recombination balance is

$$n_{\text{H}^0} \int_{\nu_0}^{\infty} \frac{4\pi J_{\nu}}{h\nu} \sigma_{\nu} d\nu = n_p n_e \alpha_{\text{B,H}}$$

L_{ν} = luminosity of the central star at frequency ν .

From the RT equation,

$$4\pi J_{\nu} = \frac{L_{\nu}}{4\pi r^2} e^{-\tau_{\nu}}, \quad \text{where } \tau_{\nu} = \int_0^r n_{\text{H}^0} \sigma_{\nu} dr$$



geometrical attenuation + radiative absorption

Integrating the balance equation over the whole volume:

$$\int_0^{\infty} \int_{\nu_0}^{\infty} \frac{L_{\nu}/h\nu}{4\pi r^2} e^{-\tau_{\nu}} n_{\text{H}^0} \sigma_{\nu} d\nu (4\pi r^2) dr = \int_0^{\infty} n_p n_e \alpha_{\text{B,H}} (4\pi r^2) dr$$

$$\int_{\nu_0}^{\infty} L_{\nu}/h\nu \left[\int_0^{\infty} e^{-\tau_{\nu}} n_{\text{H}^0} \sigma_{\nu} dr \right] d\nu = \int_0^{\infty} n_p n_e \alpha_{\text{B,H}} (4\pi r^2) dr$$


The square bracket term in the left side is

$$\int_0^\infty e^{-\tau_\nu} n_{\text{H}^0} \sigma_\nu dr = \int_0^\infty e^{-\tau_\nu} d\tau_\nu = 1$$

Then, we obtain

$$Q_0 = \int_0^\infty n_p n_e \alpha_{\text{B},\text{H}} dV,$$

where $Q_0 \equiv \int_{\nu_0}^\infty \frac{L_\nu}{h\nu} d\nu$ and $dV = 4\pi r^2 dr$




- Assuming that ***the ionization is nearly complete*** ($n_p = n_e = n_{\text{H}}$) ***within*** R_s , and nearly zero ($n_{\text{H}^0} = n_{\text{H}}$, $n_e = 0$) outside R_s , we obtain the size of the ionized region.

$$\begin{aligned} Q_0 &= n_{\text{H}}^2 \alpha_{\text{B},\text{H}} \frac{4\pi}{3} R_s^3 \\ R_s &= \left(\frac{3}{4\pi} \frac{Q_0}{\alpha_{\text{B},\text{H}} n_{\text{H}}^2} \right)^{1/3} \\ &= 3.17 \left(\frac{Q_0}{10^{49} \text{ photons s}^{-1}} \right)^{1/3} \left(\frac{n_{\text{H}}}{10^2 \text{ cm}^{-3}} \right)^{-2/3} \left(\frac{T}{10^4 \text{ K}} \right)^{0.28} [\text{pc}] \end{aligned}$$

The physical meaning of this is that ***the total number of ionizing photons emitted by the star balances the total number of recombinations within the ionized volume*** $(4\pi/3)R_s^3$, often called the Strömgren sphere. Its radius R_s is called the Strömgren radius.

- ***Opacity as a function of distance***

- We note that the medium is fully ionized within the Strömgren sphere. Thus, within the Strömgren sphere, the opacity is nearly zero. The opacity suddenly increases at the boundary of the ionized region.



- ***Mean free path***

- The mean free path of an ionizing photon is

$$\lambda_{\text{mfp}} = \frac{1}{n_H \sigma_{\text{pi}}} \sim 5 \times 10^{-4} \text{ pc} \left(\frac{n_H}{10^2 \text{ cm}^{-2}} \right)^{-1} \left(\frac{\sigma_{\text{pi}}}{6.304 \times 10^{-18} \text{ cm}^{-2}} \right)^{-1}$$

This tells us that the transition from ionized gas to neutral gas at the boundary of the H II region will occur over a distance that is very small compared to the Strömgren radius.

-
- Time Scales:
 - ***Ionization time scale:*** The Strömgren sphere analysis assumes a steady state solution. What is the time scale for approach to the steady state? Suppose that we start with a neutral region, and the ionizing source is suddenly turned on.

$$t_{\text{ioniz.}} = \frac{\text{total number of ions to be created}}{\text{number of photons supplied per unit time}}$$

$$= \frac{(4\pi/3)R_s^3 n_{\text{H}}}{Q_0} = \frac{1}{\alpha_{\text{B,H}} n_{\text{H}}} = 1.22 \times 10^3 \left(\frac{10^2 \text{ cm}^{-3}}{n_{\text{H}}} \right) \text{ [yr]}$$

- ***Recombination time scale:*** Suppose that the ionizing source suddenly turns off. The ionized region will recombine on the recombination time scale:

$$t_{\text{rec}} = \frac{1}{\alpha_{\text{B,H}} n_{\text{H}}} = 1.22 \times 10^3 \left(\frac{10^2 \text{ cm}^{-3}}{n_{\text{H}}} \right) \text{ [yr]}$$

Therefore, the recombination time scale is identical to the ionization time scale!

The ionization/recombination time scale is shorter than the main-sequence lifetime > 5 Myr for a massive star.

Ionization of Helium

- Now, what about helium?
 - Out of every 1000 atoms, there are on average 912 hydrogen atoms, 87 helium atoms and one heavy atom.
 - ▶ Looking at the photoionization cross sections for H^0 , He^0 , He^{+1} , we see that above the 24.6 eV threshold for ionizing He^0 , the photoionization cross section for helium is larger than that for hydrogen.

$$\begin{aligned}\sigma_{\text{pi},He^0} &\approx 6.5 \sigma_{\text{pi},H^0} \quad \text{at } h\nu \sim 24.6 \text{ eV} \\ &\approx 14 \sigma_{\text{pi},H^0} \quad \text{at } h\nu \sim 54.5 \text{ eV}\end{aligned}$$

- ▶ Thus, the photoionization cross section for He is ~ 10 times that of H, while the number density of He is ~ 0.1 times that of H.
- ▶ This implies that if we suddenly turn on a hot star, ***the initial photons in the range $24.6 \text{ eV} < h\nu < 54.4 \text{ eV}$ will be about as likely to photoionize a helium atom as a hydrogen atom.***
- ▶ ***In the range of $13.6 \text{ eV} < h\nu < 24.6 \text{ eV}$, on the other hand, nearly all the photons go to ionize H;*** scarcer atoms (metals like O and C) account for only a tiny fraction of the ionizations.

- ***Radiative Recombination of Helium***



$$\alpha_A(T) \approx 4.13 \times 10^{-13} Z(T_4/Z^2)^{-0.7131-0.0115 \ln(T_4/Z^2)} \quad [\text{cm}^3 \text{s}^{-1}] \quad (30 \text{ K} < T/Z^2 < 3 \times 10^4 \text{ K})$$

$$\alpha_B(T) \approx 2.54 \times 10^{-13} Z(T_4/Z^2)^{-0.8163-0.0208 \ln(T_4/Z^2)} \quad [\text{cm}^3 \text{s}^{-1}]$$



$$\alpha_{1s^2, \text{He}} = 1.54 \times 10^{-13} T_4^{-0.486} \quad [\text{cm}^3 \text{s}^{-1}] \quad (0.5 < T_4 < 2)$$

$$\alpha_{B, \text{He}} = 2.72 \times 10^{-13} T_4^{-0.789} \quad [\text{cm}^3 \text{s}^{-1}]$$

Here, $\alpha_{1s^2, \text{He}}$ is the recombination rate to the ground state $1s^2 \ ^1S_0$,
and $\alpha_{B, \text{He}}$ is the recombination rate coefficient to all states except the ground state.

Note: $\alpha_{B, \text{H}} \approx \alpha_{B, \text{He}}$ and $\alpha_{A, \text{H}} \approx \alpha_{A, \text{He}}$.

- ***Effective recombination rate coefficient for Helium***

- Note that ***the stellar LyC photons with $h\nu > 24.6 \text{ eV}$ are capable of photoionizing not only neutral helium atoms but also neutral hydrogen atoms***
- The recombinations directly to the ***ground state $1s^2 1S_0$*** of neutral helium produce photons with $h\nu > 24.6 \text{ eV}$. ***The recombination continuum photons are capable of photoionizing not only neutral helium atoms but also neutral hydrogen atoms; the fraction of these that ionize hydrogen is***

$$\begin{aligned} y &= \frac{n_{\text{H}^0} \sigma_{\text{pi}, \text{H}^0}(E)}{n_{\text{H}^0} \sigma_{\text{pi}, \text{H}^0}(E) + n_{\text{He}^0} \sigma_{\text{pi}, \text{He}^0}(E)} \\ &= \left[1 + \frac{n_{\text{He}^0}}{n_{\text{H}^0}} \frac{\sigma_{\text{pi}, \text{He}^0}(E)}{\sigma_{\text{pi}, \text{H}^0}(E)} \right]^{-1}, \quad \text{where } E \approx 24.6 \text{ eV} + kT \end{aligned}$$

$$\sigma_{\text{pi}, \text{He}^0}/\sigma_{\text{pi}, \text{H}^0} > 6.0 \text{ for } E > 24.6 \text{ eV}$$

$$y < 0.5 \text{ if } n_{\text{He}^0}/n_{\text{H}^0} > 0.16$$

This fraction contributes to the recombination of He, while the remaining fraction, $1 - y$, photoionizes He again. Therefore, in an optically thick gas, the effective radiative recombination rate coefficient for $\text{He}^+ \rightarrow \text{He}^0$ is

$$\alpha_{\text{eff}, \text{He}} = \alpha_{\text{B}, \text{He}} + y \alpha_{1s^2, \text{He}} = \alpha_{\text{A}, \text{He}} - (1 - y) \alpha_{1s^2, \text{He}}$$


At $T = 10,000 \text{ K}$,

$\alpha_{\text{B}, \text{He}} = 2.72 \times 10^{-13} \text{ [cm}^3 \text{s}^{-1}\text{]}$	\rightarrow	$\alpha_{\text{eff}, \text{He}} \approx 3.0 \times 10^{-13} \text{ [cm}^3 \text{s}^{-1}\text{]}$
$\alpha_{1s^2, \text{He}} = 1.54 \times 10^{-13} \text{ [cm}^3 \text{s}^{-1}\text{]}$		$\approx 1.2 \alpha_{\text{B}, \text{H}}$
		$y \approx 0.2$

- This is not all. Consider now the recombination to **excited levels** of He^0 , which are followed by a radiative cascade down. Most of photons produced by the cascades have $h\nu > 13.6 \text{ eV}$. **A fraction of these photons are capable of photoionizing hydrogen. Let z be this fraction.** However, note that **this fraction is not relevant to the recombination of He, but contribute to the photoionization H.**

$z \approx 0.96$ at low densities
 ≈ 0.67 at high densities

We take an intermediate value $z \approx 0.8$.



- See Section 14.3.2 and 15.5 of [Draine] for details.
- [Ryden] assumes that $z = 1$.

Note that $1.08 \mu\text{m}$ is useful to probe the planetary atmosphere

[Fig. 14.3 of Draine]

- **How many recombinations occur for He:** Suppose that we have a Strömgren sphere with the cosmic abundance ratio of helium to hydrogen $f \equiv n_{\text{He}}/n_{\text{H}} \approx 0.096$. Now define:

$$Q_0 \equiv \int_{I_{\text{H}}/h}^{\infty} \frac{L_{\nu}}{h\nu} d\nu, \quad Q_1 \equiv \int_{I_{\text{He}}/h}^{\infty} \frac{L_{\nu}}{h\nu} d\nu \quad (Q_1 < Q_0)$$

- In the very central region, the hydrogen would be fully ionized, and the helium would be all singly ionized. Even **the hottest O stars don't produce a significant number of photons with $h\nu > 54.5 \text{ eV}$** ; hence, there will be no significant amount of doubly ionized He^{+2} .

- This will result in $n_p = n_{\text{H}}$

$$n_{\text{He}^+} = n_{\text{He}} = f n_{\text{H}}$$

$$n_e = n_p + n_{\text{He}^+} = (1 + f) n_{\text{H}} \quad \text{inside the Strömgren sphere.}$$

- The volumetric rate of the hydrogen recombination is

$$\frac{dn_p}{dt} = -\alpha_{\text{B,H}} n_e n_p = -\alpha_{\text{B,H}} (1 + f) n_{\text{H}}^2$$

- The volumetric rate of He recombination is

$$\frac{dn_{\text{He}^+}}{dt} = -\alpha_{\text{eff,He}} n_e n_{\text{He}^+} = -\alpha_{\text{eff,He}} f (1 + f) n_{\text{H}}^2$$

- Comparing the two equations, we see that

$$\begin{aligned}\frac{dn_{\text{He}^+}}{dt} &= \left(\frac{\alpha_{\text{eff}, \text{He}}}{\alpha_{\text{B}, \text{H}}} \right) f \frac{dn_p}{dt} \\ &\approx (1.2)(0.096) \frac{dn_p}{dt} \\ &\approx 0.11 \frac{dn_p}{dt}\end{aligned}$$

- Thus, **for every helium recombination, we expect about 9 hydrogen recombinations.**

- Remember the recombination paths, under the Case B condition:
 - $13.6 \text{ eV} < h\nu < 24.6 \text{ eV}$: A stellar photon will ionize one H atom.
 - $h\nu > 24.6 \text{ eV}$: For a fraction of y of the photoionization followed by the **direct recombinations to the ground state**, a stellar photon will ionize one H atom. For the remaining fraction $(1 - y)$ of these, a stellar photon will ionize one He atom.
 - $h\nu > 24.6 \text{ eV}$: For the photoionization followed by **the recombinations to excited states**, a stellar photon will ionize one H atom for a fraction of z of the recombination events.
- **Number of ionized atoms:** The number of ionized helium and hydrogen, $N(\text{He}^+)$ and $N(\text{H}^+)$, within the ionized regions can be estimated by balancing recombinations and photoionizations:

$$\begin{aligned}
 N(\text{He}^+)n_e (\alpha_{\text{B},\text{He}} + y\alpha_{1s^2,\text{He}}) &= (1 - y)Q_1 \\
 N(\text{H}^+)n_e \alpha_{\text{B},\text{H}} &= (Q_0 - Q_1) + yQ_1 + N(\text{He}^+)n_e (z\alpha_{\text{B},\text{He}} + y\alpha_{1s^2,\text{He}})
 \end{aligned}$$

Contribution by the recombination to the ground state.
 Contribution by the recombination to the excited state.

stellar LyC with $h\nu > 24.5 \text{ eV}$ that ionize H

$\rightarrow N(\text{H}^+)n_e \alpha_{\text{B},\text{H}} = Q_0 - N(\text{He}^+)n_e (1 - z) \alpha_{\text{B},\text{He}}$

$\longrightarrow \frac{N(\text{He}^+)}{N(\text{H}^+)} = \frac{(1 - y)\alpha_{\text{B},\text{H}}(Q_1/Q_0)}{\alpha_{\text{B},\text{He}} + y\alpha_{1s^2,\text{He}} - (1 - y)(1 - z)(Q_1/Q_0)\alpha_{\text{B},\text{He}}}$

$$\frac{N(\text{He}^+)}{N(\text{H}^+)} \approx \frac{0.68(Q_1/Q_0)}{1 - 0.17(Q_1/Q_0)} \quad \text{for } z \approx 0.8, T = 8000\text{K}, \text{ and } y = 0.2$$

- Condition for full ionization of the He in the H⁺ Strömgren sphere:

$$\frac{N(\text{He}^+)}{N(\text{H}^+)} = \frac{n_{\text{He}}}{n_{\text{H}}} = 0.096 \rightarrow \frac{Q_1}{Q_0} \approx 0.15$$


- **Radius of the He⁺ zone:**

$$N(\text{He}^+) = \frac{4\pi}{3} R_{\text{He}}^3 n_{\text{He}}$$

$$N(\text{H}^+) = \frac{4\pi}{3} R_{\text{H}}^3 n_{\text{H}}$$

$R_{\text{He}} < R_{\text{H}}$ if $Q_1/Q_0 \lesssim 0.15$

$$\begin{aligned} \frac{R_{\text{He}}}{R_{\text{H}}} &= \left[\frac{n_{\text{H}}}{n_{\text{He}}} \frac{N(\text{He}^+)}{N(\text{H}^+)} \right]^{1/3} \\ &= \left[\frac{7.08(Q_1/Q_0)}{1 - 0.17(Q_1/Q_0)} \right]^{1/3} \end{aligned}$$




- On the main sequence, a star with spectral class O7, corresponding to effective temperature $T_{\text{eff}} = 37,000$ K, will have a critical ratio $Q_1/Q_0 \sim 0.14$.
 - ▶ For cooler ionizing stars, the ionized helium sphere will have a radius that is smaller than the radius of the ionized hydrogen sphere.
 - ▶ For stellar temperature $T_{\text{eff}} > 37,000$ K, the ionized helium sphere has the same size as the ionized hydrogen sphere, because of the limit on the abundance. The photons with $h\nu > 24.6\text{ eV}$ will be used up to ionize H.

Table 15.1 [Draine]

SpTp	M/M_{\odot}	T_{eff} (K)	$\log_{10}(Q_0/\text{s}^{-1})^b$	Q_1/Q_0^c	$\log_{10}(L/L_{\odot})^d$
O3V	58.0	44850	49.64	0.251	5.84
O4V	46.9	42860	49.44	0.224	5.67
O5V	38.1	40860	49.22	0.209	5.49
O5.5V	34.4	39870	49.10	0.204	5.41
O6V	31.0	38870	48.99	0.186	5.32
O6.5V	28.0	37870	48.88	0.162	5.23
O7V	25.3	36870	48.75	0.135	5.14
O7.5V	22.9	35870	48.61	0.107	5.05
O8V	20.8	34880	48.44	0.072	4.96
O8.5V	18.8	33880	48.27	0.0347	4.86
O9V	17.1	32830	48.06	0.0145	4.77
O9.5V	15.6	31880	47.88	0.0083	4.68
O3III	56.0	44540	49.77	0.234	5.96
O4III	47.4	42420	49.64	0.204	5.85
O5III	40.4	40310	49.48	0.186	5.73
O5.5III	37.4	39250	49.40	0.170	5.67
O6III	34.5	38190	49.32	0.158	5.61
O6.5III	32.0	37130	49.23	0.141	5.54
O7III	29.6	36080	49.13	0.129	5.48
O7.5III	27.5	35020	49.01	0.105	5.42
O8III	25.5	33960	48.88	0.072	5.35
O8.5III	23.7	32900	48.75	0.0417	5.28
O9III	22.0	31850	48.65	0.0257	5.21
O9.5III	20.6	30790	48.42	0.0129	5.15
O3I	67.5	42230	49.78	0.204	5.99
O4I	58.5	40420	49.70	0.182	5.93
O5I	50.7	38610	49.62	0.158	5.87
O5.5I	47.3	37710	49.58	0.151	5.84
O6I	44.1	36800	49.52	0.141	5.81
O6.5I	41.2	35900	49.46	0.132	5.78
O7I	38.4	34990	49.41	0.115	5.75
O7.5I	36.0	34080	49.31	0.100	5.72
O8I	33.7	33180	49.25	0.079	5.68
O8.5I	31.5	32270	49.19	0.065	5.65
O9I	29.6	31370	49.11	0.0363	5.61
O9.5I	27.8	30460	49.00	0.0224	5.57


^a After Martins et al. (2005).^b Q_0 = rate of emission of $h\nu > 13.6 \text{ eV}$ photons.^c Q_1 = rate of emission of $h\nu > 24.6 \text{ eV}$ photons.^d L = total electromagnetic luminosity.

Figure 15.5 [Draine]

Q₁/Q₀ vs. stellar temperature


Q₁/Q₀ > 0.15 is required for He to be ionized throughout the H II region, corresponding to $T_{\text{eff}} > 37,000 \text{ K}$.

Figure 2.4 [Osterbrock]



Ionization structure of two homogeneous H + He models for H II regions.

Figure 2.5 [Osterbrock]



Relative radius of He^+ zone as a function of effective temperature of exciting star.

- **Metals:** Ions that requires $E > 24.6$ eV for their formation will present only in the He^+ zone.

[Draine] **Table 15.2** Abundant Ions in H II Regions^a

Element	H II and He I zone ^b		H II and He II zone ^c	
	Ion	$h\nu$ (eV) ^d	Ion	$h\nu$ (eV) ^d
H	H II	13.60	H II	13.60
He	He I	0	He II	24.59
C	C II	11.26	C III ^e	24.38
			C IV	47.88
N	N II	14.53	N III	29.60
			N IV	47.45
O	O II	13.62	O III	35.12
Ne	Ne II	21.56	Ne III	40.96
Na	(Na II) ^f	5.14	(Na II) ^f	5.14
			Na III	47.29
Mg	Mg II	7.65	(Mg III) ^f	15.04
	(Mg III) ^f	15.04		
Al	Al III	18.83	(Al IV) ^f	28.45
Si	Si III	16.35	Si IV	33.49
			(Si V) ^f	45.14
S	S II	10.36	S III	23.33
	S III	23.33	S IV	34.83
Ar	Ar II	15.76	Ar III	27.63
			Ar IV	40.74
Ca	Ca III	11.87	Ca IV	50.91
Fe	Fe III	16.16	Fe IV	30.65
Ni	Ni III	18.17	Ni IV	35.17

^a Limited to elements X with $N_X/N_{\text{H}} > 10^{-6}$.

^b Ions that can be created by radiation with $13.60 < h\nu < 24.59$ eV.

^c Ions that can be created by radiation with $24.59 < h\nu < 54.42$ eV.

^d Photon energy required to create ion.

^e Ionization potential is just below 24.59 eV.

^f Closed shell, with no excited states below 13.6 eV.

[Page 238, Dopita]

The dominant ionization zones of the nebula for the most abundant elements and important coolants are as follows:

H I, He I : C II, N I, O I, Ne I, S II,

H II, He I : C II, (C III), N II, O II, Ne II, S II, (S III),


H II, He II : C III, (C IV), N III, O III, Ne III, S III, (S IV, S V),


H II, He III : C IV, N IV, O IV, Ne III, S V, and higher,

[Figure 5.3, Dyson]

Ionization stratification in a nebula.


(i) Low stellar temperature, (ii) High stellar temperature





[Figure 9.4, Dopita, Astrophysics of the Diffuse Universe]

The temperature, density, and ionization structure of a model H II region illuminated by a star with an effective temperature of 53,000 K. Note how the ionization structure in the heavy elements follows that of hydrogen and helium.




[Seon & Witt, 2012, ApJ, 758, 19]

Figure 4. Top: brightness profiles of $H\alpha$, $\text{He I } \lambda 5876$, $[\text{N II}] \lambda 6583$, and $[\text{S II}] \lambda 6716$ lines (in units of $\text{erg cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$) for various central ionization sources. Bottom: brightness profiles of line ratios $\text{He I}/H\alpha$, $[\text{N II}]/H\alpha$, and $[\text{S II}]/H\alpha$. Elemental abundances for WNM and hydrogen density of $n_H = 10 \text{ cm}^{-3}$ were assumed for the photoionization models. The curves from the outermost to innermost correspond to O3V to B1V stars progressively. Solid and dashed lines were alternatively used for clarification.

Recombination lines

- Recombination Radiation = Recombination Lines + Recombination Continuum
- Diagnostics using the recombination lines
 - **Temperature**: The hydrogen recombination spectrum depends on temperature T , and therefore measured line ratios can be used to estimate T .
 - **Reddening**: Measurements of the relative intensities of recombination lines with different wavelengths can be used to estimate the reddening by dust between us and the emitting region.
- ***Case A Recombination Spectrum***
 - In the optically thin limit, the power radiated per volume in the transition $nl \rightarrow n'l'$ is

$$4\pi j(nl \rightarrow n'l') = n_e n_p \frac{A(nl \rightarrow n'l') h \nu_{nl \rightarrow n'l'}}{\sum_{n''l''} A(nl \rightarrow n''l'')} \times \left[\alpha(nl) + \sum_{n''l'' (n'' > n)} \alpha(n''l'') P_A(n''l'', nl) \right]$$



↓ direct recombination ↓ cascade

Note a typo in Eq (14.7) of Draine

$P_A(n''l'', nl)$ is the Case A probability that an atom in level $n''l''$ will follow a decay path that takes it through level nl . The probabilities can be readily calculated from the known transition probabilities $A(nl \rightarrow n'l')$ using straightforward branching probability arguments.

- ***Case B Recombination Spectrum***

- The resonant absorption cross-sections for Ly α , Ly β ,... are much larger than photoionization cross sections.

$$\tau_0(\text{Ly}\alpha) = 8.02 \times 10^4 \left(\frac{15 \text{ km s}^{-1}}{b} \right) \tau(\text{Ly cont})$$

$$\tau(\text{Ly cont}) = 6.30 \times 10^{-18} \text{ cm}^2 N(\text{H})$$

- ***Any nebula that is optically thick to Lyman continuum ($E > 13.6 \text{ eV}$) will be very optical thick to all of the Lyman series ($n \rightarrow 1$) transitions.***
- Note that the cross sections for resonant absorption in the $1 \rightarrow n$ transitions becomes equal to the photoionization cross section as $n \rightarrow \infty$.

$$\tau_{\text{reson.}}(1 \rightarrow n) \geq \tau_{\text{reson.}}(1 \rightarrow \infty) = \tau_{\text{photo.}}$$

-
- **On-the-spot approximation:**
 - ▶ Therefore, under Case B condition, Lyman series photons will (immediately) be resonantly absorbed by other hydrogen atoms in the ground state. They will travel only a short distance before being reabsorbed.
 - ▶ It is helpful to think about the radiative decay and resonant reabsorption as though the photon were reabsorbed by the same atom as emitted.
 - ▶ Consider a hydrogen atom in level $n \geq 3$ (for instance, $n = 3$) . Then, Ly β , Ly γ ,... will immediately be resonantly absorbed, returning back to the initial state $n \geq 3$. After returning to the initial state, the atom will again decay one of its allowed decay paths (for instance, $3 \rightarrow 2 \rightarrow 1$ and $4 \rightarrow 2 \rightarrow 1$). The atom may emit another Lyman series photon, which will again be absorbed.
 - ▶ This process will repeat until eventually “non-Lyman transitions” + a “Ly α transition” (or “non-Lyman transitions” + 2-photon transition) occur.

For instance,




Two-photon continuum emission occur, if the repeated process eventually populates 2s state, instead of 2p.


- ▶ Under this condition, no Lyman series lines (except for Ly α) will be produced.

Case B

(1) Lyman Continuum Photons




Recombination to $n = 1$



Absorption of emitted LyC photons
by other hydrogen atoms


(2) Lyman Series Line Photons

example 1




Emission of higher series lines
(Balmer, etc) + Ly α and 2-photons

example 2




Emission of Ly γ ($n \geq 2$) photons
(Ly β , Ly γ , ...)


Resonance




Fluorescence



Absorption of Ly β photons
by other hydrogen atoms





- **Balmer lines:**
 - Under Case B condition, the rate coefficients for recombinations that result in emission of H α , H β can be approximated by

$$\alpha_{\text{eff}, \text{H}\alpha} \approx 1.17 \times 10^{-13} T_4^{-0.942 - 0.031 \ln T_4} [\text{cm}^3 \text{s}^{-1}] \quad (T_4 \equiv T/10^4 \text{ K})$$

$$\alpha_{\text{eff}, \text{H}\beta} \approx 3.03 \times 10^{-14} T_4^{-0.874 - 0.058 \ln T_4} [\text{cm}^3 \text{s}^{-1}]$$

- Emissivities of Balmer lines:

Using the statistical balance for the level population, we can obtain the emissivity. (Note that, ***in the case of hydrogen and helium, the population caused by collisional excitation is negligible.***)

Population of u state by recombination = Depopulation by radiative decay.

$$4\pi j_{ul} = n_u A_{ul} (h\nu_{ul}) = n_e n_p \alpha_{\text{eff}, u} (h\nu_{ul})$$

$$4\pi j_{\text{H}\alpha} = n_e n_p \alpha_{\text{eff}, \text{H}\alpha} h\nu_{\text{H}\alpha}$$

$$4\pi j_{\text{H}\beta} = n_e n_p \alpha_{\text{eff}, \text{H}\beta} h\nu_{\text{H}\beta}$$

- Balmer Decrement**: The ratio between Balmer lines can be used **to estimate the dust reddening.**

$$\frac{j_{\text{H}\alpha}}{j_{\text{H}\beta}} = \frac{\alpha_{\text{eff}, \text{H}\alpha}}{\alpha_{\text{eff}, \text{H}\beta}} \frac{\nu_{\text{H}\alpha}}{\nu_{\text{H}\beta}} = 2.86 T_4^{-0.068 + 0.027 \ln T_4}$$

Note: $\lambda_{\text{H}\alpha} = 6563\text{\AA}$
 $\lambda_{\text{H}\beta} = 4861\text{\AA}$

See Table 14.2 of Draine for other lines.

- Lyman α

- Let $\alpha_{\text{eff},2s}$ and $\alpha_{\text{eff},2p}$ be the effective rate coefficients for populating the 2s and 2p states. By definition, it is clear that the case B radiative recombination process must eventually take the atom to either the 2s level or the 2p level. Thus,

$$\alpha_{\text{eff},2s} + \alpha_{\text{eff},2p} = \alpha_B$$

- The fractions $f(2s) \equiv \frac{\alpha_{\text{eff},2s}}{\alpha_B} \approx \frac{1}{3}$ and $f(2p) \equiv \frac{\alpha_{\text{eff},2p}}{\alpha_B} \approx \frac{2}{3}$ are given in the following table.

T(K)	f(2s)	f(2p)
4000	0.285	0.715
5000	0.305	0.695
10000	0.325	0.675
20000	0.356	0.644

Tables 14.2 and 14.3 of [Draine]

A minor discrepancy between this and Cantalupo et al. (2008, ApJ, 672, 48):

$$f(\text{Ly}\alpha) = 0.686 - 0.106 \log(T/10^4 \text{ K}) - 0.009 (T/10^4 \text{ K})^{-0.44}$$

- Then, the emissivity for Ly α is

$$\begin{aligned} 4\pi j_{\text{Ly}\alpha} &= n_e n_p \alpha_{\text{eff},2p} h\nu_{\text{Ly}\alpha} \\ &\approx \frac{2}{3} n_e n_p \alpha_B h\nu_{\text{Ly}\alpha} \end{aligned}$$

In a high density medium ($n_e \gtrsim 1.55 \times 10^4 \text{ cm}^{-3}$), the Ly α emissivity will be increased by the collisional transition from 2s to 2p state (see 14.2.4 of [Draine]).

How many Ly α , H α , and H β photons are produced for each recombination event:

$$f(\text{Ly}\alpha) = \frac{\alpha_{\text{eff},2p}}{\alpha_B} \approx \frac{2}{3} \Rightarrow f(2p)$$

$$f(\text{H}\alpha) = \frac{\alpha_{\text{eff,H}\alpha}}{\alpha_B} = 0.452 T_4^{-0.109 - 0.003 \ln T_4}$$

$$f(\text{H}\beta) = \frac{\alpha_{\text{eff,H}\beta}}{\alpha_B} = 0.117 T_4^{-0.041 - 0.02 \ln T_4}$$

- **Radiative Recombination: Heavy Elements**

- We do not concern ourselves with the possibility that photons emitted from recombination to the ground state could be reabsorbed locally by another atom.
- That is, *we assume Case A condition when studying the recombination of heavy elements.*
- Radiative recombination of elements such as O and Ne is accompanied by emission of characteristic lines - the recombining electrons are captured into excited states, which then emit a cascade of line radiation.
- For example, radiative recombination of O III sometimes populates an excited state, resulting in O II 4462.8Å and O II 4073.79Å emission (allowed lines).
- In H II regions and planetary nebulae, these recombination lines are faint compared to the recombination lines of H, simply because of the greatly reduced abundance of heavy elements, but can nevertheless be measured.
- The abundances obtained from **recombination lines** should, in principle, agree with the abundances derived from the much stronger **collisionally excited lines**. However, it is known that *recombination lines give abundances that are larger than that estimated from collisionally excited lines. This is a puzzle that is yet to be resolved.*

Appendix: Ionization Fraction within an H II region

- Let's consider a shell between radii r and $r + dr$.
 - Number of ionizing photons within the volume = Number of Recombinations within in the volume

$$|Q(r + \Delta r) - Q(r)| = n_p n_e \alpha_B \Delta V$$

$$\frac{dQ}{dr} = -n_p n_e \alpha_B 4\pi r^2$$

$$\begin{aligned} Q(r) &= Q_0 - \int_0^r n_p n_e \alpha_B 4\pi r'^2 dr' \\ &= Q_0 \left[1 - 3 \int_0^{r/R_s} x^2 y^2 dy \right] \end{aligned} \quad \begin{aligned} \text{where } Q_0 &\equiv Q(r=0) \\ x &\equiv n_p/n_H = n_e/n_H \\ y &\equiv r/R_s \end{aligned}$$

$$R_s = \left(\frac{3}{4\pi} \frac{Q_0}{\alpha_{B,H} n_H^2} \right)^{1/3}$$

- At each point,
 - The rate of Case B recombinations per volume must be balanced by the rate of photoionization per volume:

$$\frac{Q(r)}{4\pi r^2} n_{H^0} \sigma_{\text{pi}} = n_p n_e \alpha_B$$

- This can be rewritten as

$$\frac{Q(r)}{4\pi r^2} (1-x) n_{\text{H}} \sigma_{\text{pi}} = x^2 n_{\text{H}}^2 \alpha_{\text{B}}$$

$$\frac{Q(r)}{Q_0} \frac{(4\pi/3) R_s^3 \alpha_{\text{B}} n_{\text{H}}^2}{4\pi r^2} (1-x) n_{\text{H}} \sigma_{\text{pi}} = x^2 n_{\text{H}}^2 \alpha_{\text{B}}$$

$$\frac{x^2}{1-x} = \frac{Q(r)}{Q_0} \frac{\tau_s}{3y^2}$$

where $\tau_s \equiv n_{\text{H}} \sigma_{\text{pi}} R_s$

$$= 2880 \left(\frac{Q_0}{10^{49} \text{ s}^{-1}} \right)^{1/3} \left(\frac{n_{\text{H}}}{10^2 \text{ cm}^{-3}} \right)^{1/3} \left(\frac{T}{10^4 \text{ K}} \right)^{0.28} \left(\frac{\sigma_{\text{pi}}}{2.95 \times 10^{-18} \text{ cm}^2} \right)$$

- Now, we can estimate the ionization degree x at each point r , by simultaneously solving the following equations:

$$\frac{x^2}{1-x} = \frac{Q(y)}{Q_0} \frac{\tau_s}{3y^2}$$

$$\frac{Q(y)}{Q_0} = \left[1 - 3 \int_0^y x^2 y'^2 dy' \right]$$

$$(0 \leq y = r/R_s \leq 1)$$

Heating and Cooling in H II Regions: Heating

- ***Temperature***

- $T_{\text{HII}} \sim 10,000 \text{ K}$. Observations indicate that the temperatures of H II regions are remarkably independent of the effective temperature of the central star.
- The temperature is not determined by the central star. It is ***the result of a balance between heating and cooling mechanisms*** in the ionized gas of the H II region.
- The main source of heating in an ionized nebula is photoionization.

- ***Photoionization Heating***

- When hydrogen is photoionized from its ground state, the photoelectron that is emitted carries away a kinetic energy:

$$E = h\nu - I_{\text{H}} \quad (h\nu = \text{energy of incident photon})$$

The **mean energy of the ejected electrons**, averaged over the all photoionization, is

$$\langle E \rangle = \langle h\nu \rangle - I_{\text{H}}$$

- The average energy $\langle h\nu \rangle$ of an ionizing photon must be weighted by the photoionization cross-section.

$$\langle h\nu \rangle = \frac{\int_{\nu_0}^{\infty} (4\pi J_{\nu}/h\nu)(h\nu)\sigma_{\text{pi}} d\nu}{\int_{\nu_0}^{\infty} (4\pi J_{\nu}/h\nu)\sigma_{\text{pi}} d\nu}$$

- Although stars are not blackbodies, we will use the Planck function. Because the energy of ionizing photons is $h\nu > 13.6 \text{ eV}$, we use the high-energy Wien tail with an effective temperature T_{eff} .

$$J_\nu \propto \nu^3 \exp\left(-\frac{h\nu}{kT_{\text{eff}}}\right) \quad \text{and} \quad \sigma_{\text{pi}} \propto \nu^{-3}$$

$$\begin{aligned} \langle h\nu \rangle &= \frac{h \int_{\nu_0}^{\infty} (\nu^2 e^{-h\nu/kT_{\text{eff}}}) \nu \cdot \nu^{-3} d\nu}{\int_{\nu_0}^{\infty} (\nu^2 e^{-h\nu/kT_{\text{eff}}}) \nu^{-3} d\nu} \\ &= kT_{\text{eff}} \frac{\int_{x_0}^{\infty} e^{-x} dx}{\int_{x_0}^{\infty} e^{-x} x^{-1} dx} \quad \text{Here, } x \equiv h\nu/kT_{\text{eff}} \text{ and } x_0 \equiv h\nu_0/kT_{\text{eff}} \\ &= kT_{\text{eff}} \frac{e^{-x_0}}{\int_{x_0}^{\infty} e^{-x} x^{-1} dx} = kT_{\text{eff}} \frac{e^{-x_0}}{E_1(x_0)} \end{aligned}$$

The integral in the denominator is the first exponential integral $E_1(x_0)$. Then, we obtain

$$E_1(x_0) \simeq \frac{e^{-x_0}}{x_0} \left[1 - \frac{1}{x_0} + \mathcal{O}(x_0^{-2}) \right] \quad \text{for } x_0 \gg 1$$

$$\langle h\nu \rangle \approx kT_{\text{eff}} x_0 \left(1 + \frac{1}{x_0} \right) = h\nu_0 + kT_{\text{eff}} \longrightarrow$$

Mean kinetic energy of the ejected electrons:

$$\langle E \rangle = \langle h\nu \rangle - I_{\text{H}} \approx kT_{\text{eff}}$$

-
- **Volumetric heating rate:** In photoionization equilibrium,

$$n_{\text{H}^0} \zeta_{\text{pi}} = n_e n_p \alpha_{\text{B,H}}$$

Hence, the volumetric heating rate is

$$\begin{aligned} \mathcal{G}_{\text{pi}} &= n_{\text{H}^0} \zeta_{\text{pi}} \langle E \rangle && \longleftarrow n_{\text{H}^0} \zeta_{\text{pi}} = n_e n_p \alpha_{\text{B,H}} \quad \text{and} \quad \langle E \rangle = kT_{\text{eff}} \\ &= n_{\text{H}}^2 \alpha_{\text{B,H}} kT_{\text{eff}} && \longleftarrow \alpha_{\text{B,H}} \approx 2.59 \times 10^{-13} (T_{\text{gas}}/10^4 \text{ K})^{-0.833} \quad [\text{cm}^3 \text{ s}^{-1}] \\ &\propto T_{\text{gas}}^{-0.83} T_{\text{eff}} \end{aligned}$$

Notice that *the volumetric heating rate decreases with increasing gas temperature.*

- **Necessity of the cooling mechanisms**

- ▶ An O3 main sequence star has an effective temperature $T_{\text{eff}} \sim 44,850 \text{ K}$ ($kT_{\text{eff}} \sim 3.9 \text{ eV}$), and thus the photoelectrons will have a mean energy of 3.9 eV.
- ▶ However, the free electrons in a 10,000 K nebula have a mean energy $(3/2) kT_{\text{gas}} \sim 1.3 \text{ eV}$.
- ▶ Therefore, some cooling mechanism must be reducing the average kinetic energy of the photoelectrons.

Heating and Cooling in H II Regions: Cooling

- **Main cooling sources in H II regions:**
 - Recombination Continuum and Line Emission (free-bound)
 - Thermal Bremsstrahlung (free-free)
 - Collisionally Excited Line Emission
- **Recombination Cooling:**
 - Recombination cooling occurs when electrons undergo radiative recombination with protons to form neutral hydrogen atoms. The volumetric cooling rate is then

$$\mathcal{L}_{\text{rr}} = n_e n_p \alpha_{\text{B,H}} \langle E_{\text{rr}} \rangle$$

where $\langle E_{\text{rr}} \rangle$ is the mean kinetic energy of the recombining electrons. The mean kinetic energy is obtained by weighting by cross section and integrating over the Maxwell distribution

$$\langle E_{\text{rr}} \rangle = \frac{\langle E \sigma_{\text{rr}} v \rangle_{\text{Maxwell}}}{\langle \sigma_{\text{rr}} v \rangle_{\text{Maxwell}}} = \frac{\int v^2 dve^{-E/kT_{\text{gas}}} \sigma_{\text{rr}} v E}{\int v^2 dve^{-E/kT_{\text{gas}}} \sigma_{\text{rr}} v} = \frac{\int E^2 \sigma_{\text{rr}} e^{-E/kT_{\text{gas}}} dE}{\int E \sigma_{\text{rr}} e^{-E/kT_{\text{gas}}} dE}$$

Note that $\langle E_{\text{rr}} \rangle \neq (3/2)kT_{\text{gas}}$. This is because the radiative recombination cross-section is a decreasing function of electron kinetic energy.

We will perform the integration by approximating that the radiative recombination cross-section, at about $T \sim 10^4$ K, by a power-law:

$$\sigma_{\text{rr}}(E) = \sigma_0 (E/E_0)^\gamma \quad \text{where } \gamma \approx -1.316 \text{ for Case B}$$

See Section 27.3.1 of [Draine]
for the derivation of the
power-law index.

Then, the mean energy per recombining electron (for Case B) is

$$\begin{aligned} \langle E_{\text{rr}} \rangle &= \frac{\Gamma(3 + \gamma)}{\Gamma(2 + \gamma)} kT_{\text{gas}} = (2 + \gamma) kT_{\text{gas}} & \leftarrow & \quad \Gamma(n) = \int_0^\infty x^{n-1} e^{-x} dx \\ &= 0.684 kT_{\text{gas}} \end{aligned}$$

The cooling rate from the recombination is

$$\mathcal{L}_{\text{rr}} = n_e n_p \alpha_{\text{B,H}} (\gamma + 2) kT_{\text{gas}}$$

- Gas temperature:
 - If radiative recombination were the only cooling mechanism, then the gas temperature would be found by equating the photoionization heating with the recombination cooling.

$$\mathcal{G}_{\text{pi}} = \mathcal{L}_{\text{rr}} \quad \longrightarrow \quad n_e n_p \alpha_{\text{B,H}} kT_{\text{eff}} = n_e n_p \alpha_{\text{B,H}} (\gamma + 2) kT_{\text{gas}}$$

$$T_{\text{gas}} = \frac{T_{\text{eff}}}{2 + \gamma} = \frac{T_{\text{eff}}}{0.684} = 1.462 T_{\text{eff}}$$

The resulting temperature would be ~46% higher than the effective temperature of the central star. For an O3 main sequence star with $T_{\text{eff}} = 44,900 \text{ K}$, the nebula temperature will be $T_{\text{gas}} = 66,000 \text{ K}$

This is because radiative recombination selectively removes the lower-energy free electrons (because of the higher cross section at lower energy), and thus increases the mean kinetic energy of electrons that are left without being captured.

- Hence, we need an additional cooling mechanism.
- **Free-free cooling:**
 - Bremsstrahlung cooling occurs when free electrons are accelerated by close encounters with protons or other ions, and thus emit radiation.
 - The emissivity is

$$4\pi j_{\nu}^{\text{ff}} = \frac{32\pi}{3} \left(\frac{2\pi}{3}\right)^{1/2} \frac{Z_i^2 e^6}{m_e^2 c^3} \left(\frac{m_e}{kT_{\text{gas}}}\right)^{1/2} n_i n_e g_{\text{ff}} e^{-h\nu/kT_{\text{gas}}} \quad (Z_i = 1, n_i = n_p \text{ for H})$$

where g_{ff} is the Quantum mechanical Gaunt factor.

-
- The volumetric cooling rate for a pure hydrogen gas is

$$\begin{aligned}\mathcal{L}_{\text{ff}} &= \int_0^{\infty} 4\pi j_{\nu}^{\text{ff}} d\nu \\ &= \frac{32\pi}{3} \left(\frac{2\pi}{3}\right)^{1/2} \frac{e^6}{m_e^2 h c^3} (m_e k T_{\text{gas}})^{1/2} n_p n_e \bar{g}_{\text{ff}}\end{aligned}$$

where \bar{g}_{ff} is the frequency-averaged Gaunt factor.

For temperature near $T_{\text{gas}} = 10^4$ K, a Quantum-mechanical calculation yields

$$\bar{g}_{\text{ff}} \approx 1.34 (T/10^4 \text{ K})^{0.05}$$

- The ratio between the radiative recombination cooling and free-free cooling rates is

$$\frac{\mathcal{L}_{\text{ff}}}{\mathcal{L}_{\text{rr}}} = \frac{32\pi}{3} \left(\frac{2\pi}{3}\right)^{1/2} \frac{e^6}{m_e^2 hc^3} \left(\frac{m_e}{kT_{\text{gas}}}\right)^{1/2} \frac{\bar{g}_{\text{ff}}}{(2+\gamma)\alpha_{\text{B,H}}}$$

$$\frac{\mathcal{L}_{\text{ff}}}{\mathcal{L}_{\text{rr}}} \approx 0.79 \left(T_{\text{gas}}/10^4 \text{ K}\right)^{0.37}$$

Note that both cooling mechanisms are two-body processes and thus the factors $n_e n_{\text{H}^+}$ cancel.

- Adding the free-free cooling, we can estimate the gas temperature, as follows:

$$\mathcal{G}_{\text{pi}} = \mathcal{L}_{\text{rr}} + \mathcal{L}_{\text{ff}} \longrightarrow T_{\text{eff}} = (\gamma + 2) T_{\text{gas}} \left[1 + 0.79 \left(T_{\text{gas}}/10^4 \text{ K}\right)^{0.37}\right]$$

$$\gamma + 2 = 0.684$$

Example: for an O3 main sequence star with $T_{\text{eff}} = 44,900 \text{ K}$, the nebula temperature will be $T_{\text{gas}} = 30,000 \text{ K}$ if both the radiative recombination and free-free coolings are taken into account. This temperature is still higher than that is actually observed in H II regions.

- ***Collisional excited line cooling***

- If a free electron collisionally excites an atom or ion from a lower energy level to an excited level, the energy difference between the levels is taken from the free electron's kinetic energy. If the excited atom or ion then undergoes radiative de-excitation, and if the emitted photon escapes from the nebula, then there is a net cooling of the gas.

-
- To cool from $T \sim 30,000$ K to $\sim 10,000$ K, the energy levels of the excited system must be separated by a difference $\Delta E \approx 1 - 3$ eV [$T \approx (1.2 - 3.5) \times 10^4$ K].
 - ▶ If ΔE is much lower than this value, then the photons emitted by radiative de-excitation will carry away only a small amount of energy.
 - ▶ If ΔE is much higher than this value, then only a small fraction of free electrons will have high enough energies to excite the ions or atoms.
 - ▶ In H II regions, most of the hydrogen will be ionized. Even if some He or He^+ is present, the energy of the first excited state is so far above the ground state that the rate for collisional excitation is negligible. Ly α ($\Delta E = 10.2$ eV) emission from neutral hydrogen atoms is not effective at cooling H II regions. Similarly, the first excited state of neutral helium is far too energetic ($\Delta E = 20.6$ eV) to be collisional excited.
 - ***This is where the heavy atoms such as oxygen and nitrogen play a key role in cooling H II regions.***
 - ▶ In particular, O II, N II, and O III have forbidden transitions in the 1 - 3 eV range.
 - ▶ If the collisional excitation is followed by a collisional de-excitation, the kinetic energy of the gas will be unchanged.
 - ▶ Therefore, if a collisional excitation is to result in cooling, it must be followed by a radiative de-excitation. For radiative de-excitation to dominate over collisional de-excitation, the number density of electrons must be lower than the critical density n_{crit} .
 - ▶ The critical density for these forbidden lines are indeed high compared to typical densities in an H II region.

- Calculation of the cooling rate for the collisionally excitation lines (electron impact emission lines)
 - If the collisionally excited levels are radiatively de-excited, the rate of energy loss by the gas is

$$\mathcal{L}_{ce} = \sum_X \sum_u n(X, u) \sum_{\ell < u} A_{u\ell} E_{u\ell}$$

where $E_{u\ell} \equiv E_u - E_\ell$

where the sum is over species X and excited states u .

Recall:

[population balance for two level atoms], ignoring the stimulated emission

$$n_\ell n_e k_{\ell u} = n_u (n_e k_{u\ell} + A_{u\ell})$$

$$\rightarrow \frac{n_u}{n_\ell} = \frac{n_e k_{\ell u}}{n_e k_{u\ell} + A_{u\ell}} \quad \rightarrow \quad \frac{n_u}{n_\ell} \simeq n_e \frac{k_{\ell u}}{A_{u\ell}} \quad \text{for low density.}$$

[collisional excitation & de-excitation rate coefficients]

$$k_{u\ell} = \langle \sigma_{u\ell} v \rangle = \frac{\beta}{T^{1/2}} \frac{\langle \Omega_{u\ell} \rangle}{g_u} \quad [\text{cm}^3 \text{ s}^{-1}],$$

$$k_{\ell u} = \langle \sigma_{\ell u} v \rangle = \frac{\beta}{T^{1/2}} \frac{\langle \Omega_{u\ell} \rangle}{g_\ell} e^{-(E_u - E_\ell)/kT_{\text{gas}}} \quad [\text{cm}^3 \text{ s}^{-1}]$$

$(\beta = 8.62942 \times 10^{-6})$

[emissivity] $4\pi j_\nu = n_u A_{u\ell} (E_u - E_\ell)$

[principle of detailed balance]

$$\frac{k_{\ell u}}{k_{u\ell}} = \frac{g_u}{g_\ell} e^{-(E_u - E_\ell)/kT_{\text{gas}}}$$

We need (1) $A_{u\ell}$ and (2) $\langle \Omega_{u\ell} \rangle$.
 For three or more levels, the balance equation becomes more complicated.
 See Appendix F of Draine, Table 4.1 of Lequeux, Table 9.3 & 9.4 in Draine